

**Coordination Chemistry and Applications of Versatile 4,5-Diazafluorene Derivatives**

Journal:	<i>Dalton Transactions</i>
Manuscript ID	DT-PER-09-2015-003665.R1
Article Type:	Perspective
Date Submitted by the Author:	12-Nov-2015
Complete List of Authors:	Song, Datong; University of Toronto, Department of Chemistry Annibale, Vincent; University of British Columbia, Chemistry; University of Toronto, Chemistry



Coordination Chemistry and Applications of Versatile 4,5-Diazafluorene Derivatives

Vincent T. Annibale,^{a,b} and Datong Song^{a*}

Received 00th January 20xx,
Accepted 00th January 20xx


DOI: 10.1039/x0xx00000x

www.rsc.org/

This perspective review will examine the coordination chemistry and applications of metal complexes of 4,5-diazafluorene derivatives. The versatile derivatives of 4,5-diazafluorene can serve multiple roles, and display a number of coordination modes. The ambidentate derivatives with multiple coordination sites can allow for the syntheses of coordination polymers, multimetallic, and macrocyclic complexes. In addition, certain 4,5-diazafluorene derivatives can serve as spectator ligands to support reactivity at the metal centre, or as reactive actor ligands engaging in atypical reactivity patterns. The applications of metal complexes of 4,5-diazafluorene derivatives in catalysis, photochemistry and photophysics, as well as bioinorganic chemistry are also surveyed.

Introduction:

4,5-Diazafluorene (dafH) was first reported in the late 1970s,^{1, 2} and synthesized in two steps from 1,10-phenanthroline (phen). The first step is an oxidative ring contraction of phen with permanganate in basic aqueous media giving 4,5-diazafluoren-9-one (dafo),³ and the second step is a Wolff-Kishner reduction of dafo with hydrazine monohydrate at high temperature (Scheme 1).^{4, 5} The contraction of middle ring of phen increases the distance between the two N-donors, i.e., 2.72 Å in phen⁶ and 3.05 Å for dafo⁷ and dafH.⁸ In coordination chemistry, the dafH ligand has been conventionally viewed as a 2,2'-bipyridine (bpy) derivative with a methylene linker tethering the two pyridine rings together (Figure 1). The methylene group effectively ties back the two pyridine rings, resulting in a longer N–N distance in dafH (2.82(3) Å) compared to bpy (2.63(4) Å) in complexes. Consequentially, the overlap between the nitrogen lone pairs and the metal orbitals is less effective for dafH compared to bpy (Figure 1).^{9–13} The dafo ligand has an even longer N–N distance (2.96(7) Å) since C9 is sp² hybridized in dafo.¹⁴ Although dafH derivatives were considered merely as bpy analogues in the late 1970s, they have recently gained more attention in many areas. Ono,¹⁵ and Wong,¹⁶ previously reviewed the synthesis and coordination chemistry of several dafH derivatives. The purpose of this perspective review is to highlight the versatile nature of dafH derivatives as ligands, and also to detail some of the recent advances made using this ligand family.



Scheme 1. Synthesis of dafH from 1,10-phenanthroline with numbering scheme of dafH ligand shown.




Figure 1. Comparison between bpy and dafH.¹⁷

I. Fundamental coordination chemistry of dafH derivatives

I.1. Coordination chemistry of the parent dafH and daf ligands:

Although being conventionally viewed as a bpy derivative, the dafH ligand can also be viewed as two pyridine rings fused onto a central cyclopentadiene (CpH) ring in a *syn* fashion. The methylene group of dafH is acidic, analogously the pK_s of CpH is 18, and that of fluorene is 22.6 in DMSO.¹⁸ The methylene linker of dafH may be deprotonated to form the monoanionic 4,5-diazafluorenyl (daf⁻) (Scheme 2A).

Ambidentate ligands containing multiple potential coordination sites of different nature can be used to construct linkage isomers, homo- and heteromultimetallic complexes, and complex molecular architectures through coordination-driven self-assembly. Several 4,5-diazafluorene derivatives possess multiple coordination sites, especially those derivatives that are functionalized at the C9-position discussed later. The daf⁻ ligand is potentially an ambidentate ligand with two types of metal binding sites, an N,N-chelate site, and the C-donors of the central cyclopentadienyl-like moiety. However in most examples daf⁻ binds metals with its nitrogen donors,^{19–24} without utilizing the C-donors. In addition the daf⁻ ligand is able to form zwitterionic complexes when only two N-donors

^a Davenport Chemical Research Laboratories, Department of Chemistry, University of Toronto, 80 St. George Street, Toronto, Ontario, Canada M5S 3H6. *E-mail: dsong@chem.utoronto.ca; Fax: +1 416 978 7013; Tel: +1 416 978 7014

^b Current Address: Department of Chemistry, The University of British Columbia, 2036 Main Mall, Vancouver, British Columbia, Canada V6T 1Z1.


PERSPECTIVE

Dalton Transactions

are bound to metal centre with a -1 charge localized onto the ligand backbone. A recurring theme in this review is the reactivity of the ligand backbone in a zwitterionic setup.

Our group's initial investigations into the coordination chemistry of the 4,5-diazafluorene (daf) ligand were reported in 2008.¹⁹ The reaction of Na[daf] with [Pd(PPh₃)₂Cl₂] gave a dinuclear Pd complex [Pd(daf)(PPh₃)Cl]₂, in this exceptional case the daf ligand was bound to Pd through one of the N-donors and the anionic C-donor of the ligand backbone in an η¹(σ)-fashion (Scheme 2B).¹⁹ A zwitterionic Rh(I) complex [Rh(daf)(COD)] can also be straightforwardly prepared by reacting Na[daf] with 0.5 equiv. of [Rh(COD)Cl]₂ (Scheme 2B).

An extension of the work with Rh 4,5-diazafluorene derivatives involved the syntheses of complexes with PPh₃ as an auxiliary ligand (see Scheme 2C).²⁰ The Rh(I) complex [Rh(daf)(PPh₃)₂] can be synthesized from the reaction of Na[daf] and Wilkinson's catalyst. Complex [Rh(daf)(PPh₃)₂] reacts with H₂ to give a Rh(III) dihydride complex [Rh(daf)(PPh₃)₂(H)₂], which reacts with HCl to give [Rh(dafH)(PPh₃)₂(H)₂]Cl (Scheme 2C). Alternatively, [Rh(dafH)(PPh₃)₂(H)₂]Cl can also be synthesized by reacting dafH with Wilkinson's catalyst under H₂ atmosphere, subsequent deprotonation with NaH gives [Rh(daf)(PPh₃)₂(H)₂] cleanly.



Scheme 2. A) Deprotonation of dafH to give daf.¹⁹ B) Synthesis of [Rh(daf)(COD)] and [Pd(daf)(PPh₃)Cl]₂.¹⁹ C) Interconversion between Rh 4,5-diazafluorene complexes.²⁰


Parts of Scheme 3 highlight the ambidentate nature of daf in its coordination chemistry with a {RuCp*}⁺ fragment.²⁵ The addition of neutral dafH to 0.25 equiv. of [RuCp*(μ₃-Cl)]₄ starting material gave [RuCp*(dafH)(Cl)] where the dafH ligand coordinates through the nitrogen chelate (Scheme 3).²⁵ However the addition of Na[daf] to 0.25 equiv. of [RuCp*(μ₃-Cl)]₄ gave a monomeric sandwich complex [RuCp*(daf)] where the daf ligand coordinates through the central cyclopentadienyl-like ring in an η⁵-fashion leaving the nitrogen chelate vacant (Scheme 3).²⁵ Surprisingly when the [RuCp*(μ₃-

Cl)]₄ starting material is treated with K[daf], or if the coordinated 4,5-diazafluorene ligand of [RuCp*(dafH)] is deprotonated with K[HBET₃] the self-assembly of a tetraruthenamacyclo [RuCp*(daf)]₄ occurred where the daf ligand coordinated through both the N-donors and the C-donor of the ligand backbone in an η¹(σ)-fashion (Scheme 3).²⁵ The different coordination behaviours of daf in [RuCp*(daf)]_n (n = 1 or 4), a monomeric sandwich complex and a tetraruthenamacyclo, demonstrate its ambidentate nature.


We have used 4,5-diazafluorene derivatives to synthesize heterobimetallic complexes.^{26, 27} When the salt [Na(dme)₃][Pt(daf)(Ph)₂] is treated with [Cu(IPr)Cl], a Pt(II)–Cu(I) heterobimetallic complex [(IPr)Cu(daf)Pt(Ph)₂] was obtained where C9 of daf is coordinated to the {Cu(IPr)}⁺ fragment in an η¹(π) fashion (Scheme 4).²⁶ Surprisingly when the metallation sequence is reversed and the [Cu(daf)(IPr)] complex is treated with [Pt(Ph)₂(μ-SMe₂)_n] (n = 2 or 3) the exact same Pt(II)–Cu(I) heterobimetallic complex (as opposed to an isomer) was obtained.²⁶ In the process of generating [(IPr)Cu(daf)Pt(Ph)₂] from [Cu(daf)(IPr)] the {Cu(IPr)}⁺ fragment was replaced by the {Pt(Ph)₂} unit at the N,N-chelate site and migrated to the backbone C-donor. The built-in self-correction featured in the synthesis of [(IPr)Cu(daf)Pt(Ph)₂] system allowed for a one pot synthesis to be successfully carried out (Scheme 4).

1.2. Coordination chemistry of the aryl substituted daf_{Mes}H and daf_{Mes}⁻ ligands:

A 3,6-dimesityl substituted 4,5-diazafluorene ligand (daf_{Mes}H) has been synthesized, which can also be deprotonated to give daf_{Mes}⁻.²⁵ These bulkier 4,5-diazafluorene derivatives are also ambidentate. The daf_{Mes}⁻ ligand has an N,N-chelate site and the C-donors of the central cyclopentadienyl moiety; in addition, there is also the possibility of the mesityl substituent to participate in bonding. The coordination chemistry of neutral daf_{Mes}H and monoanionic daf_{Mes}⁻ ligands toward the {RuCp*}⁺ fragment was examined (Scheme 3). When K[daf_{Mes}⁻] was added to 0.25 equiv. of [RuCp*(μ₃-Cl)]₄, a sandwich complex [RuCp*(daf_{Mes}⁻)] formed where the nitrogen chelate is vacant and the daf_{Mes}⁻ ligand is coordinated through the central C₅-ring in a η⁵-fashion (Scheme 3).²⁵ However upon adding the neutral daf_{Mes}H ligand to 0.25 equiv. [RuCp*(μ₃-Cl)]₄, [RuCp*(daf_{Mes}H)]Cl was obtained. Since the mesityl groups prevent the N-donors from coordinating to the {RuCp*}⁺ fragment, the {RuCp*}⁺ fragment coordinates to the arene (Scheme 3).²⁵ After deprotonation of [RuCp*(daf_{Mes}H)]Cl with KOtBu, a zwitterionic linkage isomer of [RuCp*(daf_{Mes}⁻)] formed, where the Ru(II) remains coordinated to the arene (Scheme 3).²⁵ Interestingly two different linkage isomers of [RuCp*(daf_{Mes}⁻)], a sandwich complex and a zwitterionic complex, were formed depending on the reaction sequence.



Scheme 3. Coordination chemistry of ambidentate 4,5-diazafluorene derivatives with a RuCp* synthon.²⁵




Scheme 4. Three synthetic routes to [(IPr)Cu(daf)Pt(Ph)₂].²⁶

A variety of heterobimetallic complexes can also be synthesized where the central cyclopentadienyl ring of 4,5-diazafluorene is coordinated to one metal centre and the nitrogen chelate is coordinated to a second metal centre, and the ligand displays either η⁵,κ²-[N,N] or η⁵,κ¹-N coordination

modes.²⁷ For example, [PtPh₂(daf)]⁻ and [PtPh₂(daf_{Mes})]⁻ can be metallated successfully with the [RuCp*(μ₃-Cl)₄] to obtain Ru(II)–Pt(II) heterobimetallic complexes [RuCp*(daf)Pt(Ph)₂] and [RuCp*(daf_{Mes})Pt(Ph)₂] (Scheme 5). These Ru(II)–Pt(II) heterobimetallic complexes can also be synthesized from the Ru sandwich complexes [RuCp*(daf)], and [RuCp*(daf_{Mes})] (Scheme 5).²⁷ The [RuCp*(daf_{Mes})] sandwich isomer with a vacant N,N-chelate can also serve as a metalloligand for the complexation with metal halides such as CuCl, FeCl₂(THF)_{1.5} and CoCl₂(THF)_{1.5} (Scheme 5).²⁷

PERSPECTIVE




Scheme 5. Synthesis of heterodinuclear Ru(II)-M complexes where the daf or daf_{Mes} ligand displayed η³,κ⁻[N,N] or η³,κ⁺-N coordination modes.²⁷

1.3. Coordination chemistry of the phosphine donor functionalized daf_{pH} and daf_p⁻ ligands:

Our group also installed a phosphine arm at the C9-position of 4,5-diazafluorene to give the daf_{pH} ligand, which can also be deprotonated to form the daf_p⁻ ligand. Both daf_{pH} and daf_p⁻ have been used to assemble head-to-tail macrocycles with {RuCp*}⁺ (Scheme 3).²⁵ We further demonstrated the transfer of daf_p⁻ from [Cu(IPr)(daf_p)_n] (n = 1 or 2) to either Rh(I) or Au(I) resulting in macrocyclic complexes (Scheme 6).²⁸ There are a few benefits to constructing these macrocyclic Rh(I) and Au(I) complexes through ligand transfer from a {Cu(IPr)}⁺ fragment, compared to the conventional synthesis by directly reacting the daf_p⁻ salt and metal chloride starting materials. The benefits include improved yields, shortened reaction time, and simplified isolation of the product as the soluble [Cu(IPr)Cl] byproduct can easily be removed by filtration. In addition, daf_{pH} and daf_p⁻ are also ambidentate ligands with phosphine and N-donor coordination sites, while daf_p⁻ can also anchor a metal in the P,C-coordination site. The daf_p⁻ ligand displayed several coordination modes, where the diazafluorenyl moiety coordinates through one or both N-donors and the P-donor with or without the participation of the C-donor of the cyclopentadienyl-like moiety.

We also have demonstrated that in the heterodinuclear complex [(NHC)Cu(daf_p)Pt(Ph)₂] the tethered phosphine of daf_p⁻ helps anchor the Pt(II) center onto the carbon site, and the Cu(I) center is bound to the N,N-chelate site (Scheme 6).²⁶ The synthesis of these Pt^{II}-Cu^I heterobimetallics is highly regioselective, and can also be performed in one-pot (Scheme 6).²⁶



Scheme 6. Synthesis of [Cu(NHC)(daf_p)] from [K(daf_p)], and the synthesis of macrocyclic complexes [Au(daf_p)₂], [Rh(daf_p)(COD)]₂ through salt metathesis, and daf_p ligand transfer from a Cu(NHC) complex.²⁶ In addition the selective syntheses of Pt(II)-Cu(I) heterobimetallic complexes.²⁶


1.4. Coordination chemistry of sulfur donor functionalized 4,5-diazafluorene derivatives:

Baudron, Hosseini, and co-workers have created ambidentate ligands with sulfur donors and demonstrated their use in the stepwise synthesis of multimetallic complexes and coordination polymers.²⁹⁻³² Both 4,5-diazafluorenyl-9-dithiolene (L¹) ligands,^{30, 32} and 4,5-diazafluorenyl-9-dithiolate (L²),^{29, 31, 32} are generated *in situ* by the removal of propionitrile S-protecting groups using NR₄OH (Scheme 7). Both ambidentate dianionic ligands provide different chelates, a dithiolate or dithiolene chelate and the 4,5-diazafluorene moiety. The homoleptic mononuclear complexes of the generic form [M(L¹)₂](NBu₄)₂ (where M = Ni(II) or Hg(II)),³⁰ and [M(L²)₂](NEt₄)₂ (M = Ni(II), Pd(II), Zn(II), or Hg(II))³¹ can be synthesized (Scheme 7). The homoleptic complexes all have a central metal which is coordinated by the dithiolate or dithiolene moiety of the ligands, leaving the 4,5-diazafluorene chelate sites vacant. These mononuclear complexes form one-dimensional coordination polymers in the presence of Na⁺ ions (Scheme 7).^{30, 31} The heterotrimetallic complex [(cyclenNi)₂Pd(L²)₂](BF₄)₂ can be prepared where the {(cyclen)Ni}²⁺ fragment is coordinated to the 4,5-diazafluorenyl moiety via the N,N-chelate (Scheme 7).³¹ The 4,5-diazafluorenyl-9-dithiolate ligand can serve as a bridging ligand through the two anionic S-donors to assemble an octanuclear Cu complex [Cu₈(L²)₆]⁴⁻ where the eight Cu centres occupy the vertices of a cube and on the periphery of the complex are the potentially chelating 4,5-diazafluorenyl moieties (Figure 2).²⁹

Heteronuclear metallamacrocycles can be constructed from L¹ and L² ligands (Scheme 8).³² The dianionic ligands are first metallated with (dppp)M¹Cl₂ (where M¹ = Pd, Pt, and dppp = 1,2-bis(diphenylphosphino)propane) which bears a diphosphine capping ligand allowing for a discrete neutral

species of the general form $[(dppp)M^1(L^1)]$ or $[(dppp)M^1(L^2)]$ to be isolated, where both L^1 and L^2 ligands coordinate to M^1 through the S,S-chelate, leaving the 4,5-diazafluorenyl moiety available for a second metal (Scheme 8).³² The addition of a second group 10 metal halide M^2X leads to the formation of head-to-tail metallamacrocycles where the formation of metal-sulfur linkages are involved in the assembly (Scheme 8).³²

The heteroleptic Ir(III) complex $[(dppp)_2(FluoDT(SPN)_2)]^+$ with the propionitrile-protected 4,5-diazafluorene-9-dithiolene proligand (where $dppp = 2-(2,4\text{-difluorophenyl})\text{pyridine}$ and $FluoDT(SPN)_2 = 4,5\text{-diaz-9-[4,5-bis(cyanoethylsulfanyl)-1,3-dithiol-2-ylidene]fluorene}$, see Scheme 7 for the structure of $FluoDT(SPN)_2$) has been used in a cascade reaction in basic media as both the reducing agent for chloroauric acid and the capping agent for the resulting gold nanoparticles.³³ In basic media the protecting groups of $[(dppp)_2(FluoDT(SPN)_2)]^+$ are cleaved resulting in the 4,5-diazafluorene-9-dithiolene (L^1) ligand which is redox-active and able to reduce Au^{3+} to Au^0 , and the anionic S-donors allow the Ir(III) capping agent to bind strongly to the surface of the Au^0 nanoparticles.³³ The coordination chemistry of the bis-thioether analogue 4,5-diaza-9-[4,5-bis(methylthio)-1,3-dithiol-2-ylidene]fluorene (L^3) with transition metals was explored.³⁴⁻⁴⁰ The solid state structure and IR spectrum and the solution UV-vis spectrum, magnetic properties have been reported for $[M(L^3)(tpa)](SbF_6)_2$ (where $tpa = \text{tris}(2\text{-pyridylmethyl})\text{amine}$). The L^3 ligand is labile and readily displaced in coordinating solvents such as acetone, or MeCN.⁴⁰



Scheme 7. Synthesis of homoleptic complexes $[M(L^1)]_2(NnBu_4)_2$ and $[M(L^1)]_2(NEt_3)_2$ and subsequent synthesis of heterotrimeric complex $[(\text{cyclen}Ni)_2Pd(L^1)_2](BF_4)_2$ and Na⁺ containing coordination polymers.³¹





Figure 2. Octanuclear Cu cluster, $[Cu_8(L^2)_6](NEt_3)_4$, composed of L^2 4,5-diazafluorenyl-9-dithiolate ligands.



Scheme 8. Heteronuclear metallamacrocycles constructed from 4,5-diazafluorenyl-9-dithiolene (L^1), and 4,5-diazafluorenyl-9-dithiolate (L^2).

1.5. Coordination chemistry of 9-hydroxy-9-alkynyl functionalized dafH derivatives:


The coordination chemistry of 9-hydroxy-9-alkynyl-4,5-diazafluorene derivatives which possess several potential sites of coordination has been explored.^{7, 41, 42} The 9-hydroxy-9-ethynyl-4,5-diazafluorene ligand can be dehydrated upon binding with a 16-electron coordinatively unsaturated species such as *in situ* generated $[M(\text{diphosphine})_2Cl]^+$ ($M = Ru, Os$)⁴¹ and $[Ru(\text{dppe})_2Cl](\text{OTf})$ precursor⁴² (Scheme 9). The resulting Ru and Os allenylidene complexes feature a vacant N,N-chelate which is a potential site for further coordination. Indeed this N,N-chelate moiety can be coordinated to a second metal centre to form Ru(II)-Ru(II), and Ru(II)-Re(I) multimetallic complexes.⁴²

When the 9-hydroxy-9-phenylethynyl-4,5-diazafluorene ligand is reacted with K_2PtCl_4 or ZnI_2 , the corresponding mononuclear complex forms with the metal bound to the N,N-

PERSPECTIVE

Dalton Transactions

chelate (Scheme 9).⁷ However the reaction of 9-hydroxy-9-phenylethynyl-4,5-diazafluorene with $\text{Co}_2(\text{CO})_8$ led to a (μ -alkyne)hexacarbonyldicobalt complex where the N,N-chelate remains unoccupied (Scheme 9).⁷



Scheme 9. Reactions of 9-hydroxy-9-alkynyl-4,5-diazafluorene.^{7, 41, 42}

1.6. Coordination chemistry of the dafo ligand:

Dafo has been used as a ligand for metals across the periodic table: Mn,⁴³ Co,^{44, 45} Ni,⁴⁶⁻⁴⁸ Cu,^{44, 48-62} Zn,⁷ Mo,⁶³ Ru,^{9, 64-73} Pd,^{17, 44, 74-87} Ag,⁸⁸⁻⁹⁰ Cd,^{62, 90, 91} Re,⁹² Ir,⁹³ Pt,^{7, 17, 81} Hg,⁴⁸ and the lanthanides^{50, 94-96}. In the vast majority of cases dafo behaves as a bidentate chelate ligand, however there are a few examples where dafo adopts either a monodentate κ^1 -N coordination mode,^{7, 17, 50, 54, 56, 88, 89} a bis-monodentate bridging coordination mode,^{17, 59, 60, 79, 81, 97} or even an η^4 -cyclopentadienone-like coordination mode in the $[\text{Cp}^*\text{Co}(\eta^4\text{-dafo})]$ complex.⁴⁵

1.7. Coordination polymers containing dafh derivatives

Recently metal-organic frameworks (MOFs) have received substantial interest due to the wide variety of potential architectures arising from different metal-ligand combinations, and the possibility of creating materials with intriguing applications. A variety of MOFs with different architectures were constructed from Zn(II) ions, a variety of aromatic polycarboxylic acid ligands and a 4,5-diazafluoren-9-oxime ligand.⁹⁸ The polynuclear Zn secondary building units (SBUs) comprising the various MOFs were modulated by 4,5-diazafluoren-9-oxime ligand, which can either coordinate to Zn in a chelating or monodentate fashion.⁹⁸

Azide containing MOFs can potentially be used as molecule-based magnets. A three-fold interpenetrating MOF of the general formula $[(\text{Mn}-\mu_{1,3}\text{-N}_3-\mu_{1,1}\text{-N}_3)_3(\text{L}^4)]$ where L^4 is the bis(bidentate) Schiff base ligand 4,5-diazafluoren-9-one azine exhibited spin-canted long-range ferromagnetic ordering.⁹⁹ The L^4 ligand served as the long links in the 3D structure while the chain of $\text{Mn}-\mu_{1,3}\text{-N}_3-\mu_{1,1}\text{-N}_3$ served as the SBU.⁹⁹ A series of isostructural tetranuclear clusters $[\text{M}_4(\text{dafo})_4(\text{N}_3)_2(\mu_{1,1}\text{-N}_3)_4(\mu_{1,1,1}\text{-N}_3)_2]$ have been synthesized with azido ligands bridging the four M^{2+} ions, and the dafo coordinated in a chelating fashion to the metal vertices of the clusters, where the metal can be Co,^{43, 100} Mn,⁴³ Cd,⁹¹ or Cu.⁴⁹

A luminescent Ag(I) one-dimensional polymer chain of the generic formula $[\text{Ag}_2(\text{L}^5)_2(\text{ClO}_4)_2]_n$ where L^5 =4,5-diaza-9,9'-spirobifluorene was structurally characterized.¹⁰¹ Both of the L^5 ligands adopt a bis-monodentate bridging mode between the two crystallographically independent Ag(I) centres where the Ag–Ag distance is 2.776(1) Å, and one of ClO_4^- ligands also bridges two adjacent Ag centres to construct the polymer chain (Figure 3).¹⁰¹




Figure 3. Coordination polymer constructed from Ag(I) and 4,5-diaza-9,9'-spirobifluorene (L^5).¹⁰¹

The fullerene C_{60} with four peripheral malonate groups can be functionalized with two 4,5-diazafluorene moieties in the *trans*-1 positions, since 4,5-diazafluorene is planar the N,N-chelates can be situated 180° relative to each other.¹⁰² The reaction between AgOTf and this large ditopic bis(4,5-diazafluorene)tetrakis(malonate) substituted fullerene ligand (L^6) gives a 1-D coordination polymer $[\text{Ag}_2(\text{L}^6)(\text{OTf})_2]_n$ where each Ag(I) centre is four-coordinate and is bound to the N,N-chelate of one of the 4,5-diazafluorene moieties, an O-donor from the OTf^- ligand, and a C-donor from a neighboring C_{60} cage in an η^1 -fashion (Figure 4).¹⁰² In the solid state of this fullerene-based coordination polymer the two antiparallel 4,5-diazafluorene moieties from two neighboring L^6 ligands engage in face-to-face π - π interactions.¹⁰² The choice of Ag(I) precursor is crucial for coordination polymer formation. For example, if AgBF_4 is used instead, a non-polymeric salt $[(\text{L}^6)(\text{Ag}(\text{toluene})_2)](\text{BF}_4)_2 \cdot \text{CH}_2\text{Cl}_2$ forms, where each Ag(I) centre is coordinated to the N,N-chelate of the 4,5-diazafluorene moieties and is bound to one η^1 -toluene and one η^2 -toluene.¹⁰²




Figure 4. A portion of the solid-state structure of the one-dimensional coordination polymer $[(\text{L}^6)(\text{AgOTf})_2]_n$, where L^6 is a bis(4,5-diazafluorene)tetrakis(malonate) substituted fullerene, toluene solvent molecules and H atoms omitted for clarity. Ag: pink, S: yellow, F: green, O: red, N: blue, C: grey.¹⁰²

A simple coordination polymer of $[\text{Cd}(\text{dafo})(\text{NCS})_2]_n$ can be prepared where each Cd centre adopts a distorted octahedral

geometry and the ambidentate NCS^- ligands bridge adjacent Cd centres through both the N and S termini.¹⁰³

Jung, Lee, and co-workers demonstrated the use of a heteroditopic bis(4,5-diazafluorenylimino)dibenzo[18]crown-6 ether based ligand which could be used to synthesize a coordination polymer gel in the presence of Zn^{2+} and Cs^+ ions.¹⁰⁴ The Zn^{2+} ion is bound by two 4,5-diazafluorenylimino moieties with a tetrahedral coordination geometry and one Cs^+ ion is sandwiched between two crown ether rings to give a highly cross-linked coordination polymer gel, where the rheological properties and microstructure are strongly dependent on the presence and concentration of Cs^+ ions.¹⁰⁴

There are several examples in the literature inorganic-organic hybrid polyoxometallates which contain Keggin-type cluster anion and metal-dafo complex cation (typically Cu, in a few examples Ag or Cd).^{50, 51, 57, 59, 60, 90, 97, 105} The metal-dafo cation within these hybrid compounds seems to direct the assembly of other supramolecular interactions within the solid state. The coordination geometry of the metal, how the Keggin cluster is coordinated to the cation, and even the number of metals can play a role. For example helical assemblies can form when dinuclear $[\text{Cu}_2(\text{dafo})_2(\text{H}_2\text{O})]^{2+}$ cations are used as a hinge-like motif to link Keggin clusters together in forming the three-dimensional framework.^{59, 60}

II. Applications of 4,5-diazafluorene derivatives


II.1. As actor ligands in reactivity chemistry.

Most of inorganic and organometallic chemistry is dominated by metal centred reactivity where the ligand is a spectator, hence the term "spectator ligand". There are emerging examples of ligand-centred, or metal-ligand cooperative reactivity where reactions with incoming substrates occur at the "actor ligands". The use of multifunctional actor ligands in small molecule activation and catalysis has gained significant interest in the chemistry community and there are few recent reviews on this topic.¹⁰⁶⁻¹¹³ There are several examples where 4,5-diazafluorene derivatives behave as actor ligands, where the majority of the ligand-based reactions occur at the reactive C9.


Rillema and co-workers have uncovered divergent chemical behaviour for coordinated dafo ligand and free dafo in reactions with nucleophiles.^{70, 71} Free dafo reacts with 2-aminopyridine in ethanol to give a Schiff-base product (Scheme 10A), however under the same reaction conditions the coordinated dafo ligand of $[\text{Ru}(\text{bpy})_2(\text{dafo})]^{2+}$ reacts with 2-aminopyridine to give a ring-opened product possessing a coordinated esterified 2,2'-bipyridine ligand (Scheme 10B).⁷¹ In another example no reaction occurs between free dafo and DBU in wet dichloromethane (Scheme 10A), while coordinated dafo in $[\text{Ru}(\text{bpy})_2(\text{dafo})]^{2+}$ reacts with DBU to give another ring-opened product (Scheme 10B).⁷⁰ The spectator metal centre plays a major role in altering the reaction pathways of dafo with nucleophiles; the driving force to form a coordinated 2,2'-bipyridine ligand from a coordinated dafo is the release of coordination-induced ring strain and the formation of shorter, stronger Ru-N bonds.^{70, 71} There are two potential nucleophilic sites in 2-aminopyridine, the amine and the pyridyl ring

nitrogen atoms; the spectator metal ion increases the electron density on the carbonyl carbon atom of dafo thus hindering the reaction with the amine nitrogen and the formation of the Schiff-base product.⁷¹ Free dafo does not react with ethylene glycol, meanwhile coordinated dafo of $[\text{Ru}(\text{bpy})_n(\text{dafo})_m]^{2+}$ ($n=1$ and $m=2$, or, $n=0$, and $m=3$) reacts to give the corresponding ketal ligand coordinated to Ru(II) which is resistant to hydrolysis with $\text{HCl}_{(\text{aq})}$ – a highly unusual feature compared with most organic ketals.⁷² Siemeling and co-workers synthesized the sandwich complex $[\text{Cp}^*\text{Co}(\eta^4\text{-dafo})]$, where the dafo ligand is coordinated through the π system leaving the $[N,M]$ -chelate vacant.⁴⁵ The $[\text{Cp}^*\text{Co}]$ fragment coordinates η^2 to each of the two six-membered dearomatized rings of dafo; this results in cyclopentadienone-like behaviour where the nucleophilicity of the oxygen atom is increased substantially versus free dafo.⁴⁵ The nucleophilic, coordinated, cyclopentadienone-like dafo of $[\text{Cp}^*\text{Co}(\eta^4\text{-dafo})]$ reacts with electrophiles such as acetyl chloride to give the *O*-acylated cobaltocenium species (Scheme 10C); in contrast free dafo does not react with acetyl chloride (Scheme 10A).⁴⁵

Our group discovered a surprising example of ligand-based reactivity: free 4,5-diazafluorene is air-stable but the coordinated 4,5-diazafluorene ligand in $[\text{Ru}(\text{dafH})(\text{PPh}_3)_2(\text{Cl})_2]$ can selectively undergo an aerobic oxidation reaction giving a coordinated dafo ligand (Scheme 11).⁶⁶ The selectivity of this ligand-based oxidation reaction is surprising since the typically oxygen-sensitive phosphine ligands are left intact.



Scheme 10. A) Reactivity of uncoordinated free dafo with 2-aminopyridine, DBU, and acetyl chloride.^{70, 71} B) Reactivity of $[\text{Ru}(\text{dafo})(\text{bpy})_2]^{2+}$ with 2-aminopyridine, and DBU.^{70, 71} C) Reactivity of $[\text{CoCp}^*(\eta^4\text{-dafo})]$ with acetyl chloride.




Scheme 11. Selective oxidation of a coordinated dafo ligand.⁶⁶

Our group also previously demonstrated an interesting example of metal-ligand cooperativity. The Ru(II) 4,5-

PERSPECTIVE


diazafluorene complex, $[\text{Ru}(\text{daf})(\text{PPh}_3)_2(\text{H})(\text{N}_2)]$, can be synthesized where the *daf* ligand possesses a central negatively charged cyclopentadienyl like moiety that remains uncoordinated and thus has unquenched basicity (Scheme 12).²¹ $[\text{Ru}(\text{daf})(\text{PPh}_3)_2(\text{H})(\text{N}_2)]$ reversibly splits dihydrogen over a long-range, between the metal centre and the backbone carbanion, a distance of ~ 5 Å, to yield complex $[\text{Ru}(\text{dafH})(\text{PPh}_3)_2(\text{H})_2]$.²¹ The π -system of the diazafluorene ligand is disrupted and restored during the forward and backwards reactions, respectively. It is also worth noting that the reversible synthesis of a metal-dinitrogen complex via the metal-hydride route, a route which circumvents the need for harsh reducing agents, is quite unique.¹¹⁴



Scheme 12. Reactions of $[\text{Ru}(\text{daf})(\text{PPh}_3)_2(\text{H})(\text{N}_2)]$ toward reversible H_2 splitting,²¹ and a reversible formal insertion of CO_2 into a *daf* C–H bond.¹¹⁵

The formation of H_2 from a coordinated 4,5-diazafluorene ligand giving a coordinated 4,5-diazafluorenyl ligand has been noted by other groups. Mach and co-workers observed the evolution of H_2 when 4,5-diazafluorene was added to $[\text{Cp}^*_2\text{Ti}(\text{Me}_3\text{SiCCSiMe}_3)]$ which gave the paramagnetic $[\text{Cp}^*_2\text{Ti}(\text{III})(\text{daf})]$ adduct (Scheme 13A).²³ Andersen and co-workers found that the stabilized biradical $[\text{Cp}^*_2\text{Yb}(\text{dafH})]$ adduct slowly eliminated H_2 to give $[\text{Cp}^*_2\text{Yb}(\text{daf})]$; in addition the dinuclear Yb complex prepared with 9,9'-bis-4,5-diaza-9H-fluorene ligand also thermally produces $[\text{Cp}^*_2\text{Yb}(\text{daf})]$ (Scheme 13B).²⁴ The mechanism for dihydrogen formation is proposed on the basis of kinetic and labelling experiments to involve the dinuclear Yb complex as an intermediate.²⁴ The oxidation states of Yb was described as being intermediate between +2 and +3 for both $[\text{Cp}^*_2\text{Yb}(\text{dafH})]$ and $[\text{Cp}^*_2\text{Yb}(\text{bpy})]$ with an equilibrium between the at least two low-lying open-shell singlet states.²⁴ Andersen and co-workers also performed calculations on 4,5-diazafluorene ligand in an attempt to understand how even though $[\text{Cp}^*_2\text{Yb}(\text{dafH})]$ and $[\text{Cp}^*_2\text{Yb}(\text{bpy})]$ have multiconfigurational ground states, they differ in reactivity where $[\text{Cp}^*_2\text{Yb}(\text{dafH})]$ eliminates H_2 , and $[\text{Cp}^*_2\text{Yb}(\text{bpy})]$ does not.²⁴ The *dafH* ligand has unpaired spin density distributed in various p_π orbitals on the nitrogen and carbon atoms.²⁴ The LUMO+1 of $2b_1$ symmetry has unpaired spin density at the 9-position carbon, this orbital is possibly responsible for the chemistry observed.²⁴ The unpaired spin density at the 9-position aids in the cleavage of the C–H allowing the formation of H_2 and a C–C bond to give the dinuclear Yb complex.

Dalton Transactions




Scheme 13. Two studies involving the formation of H_2 from a coordinated *daf* ligand. A) Isolation of $[\text{Ti}(\text{Cp}^*)_2(\text{daf})]$.²³ B) Synthesis of $[\text{Yb}(\text{Cp}^*)_2(\text{daf})]$ from stabilized biradical $[\text{Yb}(\text{Cp}^*)_2(\text{dafH})]$, or the dinuclear complex $[\text{Yb}(\text{Cp}^*)_2(\text{bdafH}_2)]$.²⁴

Our group also investigated the reactivity of the zwitterionic Ru(II) diazafluorene complex $[\text{Ru}(\text{daf})(\text{PPh}_3)_2(\text{H})(\text{N}_2)]$ toward CO_2 and uncovered an interesting example of ligand-based reactivity.¹¹⁵ At room temperature $[\text{Ru}(\text{daf})(\text{PPh}_3)_2(\text{H})(\text{N}_2)]$ selectively and reversibly undergoes a formal insertion of CO_2 into a remote ligand C–H bond to generate a monoanionic 4,5-diazafluorenyl-9-carboxylic acid ligand (*dafCO}_2\text{H}^-*) on Ru(II) (Scheme 12).¹¹⁵ The activation of CO_2 in our system occurs at the ligand backbone remote from the metal centre where the metal's role is to adjust the nucleophilicity of the ligand-based carbanion, the acidity of the C–H bond involved in proton migration, as well as the strength of the newly formed C–C bond. Given the unusual situation of having an actor ligand and a spectator metal centre, a variety of spectator metal centres were used to tune the reactivity and electronics of the actor *daf* ligand for tandem CO_2 and C–H activation.²² Since Rh(III) is isoelectronic with Ru(II), complex $[\text{Rh}(\text{daf})(\text{PPh}_3)_2(\text{H})_2]$ also reacts with CO_2 in an analogous way where CO_2 reversibly inserts into the C–H bond of the ligand backbone (Scheme 14).²²

In contrast, when the more electron-rich Rh(I) complex $[\text{Rh}(\text{daf})(\text{PPh}_3)_2]$ is placed under CO_2 , a dinuclear Rh(I) complex, $[(\text{Rh}(\text{PPh}_3)_2)_2(\text{dafCO}_2)]$, is formed where the two Rh(I) centres are bridged by a dianionic 4,5-diazafluorenyl-9-carboxylate ligand (dafCO_2^{2-}), along with the formation of free *dafH* (Scheme 14).²² The result of tuning *daf* with a more electron rich metal centre is increased basicity of the ligand-based carbanion which can deprotonate the carboxylic acid initially formed from CO_2 insertion into the ligand C–H bond. This proton transfer gives $[\text{Rh}(\text{dafH})(\text{PPh}_3)_2]^+$ and $[\text{Rh}(\text{dafCO}_2)(\text{PPh}_3)_2]$; the carboxylate of the latter replaces the *dafH* ligand of the former to yield the dinuclear product.²² As a result CO_2 is trapped by the second metal centre, and also there is no proton on O that can engage in proton transfer necessary for the decarboxylation.²² The dinuclear

[[Rh(PPh₃)₂]₂(dafCO₂)] complex was reacted with H₂ to attempt ligand-based CO₂ reduction.¹¹⁶ A series of stepwise stoichiometric reactions with H₂, NMR experiments at low temperatures with added PPh₃ or CO₂, along with ¹³C-labelling experiments were conducted in an attempt to gain some mechanistic insight.¹¹⁶ Upon the addition of a CO₂ and H₂ gas mixture to [[Rh(PPh₃)₂]₂(dafCO₂)], a mixture of the carboxylated Rh(III) complex [Rh(dafCO₂H)(PPh₃)₂(H)₂] and [Rh(PPh₃)₂(H)₂(κ²-HCO₂)] results (Scheme 14).¹¹⁶




Scheme 14. Chemistry of Rh 4,5-diazafluorenyl complexes with CO₂ and H₂.^{20, 22, 116}

It is worth noting that the carboxylic acid intermediate in the reaction between [Rh(daf)(PPh₃)₂] and CO₂ could not be isolated or even observed in NMR experiments, presumably because the highly basic carbanion in [Rh(daf)(PPh₃)₂] effected by the electron rich Rh(I) centre deprotonates the carboxylic acid to trigger the formation of the final product too quickly before the concentration of the carboxylic acid intermediate could build up. In order to observe and isolate both the kinetic and thermodynamic products of CO₂ reaction, the slightly less electron donating Cu(I) was used as the spectator metal centre.²² When CO₂ is added to [Cu(daf)(IPr)] in toluene the kinetic product [Cu(dafCO₂H)(IPr)] precipitates from solution (Scheme 15).²² Since [Cu(dafCO₂H)(IPr)] and [Cu(daf)(IPr)] are both soluble in DMSO, the reaction between these two species is readily observable when DMSO is used as the solvent for both the carboxylation of [Cu(daf)(IPr)] with CO₂ and the decarboxylation of [Cu(dafCO₂H)(IPr)] under N₂ atmosphere. In both cases, a dinuclear Cu(I) complex [(Cu(IPr))₂(dafCO₂)] and dafH are obtained as thermodynamic products (Scheme 15).²² In both the Cu(I) and Rh(I) cases the metal 4,5-diazafluorenyl and the metal 4,5-diazafluorenyl-9-carboxylic acid complexes react with each other in solution, in contrast to the less electron rich Ru(II) and Rh(III) systems.

II.2. 4,5-Diazafluorene derivatives as ligands in catalysis:


Nitrogen-donor ligands have been used extensively in oxidative aerobic organic transformations, especially given their robustness versus traditional phosphine ligands under oxidizing reaction conditions. Stahl and co-workers initially explored the use of 4,5-diazafluorene derivatives in oxidative organic reactions where O₂ is the oxidant.⁷⁶ A variety of nitrogen-donor ligands were screened in the Pd-catalyzed aerobic allylic acetoxylation of allylbenzene; most of the

ligands screened gave low yields of the cinnamyl acetate product (Table 1).⁷⁴ Conversely when 9,9-dimethyl-4,5-diazafluorene (Me₂daf)



Scheme 15. Reaction of [Cu(daf)(IPr)] with CO₂ which gives [Cu(dafCO₂H)(IPr)] as the kinetic product and a mixture of [(Cu(IPr))₂(dafCO₂)] and dafH as thermodynamic products.²²

Table 1. Nitrogen-based ligands screen in the Pd-catalyzed aerobic allylic acetoxylation of allylbenzene.



was used as the ligand a 50% yield of cinnamyl acetate was obtained, the yield was further improved to 81% when dafo was used as the ligand.⁷⁴ The structures of 4,5-diazafluorene derivatives seem to have a large impact on the catalytic results obtained relative to the other nitrogen-donor ligands tested. Both the ability of dafo to withdraw electron density through π back-bonding, and the unique ligand bite angle may play a role in the catalysis. Recently Stahl reported the mechanistic investigations where they suggest that the dafo ligand promotes the C–O reductive elimination, but further studies are needed.⁷⁷ Stahl and co-workers have also demonstrated the use of 4,5-diazafluorene derivatives as ancillary ligands in the aerobic Pd-catalyzed cross-coupling of indoles with benzene.⁷⁵ The regioselectivity for arylation at the C2- vs. the C3- position of the indole compound was dramatically affected


PERSPECTIVE

Dalton Transactions


by the identity of 4,5-diazafluorene derivative and the anionic ligand used (Scheme 16).⁷⁵

Stahl,⁷⁸ along with Zhao and Huang,⁸⁰ independently and simultaneously reported the use of the dafo ligand in Pd-catalyzed aerobic dehydrogenation to form α,β -unsaturated aldehydes, ketones, esters, and azobenzenes (Scheme 17). Typically enones and other α,β -unsaturated carbonyl compounds are prepared in stepwise protocols. The aerobic Pd-dafo catalyzed reaction is a much more efficient alternative.


Recently it has been shown that dafo is an effective ligand for the Pd-catalyzed aerobic dehydrogenative Heck reaction to couple furans and thiophenes with cinnamic acid and stilbene derivatives.⁸⁵ In addition a variety of alkenes could be coupled with ferrocene using a Pd dafo catalyst in an aerobic dehydrogenative Heck reaction (Scheme 18).⁸³ A combination of kinetics, competition and ESI-MS (to characterize catalytic intermediates) experiments suggest that dafo plays a role at each stage of the catalytic cycle,⁸⁵ i.e., the dafo influences C–H bond activation, insertion of alkenes, the stereo-selective step, and the regeneration of the catalyst with O₂.⁸⁵



Scheme 16. Pd-catalyzed aerobic coupling of indoles with benzene using 4,5-diazafluorene derivatives as ligands.⁷⁵



Scheme 17. Pd dafo catalyzed aerobic dehydrogenation to form double bonds.^{78, 80}



Scheme 18. Pd-dafo-catalyzed aerobic Heck coupling reaction of an alkene with ferrocene.⁸³

Elsevier and co-workers have looked at the influence of various nitrogen chelates on Pd catalyzed C–C bond formation reactions.⁷⁹ Zerovalent mono and binuclear palladium and platinum bis(quinone) complexes of dafoH and dafo have been prepared, where the 4,5-diazafluorene derivative either acts as a chelate, monodentate, or bridging ligand.^{17, 81, 82} In the reaction of cinnamylchloride with benzyl Grignard the regioselectivity when palladium complexes of 4,5-diazafluorene derivatives are used is extremely high for

substitution at the less substituted allylic carbon with less than 5% of homo-coupling product, a sharp contrast to when phosphine complexes are used.⁷⁹

[Rh(daf)(COD)] can catalyze the hydrogenation of olefins, however it is not as fast a catalyst when compared with [Rh(PPh₃)₃Cl] or [Ru(PPh₃)₃Cl₂], and does not hydrogenate internal olefins.¹⁹ Wilkinson's catalyst [Rh(PPh₃)₃Cl] dissociates in solution and produces hydride species under H₂. In contrast, no hydride was observed when a solution of [Rh(daf)(COD)] was exposed to H₂.


[Rh(dafH)(PPh₃)₂(H)₂]Cl was also found to be a selective olefin hydrogenation catalyst, and can even hydrogenate internal olefins and substrates with pyridyl or carbonyl groups.²⁰ The chloride counterion appears to play a role in the catalysis, i.e., if the counterion is replaced with triflate the complex is inactive towards olefin hydrogenation. Exogenous chloride anions however appear to slow the catalysis.

Recently it was shown that 1,5-dihydro-2H-cyclopenta[2,1-b:3,4-b']dipyridin-2-one (a 4,5-diazafluorene derivative with keto-enol tautomerism), L⁷, can be coordinated to Ru(II) to give [Ru(L⁷(bpy)₂)(PF₆)₂].¹¹⁷ [Ru(L⁷(bpy)₂)(PF₆)₂] was used as a photosensitizer in dye-sensitized solar cells (DSSCs) and as a catalyst for the transfer hydrogenation of ketones.¹¹⁷

The complex [MoO₂Cl₂(dafo)] was quite an active and selective olefin epoxidation catalyst when compared with other polypyridine ligands.⁶³ Of the polypyridine ligands tested the best catalysts seemed to have moderate donating capability and little steric hindrance with respect to the {MoO₂Cl₂} fragment.⁶³

Huang and co-workers reported the use of dafo as a ligand in organic transformations mediated by a gold catalyst and secondary amine that work synergistically (Scheme 19).^{118, 119}


Tri- and tetrasubstituted allenes can be synthesized in an α -vinylideneation or an α -vinylideneation/ γ -vinylideneation cascade from aldehydes and silyl-EBX.¹¹⁸ The trisubstituted allenyl aldehyde product produced in this gold-amine system is sensitive toward O₂ to yield ynone products, the result of oxidative C–C bond cleavage (Scheme 19).¹¹⁹



Scheme 19. A) α -Vinylideneation of aldehydes to give trisubstituted allenes, B) α -vinylideneation/ γ -vinylideneation cascade to give tetrasubstituted allenes. C) Aerobic C–C bond cleavage reaction with aldehydes which yields an ynone product. Both reactions are catalyzed by an Au-dafo/amine catalyst system.^{118, 119}


Stahl and coworkers have demonstrated Pd catalyzed aerobic intramolecular aryl C–H amination to give carbazole derivatives as products (Scheme 20).⁸⁶ The 1,4-dioxane solvent decomposes in the presence of O₂ to form *in situ* an alkyl peroxide that promotes the efficient Pd catalyzed aryl C–H amination.⁸⁶ The neutral dafo ligand was found to be the most efficient at promoting carbazole synthesis starting from the *N*-

benzenesulfonyl-2-aminobiphenyl test substrate giving an 80% yield, Me₂daf can also promote the reaction giving a 33% yield.⁸⁶



Scheme 20. Pd catalyzed aerobic aryl C-H amination.⁸⁶

One interesting catalytic application of 4,5-diazafluorene derivatives is the introduction of a catalytically active metal to the surface of nanoparticles to create nanocatalysts. Silica-coated magnetite nanoparticles (SMNPs) were silanated with 3-aminopropyltriethoxysilane, the exposed –NH₂ group was reacted with dafo to give the imine that was subsequently reduced to the amine (Scheme 21A).¹²⁰ The 4,5-diazafluorene-functionalized SMNPs were coordinated with Pd(II) to give a Pd-4,5-diazafluorenyl SMNPs catalyst for C(sp²)–C(sp²) cross-coupling reactions (Scheme 21B).¹²⁰ An elegant feature of this heterogeneous SMNPs nanocatalyst is that it can be separated from the reaction mixture by using an external magnet, and reused successfully in subsequent cross-coupling reactions. Using a similar protocol silica nanospheres were functionalized with a 4,5-diazafluorene-imine group, and subsequently reacted with ferric chloride.¹²¹ The silica nanosphere-based iron nanocatalyst was used to perform the one-pot coupling reaction of a terminal alkyne, dichloromethane, and amines to give propargylamines.¹²¹ In addition, the silica nanosphere-based iron catalyst could be recycled at least seven times without any appreciable loss of catalytic activity.¹²¹



Scheme 21. A) Synthesis of magnetically-separable Pd-4,5-diazafluorenyl SMNPs nanocatalyst, and B) their use as catalysts in cross-coupling reactions.¹²⁰

II.3. Photophysical and photochemical applications

II.3.1 As photosensitizers.

Ru complexes of dafH of the form [Ru(dafH)_x(bipyridine ligand)_n]²⁺ were studied as analogues to the well-known [Ru(bpy)₃]²⁺ complex because of their potential applications as photosensitizers in photochemical water-splitting processes and DSSCs. Replacing one bpy ligand in the parent complex [Ru(bpy)₃]²⁺ with a dafH ligand results in very little change in the room temperature absorption spectra, and very little change in the low temperature emission spectra (λ_{max} (77 K) = 578 nm for [Ru(bpy)₃]²⁺, λ_{max} (77 K) = 574 nm for

[Ru(dafH)(bpy)₂]²⁺ in EtOH/MeOH (4:1)).⁹⁻¹⁰ The most substantial difference is that the emission quantum yield of [Ru(bpy)₃]²⁺ in water ($\phi(298\text{ K}) = 0.042$, $\tau(298\text{ K}) = 630\text{ ns}$) is ~50 fold larger than that of [Ru(dafH)(bpy)₂]²⁺ ($\phi(298\text{ K}) = 8 \times 10^{-4}$) and the lifetime of [Ru(dafH)(bpy)₂]²⁺ was so short that it could not be determined.⁹⁻¹⁰ The dafH ligand with a longer N–N distance was found to be lower than bpy in the spectrochemical series which translated into an energetic lowering of ligand field excited states and a dramatic decrease in emission intensity approaching room temperature.⁹⁻¹⁰

A series of multinuclear Ru(II) complexes supported by multinucleating ligands possessing two or more chelating 4,5-diazafluorenyl moieties have been explored.¹²²⁻¹³⁸ The spacers between the coordinating 4,5-diazafluorenyl moieties have been varied substantially to allow for the metal–metal separation, and the degree of intramolecular charge transfer to be finely tuned. In some cases disappointingly the Ru(II) complexes had extremely short excited state lifetimes at room temperature on the order of ~0.03 ns (e.g., Ru complex of the ditopic ligand with admantyl spacer).¹²²

One of the major challenges in the field of DSSCs is to identify molecules that can efficiently harvest sunlight when incorporated into mesoscopic semiconductor thin films such as mesoporous TiO₂. Ru(II) complexes of 4,5-diazafluorene derivatives have been investigated as sensitizers for TiO₂ especially for their ability to inject electrons into TiO₂ from their excited-states. Following adsorption of [Ru(dafo)(bpy)₂]²⁺ onto TiO₂ it was noted by Heuer, Meyer and coworkers that the dafo ligand engaged in a ring-opening reaction resulting in a coordinated 3-(CO₂H)-2,2'-bipyridine ligand with the carboxylate group anchoring the complex to TiO₂.⁶⁵ The control experiment of independently anchoring the analogous Ru(II) complex of 3-(CO₂H)-2,2'-bipyridine gave comparable results; however, the excited-state injection yield for [Ru(dafo)(bpy)₂]²⁺/TiO₂ system was consistently lower.⁶⁵ Ru(II)-dafo complexes with either one or two 4,4'-dicarboxy-2,2'-bipyridine (dcbpy) ligands to anchor the complex onto TiO₂ have also been explored.^{69, 139} The thermal stability of [Ru(dafo)(dcbpy)(SCN)₂] at high temperatures has also been investigated where the final products are Ru and RuS₂; again the ring-opening of dafo seems to play a role in complex decomposition.⁶⁴ Another ligand design tested to anchor Ru(II) polypyridine complexes to TiO₂ was an ambidentate ligand (4,5-diazafluorene-9-ylidene)malonic acid (dfm), which has a 4,5-diazafluorenyl chelate with an olefin bridge to two carboxylic acid groups.¹⁴⁰ The metal-to-ligand charge transfer (MLCT) excited state lifetime of [Ru(dfm)(bpy)₂]²⁺ was extremely short in solution, yet the interfacial electron transfer to TiO₂ was efficient ($\phi_{\text{inj}} = 0.70 \pm 0.05$).¹⁴⁰ Some Ru photosensitizers with record high extinction coefficients possess 4,5-diazafluorenylidene ligands substituted with a 1,3-dithiole group ($\epsilon \geq 40,000\text{ M}^{-1}\text{cm}^{-1}$ @ 470 nm), displaying rapid and efficient charge injection to TiO₂.^{141, 142} In addition, Ru(II) complexes of 4,5-diazafluorene functionalized at the 9-position with carbazole groups,¹⁴³ aryl groups,¹⁴⁴ or bis(thiophene)pyrrolyl groups,¹³⁹ have all been used to create DSSCs based on TiO₂.

PERSPECTIVE

Dalton Transactions

II.3.2 As emitters.

Light-emitting electrochemical cells (LECs) are an attractive alternative to organic light-emitting diodes (OLEDs). LECs can be easily prepared with a solution-based process, only require a single emissive layer, use air-stable electrodes such as gold, and can operate at very low voltages with high power efficiencies.¹⁴⁵⁻¹⁴⁸ Cationic Ir(III) complexes of 4,5-diazafluorene derivatives, which are conductive and phosphorescent, can be used effectively in LECs.

Cationic Ir(III) complexes of the form $[\text{Ir}(\text{L}^n)(\text{dfppz})_2](\text{PF}_6)$ ($n=2,8,9,10,11$) with cyclometallated 1-(2,4-difluorophenyl)pyrazole (dfppz) have been prepared, where the parent 4,5-diaza-9,9'-spirobifluorene ligand (L^5) is substituted at the 3,6-positions with one (L^8) or two (L^9) pendant phenyl groups (see Figure 5 for labelling scheme).¹⁴⁸ Interestingly these pendant phenyl groups can engage in π - π stacking interactions with the dfppz ligand(s) within complexes; these interactions may limit the elongation of the Ir-ligand bonds in the excited state protecting the metal centre.¹⁴⁸

A white-light emitting LEC was created using a mixture of red- green- and blue-emitting cationic Ir(III) complexes, where the green and blue emitters have 4,5-diazafluorene derivatives disubstituted at the C9-position with gem-diaryl (L^{10}) or gem-dialkyl (L^{11}) groups (Figure 5).¹⁴⁶ Substitution at the C9-position was found to be crucial for the synthesis of Ir(III) complexes with high photoluminescence quantum yields.¹⁴⁶ Likewise, Ir(III) complexes $[\text{Ir}(\text{L}^5)(\text{ppy})_2](\text{PF}_6)$ (where ppy is a cyclometallated 2-phenylpyridine derivative) **A** and **B** (Figure 5) show high electroluminescence efficiencies: 7.1%, $22.6 \text{ Im}\cdot\text{W}^{-1}$ for orange **A**, and 7.1%, $26.2 \text{ Im}\cdot\text{W}^{-1}$ for green **B**,¹⁴⁵ owing to the spiro structural feature of L^5 , which limits self-quenching phenomena without greatly perturbing the energy gaps of the compounds.

A conceptually interesting approach toward incorporating three functions into a single molecule, a luminescent chromophore with hole and electron transport capability was explored by the Wong group. In luminescent Ir(III) complex **C** (Figure 5), the ligand L^{12} has an electron-transporting 4,5-diazafluorene site, and hole-transporting $-\text{NPh}_2$ groups.¹⁴⁷ Unfortunately the LEC device performance using this trifunctional Ir(III) complex was rather low. There is no $^3\text{MLCT}$ contribution to the lowest transition for the complex in the triplet state.¹⁴⁷

Luminescent lanthanide complexes exhibit very sharp emission bands. The design and tuning of sensitizing ligands that allow for efficient ligand-to-metal energy transfer is an area of particular interest. Luminescent lanthanide tris(β -diketonate) complexes of various 4,5-diazafluorene derivatives such as dafo,^{94-96, 149} 9,9-diaryl substituted 4,5-diazafluorene,¹⁵⁰ 4,5-diazafluorene-9-imine derivatives,¹⁵¹ and 4,5-diaza-9,9'-spirobifluorene¹⁵² have been synthesized and characterized. Several of these lanthanide complexes exhibit NIR luminescence⁹⁴⁻⁹⁶ photo- and electroluminescence,^{150, 152} and even triboluminescence.¹⁴⁹




Figure 5. Luminescent cationic Ir(III) complexes of 4,5-diazafluorene derivatives for use in solid-state light-emitting electrochemical cells.

Green phosphorescent Re(I) complexes of various 4,5-diazafluorene derivatives have been investigated for their performance in OLEDs.¹⁵³⁻¹⁵⁶ In comparing $[\text{Re}(\text{N}^*\text{N})(\text{CO})_3\text{Br}]$ type complexes of dafH and a 9,9-di(ethoxyphenyl) substituted 4,5-diazafluorene derivative, both complexes give approximately the same emission wavelength, however the complex with bulky groups at the C9 position alleviates to a large extent the self-quenching at high doping concentrations.¹⁵³ A dinuclear Re(I) complex of 9,9'-bi-4,5-diazafluorenyl can serve as a highly efficient green phosphorescent emitter in OLEDs with a maximum luminance of $2026 \text{ cd}/\text{m}^2$ and a peak current efficiency of $8.2 \text{ cd}/\text{A}$.¹⁵⁴ Non-doped devices containing $[(\text{L}^5)\text{Re}(\text{CO})_3\text{Br}]$ had outstanding performance with a further improved peak luminance of $8531 \text{ cd}/\text{m}^2$ and maximum current efficiency of $16.8 \text{ cd}/\text{A}$.¹⁵⁶

The phosphine Cu(I) complexes of 3,3'-methylene-4,4'-diphenyl-2,2'-biquinoline (mdpbq) exhibited red phosphorescence (Figure 6).¹⁵⁷ Complexes of the extremely rigid mdpbq ligand showed decent photoluminescence quantum yields in 20 wt% poly(methylmethacrylate) (PMMA) films, 0.56 and 0.43 for $[\text{Cu}(\text{mdpbq})(\text{PPh}_3)_2](\text{BF}_4)$ $[\text{Cu}(\text{mdpbq})(\text{DPEphos})](\text{BF}_4)$, respectively.¹⁵⁷ The OLEDs doped with these Cu(I) complexes gave a current efficiency up to $6.4 \text{ cd}/\text{A}$ for a multi-layer device.¹⁵⁷




Figure 6. Structures of red phosphorescent Cu(I) complexes.¹⁵⁷

Ag-dafo complexes with either carborane-based diphosphines, or simple classical diphosphine ligands of the general formula $[\text{Ag}(\text{dafo})(\text{P}-\text{P})]\text{OTf}$ are luminescent. The emissions seem independent of the phosphine ligand or the coordination environment of the Ag^+ ion.¹⁵⁸ The free dafo emits at 537 nm ($\tau = 5 \text{ ns}$), while the luminescent Ag-dafo complexes emit across the blue to orange region of the spectrum with lifetimes also in the ns range, similar to free

dafo.¹⁵⁸ The dafo ligand centred transitions seem to be responsible for the luminescence behaviour of [Ag(dafo)(P-P)]OTf and [Ag(dafo)(PPh₃)(OTf)] complexes, where the electron density at the Ag centre tuned by the different phosphine ligands modified the emission energy.¹⁵⁸

II.3.3 As harmonophores in non-linear optics.

There is great interest in utilizing coordination complexes as harmonophores with second-order non-linear optical (NLO) properties since the metal centres offer additional means of tuning the electronic properties which affect the NLO response. The metal centres could not only allow for either metal-to-ligand charge transfer (MLCT) or ligand-to-metal charge transfer (LMCT) to take place, but also modulate the energy of intraligand charge transfer (ILCT) transitions. Moreover, the metal centre is tunable in terms of the identity of the metal element, oxidation state, and coordination sphere. Complexes of 4,5-diazafluorene derivatives have found use in this field.

Ir complexes of the form [Ir(L^ΛL)(ppy)₂]ⁿ⁺ where the bidentate L^ΛL is either daFH, 9-fulleriden-4,5-diazafluorene, or cyclometallated 9-fulleriden-4-azafluorene can be used as harmonophores with second-order NLO properties.¹⁵⁹ The introduction of the fullerene moiety weakens the interaction between the cationic Ir(III) complex and the anion, which also leads to an overall increase in the NLO response with large and negative $\mu\beta_{1,907}$ values (-600 to -2190 × 10⁻⁴⁸ esu).¹⁵⁹ For the charge neutral complex with cyclometallated 9-fulleriden-4-azafluorene ligand the $\mu\beta_{1,907}$ value is lower than the cationic complexes.¹⁵⁹ In addition the second-order NLO properties of Ru(II) complexes of the forms [Ru(N^ΛN)(PPh₃)₂Cl₂] and [Ru(N^ΛN)(CO)₂Cl₂] where the N^ΛN chelate is either daFH or 9-fulleriden-4,5-diazafluorene, have also been examined.¹⁶⁰ The greater absolute values of $\mu\beta_{1,907}$ for complexes with the fullerene substituted ligand compared to those with unsubstituted daFH suggest the importance of a highly polarizable C₆₀ group.¹⁶⁰

Ru(II) bipyridine complexes of 4,5-diazafluorene-9-imine derivatives with long alkyl chains off of the imine moiety have been incorporated as surfactants into Langmuir-Blodgett thin films and have activities for second order harmonic generation that are 2.6 to 3.6 times greater than that of the organic standard (*E*)-*N*-methyl-4-(2-(4-octadecyloxyphenyl)ethenyl)pyridinium iodide.¹⁶¹ Similarly [Re(N^ΛN)(CO)₃Cl] complexes of similar 4,5-diazafluorene-9-imine derivatives can also be formed into stable Langmuir-Blodgett thin films, though a lower than expected measurement for the second-order harmonics NLO signal was observed.¹⁶²

Zn(II) complexes of the highly conjugated push-pull 4,5-diazafluorene-9-ylidene ligand family terminated with either a *N,N*-dibutylamino, or azulenyl moiety exhibit amongst the highest reported $\mu\beta_{1,907}$ value for a Zn(II) complex.¹⁶³ With respect to the free ligands, coordination to Lewis acidic Zn(II) enhances the $\mu\beta_{1,907}$ value presumably by red-shifting the ILCT transition.¹⁶³

II.3.4 As luminescent sensors.

Ru polypyridine complexes of 4,5-diazafluorene derivatives possessing a crown ether moiety attached to the ligand backbone have been synthesized. In most cases the binding with alkali metal cations was studied, where both the spectroscopic and electrochemical properties of the complex were perturbed by cation binding.^{131, 135, 137, 164, 165}

Duan, Bai, and coworkers reported [Ru(L¹³)(bpy)₂]²⁺ (where L¹³ is 4,5-diazafluorene-9-one-2,4-dinitrophenylhydrazone) as a selective chromo- and fluorogenic dual responding fluoride sensor.¹⁶⁶ Not only does the presence of F⁻ enhance the luminescence intensity but also triggers a dramatic color change from yellow to magenta. The Ru(II) complex could even be adsorbed onto paper to allow for the preparation of colorimetric testing strips for F⁻ concentration in water. Interestingly, even when the concentration of F⁻ is as low as 1 mg·L⁻¹, the color change is visible to naked eyes.

II.4. Bioinorganic chemistry of 4,5-diazafluorene derivatives:

Recently Ru(II) complexes [Ru(L¹⁴)(bpy)₂](PF₆)₂ (where L¹⁴ is a 4,5-diazafluorene-9-imine ligand with various *N*-aryl groups) have been found to be effective topical antibiotics against the bacteria *Staphylococcus aureus* which is resistant to the antibiotic methicillin.¹⁶⁷ In particular the derivative with a -OC₇H₁₅ group attached to the *N*-aryl group, which is non-toxic toward human skin keratinocyte cells, exhibits strong microbicidal and bacterial growth inhibitory effects.¹⁶⁷ One possible mechanism for how these Ru(II) complexes are active against *S. aureus* is through the generation of reactive oxygen species.¹⁶⁷

Two crystal structures of the bis-dafo Ag(I) nitrate complexes were obtained where the nitrate ion is either bound to the metal centre in the case of [Ag(dafo)₂(NO₃)],¹⁶⁸ or not in the case of [Ag(dafo)₂](NO₃)·H₂O.^{88, 168} These complexes exhibited broad spectrum antibacterial properties against six different clinically resistant strains of diabetic foot bacteria, and had a significantly lower minimum inhibitory concentration (MIC) compared with currently available commercial antibiotics.¹⁶⁸ Initial studies also showed that the silver complexes could be loaded into hydrogels for possible incorporation into wound dressings.¹⁶⁸ The *in vitro* linear dichroism studies also showed that both dafo and the silver complexes bind to calf thymus DNA, however the complexes showed significantly stronger binding to DNA compared with the free ligand.¹⁶⁸

The free 9-diazo-4,5-diazafluorene can be photochemically activated to release dinitrogen as a byproduct and generate a triplet carbene.¹⁶⁹ Matrix photolysis experiments of complex [Cu(9-diazo-4,5-diazafluorene)₂(NO₃)₂] (Figure 7) indicate the formation of a Cu(I)-L^{•+} species (S=1/2), where the radical is primarily localized at the C9-position.¹⁶⁹ Presumably the Cu(II) center is reduced by the triplet carbene formed via photolysis. In solution the nitrate ligands are labile, and other chemical or biological substrates such as DNA can coordinate to Cu. Consequently, [Cu(9-diazo-4,5-diazafluorene)₂(NO₃)₂] cleaves DNA upon irradiation with visible light under anaerobic conditions; in contrast, the free 9-diazo-4,5-diazafluorene

ligand is significantly less effective for the photocleavage of DNA.¹⁶⁹⁻¹⁷¹




Figure 7. Structure of [Cu(9-diazo-4,5-diazafluorene)₂(NO₃)₂].^{169, 170}

Conclusions

Since the first report on the synthesis of dafH, the chemistry of dafH derivatives as ligands has evolved into a vibrant field. From the fundamental chemistry point of view, the parent compounds dafH and dafo can be derivatized in many ways to achieve the desired functions. In particular, the functionalization at the C9 position has been in the spotlight for generating new ligand series. Many of the ambidentate derivatives have been utilized to construct coordination polymers, self-assembled macrocycles, and heteromultimetallic complexes with great specificity. The reactive nature of the 9-position of dafH, daf and dafo ligands gave rises to the actor-ligand behavior in the corresponding metal complexes, which distinguish dafH derivatives from bpy. From an applications point of view, dafH derivatives have found use in many areas. Their use as ancillary ligands in catalysis has been fruitful especially in transformations performed under oxidizing conditions. The photochemistry and physics of metal complexes of 4,5-diazafluorene derivatives have been studied as emitters in solid state lighting, photosensitizers in DSSCs, and harmonophores in non-linear optics. Biological studies of 4,5-diazafluorene metal complexes have included DNA binding and cleavage, and antimicrobial properties. Further research may lead to new types of fundamental reactivities as well as applications.

References

- 1 K. Kloc, J. Mlochowski and Z. Szulc, *J. Prakt. Chem.*, 1977, **319**, 959-967.
- 2 K. Kloc, Z. Szulc and J. Mlochowski, *Heterocycles*, 1978, **9**, 849-852.
- 3 I. F. Eckhard and L. A. Summers, *Aust. J. Chem.*, 1973, **26**, 2727-2728.
- 4 R. P. Thummel, F. Lefoulon and R. Mahadevan, *J. Org. Chem.*, 1985, **50**, 3824-3828.
- 5 M. J. Plater, S. Kemp and E. Lattmann, *J. Chem. Soc., Perkin Trans. 1*, 2000, 971-979.
- 6 S. Nishigaki, H. Yoshioka and K. Nakatsu, *Acta Cryst.*, 1978, **B(34)**, 875-879.
- 7 L. Maguire, C. M. Seward, S. Baljak, T. Reumann, Y. Ortin, E. Banide, K. Nikitin, H. Müller-Bunz and M. J. McGlinchey, *Eur. J. Inorg. Chem.*, 2009, 3250-3258.
- 8 M. Riklin, A. von Zelewsky, A. Bashall, M. McPartlin, A. Baysal, J. A. Connor and J. D. Wallis, *Helv. Chim. Acta*, 1999, **82**, 1666-1680.

- 9 L. J. Henderson, F. R. Fronczek and W. R. Cherry, *J. Am. Chem. Soc.*, 1984, **106**, 5876-5879.
- 10 W. M. Wacholtz, R. A. Auerbach, R. H. Schmehl, M. Ollino and W. R. Cherry, *Inorg. Chem.*, 1985, **24**, 1758-1760.
- 11 R. P. Thummel, F. Lefoulon and J. D. Korp, *Inorg. Chem.*, 1987, **26**, 2370-2376.
- 12 T. C. Streckas, H. D. Gafney, S. A. Tysoe, R. P. Thummel and F. Lefoulon, *Inorg. Chem.*, 1989, **28**, 2964-2967.
- 13 K. Maruszewski and J. R. Kincaid, *Inorg. Chem.*, 1995, **34**, 2002-2006.
- 14 D. Aguilà, E. Escribano, S. Speed, D. Talancón, L. Yermán and S. Alvarez, *Dalton Trans.*, 2009, 6610-6625.
- 15 K. Ono and K. Saito, *Heterocycles*, 2008, **75**, 2381-2413.
- 16 W. Wong, *Coord. Chem. Rev.*, 2005, **249**, 971-997.
- 17 R. A. Klein, P. Witte, R. van Belzen, J. Fraanje, K. Goubitz, M. Numan, H. Schenk, J. M. Ernsting and C. J. Elsevier, *Eur. J. Inorg. Chem.*, 1998, 319-330.
- 18 F. G. Bordwell, *Acc. Chem. Res.*, 1988, **21**, 456-463.
- 19 H. Jiang and D. Song, *Organometallics*, 2008, **27**, 3587-3592.
- 20 H. Jiang, E. Stepowska and D. Song, *Eur. J. Inorg. Chem.*, 2009, 2083-2089.
- 21 E. Stepowska, H. Jiang and D. Song, *Chem. Commun.*, 2010, **46**, 556-558.
- 22 V. T. Annibale, D. A. Dalessandro and D. Song, *J. Am. Chem. Soc.*, 2013, **135**, 16175-16183.
- 23 P. T. Witte, R. Klein, H. Kooijman, A. L. Spek, M. Polášek, V. Varga and K. Mach, *J. Organomet. Chem.*, 1996, **519**, 195-204.
- 24 G. Nocton, C. H. Booth, L. Maron and R. A. Andersen, *Organometallics*, 2013, **32**, 1150-1158.
- 25 V. T. Annibale, R. Batcup, T. Bai, S. J. Hughes and D. Song, *Organometallics*, 2013, **32**, 6511-6521.
- 26 R. Batcup, F. S. N. Chiu, V. T. Annibale, J. U. Huh, R. Tan and D. Song, *Dalton Trans.*, 2013, **42**, 16343-16350.
- 27 R. Batcup, V. T. Annibale and D. Song, *Dalton Trans.*, 2014, **43**, 8951-8958.
- 28 R. Tan, F. S. N. Chiu, A. Hadzovic and D. Song, *Organometallics*, 2012, **31**, 2184-2192.
- 29 S. A. Baudron, M. W. Hosseini and N. Kyritsakas, *New J. Chem.*, 2006, **30**, 1083-1086.
- 30 S. Baudron and M. Hosseini, *Inorg. Chem.*, 2006, **45**, 5260-5262.
- 31 S. A. Baudron, M. W. Hosseini, N. Kyritsakas and M. Kurmoo, *Dalton Trans.*, 2007, 1129-1139.
- 32 S. A. Baudron and M. W. Hosseini, *Chem. Commun.*, 2008, , 4558-4560.
- 33 G. Nasr, A. Guerlin, F. Dumur, S. A. Baudron, E. Dumas, F. Miomandre, G. Clavier, M. Sliwa and C. R. Mayer, *J. Am. Chem. Soc.*, 2011, **133**, 6501-6504.
- 34 K. Sako, M. Kusakabe and H. Tatemitsu, *Mol. Cryst. Liq. Cryst. Sci. Technol., Sect. A*, 1996, **285**, 101-106.
- 35 Q. Zhu, J. Dai, D. Jia, L. Cao and H. Lin, *Eur. J. Inorg. Chem.*, 2004, 4789-4794.
- 36 Q. Zhu, J. Dai, D. Jia, L. Cao and H. Lin, *Polyhedron*, 2004, **23**, 2259-2264.
- 37 Q. Zhu, Y. Zhang, R. Dai, G. Bian, D. Jia, J. Zhang and L. Guo, *Chem. Lett.*, 2003, **32**, 762-763.
- 38 Q. Zhu, W. Lu, Y. Zhang, G. Bian, J. Gu, X. Lin and J. Dai, *Eur. J. Inorg. Chem.*, 2008, 230-238.
- 39 Q. Zhu, L. Yu, Y. Qin, L. Huo, M. Shao and J. Dai, *CrystEngComm*, 2011, **13**, 2521-2528.
- 40 T. Culpitt, I. A. Guzei, L. C. Spencer, A. Simonson, J. S. Miller, M. R. Wimmer and K. J. Nelson, *Inorg. Chim. Acta.*, 2015, **427**, 162-167.
- 41 M. P. Cifuentes, M. G. Humphrey, G. A. Koutsantonis, N. A. Lengkeek, S. Petrie, V. Sanford, P. A. Schauer, B. W. Skelton, R. Stranger and A. H. White, *Organometallics*, 2008, **27**, 1716-1726.
- 42 O. Pélerin, C. Olivier, T. Roisnel, D. Touchard and S. Rigaut, *J. Organomet. Chem.*, 2008, **693**, 2153-2158.

- 43 J. Zhao, R. Zhao, Q. Yang, B. Hu, F. Liu and X. Bu, *Dalton Trans.*, 2013, **42**, 14509-14515.
- 44 O. Jung and C. G. Pierpont, *Inorg. Chem.*, 1994, **33**, 2227-2235.
- 45 U. Siemeling, I. Scheppelmann, B. Neumann, H. G. Stammler and W. W. Schoeller, *Organometallics*, 2004, **23**, 626-628.
- 46 D. Li and H. Zeng, *Appl. Organomet. Chem.*, 2013, **27**, 89-97.
- 47 D. Li and Z. Tao, *J. Coord. Chem.*, 2013, **66**, 4237-4254.
- 48 B. Machura, M. Wolff, J. Palion, A. Świtlicka, I. Nawrot and K. Michalik, *Struct. Chem.*, 2011, **22**, 1053-1064.
- 49 B. K. Babu, A. R. Biju, S. Sunkari, M. V. Rajasekharan and J. P. Tuchagues, *Eur. J. Inorg. Chem.*, 2013, 1444-1450.
- 50 X. Chen, D. Wang, Q. Luo and R. Wang, *J. Coord. Chem.*, 2009, **62**, 3895-3904.
- 51 F. Dong, L. Che, L. Li and F. Luo, *Transit. Met. Chem.*, 2011, **36**, 125-130.
- 52 P. Kulkarni, S. Padhye and E. Sinn, *Inorg. Chim. Acta*, 2001, **321**, 193-199.
- 53 P. Kulkarni, S. Padhye, E. Sinn, C. E. Anson and A. K. Powell, *Inorg. Chim. Acta*, 2002, **332**, 167-175.
- 54 D. Li, J. Tian, W. Gu, X. Liu, H. Zeng and S. Yan, *J. Inorg. Biochem.*, 2011, **105**, 894-901.
- 55 G. Li, N. Liu, S. Liu and S. Zhang, *Electrochim. Acta*, 2008, **53**, 2870-2876.
- 56 B. Li, B. Li, X. Zhu and Y. Zhang, *Inorg. Chem. Commun.*, 2003, **6**, 1304-1306.
- 57 J. Meng, X. Wang, E. Wang, H. Fu and Y. Zhong, *Transit. Met. Chem.*, 2009, **34**, 361-366.
- 58 S. Menon and M. V. Rajasekharan, *Inorg. Chem.*, 1997, **36**, 4983-4987.
- 59 H. Pang, J. Sha, J. Peng, A. Tian, C. Zhang, P. Zhang, Y. Chen, M. Zhu and Y. Wang, *Inorg. Chem. Commun.*, 2009, **12**, 735-738.
- 60 H. Pang, C. Zhang, J. Peng, Y. Wang, J. Sha, A. Tian, P. Zhang, Y. Chen, M. Zhu and Z. Su, *Eur. J. Inorg. Chem.*, 2009, 5175-5180.
- 61 M. N. Patel, B. S. Bhatt and P. A. Dosi, *Chem. Biodivers.*, 2012, **9**, 2810-2824.
- 62 S. Shao, Y. Zhu, K. Ma, H. Zhao and Y. Qiu, *J. Coord. Chem.*, 2013, **66**, 2702-2711.
- 63 A. Günyar and F. E. Kühn, *J. Mol. Catal. A: Chem.*, 2010, **319**, 108-113.
- 64 F. M. Emen, K. Ocakoglu and N. Kulcu, *J. Therm. Anal. Calorim.*, 2012, **110**, 799-805.
- 65 W. B. Heuer, H. Xia, M. Abrahamsson, Z. Zhou, S. Ardo, A. A. Narducci Sarjeant and G. J. Meyer, *Inorg. Chem.*, 2010, **49**, 7726-7734.
- 66 H. Jiang, E. Stepowska and D. Song, *Dalton Trans.*, 2008, 5879-5881.
- 67 K. Ocakoglu, E. Harputlu, P. Guloglu and S. Erten-Ela, *Inorg. Chem. Commun.*, 2012, **24**, 118-124.
- 68 A. M. Pyle, J. P. Rehmann, R. Meshoyrer, C. V. Kumar, N. J. Turro and J. K. Barton, *J. Am. Chem. Soc.*, 1989, **111**, 3051-3058.
- 69 W. K. Seok, A. K. Gupta, S. Roh, W. Lee and S. Han, *Bull. Korean Chem. Soc.*, 2007, **28**, 1311-1316.
- 70 Y. Wang and D. P. Rillema, *Inorg. Chem. Commun.*, 1998, **1**, 27-29.
- 71 Y. Wang and D. P. Rillema, *Tetrahedron Lett.*, 1997, **38**, 6627-6630.
- 72 Y. Wang, W. Perez, G. Y. Zheng and D. P. Rillema, *Inorg. Chem.*, 1998, **37**, 2051-2059.
- 73 G. Yang, L. Ji, X. Zhou and Z. Zhou, *Transit. Met. Chem.*, 1998, **23**, 273-276.
- 74 A. N. Campbell, P. B. White, I. A. Guzei and S. S. Stahl, *J. Am. Chem. Soc.*, 2010, **132**, 15116-15119.
- 75 A. N. Campbell, E. B. Meyer and S. S. Stahl, *Chem. Commun.*, 2011, **47**, 10257-10259.
- 76 A. N. Campbell and S. S. Stahl, *Acc. Chem. Res.*, 2012, **45**, 851-863.
- 77 T. Diao and S. S. Stahl, *Polyhedron*, 2014, **84**, 96-102.
- 78 T. Diao, T. J. Wadzinski and S. S. Stahl, *Chem. Sci.*, 2012, **3**, 887-891.
- 79 C. J. Elsevier, *Coord. Chem. Rev.*, 1999, **185-6**, 809-822.
- 80 W. Gao, Z. He, Y. Qian, J. Zhao and Y. Huang, *Chem. Sci.*, 2012, **3**, 883-886.
- 81 R. A. Klein, C. J. Elsevier and F. Hartl, *Organometallics*, 1997, **16**, 1284-1291.
- 82 R. A. Klein, R. van Belzen, K. Vrieze, C. J. Elsevier, R. P. Thummel, J. Fraanje and K. Goubitz, *Collect. Czech. Chem. Commun.*, 1997, **62**, 238-256.
- 83 M. Piotrowicz and J. Zakrzewski, *Organometallics*, 2013, **32**, 106-109.
- 84 J. W. Sprengers, M. de Greef, M. A. Duin and C. J. Elsevier, *Eur. J. Inorg. Chem.*, 2003, 3811-3819.
- 85 A. Vasseur, C. Laugel, D. Harakat, J. Muzart and J. Le Bras, *Eur. J. Org. Chem.*, 2015, 944-948.
- 86 A. B. Weinstein and S. S. Stahl, *Catal. Sci. Technol.*, 2014, **4**, 4301-4307.
- 87 Y. Xia, Z. Miao, F. Wang, H. Yao, M. Cui, Y. Ma, Z. Qi and Y. Sun, *J. Organomet. Chem.*, 2015, **779**, 81-85.
- 88 A. R. Biju and M. V. Rajasekharan, *Polyhedron*, 2008, **27**, 2065-2068.
- 89 A. A. Massoud, Y. M. Gohar, V. Langer, P. Lincoln, F. R. Svensson, J. Jänis, S. T. Gårdebjer, M. Haukka, F. Jonsson, E. Aneheim, P. Löwenhielm, M. A. M. Abu-Youssef and L. R. Öhrström, *New J. Chem.*, 2011, **35**, 640-648.
- 90 J. Meng, X. Wang, Y. Ma, E. Wang and X. Xu, *J. Coord. Chem.*, 2008, **61**, 2853-2860.
- 91 B. Machura, I. Nawrot and K. Michalik, *Polyhedron*, 2012, **31**, 548-557.
- 92 B. Machura, M. Wolff and A. Świtlicka, *Inorg. Chem. Commun.*, 2011, **14**, 17-21.
- 93 B. Tong, F. Wu, Q. Mei and Q. Zhang, *Z. Naturforsch.*, 2010, **65**, 511-515.
- 94 S. Dang, J. Yu, X. Wang, Z. Guo, L. Sun, R. Deng, J. Feng, W. Fan and H. Zhang, *J. Photochem. Photobiol. A*, 2010, **214**, 152-160.
- 95 S. Dang, J. Yu, X. Wang, L. Sun, R. Deng, J. Feng, W. Fan and H. Zhang, *J. Lumin.*, 2011, **131**, 1857-1863.
- 96 S. Dang, J. Yu, J. Yu, X. Wang, L. Sun, J. Feng, W. Fan and H. Zhang, *Mater. Lett.*, 2011, **65**, 1642-1644.
- 97 A. Tian, Z. Han, J. Peng, J. Ying, J. Sha, B. Dong, J. Zhai and H. Liu, *Inorg. Chim. Acta*, 2008, **361**, 1332-1338.
- 98 K. Shao, Y. Zhao, Y. Lan, X. Wang, Z. Su and R. Wang, *CrystEngComm*, 2011, **13**, 889-896.
- 99 C. Liu, S. Gao, D. Zhang, Y. Huang, R. Xiong, Z. Liu, F. Jiang and D. Zhu, *Angew. Chem. Int. Ed.*, 2004, **43**, 990-994.
- 100 J. Cano, F. A. Mautner, C. Berger, R. C. Fischer and R. Vicente, *Polyhedron*, 2013, **50**, 240-245.
- 101 C. Wang, C. Yang, S. Tseng, S. Lin, T. Wu, M. Fuh, G. Lee, K. Wong, R. Chen, Y. Cheng and P. Chou, *Inorg. Chem.*, 2004, **43**, 4781-4783.
- 102 P. Peng, F. Li, F. L. Bowles, V. S. P. K. Neti, A. J. Metta-Magana, M. M. Olmstead, A. L. Balch and L. Echegoyen, *Chem. Commun.*, 2013, **49**, 3209-3211.
- 103 A. Pal, B. Biswas, M. Mitra, C. S. Purohit, C. Lin and R. Ghosh, *J. Chem. Sci.*, 2014, **126**, 717-725.
- 104 E. Noh, S. Park, S. Kang, J. Y. Lee and J. H. Jung, *Chem. Eur. J.*, 2013, **19**, 2620-2627.
- 105 S. Wang, Z. Sun, C. Zhang, L. Ni, C. Wang, Y. Gao, L. Lv, J. Chang and W. Hao, *Inorg. Chem. Commun.*, 2014, **41**, 47-50.
- 106 B. Askevold, H. W. Roesky and S. Schneider, *ChemCatChem*, 2012, **4**, 307-320.
- 107 S. Schneider, J. Meiners and B. Askevold, *Eur. J. Inorg. Chem.*, 2012, , 412-429.
- 108 C. Gunanathan and D. Milstein, *Acc. Chem. Res.*, 2011, **44**, 588-602.

- 109 C. Gunanathan and D. Milstein, *Top. Organomet. Chem.*, 2011, **37**, 55-84.
- 110 J. I. van der Vlugt, *Eur. J. Inorg. Chem.*, 2012, , 363-375.
- 111 J. I. van der Vlugt and J. N. H. Reek, *Angew. Chem. Int. Ed.*, 2009, **48**, 8832-8846.
- 112 R. H. Crabtree, *New J. Chem.*, 2011, **35**, 18-23.
- 113 V. T. Annibale and D. Song, *RSC Adv.*, 2013, **3**, 11432-11449.
- 114 J. Ballmann, R. F. Munhá and M. D. Fryzuk, *Chem. Commun.*, 2010, **46**, 1013-1025.
- 115 V. T. Annibale and D. Song, *Chem. Commun.*, 2012, **48**, 5416-5418.
- 116 V. T. Annibale and D. Song, *Organometallics*, 2014, **33**, 2776-2783.
- 117 A. Baysal, M. Aydemir, F. Durap, S. Özkar, L. T. Yildirim and Y. S. Ocak, *Polyhedron*, 2015, **89**, 55-61.
- 118 Z. Wang, X. Li and Y. Huang, *Angew. Chem. Int. Ed.*, 2013, **52**, 14219-14223.
- 119 Z. Wang, L. Li and Y. Huang, *J. Am. Chem. Soc.*, 2014, **136**, 12233-12236.
- 120 M. A. Zolfigol, T. Azadbakht, V. Khakyzadeh, R. Nejatyami and D. M. Perrin, *RSC Adv.*, 2014, **4**, 40036-40042.
- 121 R. K. Sharma, S. Sharma and G. Gaba, *RSC Adv.*, 2014, **4**, 49198-49211.
- 122 L. De Cola, V. Balzani, F. Barigelletti, L. Flamigni, P. Belser and S. Bernhard, *Recl. Trav. Chim. Pays-Bas*, 1995, **114**, 534-&.
- 123 R. N. Warrener, A. B. B. Ferreira, A. C. Schultz, D. N. Butler, F. R. Keene and L. S. Kelso, *Angew. Chem. Int. Ed. Engl.*, 1996, **35**, 2485-2487.
- 124 R. N. Warrener, A. C. Schultz, M. A. Houghton and D. N. Butler, *Tetrahedron*, 1997, **53**, 3991-4012.
- 125 Y. Wang, W. J. Perez, G. Y. Zheng, D. P. Rillema and C. L. Huber, *Inorg. Chem.*, 1998, **37**, 2227-2234.
- 126 F. Cheng, C. He, H. Yin and N. Tang, *Transit. Met. Chem.*, 2013, **38**, 259-265.
- 127 F. Cheng, C. He, H. Yin, N. Tang and N. Hou, *Z. Anorg. Allg. Chem.*, 2013, **639**, 1284-1290.
- 128 F. Cheng, J. Chen, F. Wang, N. Tang and L. Chen, *J. Coord. Chem.*, 2012, **65**, 205-217.
- 129 F. Cheng, J. Chen, C. Sun, N. Tang and L. Chen, *Z. Anorg. Allg. Chem.*, 2011, **637**, 160-166.
- 130 F. Cheng, J. Chen, F. Wang, N. Tang and L. Chen, *Transit. Met. Chem.*, 2011, **36**, 573-578.
- 131 F. Cheng, N. Tang, J. Chen, F. Wang and L. Chen, *Z. Naturforsch. (B)*, 2011, **66**, 923-929.
- 132 F. Cheng, N. Tang, J. Chen, L. Chen, L. Jia and G. Chen, *Inorg. Chem. Commun.*, 2010, **13**, 258-261.
- 133 F. Cheng, L. Chen, G. Bo and N. Tang, *Inorg. Chem. Commun.*, 2009, **12**, 227-230.
- 134 F. Cheng, L. Chen, J. Zhang, J. Wang and N. Tang, *Inorg. Chem. Commun.*, 2009, **12**, 728-730.
- 135 F. Cheng and N. Tang, *Inorg. Chem. Commun.*, 2008, **11**, 939-942.
- 136 F. Cheng and N. Tang, *Inorg. Chem. Commun.*, 2008, **11**, 243-245.
- 137 F. Cheng, N. Tang and L. Chen, *Z. Anorg. Allg. Chem.*, 2008, **634**, 1608-1612.
- 138 F. Cheng, C. He, L. Yao, F. Wang and N. Tang, *J. Coord. Chem.*, 2015, **68**, 704-716.
- 139 K. Ocakoglu, S. Sogut, H. Sarica, P. Guloglu and S. Erten-Ela, *Synth Met*, 2013, **174**, 24-32.
- 140 W. B. Heuer, H. Xia, W. Ward, Z. Zhou, W. H. Pearson, M. A. Siegler, A. A. Narducci Sarjeant, M. Abrahamsson and G. J. Meyer, *Inorg. Chem.*, 2012, **51**, 3981-3988.
- 141 A. Staniszewski, W. B. Heuer and G. J. Meyer, *Inorg. Chem.*, 2008, **47**, 7062-7064.
- 142 M. Abrahamsson, J. H. J. Hedberg, H. Becker, A. Staniszewski, W. H. Pearson, W. B. Heuer and G. J. Meyer, *ChemPhysChem*, 2014, **15**, 1154-1163.
- 143 R. Sivakumar, A. Manivel, M. Meléndrez, J. Martínez-Oyanedel, M. Bunster, C. Vergara and P. Manidurai, *Polyhedron*, 2015, **87**, 135-140.
- 144 K. Ono, H. Tanaka, M. Shiozawa, T. Motohiro, S. Kunikane and K. Saito, *Chem. Lett.*, 2007, **36**, 892-893.
- 145 H. Su, F. Fang, T. Hwu, H. Hsieh, H. Chen, G. Lee, S. Peng, K. Wong and C. Wu, *Adv. Funct. Mater.*, 2007, **17**, 1019-1027.
- 146 H. Su, H. Chen, F. Fang, C. Liu, C. Wu, K. Wong, Y. Liu and S. Peng, *J. Am. Chem. Soc.*, 2008, **130**, 3413-3419.
- 147 H. Chen, K. Wong, Y. Liu, Y. Wang, Y. Cheng, M. Chung, P. Chou and H. Su, *J. Mater. Chem.*, 2011, **21**, 768-774.
- 148 H. Chen, W. Hung, S. Chen, T. Wang, S. Lin, S. Chou, C. Liao, H. Su, H. Pan, P. Chou, Y. Liu and K. Wong, *Inorg. Chem.*, 2012, **51**, 12114-12121.
- 149 X. Chen, X. Zhu, Y. Xu, S. S. S. Raj, S. Öztürk, H. Fun, J. Ma and X. You, *J. Mater. Chem.*, 1999, **9**, 2919-2922.
- 150 Z. Liu, F. Wen and W. Li, *Thin Solid Films*, 2005, **478**, 265-270.
- 151 K. Z. Wang, L. H. Gao, C. H. Huang, G. Q. Yao, X. S. Zhao, X. H. Xia, J. M. Xu and T. K. Li, *Solid State Commun.*, 1996, **98**, 1075-1079.
- 152 S. Wang, J. Zhang, Y. Hou, C. Du and Y. Wu, *J. Mater. Chem.*, 2011, **21**, 7559-7561.
- 153 X. Li, D. Zhang, W. Li, B. Chu, L. Han, T. Li, Z. Su, J. Zhu, S. Wu, Y. Chen, P. Lei, Z. Hu and Z. Zhang, *Synth Met*, 2009, **159**, 1340-1344.
- 154 X. Li, D. Zhang, W. Li, B. Chu, L. Han, T. Li, Z. Su, J. Zhu, Y. Chen, Z. Hu, P. Lei and Z. Zhang, *Opt. Mater.*, 2009, **31**, 1173-1176.
- 155 X. Li, D. Zhang, H. Chi, G. Xiao, Y. Dong, S. Wu, Z. Su, Z. Zhang, P. Lei, Z. Hu and W. Li, *Appl. Phys. Lett.*, 2010, **97**, 263303.
- 156 X. Li, H. Chi, G. Lu, G. Xiao, Y. Dong, D. Zhang, Z. Zhang and Z. Hu, *Org. Electron.*, 2012, **13**, 3138-3144.
- 157 Q. Zhang, J. Ding, Y. Cheng, L. Wang, Z. Xie, X. Jing and F. Wang, *Adv. Funct. Mater.*, 2007, **17**, 2983-2990.
- 158 O. Crespo, M. C. Gimeno, A. Laguna, R. Marriott, J. M. Sáez-Rocher and M. D. Villacampa, *Dalton Trans.*, 2014, **43**, 12214-12220.
- 159 C. Dragonetti, A. Valore, A. Colombo, S. Righetto, G. Rampinini, F. Colombo, L. Rocchigiani and A. Macchioni, *Inorg. Chim. Acta*, 2012, **382**, 72-78.
- 160 A. Valore, M. Balordi, A. Colombo, C. Dragonetti, S. Righetto, D. Roberto, R. Ugo, T. Benincori, G. Rampinini, F. Sannicolò and F. Demartin, *Dalton Trans.*, 2010, **39**, 10314-10318.
- 161 B. W. Chu and V. W. Yam, *Inorg. Chem.*, 2001, **40**, 3324-3329.
- 162 V. W. W. Yam, K. Wang, C. Wang, Y. Yang and K. Cheung, *Organometallics*, 1998, **17**, 2440-2446.
- 163 A. Colombo, C. Dragonetti, S. Righetto, D. Roberto, A. Valore, T. Benincori, F. Colombo and F. Sannicolò, *J. Mater. Chem.*, 2012, **22**, 19761-19766.
- 164 K. Sako, T. Kakehi, S. Nakano, H. Oku, X. F. Shen, T. Iwanaga, M. Yoshikawa, K. Sugahara, S. Toyota, H. Takemura, T. Shinmyozu, M. Shiotsuka and H. Tatemitsu, *Tetrahedron Lett.*, 2014, **55**, 749-752.
- 165 I. Erden, A. Erdoğan, N. Demirhan and U. Avciata, *Transit. Met. Chem.*, 2008, **33**, 439-442.
- 166 Z. Lin, Y. Zhao, C. Duan, B. Zhang and Z. Bai, *Dalton Trans.*, 2006, , 3678-3684.
- 167 P. Lam, G. Lu, K. Hon, K. Lee, C. Ho, X. Wang, J. C. O. Tang, K. Lam, R. S. M. Wong, S. H. L. Kok, Z. Bian, H. Li, K. Lee, R. Gambari, C. Chui and W. Wong, *Dalton Trans.*, 2014, **43**, 3949-3957.
- 168 A. A. Massoud, Y. M. Gohar, V. Langer, P. Lincoln, F. R. Svensson, J. Janis, S. T. Gardebjer, M. Haukka, F. Jonsson, E. Aneheim, P. Lowenhielm, M. A. M. Abu-Youssef and L. R. Ohrstrom, *New J. Chem.*, 2011, **35**, 640-648.
- 169 B. J. Kraft, H. J. Eppley, J. C. Huffman and J. M. Zaleski, *J. Am. Chem. Soc.*, 2002, **124**, 272-280.

Dalton Transactions

PERSPECTIVE

170 H. J. Eppley, S. M. Sato, A. D. Ellington and J. M. Zaleski, *Chem. Commun.*, 1999, 2405-2406.

171 H. J. Eppley, V. J. Isada, S. M. Lato, J. C. Huffman, A. D. Ellington and J. M. Zaleski, *J. Inorg. Biochem.*, 1999, **74**, 124-124.

This review article highlights the versatile nature of 4,5-diazafluorene derivatives as ligands, and details some recent advances made using this ligand family.

