



Synthesis and Characterization of [Fe(NCCH₃)₆][cis-Fe(InX₃)₂(CO)₄] (X = Cl, Br, I) Containing Two Terminal Indium Fragments

Journal:	<i>Dalton Transactions</i>
Manuscript ID	DT-COM-09-2015-003745.R1
Article Type:	Communication
Date Submitted by the Author:	01-Dec-2015
Complete List of Authors:	Itazaki, masumi; Osaka City University, Chemistry Ito, Masaki; Osaka City University, Chemistry Nakashima, Satoru; Hiroshima University, Natural Science Center for Basic Research and Development Nakazawa, Hiroshi; Osaka City University, Chemistry

Synthesis and Characterization of $[\text{Fe}(\text{NCCH}_3)_6][\text{cis-Fe}(\text{InX}_3)_2(\text{CO})_4]$ ($\text{X} = \text{Cl}, \text{Br}, \text{I}$) Containing Two Terminal Indium Fragments[†]

Masumi Itazaki,^{*a} Masaki Ito,^a Satoru Nakashima^b and Hiroshi Nakazawa^{*a}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Iron complexes, $[\text{Fe}(\text{NCCH}_3)_6][\text{cis-Fe}(\text{InX}_3)_2(\text{CO})_4]$ ($\text{X} = \text{Cl}$: **1, Br : **2**, I : **3**) containing two terminal indium fragments, were synthesized. Their structures were determined using X-ray analysis. The bonding mode of the Fe–In bonds in the anionic part was investigated using ⁵⁷Fe Mössbauer and IR spectroscopy. These complexes represent the first example of transition metal complexes containing two terminal indium fragments.**

The chemistry of complexes in which transition metals are involved in a Z-type interaction with group 13 elements has been developed rapidly, and interesting reactivity has been reported.^{1,2} A Z-type interaction ($\text{M} \rightarrow \text{E}$, M = transition metal and E = group 13 element) is formed by the donation of two electrons from electron-rich M to electron-deficient EX_3 . In group 13 elements, syntheses and reactivities of $\text{M} \rightarrow \text{borane}$ complexes are well investigated, whereas much less information is available on $\text{M} \rightarrow \text{indane}$ complexes; however, different bonding fashions between M and In have been reported (Fig. 1).^{3–8}




Fig. 1 Various coordination modes of M–In complex.

In 1942, Hieber and Teller reported the first M–In complex $[\text{In}\{\text{Co}(\text{CO})_4\}_3]$ having M–In σ bonds.⁹ Since then, a large number of transition metal complexes containing M–In σ bond(s) have been synthesized. We also reported the bridging indylene ($\text{W}–\text{In}–\text{W}$)

complex with tungsten $[\{\text{Cp}(\text{CO})_3\text{W}\}_2\text{InCl}(\text{py})]$.¹⁰ However, only a few examples of indane complexes ($\text{M} \rightarrow \text{InX}_3$, $\text{X} = \text{Cl}, \text{Br}, \text{I}$) have been reported, and group 6 transition metal complexes have been crystallographically authenticated.¹¹ In addition, the group of Bourissou synthesized a palladium indane complex $[(\text{TPI})\text{Pd}]$ (TPI = triphosphine-indane).¹² Only three examples of iron indane complexes ($\text{Fe} \rightarrow \text{In}(\text{III})$) have been reported so far (Fig. 2).¹³






Fig. 2 Examples of previously reported Fe→In(III) complexes.

Previously, we reported the first ruthenium indane complexes $[\text{Ru}(\text{NCMe})_3(\text{CO})_2(\text{InX}_3)]$ ($\text{X} = \text{Cl}, \text{Br}$) and their reactivity with PPh_3 (eqn (1)).¹⁴ In this paper, the synthesis and characterization of iron complexes containing two terminal indium fragments are described, and the bonding mode of the two Fe–In bonds based on ⁵⁷Fe Mössbauer and IR data are discussed.



Triiron dodecacarbonyl $\text{Fe}_3(\text{CO})_{12}$ was treated with 3 equiv of InX_3 ($\text{X} = \text{Cl}, \text{Br}, \text{I}$) in acetonitrile at 80 °C for 30 min to produce an iron complex with two InX_3 groups, $[\text{Fe}(\text{NCMe})_6][\text{cis-Fe}(\text{InX}_3)_2(\text{CO})_4]$ ($\text{X} = \text{Cl}$: **1**, Br : **2**, I : **3**), in high yields (Scheme 1).



Scheme 1 Synthesis of $[\text{Fe}(\text{NCCH}_3)_6][\text{cis-Fe}(\text{InX}_3)_2(\text{CO})_4]$ ($\text{X} = \text{Cl}, \text{Br}, \text{I}$).


^a Department of Chemistry, Graduate School of Science, Osaka City University, 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan

^b Natural Science Center for Basic Research and Development, Hiroshima University, 1-4-2 Kagamiyama, Higashi-Hiroshima 739-8526, Japan

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

[†] Dedicated to Professor Kohtaro Osakada on the occasion of his 60th birthday

Complex **2** was obtained in 88% yield in the reaction of $\text{Fe}_2(\text{CO})_9$ with 2 equiv of InBr_3 in acetonitrile at 80 °C for 1 h. However, when $\text{Fe}(\text{CO})_5$ was used instead of $\text{Fe}_3(\text{CO})_{12}$ or $\text{Fe}_2(\text{CO})_9$, **2** was not formed (Scheme 2).



Scheme 2 Reaction of InBr_3 with $\text{Fe}_2(\text{CO})_9$ or $\text{Fe}(\text{CO})_5$.

The $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of **1** at 25 °C showed a broad singlet at δ 216.03 corresponding to the carbonyl groups. This signal split into two singlets (δ 218.38, 219.87) at -40 °C, indicating that **1** in solution shows fluxional behavior.

The electric charge of $[\text{Fe}(\text{NCMe})_6]$ is known to be 2+,^{15,16} therefore, the electric charge of $\text{cis-}[\text{Fe}(\text{InX}_3)_2(\text{CO})_4]$ should be 2-. As **1-3** were prepared from $\text{Fe}_3(\text{CO})_{12}$, charge disproportionation among neutral $\text{Fe}(0)$ species took place. Previously, Scheer *et al.* reported the synthesis of the analogous iron complex $[\text{Na}(\text{OEt})_2][\text{Fe}(\text{GaCl}_3)_2(\text{CO})_4]$.¹⁷ However, this complex was not synthesized by charge disproportionation but by the reaction of 2 equiv of GaCl_3 with the dianion iron complex $\text{Na}_2[\text{Fe}(\text{CO})_4]$.

The structures of **1-3** were determined by X-ray crystallography. The molecular structures of the anionic part of **1-3**, *i.e.*, $\text{cis-}[\text{Fe}(\text{InX}_3)_2(\text{CO})_4]^{2-}$ ($\text{X} = \text{Cl}, \text{Br}, \text{I}$), are shown in Figs. 3–5, respectively. Two independent molecules of **2** crystallized in the unit cell. As these are basically the same, only one anionic part, *i.e.*, $\text{cis-}[\text{Fe}(\text{InBr}_3)_2(\text{CO})_4]^{2-}$ (Fe1 molecule), is shown in Fig. 4. In all cases, the Fe atom for the cationic part is coordinated by six MeCN ligands, and the Fe atom for the anionic part is coordinated by two indium ligands and four terminal CO ligands. These InX_3 ligands are in a mutual *cis* position. The Fe–In bond distances (2.5817(5), 2.5896(6) Å for **1**, 2.5861(13)–2.6039(11) Å for **2**, and 2.6001(7), 2.6233(7) Å for **3**) are longer than those of previously reported Fe– InX_3 ($\text{X} = \text{Cl}, \text{I}$) complexes (2.517(2) Å for Cl,^{13b} 2.5275(9) Å for I^{13a}). The distances of Fe–C bonds *trans* to In (1.805(4), 1.811(4) Å for **1**, 1.782(9)–1.816(8) Å for **2**, and 1.805(5), 1.801(5) Å for **3**) are similar to those of Fe–C bonds *trans* to CO (1.802(4), 1.782(4) Å for **1**, 1.786(9)–1.812(7) Å for **2**, and 1.795(5), 1.804(5) Å for **3**). In contrast to the Ru case, the InX_3 ligands of **1-3** do not show a greater *trans*-influence than CO.¹⁴ The interaction between Fe and InX_3 in **1-3** seems to be weaker than that in the previously reported indane iron complex, because the In–X bond distances for **1-3** ($\text{X} = \text{Cl}$: 2.3865(9)–2.4153(10) Å for **1**, $\text{X} = \text{I}$: for 2.7413(6)–2.7956(5) Å for **3**) are shorter than those for the indane iron complex ($\text{X} = \text{Cl}$: 2.432(3)–2.469(3) Å,^{13b} $\text{X} = \text{I}$: for 2.7942(6)–2.8060(6) Å^{13a}).




Fig. 3 ORTEP drawing of $\text{cis-}[\text{Fe}(\text{InCl}_3)_2(\text{CO})_4]^{2-}$ anionic part of **1** in the crystal (30% thermal ellipsoidal plots). All hydrogen atoms are omitted for clarity. Selected bond lengths (Å) and angles (°): Fe1–In1 2.5896(6), Fe1–In2 2.5817(5), Fe1–C1 1.805(4), Fe1–C2 1.811(4), Fe1–C3 1.802(4), Fe1–C4 1.782(4), C1–O1 1.133(5), C2–O2 1.129(5), C3–O3 1.139(4), C4–O4 1.148(5), In1–Cl1 2.3965(10), In1–Cl2 2.3996(10), In1–Cl3 2.4153(10), In2–Cl4 2.4025(10), In2–Cl5 2.4089(10), In2–Cl6 2.3865(9), In1–Fe1–In2 92.930(17).




Fig. 4 ORTEP drawing of $\text{cis-}[\text{Fe}(\text{InBr}_3)_2(\text{CO})_4]^{2-}$ anionic part (Fe1 molecule) of **2** in the crystal (30% thermal ellipsoidal plots). All hydrogen atoms are omitted for clarity. Selected bond lengths (Å) and angles (°): Fe1–In1 2.6039(11), Fe1–In2 2.5883(10), Fe1–C1 1.782(9), Fe1–C2 1.816(8), Fe1–C3 1.812(7), Fe1–C4 1.794(8), C1–O1 1.134(10), C2–O2 1.121(9), C3–O3 1.125(8), C4–O4 1.126(9), In1–Br1 2.5456(9), In1–Br2 2.5532(10), In1–Br3 2.5655(10), In2–Br4 2.5496(11), In2–Br5 2.5539(12), In2–Br6 2.5369(11), In1–Fe1–In2 93.38(3) for Fe1 molecule. Fe2–In3 2.5861(13), Fe2–In4 2.5910(12), Fe2–C5 1.799(11), Fe2–C6 1.802(8), Fe2–C7 1.786(9), Fe2–C8 1.791(9), C5–O5 1.132(12), C6–O6 1.134(10), C7–O7 1.140(10), C8–O8 1.144(10), In3–Br7 2.5576(11), In3–Br8 2.5504(13), In3–Br9 2.5364(14), In4–Br10 2.5314(14), In4–Br11 2.5505(15), In4–Br12 2.5351(13), In3–Fe2–In4 91.77(4) for Fe2 molecule.




Fig. 5 ORTEP drawing of $\text{cis-}[\text{Fe}(\text{InI}_3)_2(\text{CO})_4]^{2-}$ anionic part of **3** in the crystal (30% thermal ellipsoidal plots). All hydrogen atoms are omitted for clarity. Selected bond lengths (Å) and angles (°): Fe1–In1 2.6001(7), Fe1–In2 2.6233(7), Fe1–C1 1.805(5), Fe1–C2 1.801(5), Fe1–C3 1.795(5), Fe1–C4 1.804(5), C1–O1 1.139(6), C2–O2 1.136(6), C3–O3 1.147(6), C4–O4 1.133(6), In1–I1 2.7493(6), In1–I2 2.7956(5), In1–I3 2.7512(6), In2–I4 2.7607(5), In2–I5 2.7845(6), In2–I6 2.7413(6), In1–Fe1–In2 95.33(2).

Next, we focused on the Fe–In bonding mode in the anionic part $cis\text{-}[\text{Fe}(\text{InX}_3)_2(\text{CO})_4]^{2-}$. Three possible combinations concerning Fe–In bonds for $[\text{Fe}(\text{InX}_3)_2(\text{CO})_4]^{2-}$ are conceivable (Fig. 6): (i) two Z-type bonds (electron donation from $[\text{Fe}(\text{CO})_4]^{2-}$ to two InX_3 serving as a Lewis acid), (ii) two covalent bonds (two $\text{Fe}\text{-}\text{InX}_3$, that is, two indate portions), (iii) one Z-type bond and one covalent bond (a situation between (i) and (ii)). The formal oxidation number of Fe is considered to be –II for (i), 0 for (ii), and –I for (iii).




Fig. 6 Three possible structures for the anionic part $cis\text{-}[\text{Fe}(\text{InX}_3)_2(\text{CO})_4]^{2-}$.




Fig. 7 ^{57}Fe Mössbauer spectrum of solid **1** recorded at 80 K. The black dot marks are the raw data, and the solid lines were obtained from least-squares fits of the spectrum with two doublets, Fe^{2+} component (blue, 40%) $\delta = 1.28$ mm/s, $\Delta E_Q = 0.55$ mm/s and Fe^0 component (red, 60%) $\delta = -0.03$ mm/s, $\Delta E_Q = 0.24$ mm/s.

The ^{57}Fe Mössbauer spectrum of solid **1**, collected at 80 K, is shown in Fig. 7. Complex **1** displays two doublets with $\delta = 1.28$ mm/s, $\Delta E_Q = 0.55$ mm/s (blue, 40% of Fe) and $\delta = -0.03$ mm/s, $\Delta E_Q = 0.24$ mm/s (red, 60% of Fe). The area ratio of the present sample is 40:60 at 80 K, which is different from the expected ratio. We measured the same sample at 300 K. The ratio (40:60) at 80 K changed to 27:73 at 300 K (see Figure S1). The intensity of Mössbauer absorption changes depending on the molecular vibration. The results suggest that the site of 40% at 80 K is easier to vibrate even at 80 K. The expected ratio would be observed at lower temperature than 80 K. The δ value of the former signal (1.28 mm/s) is in the range of the reported values for $\text{Fe}(\text{II})$ species ($\delta = 0.8\text{--}1.4$ mm/s).¹⁸ Thus, the doublet can be reasonably assigned to $[\text{Fe}(\text{NCMe})_6]^{2+}$. The δ value of the latter signal (-0.03 mm/s) is close to that of an $\text{Fe}(\text{0})$ complex, such as $\text{Fe}(\text{CO})_5$ (0.009 mm/s),¹⁹ and is quite different from that of an $\text{Fe}(\text{–II})$ complex, such as $\text{Na}_2\text{Fe}(\text{CO})_4$ (-0.251 mm/s).²⁰ The IR spectra of **1–3** show ν_{CO} absorptions at 1984, 2030, 2076 cm^{-1} for **1**, 1984, 2016, 2076 cm^{-1} for **2**, and 1988, 2009, 2064 cm^{-1} for **3**. The ν_{CO} absorptions of **1–3** are similar to that of the zero-valent iron complex $\text{Fe}(\text{CO})_5$ (2000, 2022 cm^{-1})²¹ and far from the ν_{CO} absorption of the dianion iron complex $\text{Na}_2[\text{Fe}(\text{CO})_4]$ (1730 cm^{-1}).²²

The data obtained from the Mössbauer and IR spectra suggest that the formal oxidation number of the Fe atom in $cis\text{-}[\text{Fe}(\text{InX}_3)_2(\text{CO})_4]^{2-}$ is 0 (zero). In addition, there are no large difference between two Fe–In distances and among six In–X distances in **1–3**. Therefore, (ii) in Fig. 6 seems to be an appropriate expression for $cis\text{-}[\text{Fe}(\text{InX}_3)_2(\text{CO})_4]^{2-}$.

In summary, we synthesized and characterized the first iron complexes $[\text{Fe}(\text{NCMe})_6][cis\text{-}\text{Fe}(\text{InX}_3)_2(\text{CO})_4]$ (X = Cl: **1**, Br: **2**, I: **3**) containing two terminal indium fragments. In solid state and in solution state at -40 °C, two InX_3 fragments are situated in a *cis* position. However, they show *cis/trans* fractional behavior in solution at room temperature. The ^{57}Fe Mössbauer and IR data suggest that $\text{Fe}^0(\text{CO})_4$ has two indate ($\text{Fe}\text{-}\text{InX}_3$) portions.

This work was supported by a Grant-in-Aid for Scientific Research from JSPS (Category C, No. 25410073 (M.I.)), by a Challenging Exploratory Research Grant (No. 25620048 (H.N. and M.I.)), and by a Grant-in Aid for Scientific Research on Innovation Area “Stimuli-responsive Chemical Species” (No. 25109538 (H.N.)) from MEXT, Japan.

Notes and references

- (a) H. Braunschweig, R. D. Dewhurst, F. Hupp, C. Kaufmann, A. K. Phukan, C. Schneider and Q. Ye, *Chem. Sci.*, 2014, **5**, 4099; (b) H. Kameo and H. Nakazawa, *Chem. Asian J.*, 2013, **8**, 1720; (c) J. Bauer, H. Braunschweig, and R. D. Dewhurst, *Chem. Rev.*, 2012, **112**, 4329; (d) G. R. Owen, *Chem. Soc. Rev.*, 2012, **41**, 3535; (e) A. Amgoune and D. Bourissou, *Chem. Commun.*, 2011, **47**, 859; (f) H. Braunschweig and R. D. Dewhurst, *Dalton Trans.*, 2011, **40**, 549; (g) H. Braunschweig, R. D. Dewhurst and A. Schneider, *Chem. Rev.*, 2010, **110**, 3924; (h) I. Kuzu, I. Krummenacher, J. Meyer, F. Armbruster and F. Breher, *Dalton Trans.*, 2008, 5836.
- Especially, the chemistry of metal-only Lewis pairs have received a lot of attention in recent years. For example, see: (a) P. J. Malinowski and I. Krossing, *Angew. Chem. Int. Ed.*, 2014, **53**, 13460; (b) H. Braunschweig, R. D. Dewhurst, F. Hupp and C. Schneider, *Chem. Commun.*, 2014, **50**, 15685;
- (a) C. Genel, T. Steinke, M. Cokoja, A. Kempter and R. A. Fischer, *Eur. J. Inorg. Chem.*, 2004, 4161; (b) R. Murugavel and V. Chandrasekhar, *Angew. Chem., Int. Ed.*, 1999, **38**, 1211; (c) R. A. Fischer and J. Weiß, *Angew. Chem., Int. Ed.*, 1999, **38**, 2831.
- S. González-Gallardo, T. Bollermann, R. A. Fischer and R. Murugavel, *Chem. Rev.*, 2012, **112**, 3136.
- M. Asay, C. Jones and M. Driess, *Chem. Rev.*, 2011, **111**, 354.
- C. Boehme, J. Uddin and G. Frenking, *Coord. Chem. Rev.*, 2000, 249.
- R. J. Baker and C. Jones, *Coord. Chem. Rev.*, 2005, 1857.
- Group 13 Chemistry I: Fundamental New Developments. Structure & Bonding*, ed. by H. W. Roesky and D. A. Atwood, Springer-Verlag: Berlin, Heidelberg, New York, 2002, Vol. 103.
- W. Hieber and U. Teller, *Z. Anorg. Allg. Chem.*, 1942, **249**, 43.
- H. Nakazawa, M. Itazaki and M. Owaribe, *Acta Crystallogr.*, 2005, **E61**, m945.
- (a) P. Rutsch, G. Renner, G. Huttner and S. Z. Sandhoefner, *Naturforsch.*, 2002, **57b**, 757; (b) B. Nuber, W. Schatz and M. L. Ziegler, *Z. Naturforsch.*, 1990, **45b**, 508; (c) L. M. Clarkson, W. Clegg, N. C. Norman, A. J. Tucker and P. M. Webster, *Inorg. Chem.*, 1988, **27**, 2653.
- E. J. Derrah, M. Sircoglou, M. Mercy, S. Ladeira, G. Bouhadir, K. Miqueu, L. Maron and D. Bourissou, *Organometallics*, 2011, **30**, 657.

COMMUNICATION

Journal Name

- 13 (a) S. Aldridge, D. L. Kays, N. R. Bunn, N. D. Coombs and L. L. Ooi, *Main Group Met. Chem.*, 2005, **28**, 201; (b) M. Esser, B. Neumüller, W. Petz, J. Uddin and G. Frenking, *Z. Anorg. Allg. Chem.*, 2000, **626**, 915; (c) J. K. Ruff, *Inorg. Chem.*, 1968, **7**, 1499.
- 14 M. Itazaki, M. Ito and H. Nakazawa, *Eur. J. Inorg. Chem.*, 2015, 2033.
- 15 D. C. Boyd, D. A. Bohling and K. R. Mann, *J. Am. Chem. Soc.*, 1985, **107**, 1641.
- 16 H. Wasada, Y. Wasada-Tsutsui, T. Hashimoto and S. Funahashi, *Int. J. Quantum Chem.*, 2009, **109**, 2208.
- 17 E. Leiner, O. Hampe and M. Scheer, *Eur. J. Inorg. Chem.*, 2002, 584.
- 18 J. Silver and R. Withnall, *Chem. Rev.*, 2004, **104**, 2833.
- 19 R. H. Herber, R. B. King and G. K. Wertheim, *Inorg. Chem.*, 1964, **3**, 101.
- 20 N. E. Erickson and A. W. Fairhall, *Inorg. Chem.*, 1965, **4**, 1320.
- 21 K. Noack and M. Ruch, *J. Organomet. Chem.*, 1969, **17**, 309.
- 22 H. Strong, P. J. Krusic and J. S. Filippo, *Inorg. Synth.*, 1986, **24**, 157.

A graphical and textual abstract

Synthesis and Characterization of $[\text{Fe}(\text{NCCH}_3)_6][\text{cis-Fe}(\text{InX}_3)_2(\text{CO})_4]$ ($\text{X} = \text{Cl}, \text{Br}, \text{I}$)
Containing Two Terminal Indium Fragments

Masumi Itazaki,^{*a} Masaki Ito,^a Satoru Nakashima^b and Hiroshi Nakazawa^{*a}

The first example of iron complexes containing two Fe–In covalent bonds, $[\text{Fe}(\text{NCCH}_3)_6][\text{cis-Fe}(\text{InX}_3)_2(\text{CO})_4]$ ($\text{X} = \text{Cl}$: **1**, Br : **2**, I : **3**), were synthesized and the bonding mode of the Fe–In bonds in the anionic part was investigated using ^{57}Fe Mössbauer and IR spectroscopy.

