

Dalton Transactions

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

A new Metal-Organic Polymeric system capable of stimuli-responsive controlled release of the drug ibuprofen.

Received 00th January 20xx,
Accepted 00th January 20xx

Ana Belén Lago^{a*}, Arantxa Pino-Cuevas^a, Rosa Carballo^a, Ezequiel M. Vázquez-López^a

DOI: 10.1039/x0xx00000x

A series of new zinc compounds has been prepared easily under mild synthetic conditions with the drug ibuprofen incorporated as a ligand to form different coordination polymers. The host materials have a high drug content and exhibit high stability, low cytotoxicity, good biodegradability and high biocompatibility. The ion exchange mechanism involved in the ibuprofen release process and the pH-controlled drug release have been studied.

www.rsc.org/

Introduction

The search for new medical strategies with high therapeutic efficiencies and reduced adverse side-effects remains a challenge for the scientific community. The development of new drug delivery systems (DDS) to generate new solid forms of active pharmaceutical ingredients (APIs) has attracted increased interest in pharmaceutical materials science. A wide range of different architectures have been employed in an effort to optimize drug therapeutics.¹ Various solid-phase carriers, such as polymeric resins or metal organic frameworks (MOFs), have emerged as promising platforms for drug delivery systems.² Many of the drawbacks associated with small molecule drugs can be circumvented through the use of these novel systems.

Ion exchange resins are cross-linked polymers that incorporate compounds and these systems have been widely studied in drug delivery research. Numerous ion-exchange resins are commercially available and they allow some control over the rate of drug release.³ The ionic strength and pH at the site of delivery play key roles in the liberation of immobilized drugs from resins. This class of compounds can be considered as a relevant example in the search for new dosage forms.

Hybrid materials composed of metal ions and organic bridging ligands exhibit promising characteristics as drug carriers, including intrinsic biodegradability as a result of the relatively labile metal-ligand bonds and high drug-loading capacity.⁴ Moreover, the addition of metal ions to the polymeric system allows changes in the physicochemical and biological properties of the material.⁵ One particular approach in this field is the direct coupling of the biomolecule as the linker with the

metal to prepare MOF compounds.⁶ The release of the active molecule can be achieved by desorption, anionic exchange or through biodegradation of the material.

An alternative approach consists of the direct coordination of the biomolecule to the metal ion connector but with the coordination polymer built up within a different biodegradable polydentate linker.⁷ As ion-exchange resins, these coordination polymers will contain two principal parts: a structural unit consisting of a polymeric chain and a functional section, which is the ion-active site group.

An important goal in developing materials with pharmaceutical applications is to design biocompatible carriers with little or zero toxicity. In this sense, in the design of coordination polymers it seems appropriate to include bioactive metal cations such as zinc(II), which is an essential nutrient for life and can also be used as an antibacterial growth agent.⁸ Moreover, stimuli-responsive controlled-release systems, which undergo physical or chemical changes in response to small changes in the environmental conditions, are highly desired. In comparison to sustained-release systems, these dynamic systems can achieve a site-selective, controlled-release pattern, which improves the therapeutic efficacy.⁹ However, it remains a challenge to find more convenient ways to achieve a fully controllable drug-release system under normal physiological conditions.¹⁰

Herein, we describe a zinc coordination polymer with a fully controlled architecture as a starting material for ibuprofen drug delivery. The choice of ibuprofen as the API is due to its moderate size, poor solubility, its relatively simple molecular geometry, and its major relevance as a typical test molecule in controlled drug delivery experiments.¹¹ Furthermore, in the field of coordination polymer research our group has previously investigated the ability of the ditopic linker ligand bis(4-pyridylthio)methane (SCS) to form polymeric compounds.¹²

One of the most attractive features of these new compounds is the simplicity of their synthesis and the straightforward loading process, which is achieved in one-pot and in a controlled way. The mechanism involved in the drug delivery process is

^a Departamento de Química Inorgánica, Facultad de Química, Instituto de Investigación Biomédica (IBI)-Universidad de Vigo, 36310 Vigo, Galicia (Spain).

*Corresponding author: A. B. Lago, e-mail: ablago@uvigo.es

† Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

controlled by anion exchange and this was also studied. These features were of sufficient interest to attract us to study new coordination polymers for applications in materials science and, in particular, as drug delivery systems. The synthesis, characterization, cytotoxicity and release behavior of this system was evaluated. This system could also be regarded as a model for other drugs that contain carboxylic acid groups capable of anionic exchange.

Experimental

Materials and physical measurements

Bis(4-pyridylthio)methane (SCS) was prepared as described previously.¹³ The other starting materials and solvents were obtained commercially and were used as supplied. Ibuprofen was used as a racemic mixture. Elemental analyses (C, H, N, S) were carried out on a Fisons EA-1108 microanalyser. IR spectra were recorded from KBr discs (4000–400 cm⁻¹) on a Jasco FT/IR-6100 spectrophotometer. ¹H-NMR spectra were obtained on a Bruker AMX 400 spectrometer. Mass spectra (ESI⁺) were recorded on a Hewlett-Packard 5989A spectrometer. UV-Vis spectra were recorded on a Jasco V-670 spectrophotometer. TGA was performed on a SETSYS Evolution Setaram thermogravimetric analyser in a flow of N₂ at a heating rate of 10 °C min⁻¹.

Synthesis of complexes

[Zn(ibu)₂(SCS)] (1)

A solution of SCS (0.427 mmol) in ethanol/water (2:1) was added to a solution of zinc(II) acetate dihydrate, Zn(CH₃CO₂)₂·2H₂O (0.427 mmol) in the same solvent mixture (10 mL). A solution of ibuprofen (Hibu) (0.853 mmol) in the same solvent mixture was added slowly to the above colorless solution. The suspension was stirred and heated under reflux for 8 h. The resulting white precipitate was filtered off, washed with water and ethanol, and dried under vacuum.

Yield: 231 mg (80%). Anal. found: C, 61.93; H, 6.24; N, 3.96; S, 8.71 C₃₇H₄₂N₂S₂O₄Zn requires: C, 62.35; H, 5.98; N, 3.96; S, 9.05. ¹H-NMR (400 MHz, [D₆]-DMSO, 25 °C): δ 0.84–0.86 (d, 12H, C12-H3, C11-H3, ibu); 1.28–1.30 (d, 6H, C3-H3, ibu); 1.77–1.80 (q, 2H, C11-H, ibu); 2.33–2.39 (d, 4H, C10-H2, ibu); 3.47–3.49 (dd, 2H, C2-H2, ibu); 4.99 (s, 2H, C14-H2, SCS); 7.02–7.04 (d, 4H, C5arom, C7arom, ibu); 7.17–7.19 (d, 4H, C6arom, C8arom, ibu); 7.38–7.40 (d, 4H, C15arom, C16arom, SCS); 8.41–8.43 (d, 4H, C17arom, C18arom, SCS). IR (KBr): ν(cm⁻¹) = 2965m, 2927m, 2868m (CH stretching), 1625s (ν_{asym}(OCO)), 1595s (ν (CC) + ν (CN)), 1374m, 1362 (ν_{sym}(OCO)), 1244m (ω (CH,CH₂)), 1063m (ν (CH) + δ (C-H)), 740m, 717 (ν (CS)). MS-ESI: *m/z* (%) = 235 (49) [SCS+H]⁺, 475 (100) [Zn(ibu)₂+H]⁺, 786 (51) [Zn(ibu)₂(SCS)(dmsO)]⁺

[ZnCl(ibu)(SCS)] (2)

A solution of SCS (0.427 mmol) in ethanol/water (2:1) was added to a solution of ZnCl₂ (0.427 mmol) in the same solvent mixture (10 mL). A white precipitate formed immediately and the suspension was stirred for 10 min. A solution of Hibu

(0.427 mmol) and NaOH (0.475 mmol) in the same solvent mixture (10 mL) was added to the suspension. The suspension was stirred for 16 h. The resulting white precipitate was filtered off, washed with ethanol and water and dried under vacuum. Colorless single crystals were obtained by slow evaporation at rt (1 week) of the mother liquor and better quality crystals were obtained by slow diffusion in a test tube of a solution of SCS (0.025 mmol), Hibu (0.025 mmol) and NaOH (0.025 mmol) in ethanol (10 mL) and a solution of ZnCl₂ (0.025 mmol) in water (10 mL) at rt. The powder X-ray diffraction data were compared to those simulated from the crystal structure and showed that the crystal is representative of the bulk material. Data for **1** from the first synthetic procedure:

Yield: 216 mg (95%). Anal. found: C, 53.22; H, 4.77; N, 5.35; S, 11.97 C₂₄H₂₆N₂S₂O₂ClZn requires: C, 53.44; H, 4.86; N, 5.19; S, 11.89. ¹H-NMR (400 MHz, [D₆]-DMSO, 25 °C): δ 0.84–0.86 (d, 6H, C12-H3, C11-H3, ibu); 1.30–1.32 (d, 3H, C3-H3, ibu); 1.77–1.81 (q, 1H, C11-H, ibu); 2.38–2.40 (d, 2H, C10-H2, ibu); 3.48–3.53 (dd, 1H, C2-H, ibu); 4.99 (s, 2H, C14-H2, SCS); 7.03–7.05 (d, 2H, C5arom, C7arom, ibu); 7.18–7.20 (d, 2H, C6arom, C8arom, ibu); 7.39–7.40 (d, 4H, C15arom, C16arom, SCS); 8.41–8.43 (d, 4H, C17arom, C18arom, SCS). IR (KBr): ν(cm⁻¹) = 2954m, 2912m, 2866m (CH stretching), 1597s (ν_{asym}(OCO), ν (CC) + ν (CN)), 1381m (ν_{sym}(OCO)), 1233w (ω (CH,CH₂)), 1064m (ν (CH) + δ (C-H)), 729m (ν (CS)). MS-ESI: *m/z* (%) = 235 (100) [SCS+H]⁺, 475 (40) [Zn(ibu)₂+H]⁺, 371 (5) [ZnCl₂(SCS)+H]⁺, 786 (9) [Zn(ibu)₂(SCS)(dmsO)]⁺

[Zn(ibu)₂(H₂O)₂] (3)

This compound was previously described with a different synthetic procedure without its structure being elucidated by single crystal X-ray diffraction.¹⁴

ZnCl₂ (0.418 mmol), Hibu (0.418 mmol), NaOH (0.418 mmol) and ethanol/water (2:1, 20 mL) were added to a high pressure vessel, which was placed in a microwave oven. The power input was limited by temperature T_{lim} = 90 °C and power P_{lim} = 500 W. For heating up to T_{lim} a time of 5 min was used and the temperature was maintained for 5 min at 90 °C with a stirring rate of 400 rpm. Under these conditions the pressure reached about 4.0 bar. Cooling was carried out over 15 min (pressurized air). The resulting white precipitate was filtered off, washed with ethanol and water and dried under vacuum.

Colorless single crystals were obtained by slow diffusion in a test tube of a solution of Hibu (0.250 mmol) and NaOH (0.375 mmol) in ethanol (10 mL) and a solution of ZnCl₂ (0.250 mmol) in water (10 mL) at rt. The powder X-ray diffraction data were compared to those simulated from the crystal structure and showed that the crystal was representative of the bulk material.

Yield: 76 mg (71%). Anal. found: C, 61.05; H, 7.56; C₂₆H₃₄O₄Zn·2H₂O requires: C, 61.24; H, 7.12. ¹H-NMR (400 MHz, [D₆]-DMSO, 25 °C): δ 0.84–0.86 (d, 6H, C12-H3, C11-H3); 1.28–1.30 (d, 3H, C3-H3); 1.75–1.82 (q, 1H, C11-H); 2.38–2.39 (d, 2H, C10-H2); 3.46–3.51 (dd, 1H, C2-H); 7.02–7.04 (d, 2H, C5arom, C7arom); 7.17–7.19 (d, 2H, C6arom, C8arom); IR (KBr): ν(cm⁻¹) = 2952m, 2921m, 2867m (CH

stretching), 1623m ($\nu(\text{OC})_{\text{Hbonded}}$), 1541vs ($\nu_{\text{asym}}(\text{OCO})$), 1510s, 1458 ($\nu(\text{CC}) + \beta(\text{HCH})$), 1410s ($\nu_{\text{sym}}(\text{OCO})$), 1258m ($\nu(\text{OC}) + \beta(\text{HCH}) + \tau(\text{HCCC})$), 1067m ($\beta(\text{HCC}) + \beta(\text{CCC})$).

[ZnCl₂(SCS)] (4)

A solution of SCS (0.210 mmol) in ethanol (10 mL) was added to a solution of ZnCl₂ (0.210 mmol) in water (10 mL). The solution was heated under reflux for 9 h and stirred for 2 d. The resulting white precipitate was filtered off, washed with ethanol and water and dried under vacuum. Colorless single crystals of the monomeric species [ZnCl₂(SCS)₂] (single crystals 5) were obtained by slow evaporation at rt (1 month) of the mother liquor. Colorless single crystals of the polymeric structure [ZnCl₂(SCS)] (single crystals 4) were obtained by slow diffusion in a test tube of a solution of SCS (0.210 mmol) in ethanol (10 mL) and a solution of ZnCl₂ (0.210 mmol) in water (10 mL) at rt over 3 days. Data from the first synthetic procedure:

Yield: 50 mg (63%). Anal. found: C, 35.87; H, 2.62; N, 7.46; S, 17.30. C₁₁H₁₀N₂S₂Cl₂Zn requires: C, 35.64; H, 2.71; N, 7.56; S, 17.30. ¹H-NMR (400 MHz, [D₆]-DMSO, 25 °C): 5.00 (s, 2H, C14-H2, SCS); 7.40–7.41 (dd, 4H, C15arom, C16arom, SCS), 8.41–8.43 (dd, 4H, C17arom, C18arom, SCS). IR (KBr): $\nu(\text{cm}^{-1}) = 1594\text{s} (\nu(\text{CC}) + \nu(\text{CN}))$, 1214m ($\omega(\text{CH}, \text{CH}_2)$), 1060m ($\nu(\text{CH}) + \delta(\text{C-H})$), 725s ($\nu(\text{CS})$). MS-ESI: m/z (%) = 235 (100) [SCS+H]⁺, 567 (3) [ZnCl(SCS)₂]⁺, 705 (40) [Zn₂Cl₃(SCS)₂]

The labeling scheme for Hibu and SCS are shown in Figure S1B

Cytotoxicity study

The cytotoxicity experiments for SCS, 1, 2, 3 and 4 were carried out with cell line MRC-5 (non-tumor human fibroblast). Cells were cultured in Eagle's Minimum Essential Medium supplemented with 10% FBS (fetal bovine serum) at 37 °C in a humidified atmosphere of 95% of air and 5% CO₂.

Growth inhibition induced by the compounds was evaluated by a colorimetric method based on the tetrazolium salt MTT ([3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide]), which is reduced by living cells to yield formazan, which is purple. Cells were seeded in 96-well plates at a density of 1×10^4 cells and left to incubate for 24 h. After this time, DMSO solutions of the compounds were added to each well in such a way that the same proportion of DMSO was maintained in each well. Incubation was performed at 37 °C/5% CO₂ for 7 days. At the end of the incubation period, 10 μL of MTT solution (5 mg/mL in PBS) was added and the solutions were incubated for 4 h at 37 °C/5% CO₂. After the addition of 100 μL of SDS (10% in 0.01 M HCl) the solutions were again incubated for 12–14 h. Finally, the cell viability was evaluated by measuring the absorbance at 595 nm using a plate spectrophotometer (Tecan Ultra Evolution). The cell viability was calculated by dividing the absorbance of each well by that of the control wells (cells treated with medium containing 1% DMSO). Each experiment was repeated at least three times. The results of these experiments are shown in Table 1.

Table 1

Compound	Concentration (μM)	% growth inhibition
SCS	100	32 \pm 1
[Zn(ibu) ₂ (SCS)] (1)	100	44 \pm 2
[Zn(Cl)(ibu)(SCS)] (2)	100	49 \pm 1
[Zn(ibu)(H ₂ O) ₂] (3)	100	17 \pm 3
[ZnCl ₂ (SCS)] (4)	100	39 \pm 1

Liberation studies and degradation of the compounds

The presence of ibuprofen as a solid or in solution was detected by FTIR, NMR and UV-Vis techniques. The molar ratio of SCS to ibu⁻, i.e., 1:2 in (1) and 1:1 in (2), is crucial to characterize the solid compounds and also to gain an insight into the liberation and/or degradation processes of the compounds. ¹H-NMR and elemental analysis studies were performed at regular intervals in order to shed light on the SCS:ibu⁻ molar ratio, both in the solid state and in solution. UV spectrophotometry was carried out at room temperature on aqueous or PBS solutions in order to confirm the presence of SCS or ibu⁻ in solution. Three absorption bands were observed in the spectrum of ibuprofen in aqueous solution: one intense band at 220 nm and two weak bands at high concentrations at 264 and 272 nm.¹⁵ Bands were observed at the same wavelength for a solution of the sodium salt of ibuprofenate and [Zn(ibu)₂(H₂O)₂] (3). The absorption maximum of SCS was observed at 265 nm along with a lower intensity band at 212 nm. These bands experience bathochromic shifts upon protonation (*vide infra*). The UV-Vis spectra of polymeric compounds 1 and 2 show bands due to the two ligands but the concentration of each analyte could not be determined by this method due to overlap of the bands. Quantitative NMR (qNMR) was used to quantify the Ibu⁻ and SCS released with D₂O as solvent. Solutions with known concentrations in DMSO were used as the inputs for quantification of the ligand concentrations in solution (Tables S6, S7, S1).

Synthesis and Characterization

In order to load ibuprofen into coordination polymers in a one-pot synthesis, either an ibuprofen solution or ibuprofen previously deprotonated with NaOH was added to the corresponding metal salt and the organic SCS linker to obtain two mixed-ligand compounds [Zn(ibu)₂(SCS)] (1) and [ZnCl(ibu)(SCS)] (2). Moreover, single-ligand compounds [Zn(ibu)₂(H₂O)₂] (3) and [ZnCl₂(SCS)] (4) were also isolated (Scheme 1). Our interest in compounds 3 and 4 was to understand fully the potential products that result from the degradation of the polymeric compounds 1 and 2, as well as to ascertain all the possible combined compounds from the building units that participate in the formation of the polymeric ibuprofen systems. Polymeric compounds were isolated as pure phases in excellent yields (80% for compound 1 and 95% for compound 2) from reactions carried out under mild conditions using water and ethanol solvents. The *in situ* synthesis and loading at room temperature of the drug into the final

bidentate N,N-divergent SCS ligand with zinc metallic units would result in the formation of very stable monodimensional neutral chains through coordination bonds.

The zinc centers are arranged in a tetrahedral geometry and are coordinated to different atoms (Cl, N or O) with Zn–O and Zn–N distances of around 2 Å and slightly longer distances for the Zn–Cl interactions (around 2.2 Å). When ibuprofen molecules are present, the metal centers can increase their coordination number due to weaker interactions established with oxygen atoms of the carboxylate groups (**2** and **3**), which results in an anisobidentate coordination. Thus, the coordination sphere around the zinc metal cations in **3** can be described as 4+2 and it consists of four oxygen atoms from water and carboxylate at expected distances and the other oxygen atom of the anisobidentate chelate with longer Zn···O distances (around 2.9 Å). In SCS-polymeric compounds the metal centers are coordinated to two nitrogen atoms and two chloride atoms in **4** and to one chloride and one ibuprofen oxygen atom in **2**. In compound **2** the zinc is also coordinated to other ibuprofen oxygen atoms with longer bond distances (2.561 Å). The Addison parameter (τ) of 0.94 suggests that the coordination sphere around the zinc atom adopts an almost ideal trigonal bipyramidal geometry, with chloride and oxygen atoms in the apical positions. In Zn-HSCSH the zinc cation is tetraordinated by chloride ions to form the $[\text{ZnCl}_4]^{2-}$ anion.




Figure 2. View of the structures of **2** (A); **4** (B); **3** (C); **5** (D) and Zn-HSCSH (E) showing their representations in this work.

On the basis of these observations, the stability of the coordination environment and the lability of the coordination interactions provide the required flexibility to the coordination environment of the metal center. This characteristic is essential to achieve the anionic exchange and therefore the delivery of ibuprofen. Anionic ibuprofen molecules are pendant ligands in the coordination polymers formed with two coordination sites that can be substituted in **1** and one coordination site in **2**. In the polymeric complexes ibuprofenate molecules are the most labile ligands and they can be easily substituted by other anionic ligands.

The flexible twisted SCS molecule has proven to be a suitable ligand that gives rise to polymerization through coordination bonds.¹² Each SCS ligand adopts a μ_2 -bridging mode to connect the zinc centers with Zn···Zn distances of 12.695 Å and 12.657

Å and N···N distances of 9.399 Å and 9.363 Å in **2** and **4**, respectively.

Compounds **2** and **4** can be described as chain-type neutral structures produced by the zinc metal centers and SCS ligands. The anionic ligands can be chloride atoms or ibuprofenate molecules and these are arranged perpendicular to the growth of the chains, with the ligands of the same nature oriented to the same side (**2**). However, the expected polymerization could be disrupted – as observed in **5** and also in acidic media (*vide infra*).

Compounds **5** and **4** are the monomeric and the polymeric structures, respectively, obtained in the synthesis of $[\text{ZnCl}_2(\text{SCS})]$ (**4**). The less common κN terminal mode of the SCS ligand is found in **5**. In accordance with this situation, the structural parameters of the SCS ligand are closer to those of the free ligand with intramolecular N···N distances of 8.155 and 9.070 Å.

The values of the C–S–C–S torsion angles are 179.10/84.75° in **4** and 178.34/85.32° in **2**, i.e., far from that in the free ligand [73.73(7)°]. These torsion angles in the monodentate terminal ligand (**5**) are also shorter and have closer values to those in the free molecule; 60.50/62.34° and 76.91/64.29°.

Crystals of Zn-HSCSH obtained in acidic media consist of $[\text{ZnCl}_4]^{2-}$ anions surrounding the $[\text{H}_2\text{-SCS}]^{2+}$ cations that result from protonation of the SCS ligand. Isolation of this species provides evidence that the acidic medium can be used to control the degree of polymerization in this system. Protonation of SCS leads to slight changes in the distances and angles related to pyridine rings but appreciable differences in the torsion angle, as observed previously for the analogous structure of $[\text{H}_2\text{-SCS}][\text{CuCl}_4]$.¹⁹ The torsion angles of the C–S–C–S bridge, with values of 174.90/85.73 and 164.67/81.25°, are the largest angles observed in this work and they correspond to the less common TG conformation.²⁰ In accordance with this situation, the intramolecular N···N distances are 9.509 and 8.671 Å.

The zinc-ibuprofen compound **3** crystallizes in the chiral non-centrosymmetric space group C_2 with the metal atom lying in a 2-fold axis. This compound crystallized with two water molecules and it is a neutral mononuclear complex with two enantiopure ibuprofen ligands acting in a monodentate manner. However, the determination of the absolute structure on a single crystal is not representative of the absolute structure of all crystals in the batch.

Release of ibuprofen anion by ion-exchange

In polymeric compounds **1** and **2** ibuprofenate is the most labile ligand and it can be substituted by other anionic ligands. Both compounds are able to undergo exchange reactions when placed in contact with a solution containing anions that are removed from solution through exchange with the ibuprofen anions coordinated to the polymers.

The release of ibuprofen was investigated under simulated physiological conditions in phosphate buffered solution (PBS, pH = 7.4). The release profiles in NaCl solutions were also studied. Compound **1** was soaked in an NaCl solution and a transformation to compound **2** was observed due to the

exchange of 30 wt%, which corresponds to the replacement of sodium ibuprofen. In this way, compound **1** exhibits a phase transition to **2** induced by an anionic exchange process. The structural transformation was confirmed by X-ray powder diffraction experiments carried out before and after the treatment (Figure S6, SI). Anion exchange could be detected in the remaining solid powder by determination of the final SCS:ibu⁻ molar ratio (1:1), which corresponds to [ZnCl(ibu)(SCS)] (**2**). This study was carried out by elemental analysis, NMR and/or IR and techniques (Section 4.1, SI). The UV-Vis spectra of the resulting crude solution showed the typical bands of the ibu⁻ molecule at high concentrations. Dilution of this solution allowed us to detect the characteristic band at 220 nm (Figure S6, SI).




Figure 3. Schematic representation of the anionic exchange process.

Two different treatments were performed in order to study the behavior of **2** in sodium chloride solutions; mild conditions involved stirring the mixture for 21 days and harsher conditions involved heating under reflux for 12 hours. Degradation of **2** was observed under both sets of conditions since both ibu⁻ and SCS species were detected in solution (Figure S7, SI). The crystalline integrity of the framework was maintained throughout the process, as evidenced by PXRD and the 1:1 molar ratio (SCS:ibu⁻) confirmed by NMR. Nevertheless, it was estimated by weight difference that 50% of the initial weight was lost under reflux conditions. Moreover, NaCl was identified in the bulk sample by powder diffraction. A significant decrease in the total percentage of organic elements (C, H, N, S) was also observed in the remaining powders and this loss was more marked after the stirring treatment. This observation indicates the presence of inorganic salts and that the drug release process is accompanied by gradual decomposition of the initial compound **2**.

The behavior exhibited by this system could be suitable for the administration of a high dose in an initial stage since the first step has a high exchange capacity and the resulting compound (**2**), which has a lower drug loading, shows a slower release rate along with decomposition of the material. This level of degradation was not observed in pure water. The possible decomposition of **2** was further evaluated by experiments in PBS solutions with D₂O as solvent.

The release of ibuprofen from **1** and **2** was investigated by ligand exchange reactions in simulated physiological PBS

buffers at r.t. (Figure 3). In this medium, the release process for the coordinated drug is triggered by phosphate anions due to the preferential coordination of PO₄³⁻ anions with zinc cations. In order to determine the release profile, the polymeric samples **1** and **2** were placed in 0.1 M PBS buffer and the drug release was monitored by UV-Vis analysis (Figures S8, S9, SI). At each time point, aliquots of buffer were removed from the buffer/polymer suspensions and replaced with fresh buffer. The aliquot was then analysed by UV-Vis spectrophotometry. The results showed that the UV intensity of the ibu⁻ (band at 221 nm) was the most intense and it remained practically constant. Compounds were also soaked under continuous stirring followed by solution removal after eight days, when the solvents were separated from the solids and they were completely replaced with fresh PBS buffer solution. The intensity of the ibuprofen UV-band (221 nm) increased significantly after the complete change of the solution. As fresh PBS buffer solution was added the intensity of the UV band of the SCS ligand (at 264 nm) increased until it became more intense than the band related to ibu⁻ (at least six replacements were required). Similar behavior was found for **2**, albeit with less pronounced changes in intensity of the UV bands (Figure S9, SI).

The diffraction patterns of the solids remaining after the treatments discussed above show that both systems are crystalline and correspond to a pattern due to the same unidentified structure. SEM images recorded after anionic exchange also showed that the morphology changed completely after the treatment (Figure 1). A significant absorption band at around 1050 cm⁻¹ due to phosphate was observed in the IR spectrum of the residue after drug release (Figure S11, SI). This suggests the involvement of the phosphate anions in the drug release process through the formation of a compound with Zn²⁺, SCS and PO₄³⁻ anions.

Different experiments were carried out in order to shed light on the release mechanism and to quantify the ibuprofen in solution (Section 4, SI). The decomposition of polymeric carrier **2** was further clarified by NMR studies of the remaining solutions in PBS buffer using D₂O as solvent. The use of D₂O instead of water allowed an in-depth study by NMR spectroscopy of the mechanism of ibuprofen delivery. Interestingly, a molar ratio of 8:1 (ibu⁻:SCS) was obtained when the equilibrium was reached and this remained constant during one month of stirring. The ibu⁻ concentration in the solution at this molar ratio was 2.7·10⁻³ M (0.34·10⁻³ M of SCS) and this was stable over time. When the complete replacement of solvent had been achieved the concentration of ibu⁻ was 2.25·10⁻³ M in the first three batches and this started to decrease after the fourth replacement (Table S7, SI).

The release of SCS ligand indicated that the drug release process is accompanied by gradual decomposition of the polymeric compounds in the last stage. This process resulted in powders in which the main components were Zn²⁺ and PO₄³⁻ anions. This point is also advantageous since degradation of the drug carrier is generally required in clinical applications.

Release of ibuprofen in acidic media

The behavior of **1** and **2** in acidic media was examined. Both compounds were soaked in water with different amounts of 0.1 M HCl (Figure 4). The pH was measured after 30 minutes and 24 hours of treatment although significant variation in this parameter was not observed. The final values were in the range 4.6–4.9 after the addition of 100 μL of 0.1 M HCl, 4.1–4.5 after 300 μL and around 2.2–2.7 after 1000 μL of 0.1 M HCl. Pristine compounds dissolved completely in more acidic media (pH between 2.2–2.7) with ibuprofen adhering to the walls and floating in the vial due to its insolubility in water. Characterization of the final solid phases after the treatment (pH between 4–5) showed that in all cases the final product corresponded to the original material together with some insoluble ibuprofen (Section 5, SI). Moreover, under much more acidic conditions (pH around 2) the materials were totally degraded and single crystals of $[\text{H}_2\text{SCS}][\text{ZnCl}_4]$ were obtained from compounds **1** and **2**.




Figure 4. Schematic representation of the degradation process in acidic media.

The UV-Vis spectra of the solutions under different pH conditions are shown in Figures S14 and S15. Protonation of the SCS ligand induced a bathochromic shift in the band at 265 nm to 296 nm (Figure S13, SI). The addition of acid led to a progressive increase in the intensity of the band at 296 nm and this change was accompanied by a decrease in the absorption bands at 212 and 265 nm. The band at 212 nm was shifted steadily upon protonation to 222 nm. This behavior was observed for both compounds. According to the observed trends, one would expect that in more acidic media the amount of $[\text{SCS-H}_2]^{2+}$ species in solution would be higher than in the less acidic media.

The acid dissociation constant (pKa) of ibuprofen (4.91–5.2) means that the vast majority of ibuprofen molecules are present as undissociated ibuprofen under the acidic conditions used and this compound is in the solid state due to its low solubility in water.

The rate of degradation of **1** was further elucidated by NMR studies of the resulting solutions when a similar experiment was performed in D_2O rather than water (Table S10, SI). The ability to protonate both pyridine groups under these conditions has been studied previously¹⁹ and, in this case, the protonation is responsible for the dissolution and degradation of the initial compound. Protonation of SCS led to a significant increase in the solubility of the ligand in water and the NMR signals of the pyridine rings were shifted to lower field (Table S10, SI). The amount of $[\text{SCS-H}_2]^{2+}$ species in solution increased considerably upon acidification until complete degradation of the initial polymeric compound occurred along with the release of ibuprofen. The $[\text{SCS-H}_2]^{2+}:\text{ibu}^-$ molar ratio in solution could

be estimated by analyzing the relative intensities of the NMR signals – approximately 2:1 at pH \sim 5 and approximately 13:1 at pH \sim 4 – and ibuprofen was not detected in solution in the much more acidic conditions. Moreover, under these conditions single crystals of $[\text{H}_2\text{SCS}][\text{ZnCl}_4]$ were isolated from the solutions.

SEM images of the recovered products show evidence for cracking of the crystals, with visible changes in morphology under slightly acidic conditions (Figure 1) as the roughness of the materials increased with the level of disintegration.

Cytotoxicity

The cytotoxicity levels against normal cells (MRC-5, non-tumor human fibroblasts) of SCS, **1**, **2**, **3** and **4** were determined. The results of these experiments are included in Table 1. All of the compounds showed low growth inhibition at the maximum concentration assayed ($10 \pm 3\%$ for 100 μM) and the growth inhibition curve was not calculated. These results suggest that the introduction of high concentrations of SCS and Zn compounds barely changes the cytotoxicity profile of SCS and Zn compounds on MRC-5 cells. The cytotoxicity of the polymeric compounds **1** and **2** are related to the toxicity arising from the interactions between ibuprofen, the SCS ligand and Zn ions of the polymeric compounds with MRC-5 cells.

Conclusion

The studies described above provide a new strategy for the development of metal-drug coordination polymers constructed with biomolecular building blocks as potential solid forms of active pharmaceutical ingredients. The nature of these polymeric compounds for the storage and release of the ibuprofen drug molecule was investigated. These systems have high drug contents and high thermal and aqueous stability.

The different compositions and coordination environments of the metal centers in the two polymeric carriers (**1** and **2**) are associated with two distinct drug release stages. The unique properties of the coordination bonds, i.e., thermodynamic stability and lability, play a crucial role in this behavior. The drug release in the initial stage may be regulated by the possibility of anionic exchange in the media (note that **1** is stable in distilled water). In this way, a second stage is reached and this involves polymeric compound **2**. Release of ibuprofen in the last stage is a slower process that occurs along with decomposition of the material. An MTT cytotoxicity assay with SCS, **1**, **2**, **3** and **4** at the maximum concentration tested showed that the new functional materials possess negligible cytotoxicity. These biocompatible polymers appear to be innovative tools with the potential for drug delivery.

Acknowledgements

Financial support from ERDF (EU), MEC (Spain) and the Xunta de Galicia (Spain) (research projects CTQ2010-19386/BQU and 10TMT314002PR) are gratefully acknowledged. A. B. L thanks the Xunta de Galicia for a postdoctoral contract under the ÁngelesAlvariño Program


References

- 1 (a) H. Wang, T.-L. Hu, R.-M. Wen, Q. Wang, X.-H. Bu, *J. Mater. Chem. B.*, 2013, **1**, 3879–3882; (b) B. Lou, F. He, *New, J. Chem.*, 2013, **37**, 309–316.
- 2 (a) V. Aanad, R. Kandarapu, S. Garg, *Drug Discov. Today*, 2001, **6**, 905–914; (b) P. Horcajada, R. Gref, T. Baati, P. K. Allan, G. Maurin, P. Couvreur, G. Férey, R. E. Morris, C. Serre, *Chem. Rev.*, 2012, **112**, 1232–1268.
- 3 X. Guo, R.-K. Chang, M. A. Hussain, *J. Pharm. Sci.*, 2009, **98**, 3886–3902.
- 4 R. C. Huxford, J. D. Roca, W. Lin, *Curr. Opin. Chem. Biol.* 2010, **14**, 262–268.
- 5 L. A. Belfiore, M. P. M. Curdie, P. K. Das, *Polymer*, 2001, **42**, 9995–10006.
- 6 (a) A. Paloic, A. Puskaric, M. Mazaj, E. Zunkovio, N. Z. Logar, J. Bronic, *J. Solid State Chem.*, 2015, **225**, 59–64; (b) S. R. Miller, D. Heurtaux, T. Baati, P. Horcajada, J.-M. Grenèche, C. Serre, *Chem. Commun.*, 2010, **46**, 4526–4528.
- 7 (a) J. An, S. Geib, N. L. Rosi, *J. Am. Chem. Soc.*, 2009, **131**, 8376–8377; (b) N. Motakef-Kazemi, S. A. Shojaosadaati, A. Morsali, *Micropor. Mesopor. Mat.*, 2014, **186**, 73–79.
- 8 A. S. Prasad, *Mol. Med.*, 2008, **14**, 353–357.
- 9 S. Binauld, M. H. Stenzel, *Chem. Commun.*, 2013, **49**, 2082–2102.
- 10 L. L. Tan, H. L., Y. C. Qiu, D. X. Chen, X. Wang, R. Y. Pan, Y. Wang, S. X. A. Zhang, B. Wang, Y. W. Yang, *Chem. Sci.*, 2015, **6**, 1640–1644.
- 11 Q. Hu, J. Yu, M. Liu, A. Liu, Z. Dou, Y. Yang, *J. Med. Chem.*, 2014, **57**, 5679–5685.
- 12 (a) A. B. Lago, R. Carballo, S. Rodríguez-Hermida, E.M. Vázquez-López, *Cryst. Eng. Comm.*, 2013, **15**, 1563–1570. (b) A. B. Lago, R. Carballo, O. Fabelo, N. Fernández-Hermida, F. Lloret, E. M. Vázquez-López, *Cryst. Eng. Comm.*, 2013, **15**, 10550–10562; (c) R. Carballo, B. Covelo, N. Fernández-Hermida, E. García-Martínez, A. B. Lago, E. M. Vázquez-López, *Polyhedron*, 2009, **28**, 923–932; (d) R. Carballo, B. Covelo, N. Fernández-Hermida, E. García-Martínez, A. B. Lago, E. M. Vázquez-López, *Polyhedron*, 2008, **27**, 3247–3254.
- 13 R. Carballo, B. Covelo, E. García-Martínez, A. B. Lago, E. M. Vázquez-López, *Z. Anorg. Allg. Chem.*, 2007, **633**, 780–782.
- 14 B. Tita, M. Stefanescu, D. Tita, *Rev. Chim.*, 2011, **62**, 1060–1064; (b) C. Dendrinou-Samara, G. Tsotsou, L. V. Ekateriniadou, A. H. Kortsaris, C. P. Raptopoulou, A. Terzis, D. A. Kyriakidis, D. P. Kessissoglou, *J. Inorg. Biochem*, 1998, **71**, 171–179.
- 15 (a) D.S. Wishart, C. Knox, A.C. Guo, S. Shrivastava, M. Hassanali, P. Stothard, Z. Chang, J. Woolsey, *Nucleic Acids Res.* **2006**, 34 (Database issue):D668-72. (b) A. M. Kumar, A. Swathi, D. Supriya, V. V. L. N. Prasad, P. V. Diwan, *Am. J. Pharm. Tech. Res.*, 2012, **2**, 541–547.
- 16 (a) R. Carballo, B. Covelo, N. Fernández-Hermida, E. García-Martínez, A. B. Lago, E. M. Vázquez-López, *Cryst. Growth Des.*, 2008, **8**, 995–1004; (b) R. Carballo, B. Covelo, N. Fernández-Hermida, E. García-Martínez, A. B. Lago, E. M. Vázquez-López, *J. Mol. Struct.*, 2008, **892**, 427–432. (c) R. Carballo, B. Covelo, N. Fernández-Hermida, E. García-Martínez, A. B. Lago, E. M. Vázquez-López, *Polyhedron*, 2007, **26**, 780–782.
- 17 (a) M. L. Vueba, M. E. Pina, L. A. E. Batista de Carvalho, *J. Pharm. Sci.*, 2008, **97**, 845–859. (b) S. Bonora, A. Pisi, S. Ottani, D. Cesini, A. Maris, M. D. Foggia, *Vib. Spectrosc.*, 2014, **73**, 45–55
- 18 (a) A. B. Lago, R. Carballo, S. Rodríguez-Hermida, E. M. Vázquez-López, *Cryst. Growth Des.*, 2014, **14**, 3096–3109; (b) K. Nakamoto, *Infrared and Raman Spectra of Inorganic and Coordination Compounds*, 4th ed., J. Wiley & Sons, New York, 1986.
- 19 A. B. Lago, R. Carballo, E. García-Martínez, E. M. Vázquez López, *Cryst. Growth Des.*, 2011, **11**, 59–68.
- 20 (a) M. Korn, A. Haas, H. Oberhammer, *Chem. Ber.*, 1995, **128**, 461–464; (b) E. M. Page, D. A. Rice, K. Aersset, K. Hagen, A. R. J. Genge, *J. Phys. Chem. A.*, 2000, **104**, 6672–6676.

Graphical Abstract

A new Metal-Organic Polymeric system capable of stimuli-responsive controlled release of the drug ibuprofen.

Ana Belén Lago, Arantxa Pino-Cuevas, Rosa Carballo, Ezequiel M. Vázquez-López.
DOI:



New polymeric zinc coordination compounds capable of release ibuprofen drug.