

# Dalton Transactions

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



## Synthetic, Structural, and Computational Investigations of N-alkyl Benzo-2,1,3-Selenadiazolium Iodides and their Supramolecular Aggregates

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Lucia M. Lee,<sup>a</sup> Victoria B. Corless,<sup>a</sup> Michael Tran,<sup>a</sup> Hilary Jenkins,<sup>a</sup> James F. Britten<sup>a</sup> and Ignacio Vargas-Baca<sup>a\*</sup>

Despite their versatility, the application of telluradiazoles as supramolecular building blocks is considerably constrained by their sensitivity to moisture. Albeit more robust, their selenium analogues form weaker supramolecular interactions. These, however, are enhanced when one nitrogen atom is bonded to an alkyl group. Here we investigate general methods for the synthesis of such derivatives. Methyl, *iso*-propyl and *tert*-butyl benzo-2,1,3-selenadiazolium cations were prepared by direct alkylation or cyclo-condensation of the alkyl-phenylenediamine with selenous acid. While the former reaction only proceeds with the primary and tertiary alkyl iodides, the latter is very efficient. Difficulties reported in earlier literature are attributable to the formation of adducts of benzoselenadiazole with its alkylated cations and side reactions initiated by aerobic oxidation of iodide. However, the cations themselves are resilient to oxidation and stable in acidic to neutral aqueous medium. X-ray crystallography was used in the identification and characterization of the following compounds:  $[\text{C}_6\text{H}_4\text{N}_2(\text{R})\text{Se}]^+\text{X}^-$ , (R =  $\text{CH}(\text{CH}_3)_2$ ,  $\text{C}(\text{CH}_3)_3$ ; X = I<sup>-</sup>, I<sub>3</sub><sup>-</sup>),  $[\text{C}_6\text{H}_4\text{N}_2(\text{CH}_3)\text{Se}]^+\text{I}^-$ , and  $[\text{C}_6\text{H}_4\text{N}_2\text{Se}][\text{C}_6\text{H}_4\text{N}_2(\text{CH}_3)_2\text{Se}]_2\text{I}_2$ . Formation of Se $\cdots$ N secondary bonding interactions (chalcogen bonds) was only observed in the last structure as anion binding to selenium is a strong competitor. The relative strengths of those forces and the structural preferences they enforce were assessed with DFT-D3 calculations supplemented by AIM analysis of the electron density.

### Introduction


While hydrogen bonding and the coordination of metal ions are preponderant in supramolecular chemistry, other interactions are receiving increasing interest in this field. A notable example is halogen bonding,<sup>1-3</sup> a special case of the phenomenon termed secondary bonding<sup>4</sup> that is recurrent in the structures of compounds of the heavy p-block elements. The potential of secondary bonding in supramolecular chemistry is well exemplified by the derivatives of the 1,2,5-telluradiazole ring (**1**, Chart 1). Their molecules usually associate by two antiparallel Te $\cdots$ N secondary bonding interactions forming the  $[\text{Te-N}]_2$  supramolecular synthon,<sup>5</sup> a virtual four-membered ring (Scheme 1, E = Te).<sup>6</sup> In the absence of substantial steric hindrance, each molecule can simultaneously form the  $[\text{Te-N}]_2$  supramolecular synthon twice building in this way supramolecular ribbons.<sup>7</sup> The driving force for association of these molecules is strong enough to overcome moderate steric repulsion by allowing structural distortion of the ribbons. This effect induces properties such as chromotropism<sup>8</sup> and second-harmonic generation<sup>9</sup> which are of interest for practical

applications. In addition to the self-association of 1,2,5-telluradiazoles through Te $\cdots$ N interactions, other studies have also evaluated their affinity to Lewis bases in both solid<sup>10</sup> and solution.<sup>11-13</sup> In spite of their versatility, prospects of widespread application of telluradiazoles are limited by their sensitivity to moisture.



1 E = Te  
2 E = Se  
3 E = S

Chart 1



E = S, Se, Te

Scheme 1

The lighter members of the chalcogenadiazole family (**2,3**, Chart 1) are much more robust but SBIs centred on S and Se atoms are weaker than those formed by Te<sup>14</sup> and instances of supramolecular association are less frequent in such cases. However, crystallographic evidence suggests that attachment of Lewis acid (a transition metal ion<sup>15,16</sup> or a borane<sup>17</sup>), on one nitrogen of the heterocycle strengthens the secondary interactions made by Se and Te, shortening the corresponding

<sup>a</sup> Department of Chemistry and Chemical Biology, McMaster University, 1280 Main St. W. Hamilton, ON, Canada L8S 4M1. E-mail: vargas@chemistry.mcmaster.ca  
Electronic Supplementary Information (ESI) available: Crystallographic details for  $[\text{4}]_2[\text{4-H}]_2[\text{I}][\text{I}_3]$  and  $[\text{4}]_2(\text{C}_6\text{H}_4(\text{NH}_2)_2\text{H}^+)[\text{I}]_2$ ; Cartesian coordinates of all calculated structures. See DOI: 10.1039/x0xx00000x

distances. Such effect is also strong in the cations formed by N-alkylation and provides an explanation for the peculiar trend in reduction potentials of the N-methyl-benzo-2,1,3-chalcogenadiazolium cations in acetonitrile.<sup>18</sup> Such evidence of molecular association in solution through SBIs is consistent with the observation of  $[M_2H]^+$  supramolecular aggregates in the mass spectrum of benzo-2,1,3-telluradiazoles<sup>19</sup> as well as calculated interaction energies and AIM parameters.<sup>12,18</sup>

N-alkyl substituted selenadiazolium cationic rings could therefore be a convenient alternative to telluradiazoles in supramolecular architecture, provided that they are reasonably stable and the preparation methods are flexible enough to accommodate multiple variations in the structure. The most straightforward synthetic method would be the direct alkylation of the heterocycle in **4** (Chart 2). However, early reports of this approach describe very inefficient reactions plagued by multiple by-products with most common alkylating reagents.<sup>20,21</sup> Clean and efficient alkylations have been reported rather recently using very reactive agents such as  $Me-OS(O)_2-R'$  ( $R' = OMe, C_6H_2(NO_2)_2-2,4,$ <sup>20</sup>  $F_3C-SO_3-CH_3$ <sup>18</sup> or  $(CH_3)_3O^+BF_4^-$ <sup>22</sup>), but these methods are applicable with only a handful of alkyl groups. Alternatively, the alkyl group could be attached to nitrogen before formation of the heterocycle. This approach was actually reported in an early publication<sup>20,21</sup> but its products were not structurally authenticated. A third route to N-alkyl selenadiazolium cations, also limited in flexibility, consists of the spontaneous de-alkylation of one N nitrogen atom during the reaction of N,N'-bis(tert-butyl)-diazabutadiene with  $SeCl_4$ .<sup>23</sup>




Chart 2

As we are interested in using the N-alkylated selenadiazolium cations as supramolecular building blocks, in this report we compare the two most flexible synthetic methods for the preparation of the derivatives with primary, secondary and tertiary alkyl groups (**5**, Chart 2). We examine the crystal structures of their iodide salts, the affinity of the cations for halide anions, and spectroscopic properties useful for the characterization of these species in solution.

## Experimental

**Materials and General Procedures.** Selenium dioxide, iodomethane were used as received from commercial suppliers. The N-alkyl 1,2-phenylenediamines were prepared from the reaction of the corresponding commercially available primary

amines with 1-fluoro-2-nitrobenzene, followed by reduction with  $H_2$  catalyzed by Pd/C.<sup>24-26</sup> Benzo-2,1,3-selenadiazole was prepared by the reaction of 1,2-phenylenediamine with selenium dioxide according to literature.<sup>27</sup> Anhydrous-grade solvents were used without further purification.

**Instrumentation.**  $^1H$ ,  $^{13}C$  and  $^{77}Se$  NMR spectra were recorded on Bruker 200 and Bruker 600 spectrometers at ambient temperature. Chemical shifts are reported in  $\delta$  values (parts per million) with respect to the resonances of tetramethylsilane for  $^1H$  and  $^{13}C$  and with respect to the line of  $Me_2Se$  for the chemical shift of  $^{77}Se$ . In the last case, spectra were obtained using  $^1H$ - $^{77}Se$  HMBC unless otherwise stated. IR spectra were recorded on polyethylene pellets on a Nicolet 6700 FT-IR spectrometer with a resolution of  $4\text{ cm}^{-1}$ . Raman spectra were obtained with a Renishaw inVia spectrometer exciting at 785 nm, 30 mW, averaging  $10 \times 10s$  scans. Low- and high-resolution electrospray spectral analyses were performed using Micromass GCT spectrometer and a Micromass Quattro Ultima for electron ionization spectra. In each case, the sample was introduced into the ionization chamber in a shortened borosilicate glass capillary on a probe rod. The samples were then heated and the temperature range that displayed the most intense parent ion peaks was used for spectral acquisition. A 70 eV electron stream ionized the sample and the positively charged ions were identified by a time-of-flight detector. A Cary 50 UV-vis spectrometer and its Cary Win-UV software package were used to acquire the spectra. Additional processing was carried out using the GramsAI suite (version 8.0). Melting points were measured on a Thomas-Hoover melting point apparatus and are reported uncorrected. Elemental analysis was performed by the Science Centre of London Metropolitan University, 29 Hornsey Road, London, UK, N7 7DD.

All single crystal X-ray diffraction samples were mounted with paratone oil at the tip of a glass fibre installed on a MiTeGen goniometer head, and kept under a cold stream of nitrogen while on the diffractometer. Data were collected at 100 K on a Bruker APEX2 diffractometer, using Mo K  $\alpha$  radiation ( $\lambda = 0.71073\text{ \AA}$ ) and outfitted with an Oxford cryostream low-temperature accessory.  $\Omega$  and  $\phi$  scans were collected in  $0.5^\circ$  steps with a crystal to detector distance of 4.954 cm. The preliminary unit cell parameters were determined using a minimum of 50 frames from three different orientations, and final cell refinement after integration in SAINT. Data were corrected for absorption and scaled using face-indices as well as redundant data in SADABS. Crystal structures were solved using SHELXL and most structures were refined by full-matrix least square of all F2 values with the WinGX package.

## Syntheses

$(C_6H_4N_2Se)[C_6H_4N(CH_3)Se]_2[I]_2$  (**4**) $[5a]_2[I]_2$ ). Benzo-2,1,3-selenadiazole (0.1 g, 0.547 mol) and neat iodomethane (0.35 mL, 0.602 mmol) were mixed with stirring under nitrogen. The mixture became purple and gave a microcrystalline red powder. Yield 60%. The product was washed with hexanes and dried under vacuum. Crystals suitable for single X-ray diffraction were grown by slow diffusion of the reagents dissolved in ethanol.  $^1H$  NMR ( $d_6$ -DMSO, 600 MHz):  $\delta$  4.13 (s, 3H,  $CH_3$ ),

7.54, 7.82, 7.96 (m, 6H, Ar-CH) ppm.  $^{13}\text{C}$  NMR ( $d_6$ -DMSO, 600 MHz):  $\delta$  159.74 156.40 150.622 131.65 126.43 125.88 116.91 (Ar-CH) 39.42 ( $\text{CH}_3$ ) ppm.  $^{77}\text{Se}$  NMR ( $d_6$ -DMSO, 600 MHz):  $\delta$  1494.0 ( $[\text{C}_6\text{H}_4\text{N}(\text{NCH}_3)\text{Se}]^+$ ), 1529.7 ( $\text{C}_6\text{H}_4\text{N}_2\text{Se}$ ) ppm. IR ( $\text{cm}^{-1}$ ): 3602 (m), 2854(b), 2341 (m), 2150 (vw), 2018 (w), 1982 (w), 1949 (w), 1812 (w), 1600 (w), 1529 (m), 1515 (m), 1464(sh), 1134 (vw), 1044(m), 905(vw), 730 (sh) 720 (sh). Mp. 148–150 °C (d). E.A. % calcd. for  $\text{C}_{20}\text{H}_{18}\text{I}_2\text{N}_6\text{Se}_3$ : C 28.83, H 2.18, N 10.09, ; found C 28.72, H 2.09, N 9.98.

**[ $\text{C}_6\text{H}_4\text{N}(\text{NCH}_3)\text{Se}][\text{I}]$  (**[5a][I]**). **Method A.** A sample of **4[5a]<sub>2</sub>[I]<sub>2</sub>** (0.02 g, 0.109 mmol) was dissolved in methanol (1.6 mL) with stirring at 42–46 °C. The colour of the solution slowly changed from dark green to bright red. Cooling to room temperature produced red crystals which were recovered by filtration. Yield 17%. **Method B.** To a solution of N-methylphenylenediamine (0.11 mL, 1.023 mmol) and  $\text{H}_2\text{SeO}_3$  (0.1319 g, 1.023 mmol) in anhydrous ethanol, trifluoroacetic acid (0.07 mL, 1.023 mmol) was added dropwise under nitrogen. The reaction mixture was allowed to stir at room temperature for 20 min, time after which a solution of NaI (0.1533 g, 1.023 mmol) in ethanol (10 mL) was added with vigorous stirring, a dark red solid precipitate formed and was separated by filtration, washed with hexanes, dried under vacuum and stored under nitrogen. Yield 22%.  $^1\text{H}$  NMR ( $\text{D}_2\text{O}$ , 600MHz):  $\delta$  4.60 (s, 3H,  $\text{CH}_3$ ), 7.78, 7.96 (m, 4H, Ar-CH) ppm.  $^{13}\text{C}$  NMR ( $d_6$ -DMSO, 600 MHz):  $\delta$  156.36, 150.73, 136.53, 129.95, 125.02, 116.46 (Ar-C), 38.35 ( $\text{CH}_3$ ) ppm.  $^{77}\text{Se}$  NMR ( $d_6$ -DMSO, 600MHz):  $\delta$  1490.4 ppm. LR ESI-MS:  $m/z$  199.0 ( $\text{M}^+$ ). HR ESI-MS:  $m/z$  198.9774 ( $\text{M}^+$ ) (cf. calc. for  $\text{C}_7\text{H}_7\text{N}_2\text{Se}$  198.9774). IR ( $\text{cm}^{-1}$ ): 3606 (m), 3219 (w), 2913 (s), 2866 (b), 2345 (m), 2018 (w), 1956 (w), 1811(w), 1536 (w), 1470 (m), 1305 (w), 1167 (vw), 1135(m), 807 (w), 744(sh), 730(sh), 720 (sh). UV-Vis  $\lambda_{\text{max}}$  = 345 nm,  $\epsilon$  = 664.6  $\text{L mol}^{-1}\text{cm}^{-3}$ . Mp. 184–188 °C (d).**

**[ $\text{C}_6\text{H}_4\text{N}(\text{NCH}(\text{CH}_3)_2)\text{Se}_2[\text{I}]$  (**[5b][I]**). To a solution of N-isopropyl-benzene-1,2-diamine and  $\text{H}_2\text{SeO}_3$  in anhydrous ethanol, trifluoroacetic acid was added. The reaction mixture was allowed to stir at room temperature for 20 min under nitrogen. Upon the addition of NaI, the product immediately precipitated. The dark red solid was washed with hexane, dried and stored under a nitrogen atmosphere. Yield 30%.  $^1\text{H}$  NMR ( $\text{CD}_2\text{Cl}_2$ , 600 MHz):  $\delta$  7.99 (dd, 1H,  $J$  = 6, 12Hz, Ar-CH) 5.60 (sept,  $J$  = 6Hz, 1H, NH-CH) 1.87 (d, 6H,  $J$  = 7 Hz,  $\text{CH}_3$ ) ppm.  $^{13}\text{C}$  NMR ( $\text{CD}_2\text{Cl}_2$ , 600 MHz):  $\delta$  138.18 (Ar-C), 131.42 (Ar-C), 125.34 (Ar-C), 115.71 (Ar-C) 57.87 (NH-CH), 24.22 ( $\text{CH}_3$ ) ppm.  $^{77}\text{Se}$  NMR ( $d_6$ -DMSO, 600MHz):  $\delta$  1459.5 ppm. LR ESI-MS:  $m/z$  227.0 ( $\text{M}^+$ ). HR ESI-MS:  $m/z$  227.0095 ( $\text{M}^+$ ) (cf. calc. for  $\text{C}_9\text{H}_{11}\text{N}_2\text{Se}$  227.0088). IR ( $\text{cm}^{-1}$ ): 3605 (w), 3374 (m), 2848 (b), 2329 (w), 2010 (vw), 1952 (w) 1980 (vw), 1842 (vw), 1896 (vw), 1809 (vw), 1720 (vw), 1523 (w), 1473 (sh), 1369 (m), 1314 (m), 1176 (sh), 1164 (sh), 1142 (m), 1121 (m), 823 (w), 757 (vs), 730 (vs), 719 (vs). UV-Vis  $\lambda_{\text{max}}$  = 343 nm,  $\epsilon$  = 428.5  $\text{L mol}^{-1}\text{cm}^{-3}$ . Mp. 165–168 °C (d).**

**[ $\text{C}_6\text{H}_4\text{N}(\text{NC}(\text{CH}_3)_3)\text{Se}][\text{I}]$  (**[5c][I]**). Using the procedure above described for **[5a][I]** and **[5b][I]**, the compound was obtained as a dark red crystalline material.  $^1\text{H}$  NMR ( $d_6$ -DMSO, 600MHz):  $\delta$  8.30(d, 1H, Ar-CH) 8.08(d, 1H, Ar-CH) 7.97 (dd, 1H, Ar-CH) 7.79 (dd, 1H, Ar-CH) 1.97 (9H, ( $\text{CH}_3$ )<sub>3</sub>).  $^{13}\text{C}$  NMR ( $d_6$ -**

DMSO, 600 MHz):  $\delta$  158.46 148.03 (Ar-C) 136.44 (Ar-C) 129.51 (Ar-C) 126.17 (Ar-C) 118.40 (Ar-C) 67.28 (N-CH) 29.66 ( $\text{CH}_3$ )<sub>3</sub> ppm.  $^{77}\text{Se}$  NMR ( $d_6$ -DMSO, 600MHz):  $\delta$  1454.7 ppm. LR ESI-MS:  $m/z$  241.0 ( $\text{M}^+$ ) HR ESI-MS:  $m/z$  241.0248 ( $\text{M}^+$ ) (cf. calc. for  $\text{C}_{10}\text{H}_{13}\text{N}_2\text{Se}$  241.0244). IR ( $\text{cm}^{-1}$ ) 3647 (vw), 3604 (w), 2793 (b), 2340 (w), 2150 (vw), 2108 (w), 1859 (vw), 1602 (w), 1526 (w), 1472 (sh), 1371 (m), 1302 (m), 1180 (w), 1132 (w), 954 (vw), 872 (vw), 844 (vw), 744(sh), 719 (sh). UV-Vis  $\lambda_{\text{max}}$  = 330 nm,  $\epsilon$ =282.5  $\text{L mol}^{-1}\text{cm}^{-3}$ . Mp. 130–132 °C (d).

**Computational Methods.** All the structures considered in this study were fully optimized using the ADF DFT package (versions 2013.01–20140.1).<sup>28,29</sup> The Adiabatic Local Density Approximation (ALDA) was used for the exchange-correlation kernel<sup>30,31</sup> and the differentiated static LDA expression was used with the Vosko-Wilk-Nusair parameterization.<sup>32,33</sup> Calculation of model geometries was gradient-corrected with the exchange and correlation functionals of Perdew, Burke and Ernzerhof (PBE)<sup>34</sup> and applying the zeroth-order regular approximation (ZORA)<sup>34–38</sup> formalism with the specially adapted triple- $\zeta$  all-electron plus one-polarization-function basis sets. The contribution of dispersion was modelled with Grimme's correction.<sup>39</sup> Analytical frequency calculations were performed to ensure that each geometry was at an energy minimum.<sup>40,41</sup> The TD-DFT calculation was performed from the optimized geometry of  $[\text{C}_6\text{H}_4\text{N}(\text{NCH}_3)\text{Se}]^+$  using the statistical average of potentials model for exchange and correlation (SAOP)<sup>42–44</sup> and a basis set of quadruple-zeta quality plus polarization.

## Results and discussion

**Syntheses by direct alkylation.** Although benzo-2,1,3-selenadiazole is a weak base, it is readily protonated by strong acids.<sup>45</sup> For this reason, the purported inefficiency<sup>20,21</sup> of its reaction with primary alkyl-iodides is unusual and merited reinvestigation using a more reactive tertiary iodide. Initial attempts at bench-top alkylation with  $(\text{CH}_3)_3\text{CI}$  in a hot toluene solution yielded a complex mixture, as made evident by the  $^1\text{H}$  NMR spectrum. Slow evaporation of the solution yielded a mixture of crystals with multiple morphologies; three of which were identified by single-crystal X-ray diffraction: **[5c][I]<sub>3</sub>**, **[4]<sub>2</sub>[4-H]<sub>2</sub>[I]<sub>3</sub>** and **[4]<sub>2</sub>( $\text{C}_6\text{H}_4(\text{NH}_2)_2\text{H}^+$ )<sub>2</sub>[I]<sub>2</sub>**. The first species does contain the N-tert-butyl benzo-selenadiazolium cation (**5c**), which demonstrates that the alkylation reaction does proceed. However, the presence of the triiodide anion, plus protonated phenylenediamine and benzoselenadiazole point to the actual problem with this experiment: the iodide ion is likely oxidized by atmospheric oxygen to  $\text{I}_3^-$  which then halogenates the solvent or benzoselenadiazole to generate HI. Crystals of **[5c][I]** were obtained when the reaction was carried out under an atmosphere of nitrogen, albeit in low yield. Similarly, the alkylation reaction proceeded with neat  $\text{CH}_3\text{I}$  in anaerobic conditions, however, in this case the alkylation is incomplete due to the formation of the crystalline **4[5a]<sub>2</sub>[I]<sub>2</sub>**, which was identified by X-ray diffraction. The salt **[5a][I]** was obtained from **4[5a]<sub>2</sub>[I]<sub>2</sub>** by recrystallization from dilute solutions.

Interestingly, *iso*-propyl iodide was completely unreactive towards benzoselenadiazole under the same conditions. While the reaction of *tert*-butyl iodide is likely to proceed by a SN<sub>1</sub> mechanism, it is likely that the primary and secondary iodides prefer a SN<sub>2</sub> mechanism but steric hindrance prevents the reaction of the *iso*-propyl halide.

**Condensation of selenous acid with N-alkylated phenylenediamines.** Even in aqueous medium, benzoselenadiazole is readily formed by the reaction of H<sub>2</sub>SeO<sub>3</sub> and phenylenediamine. N-substituted *ortho*-diamino benzenes were reported to undergo an analogous reaction in glacial acetic producing the benzoselenadiazolium cations, which are isolated by precipitation with sodium halide.<sup>46,47</sup> This procedure works

as expected but we found more convenient to carry out the reaction in a mixture of ethanol and trifluoroacetic acid in order to avoid using the less volatile acetic acid. Structural characterization by X-ray diffraction did confirm the identity of the products. We also found that exposure of the cations in solution to the atmosphere resulted in iodide oxidation and crystallization of the corresponding triiodide salts, which was verified by the strong scattering band at 111 cm<sup>-1</sup> in the Raman spectrum<sup>48</sup> or crystallographic analysis.

**X-ray crystallography.** Crystallographic and refinement data for the salts of the three cations are presented in Table 1; selected distances and angles in each crystal structure are provided in Table 2.

Table 1. – Crystallographic and Refinement Parameters for 4[5a]<sub>2</sub>[I]<sub>2</sub> and the salts of 5a-c.

	4[5a] <sub>2</sub> [I] <sub>2</sub>	[5a][I]	[5b][I <sub>3</sub> ]	[5b][I]	[5c][I <sub>3</sub> ]	[5c][I]
Chemical formula	C <sub>20</sub> H <sub>18</sub> I <sub>2</sub> N <sub>6</sub> Se <sub>3</sub>	C <sub>7</sub> H <sub>10</sub> N <sub>2</sub> SeI	C <sub>9</sub> H <sub>11</sub> N <sub>2</sub> SeI <sub>3</sub>	C <sub>9</sub> H <sub>11</sub> N <sub>2</sub> SeI	C <sub>10</sub> H <sub>13</sub> N <sub>2</sub> SeI <sub>3</sub>	C <sub>10</sub> H <sub>13</sub> N <sub>2</sub> SeI
CCDC	1424372	1424371	1424373	1424374	1424370	1434655
Formula mass	833.08	325.01	606.86	353.06	620.9	367.08
Crystal system	monoclinic	monoclinic	monoclinic	monoclinic	monoclinic	orthorhombic
a, Å	15.986(1)	6.471(1)	11.317(2)	16.084(1)	7.668(1)	11.668(1)
b, Å	11.671(1)	13.216(2)	9.818(2)	10.525(1)	10.014(2)	7.699(1)
c, Å	12.743(1)	10.474(2)	13.569(2)	6.597(1)	20.949(4)	13.253(1)
α, °	90	90	90	90	90	90
β, °	92.566(3)	92.426(4)	105.047(3)	90	95.631(3)	90
γ, °	90	90	90	90	90	90
Unit cell volume, Å <sup>3</sup>	2375.2(2)	894.9(3)	1456.0(4)	1116.8(2)	1600.7(5)	1190.6(1)
Temperature, K	100.15	100.15	100.15	100.15	100.15	296
Space group	C 1 2/c1	P 21/n	P 1 21/n 1	P n a 21	P 21/c	P n m a
Z	4	4	4	4	4	4
No. of reflections measured	2700	12698	19082	16547	8761	26960
No. independent reflections	2213	2521	4908	3306	4094	3066
Final R <sub>1</sub> values (I > 2σ(I))	0.0404	0.0199	0.0261	0.0229	0.0381	0.0331
Final wR(F <sup>2</sup> ) values (I > 2σ(I))	0.0685	0.0385	0.0447	0.0419	0.0759	0.0828
Final R <sub>1</sub> values (all data)	0.0550	0.0262	0.0399	0.0276	0.0682	0.0413
Final wR(F <sup>2</sup> ) values (all data)	0.0733	0.0404	0.0481	0.0434	0.0861	0.0847

$$R_1 = \frac{\sum |F_{obs} - F_{calc}|}{\sum |F_{obs}|} \text{ and } wR(F_2) = \frac{(\sum w|F_{obs} - F_{calc}|^2)^{1/2}}{\sum wF_{obs}^2}$$

The crystal structure of 4[5a]<sub>2</sub>[I]<sub>2</sub> consists of a molecule of 4 bound to two 5a, conforming a pseudo-trimer assembled by two asymmetric [Se-N]<sub>2</sub> supramolecular synthons (Figure 1) with asymmetric SBI distances of 2.576(1) and 2.934(1) Å. This aggregate is analogous to the product formed by the reaction of benzoselenadiazole with [(CH<sub>3</sub>)<sub>3</sub>O][BF<sub>4</sub>]<sup>22</sup>, in which the Se...N and Se...F secondary bonding distances are 2.573(2) and 2.966(6) Å and 2.970(4) Å respectively. In this arrangement,

the molecules of 4 and 5a are nearly coplanar with a deviation angle of 5.06°. This structure also features Se-I short distances at 3.528(1) Å and 3.831(1) Å. Both iodides sit out of the average plane of the aggregate forming a virtual four-membered ring (i.e. the [Se I]<sub>2</sub> supramolecular synthon) with the Se atoms of the 5a molecules of two neighbouring pseudotrimers.




## Dalton Transactions

## ARTICLE


Table 2. Selected Bond Distances and Angles for **4**[**5a**]<sub>2</sub>[I]<sub>2</sub> and the salts of **5a-c**.

Compound	<b>4</b> [ <b>5a</b> ] <sub>2</sub> [I] <sub>2</sub>	[ <b>5a</b> ][I]	<b>5b</b> [I <sub>3</sub> ]	[ <b>5b</b> ][I]	[ <b>5b</b> ][I <sub>3</sub> ]	<b>5c</b> I
Se-N1	1.856(5)	1.785(2)	1.766(3)	1.774(3)	1.742(5)	1.749(2)
Se-N2	1.777(5)	1.875(2)	1.857(3)	1.881(2)	1.841(4)	1.864(2)
N1-C1	1.339(7)	1.327(3)	1.333(4)	1.333(4)	1.349(8)	1.343(3)
N2-C6	1.336(7)	1.329(3)	1.338(4)	1.338(4)	1.347(6)	1.339(3)
C1-C6	1.443(7)	1.440(3)	1.444(4)	1.439(4)	1.414(7)	1.442(4)
C1-C2	1.432(8)	1.431(3)	1.355(5)	1.428(6)	1.414(7)	1.421(4)
C2-C3	1.366(8)	1.351(4)	1.426(5)	1.355(6)	1.342(9)	1.361(3)
C3-C4	1.428(8)	1.431(4)	1.426(5)	1.360(4)	1.400(9)	1.417(4)
C4-C5	1.371(8)	1.359(3)	1.353(4)	1.427(5)	1.345(9)	1.368(4)
C5-C6	1.428(8)	1.422(3)	1.422(5)	1.425(5)	1.418(1)	1.421(3)
N1-Se-N2	88.7(2)	88.16(9)	89.6(1)	88.6(2)	90.4(4)	90.2(1)
Se-N1-C1	111.3(4)	111.2(2)	110.7(2)	111.0(3)	109.7(7)	110.3(2)
Se-N2-C6	111.6(4)	111.4(1)	110.6(2)	111.0(3)	109.8(7)	110.2(2)
Se-N2-C7	124.0(4)	124.9(2)	124.4(2)	124.7(3)	122.4(6)	122.4(2)
C6-N2-C7	124.4(5)	123.7(2)s	125.1(3)	124.3(3)	127.8(8)	127.5(2)

Figure 1. Arrangements of molecules and ions in the crystal **4**[**5a**]<sub>2</sub>[I]<sub>2</sub>. Selected distances: Se<sup>1</sup>-N<sup>1</sup>: 2.573(4) Å and 2.5937 Å.

In clear contrast with the structures of the pseudo-trimers,<sup>22</sup> and the dimers observed for [**5a**][CF<sub>3</sub>SO<sub>3</sub>]<sup>18</sup> and [C<sub>2</sub>H<sub>2</sub>N(NC(CH<sub>3</sub>)<sub>3</sub>)Se][GaCl<sub>4</sub>]<sup>49</sup>, the crystal structures of the iodide and triiodide salts of the alkylated cations **5a-c** feature no association of the heterocycles to each other. In all cases, strong Se-anion interactions prevent formation of the Se<sup>⋯</sup>N SBIs (Figures 2 and 3).

The triiodide anions in the crystals of [**5b**][I<sub>3</sub>] and [**5c**][I<sub>3</sub>] engage in two interactions with the selenium atom. The shortest (3.249(1), 3.309(1) Å, respectively) is with the terminal iodine atom I1; the contact to the middle I2 atom is longer (3.626(1), 3.727(1) Å, respectively), cf. the sum of van der Waals radii 3.88 Å. In the case of [**5b**][I<sub>3</sub>] the terminal iodine atom I3 is in close proximity (3.491(3) Å, cf. the sum of van der Waals radii 3.53 Å) to the nitrogen atom of a neighbouring **5b** cation, the nearly linear geometry of this arrangement is indicative of a halogen bond. Consequently, the I2-I3 bond is longer (2.879(1) Å) in [**5b**][I<sub>3</sub>] than in [**5c**][I<sub>3</sub>] (2.829(1) Å).

Figure 2. Crystal structures of (a) [**5b**][I<sub>3</sub>] and (b) [**5c**][I<sub>3</sub>]. Selected bond distances (a) Se<sup>⋯</sup>I<sup>1</sup>: 3.249(1) Å, Se<sup>⋯</sup>I<sup>2</sup>: 3.626(5) Å; (b) Se<sup>⋯</sup>I<sup>1</sup>: 3.309(1) Å, Se<sup>⋯</sup>I<sup>2</sup>: 3.727(1) Å.

The arrangement of the ions and the Se<sup>⋯</sup>I interactions in the [**5a-c**][I] lattices are influenced the size of the alkyl group (Figure 3). The small methyl group in [**5a**][I] confines the Se<sup>⋯</sup>I interactions to the plane of the cation forming iodide-bridged dimers that conform the [Se I]<sub>2</sub> supramolecular synthon with short (3.178 (1) Å) and long (3.610 (1) Å) SBIs. Similar binding of anions to Se has been observed in related species. For example, a virtual four-membered ring consisting of Se and Cl atoms is formed in the crystal structure of [C<sub>2</sub>H<sub>2</sub>NN(H)Se][Cl] (2.900(1) Å and (3.075 (1) Å).<sup>45</sup> The structure of [C<sub>2</sub>H<sub>2</sub>N(NC(CH<sub>3</sub>)<sub>3</sub>)Se][Cl] also features a planar selenadiazole ring with one Se-Cl bond distance of 2.605 (1) Å.<sup>23</sup> In contrast, the GaCl<sub>4</sub><sup>-</sup> salt of the same selenadiazole heterocycle does not display any Se-Cl short interactions, instead it contains the [Se-N]<sub>2</sub> supramolecular synthon.<sup>23</sup> The larger alkyl groups favour interlayer interactions (eminently electrostatic) in [Se I]<sub>∞</sub> supramolecular chains. The Se-I distances are 3.043 (1) Å and 3.696 (1) Å in the structure of [**5b**][I] and 3.147 (1) and 3.880 (1) Å in [**5c**][I].




Figure 3. Packing arrangement in the crystal structures of the iodide salts of (a) **5a** (b) **5b** and (c) **5c**.

**Spectroscopy in solution.** The iodide salts of **5a-c** are stable in air. The compounds are slightly soluble in water at room temperature, their solubility decreases with the size of the R group. In fact, it is possible to recrystallize **5a** from hot water. Aqueous solutions of the cations have a characteristic red-brown colour, their UV-vis absorption spectrum displays prominent maxima in the range of 330 - 343 nm and a weaker band between 440 - 460 nm. These spectra are in excellent agreement with the TD-DFT calculation for the **5a**, which attributes the bands to the LUMO  $\leftarrow$  HOMO and LUMO  $\leftarrow$  HOMO-1 excitations (Figure 4). The UV-vis spectra do not change from neutral to acidic pH, but hydrolysis ensues in basic medium. Given the modest solubility of the **5a-c** iodides, their  $^{77}\text{Se}$  NMR chemical shifts are best obtained through  $^1\text{H}$ - $^{77}\text{Se}$  heteronuclear multiple bond correlation (HMBC) experiments. The  $^{77}\text{Se}$   $\delta$  values (1455-1490 ppm) appear at slightly lower frequency than the 1515 ppm reported for  $[\text{C}_6\text{H}_4\text{N}(\text{NCH}_3)\text{Se}][\text{CF}_3\text{SO}_3]$  in acetonitrile.<sup>18</sup> For the pseudo-trimer **4**[**5a**]<sub>2</sub>[I]<sub>2</sub>, two resonances were observed with frequencies that correspond to those of the constituting neutral and cationic molecules, suggesting that the aggregate dissociates in solution.




Figure 4. Experimental (left) and calculated (right) UV-vis spectra of [**5a**][I]. Frontier orbitals shown as an inset in the calculated spectrum.

**DFT calculations.** As the crystal structures show there is competition between the formation of the  $[\text{Se-N}]_2$  supramolecular synthon and binding of the anion to selenium, the relative strengths of these interactions were evaluated using DFT and were extended to include the chloride and bromide

salts. For computational expediency, GGA calculations were performed with the PBE functional, supplemented by dispersion and relativistic corrections. Model structures were optimized from the crystallographic coordinates and expanded to include hypothetical isomers and the analogues with chloride and bromide anions. Whenever an experimental structure was available, bond lengths were reproduced within 0.01 Å, the largest deviations correspond to the intramolecular Se-N distances. Selenium-centred secondary bond distances were reproduced within 0.01 Å to nitrogen and 0.50 Å to iodide. The large deviation in the latter case is likely due to the additional interactions of the anion in the lattices. Interaction energies were evaluated using the transition-state method<sup>50</sup> that partitions the energy of interaction between two molecules or fragments in a hypothetical process in which the constituting units are first calculated individually, brought to their equilibrium positions without orbital mixing, and finally the electron density is relaxed by the interaction of fragment orbitals. The change from the first to the second step is regarded as the *total steric interaction* that results from the sum of the electrostatic ( $E_{\text{Elstat}}$ ) and Pauli-repulsion ( $E_{\text{Pauli}}$ ). The third step gives the orbital interaction energy ( $E_{\text{Orb}}$ ). Dispersion is treated as a separate contribution ( $E_{\text{Disp}}$ ). The total interaction energy is given by the sum of all contributions (Eq. 1).

$$E_{\text{Interaction}} = E_{\text{Elstat}} + E_{\text{Pauli}} + E_{\text{Orb}} + E_{\text{Disp}} \quad (1)$$




Figure 5. Partition schemes used in the DFT-D3 fragment interaction analyses. Each fragment (region) is identified with a distinct colour.

Three partitioning schemes (Figure 5, i-iii) were used to evaluate the secondary interactions within the pseudo-trimer **4**[**5a**]<sub>2</sub>[I]<sub>2</sub> the results are summarized in Table 3. In this analysis the average interaction energy of the  $[\text{Se-N}]_2$  supramolecular synthon was 56.4 kJ/mol. Formation of the interactions in iii and ii could be regarded as steps in the assembly of the

pseudotrimer and the calculations show that the interaction energies are nearly additive.

Table 3. Contributions to the energies (kJ/mol) of fragment interaction in  $4[\mathbf{5a}]_2[\text{I}]_2$  calculated under partition schemes i-iii.

Partition Scheme	$E_{\text{Int}}$	$E_{\text{Elstat}}$	$E_{\text{Pauli}}$	$E_{\text{Orb}}$	$E_{\text{Disp}}$
i	-112.05	-294.18	416.15	-203.76	-30.26
ii	-55.54	-145.38	209.63	-102.26	-17.54
iii	-57.65	-147.58	208.39	-103.30	-15.16

In the case of the  $[\mathbf{5a}][\text{X}]$  ( $\text{X} = \text{Cl}, \text{Br}, \text{I}$ ) ion pairs, two geometries were considered, *cis* and *trans* to the alkylated nitrogen atom (Figure 5, iv and v). The results, summarized in Table 4, indicate that in all three cases the *trans* structure is preferred. This is consistent with the results of previous calculations of the electrostatic potential maps of the borane adducts of chalcogenadiazoles which showed that the most prominent  $\sigma$  hole on the chalcogen is opposite to the substituted nitrogen.<sup>12</sup>

Table 4. Contributions to the energy of interaction (kJ/mol) between the halide anions and  $\mathbf{5a}$  in *cis* and *trans* geometries of  $[\mathbf{5a}][\text{X}]$  ( $\text{X} = \text{Cl}, \text{Br}, \text{I}$ ) ion pairs.

Partition Scheme	X	$E_{\text{Int}}$	$E_{\text{Elstat}}$	$E_{\text{Pauli}}$	$E_{\text{Orb}}$	$E_{\text{Disp}}$
iv	Cl	-520.8	-545.8	284.0	-255.2	-3.8
	Br	-492.8	-512.0	263.2	-239.6	-4.4
	I	-461.4	-470.7	232.7	-217.7	-5.7
v	Cl	-543.1	-558.9	375.5	-357.2	-2.5
	Br	-516.1	-521.0	345.1	-337.3	-2.9
	I	-487.5	-475.8	310.8	-319.1	-3.4

The dimerization energies the  $[\mathbf{5a}][\text{X}]$  ion pairs were evaluated in two possible geometries (Figure 5, vi and vii), the results are compiled in Table 5. While dimer vi is formed from the *cis* (iv) structures and contains the  $[\text{Se-N}]_2$  supramolecular synthon, the *trans* (v) models form the  $[\text{Se X}]_2$  supramolecular synthon in dimer vii. The energy of interaction is more negative in latter case, which indicates that the structure of the dimer of the methyl cation experimentally observed in the crystal of  $[\mathbf{5a}][\text{I}]$  salt is favoured by the greater stability of the *trans* geometry of the ion pair. The results also show that the dimerization of the ion pairs  $[\mathbf{5a}][\text{I}]$  is stronger than the interaction of the ion pair with the neutral molecule of  $\mathbf{4}$ ; however, the overall sum of interactions favours the formation of the pseudo-trimer.

Table 5. Calculated contributions to the binding energies for the dimers of the  $[\mathbf{5a}][\text{X}]$  ( $\text{X} = \text{Cl}, \text{Br}, \text{I}$ ) ion pairs.

Partition Scheme	X	$E_{\text{Int}}$	$E_{\text{Elstat}}$	$E_{\text{Pauli}}$	$E_{\text{Orb}}$	$E_{\text{Disp}}$
vi	Cl	-61.2	-60.5	51.4	-39.6	-12.3
	Br	-64.3	-68.2	66.23	-48.6	-13.7
	I	-63.7	-75.2	87.1	-61.0	-14.7
vii	Cl	-57.2	-172.6	242.3	-110.7	-16.2
	Br	-65.4	265.1	-187.8	-124.8	-17.8
	I	-71.6	298.5	-206.8	-144.0	-19.3

The pseudo-trimer structures  $4[\mathbf{5a}]_2[\text{X}]_2$  ( $\text{X} = \text{I}, \text{BF}_4$ ) are especially interesting because of their two distinct  $\text{Se}\cdots\text{N}$  SBI

distances. In order to assess the relative strength of each SBI, a bond-energy decomposition analysis was performed using the *extended-transition-state* method from the *natural orbitals for chemical valence* (ETS-NOCV)<sup>51</sup>. In this method the orbital component to the total bonding energy between fragments is calculated as in Eq. 2.

$$\Delta E_{\text{Orb}} = \sum_k \Delta E_{\text{orb}}^k = \sum_{k=1}^{M/2} v_k [-F_{-k,-k}^{\text{TS}} + -F_{k,k}^{\text{TS}}] \quad (2)$$

The terms  $-F_{-k,-k}^{\text{TS}}$  are diagonal transition-state Kohn-Sham matrix elements corresponding to NOCV's with eigenvalues  $-v_k$  and  $v_k$  respectively. This particular orbital interaction term provides both a qualitative and quantitative picture of the electronic density reorganization in the  $[\text{Se-N}]_2$  supramolecular synthons. Each of the fragment interaction calculations (i-iii) shows that the shorter  $\text{Se}\cdots\text{N}$  SBI, that opposite to the quaternary nitrogen, is stronger (-13.0 to -16.9 kcal/mol) than the longer (-2.4 to -7.5 kcal/mol).

Table 6. Electron density and its Laplacian at the Bond Critical Points of the  $\text{Se}\cdots\text{X}$  and  $\text{Se}\cdots\text{N}$  supramolecular interactions in  $4[\mathbf{5a}]_2[\text{I}]_2$ .

AIM Parameter at BCP	$\text{Se}\cdots\text{N}$		$\text{Se}\cdots\text{I}$
	long	short	
$\rho$ ( $\text{e}/\text{\AA}^3 \times 10^2$ )	3.81	8.24	3.69
$\nabla^2\rho$ ( $\text{e}/\text{\AA}^3 \times 10^2$ )	2.38	6.67	4.67

The  $\text{Se}\cdots\text{N}$  interactions in  $4[\mathbf{5a}]_2[\text{I}]_2$  were also examined through Bader's theory of atoms in molecules (AIM).<sup>52</sup> Previous studies<sup>12</sup> applied AIM to the examination the Lewis acidity and basicity of  $\mathbf{1-3}$ ; their results highlighted the ambiphilic character of the chalcogenadiazoles molecules and showed that the attachment of a Lewis acid to one nitrogen of the heterocycle strengthens the  $[\text{E-N}]_2$  supramolecular synthon formed by the other nitrogen. The calculated electron density ( $\rho(r)$ ) and its Laplacian ( $\nabla^2\rho(r)$ ) at the bond critical points (BCPs) of  $4[\mathbf{5a}]_2[\text{I}]_2$  are given in Table 6. Although the difference of  $\text{Se}\cdots\text{N}$  distances is small, the AIM parameters suggest that they are very different in strength as there is indication that the energy of interaction scales linearly with the BCP density.

For comparison, the corresponding BCP parameters of the  $\text{Se}\cdots\text{I}$  interaction in the pseudo-trimer are included in Table 6. It is interesting that while the BCP electron density is smaller, its Laplacian is intermediate between the values for the  $\text{Se}\cdots\text{N}$  SBIs. The comparison was further extended by calculating the BCP parameters of the dimers of  $[\mathbf{5a}][\text{X}]$  ( $\text{X} = \text{Cl}, \text{Br}, \text{I}$ ) in the two possible conformations considered under partition schemes vi and vii. The results are compiled in Table 7. All these values suggest that the  $\text{Se}\cdots\text{X}$  and  $\text{Se}\cdots\text{N}$  interactions are of similar strength. In some cases, the calculated BCP parameters approach those of the  $\text{Te}\cdots\text{N}$  SBI in the neutral dimer of 1,2,5-telluradiazole ( $4.20 \text{ e}/\text{\AA}^3 \times 10^{-2}$  and  $10.9 \text{ e}/\text{\AA}^5 \times 10^{-2}$ )<sup>12</sup>.



Table 7. Electron density and its Laplacian at the Bond Critical Points of the Se<sup>••</sup>X interactions in the dimers of the [5a][X] (X= Cl, Br, I) ion pairs.

Structure	X	AIM Parameter at BCP	
		$\rho$ (e/Å <sup>3</sup> x 10 <sup>2</sup> )	$\nabla^2\rho$ (e/Å <sup>5</sup> x 10 <sup>2</sup> )
vi <sup>a</sup>	Cl	6.67	7.95
		1.11	2.91
	Br	5.73	5.93
		1.82	2.51
	I	4.79	4.13
		1.19	2.37
vii	Cl	3.45	8.07
	Br	3.71	8.41
	I	3.91	8.67

a. Short (top) and long (bottom) Se<sup>••</sup>X interactions.

## Conclusions

As these investigations show, the most convenient method for the preparation of N-alkyl benzoselenadiazolium cations is the condensation of selenous acid with N-substituted phenylenediamines. Such intermediates can be conveniently obtained from o-fluoronitrobenzene and primary amines. As a wide variety of aromatic diamines are commercially available or readily prepared, this approach will certainly enable the preparation of a diversity of polycations that will be useful as building blocks for supramolecular structures, including species capable of forming large supramolecular aggregates. However, the N-alkyl benzoselenadiazolium cations have a strong affinity for halide anions, therefore the anions must be replaced for non-coordinating ions in order to enable the self-assembly of the building blocks through Se<sup>••</sup>N supramolecular interactions. In contrast to the telluradiazoles, and as a great advantage, the cations are not only stable in air but also in neutral to acidic aqueous media.

## Acknowledgements

The financial support of the Natural Science Engineering Research Council of Canada (NSERC, DG-IVB, PGSD-LML) and the Ontario Graduate Scholarship program (LML) is gratefully acknowledged. Portions of this work were made possible by the facilities of the Shared Hierarchical Academic Research Computing Network (SHARCNET: www.sharcnet.ca) and Compute/Calcul Canada.

## Notes and references

- G. R. Desiraju, P. S. Ho, L. Kloo, A. C. Legon, R. Marquardt, P. Metrangolo, P. Politzer, G. Resnati and K. Rissanen, *Pure Appl. Chem.*, 2013, **85**, 1711–1713.
- P. Politzer, J. S. Murray and T. Clark, *Phys. Chem. Chem. Phys.*, 2013, **15**, 11178–11189.
- A. Priimagi, G. Cavallo, P. Metrangolo and G. Resnati, *Acc. Chem. Res.* 2013, **46**, 2686–2695.
- G. R. Desiraju, *Chem. Commun.*, 1997, 1475–1482.
- A. Nangia and G. R. Desiraju, *Top. Curr. Chem.*, 1999, **198**, 57–95.

- A. F. Cozzolino, P. J. W. Elder and I. Vargas Baca, *Coord. Chem. Rev.*, 2011, **255**, 1426–1438.
- A. F. Cozzolino, J. F. Britten and I. Vargas Baca, *Cryst. Growth Des.*, 2006, **6**, 181–186.
- A. F. Cozzolino, P. S. Whitfield and I. Vargas Baca, *J. Am. Chem. Soc.*, 2010, **132**, 17265–17270.
- A. F. Cozzolino and I. Vargas Baca, *Cryst. Growth Des.*, 2011, **11**, 668–677.
- N. A. Semenov, A. V. Lonchakov, N. A. Pushkarevsky, E. A. Suturina, V. V. Korolev, E. Lork, V. G. Vasiliev, S. N. Konchenko, J. Beckmann, N. P. Gritsan and A. V. Zibarev, *Organometallics*, 2014, **33**, 4302–4314.
- G. E. Garrett, G. L. Gibson, R. N. Straus, D. S. Seferos and M. S. Taylor, *J. Am. Chem. Soc.*, 2015, **137**, 4126–4133.
- A. F. Cozzolino, P. J. W. Elder, L. M. Lee and I. Vargas Baca, *Can. J. Chem.*, 2013, **91**, 338–347.
- P. Singh, S. Sharma, H. B. Singh and R. J. Butcher, *Proc. Natl. Acad. Sci., India, Sect. A Phys. Sci.*, 2014, **84**, 269–280.
- A. F. Cozzolino, I. Vargas Baca, S. Mansour and A. H. Mahmoudkhani, *J. Am. Chem. Soc.*, 2005, **127**, 3184–3190.
- C. J. Milios, P. V. Ioannou, C. P. Raptopoulou and G. S. Papaefstathiou, *Polyhedron*, 2009, **28**, 3199–3202.
- G. Mukherjee, P. Singh, C. Ganguri, S. Sharma, H. B. Singh, N. Goel, U. P. Singh and R. J. Butcher, *Inorg. Chem.*, 2012, **51**, 8128–8140.
- A. F. Cozzolino, A. D. Bain, S. Hanhan and I. Vargas Baca, *Chem. Commun.*, 2009, 4043–4045.
- M. Risto, R. W. Reed, C. M. Robertson, R. Oilunkaniemi, R. S. Laitinen and R. T. Oakley, *Chem. Commun.*, 2008, 3278–3280.
- A. F. Cozzolino, G. Dimopoulos-Italiano, L. M. Lee and I. Vargas Baca, *Eur. J. Inorg. Chem.*, 2013, **2013**, 2751–2756.
- A. J. Nunn and J. T. Ralph, *J. Chem. Soc.*, 1965, 6769–6777.
- A. J. Nunn and J. T. Ralph, *J. Chem. Soc., C*, 1966, 1568–1570.
- G. Berionni, B. Pegot, J. Marrot and R. Goumont, *CrystEngComm*, 2009, **11**, 986–988.
- J. L. Dutton, J. J. Tindale, M. C. Jennings and P. J. Ragogna, *Chem. Commun.*, 2006, 2474–2476.
- P. Zhang, E. A. Terefenko, J. Bray, D. Deecher, A. Fensome, J. Harrison, C. Kim, E. Koury, L. Mark, C. C. McComas, C. A. Mugford, E. J. Trybulski, A. T. Vu, G. T. Whiteside and P. E. Mahaney, *J. Med. Chem.*, 2009, **52**, 5703–5711.
- M. Chakrabarty, S. Karmakar, R. Mukherjee, S. Arima and Y. Harigaya, *Monatsh. Chem.*, 2008, **140**, 375–380.
- Y. Jahng and M. A. F. M. Rahman, *Synth. Commun.*, 2006, **36**, 1213–1220.
- I. W. Harvey, M. D. McFarlane, D. J. Moody and D. M. Smith, *J. Chem. Soc., Perkin Trans. 1*, 1988, 1939–1943.
- G. te Velde and F. M. Bickelhaupt, *J. Comp. Chem.* 2001 **22**, 931–967.
- C. Fonseca Guerra, J. G. Snijders, G. te Velde and E. J. Baerends, *Theo. Chem. Acc.* 1998, **99**, 391–403.
- S. Van Gisbergen, J. Snijders and E. Baerends, *Phys. Rev. Lett.*, 1997, **78**, 3097–3100.
- S. J. A. van Gisbergen, J. G. Snijders and E. J. Baerends, *J. Chem. Phys.*, 1998, **109**, 10644–10656.
- S. H. Vosko, L. Wilk and M. Nusair, *Can. J. Phys.*, 1980, **58**, 1200–1211.
- J. P. Perdew, K. Burke and M. Ernzerhof, *Phys. Rev. Lett.*, 1996, **77**, 3865–3868.
- E. Van Lenthe and E. Baerends, *J. Chem. Phys.*, 1993, **99**, 4597–4610.
- E. Van Lenthe, E. J. Baerends and J. G. Snijders, *J. Chem. Phys.*, 1994, **101**, 9783–9792.
- E. van Lenthe, A. Ehlers and E.-J. Baerends, *J. Chem. Phys.*,

- 1999, **110**, 8943.
- 37 E. Van Lenthe, J. G. Snijders and E. J. Baerends, *J. Chem. Phys.*, 1996, **105**, 6505.
- 38 E. Van Lenthe and R. Van Leeuwen, *Int. J. Quant. Chem.*, 1996, **57**, 281-293.
- 39 S. Grimme, J. Antony, S. Ehrlich and H. Krieg, *J. Chem. Phys.*, 2010, **132**, 154104.
- 40 L. Fan and T. Ziegler, *J. Chem. Phys.*, 1992, **96**, 9005-9012.
- 41 L. Fan and T. Ziegler, *J. Phys. Chem.*, 1992, **96**, 6937-6941.
- 42 O. V. Gritsenko, P. R. T. Schipper and E. J. Baerends, *Chem. Phys. Lett.*, 1999, **302**, 199-207.
- 43 O. V. Gritsenko, P. R. T. Schipper and E. J. Baerends, *Int. J. Quant. Chem.*, 2000, **76**, 407-419.
- 44 P. Schipper and O. V. Gritsenko, *J. Chem Phys.*, 2000, **112**, 13, 1344-1352.
- 45 L. M. Lee, P. J. W. Elder, A. F. Cozzolino, Q. Yang and I. Vargas Baca, *Main Group Chem.*, 2010, **9**, 117-133.
- 46 G. I. Ereemeeva, B. K. Strelets and L. S. Efros, *Khim. Geterotsikl. Soedin.*, 1975, 276-277.
- 47 G. I. Ereemeeva, Y. I. Akulin, T. N. Timofeeva, B. K. Strelets and L. S. Efros, *Khim. Geterotsikl. Soedin.*, 1982, 1129-1130.
- 48 L. Andrews, E. S. Prochaska and A. Loewenschuss, *Inorg. Chem.*, 1980, **19**, 463-465.
- 49 J. L. Dutton and P. J. Ragoon, *Inorg. Chem.*, 2009, **48**, 1722-1730.
- 50 T. Ziegler and A. Rauk, *Theo. Chem. Acc.*, 1977, **46**, 1-10.
- 51 M. P. Mitoraj, A. Michalak and T. Ziegler, *J. Chem. Theo. Comput.*, 2009, **5**, 962-975.
- 52 R. F. W. Bader, *Acc. Chem. Res.*, 1985, **18**, 9-15.

## Synthetic, Structural, and Computational Investigations of N-alkyl Benzo-2,1,3-Selenadiazolium Iodides and their Supramolecular Aggregates

Lucia M. Lee, Victoria B. Corless, Michael Tran, Hilary Jenkins, James F. Britten and Ignacio Vargas-Baca\*

### Graphical Abstract

Alkyl benzo-2,1,3-selenadiazolium cations are conveniently prepared by cyclo-condensation of N-alkyl-phenylenediamines with selenous acid

