

# Dalton Transactions

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



## White phosphorus activation by a Th(III) complex†

Alasdair Formanuk,<sup>a</sup> Fabrizio Ortu,<sup>a</sup> Reece Beekmeyer,<sup>b</sup> Andrew Kerridge,<sup>c</sup> Ralph W. Adams<sup>a</sup> and David P. Mills<sup>a,\*</sup>

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

**[Th(Cp'')<sub>3</sub>] (Cp'' = {C<sub>5</sub>H<sub>3</sub>(SiMe<sub>3</sub>)<sub>2</sub>-1,3}) activates P<sub>4</sub> to give [{Th(Cp'')<sub>3</sub>]<sub>2</sub>(μ-η<sup>1</sup>:η<sup>1</sup>-P<sub>4</sub>)] (1), which has an unprecedented cyclo-P<sub>4</sub> binding mode. DFT studies were performed on a model of 1 to probe the bonding in this system.**

The inherent reactivity of low oxidation state early metal complexes has been widely exploited in small molecule activation chemistry for the generation of useful chemicals from abundant feedstocks (e.g. CO, CO<sub>2</sub>, N<sub>2</sub>).<sup>1</sup> In this field, white phosphorus activation by early d-transition metal complexes has been extensively investigated<sup>2</sup> as the direct preparation of organophosphorus reagents from elemental phosphorus would skip the protracted chlorination and derivatisation steps used industrially.<sup>3,4</sup> Whilst U(III) small molecule activation chemistry has also started to flourish,<sup>5</sup> the analogous Th(III) chemistry currently lags behind, which can be attributed to the paucity of Th(III) complexes.<sup>6,7</sup> Th(III) chemistry is difficult to access as there are relatively large standard reduction potentials to overcome [E<sup>Θ</sup> Th(IV) → Th(III) −3.7 V; cf. U(IV) → U(III) −0.6 V],<sup>8</sup> and there are no synthetically useful Th(III)X<sub>3</sub> (X = halide) starting materials for salt metathesis methodologies, whereas synthetic routes to U(III) complexes from U<sub>3</sub> are well-developed.<sup>9</sup>


The reactivity of Th(III) complexes has previously been limited to the serendipitous reaction of Lappert's [Th(Cp'')<sub>3</sub>] with trace H<sub>2</sub>O/O<sub>2</sub> to give [Th(Cp'')<sub>3</sub>]<sub>2</sub>(μ-O),<sup>6c</sup> its purposeful oxidation with Bu<sup>+</sup>Cl to afford [Th(Cp'')<sub>3</sub>(Cl)],<sup>6c,10</sup> the reaction of [Th{C<sub>5</sub>Me<sub>4</sub>H}<sub>3</sub>] with TEMPO (2,2,6,6-tetramethyl-piperidin-1-oxyl radical) to give [Th{C<sub>5</sub>Me<sub>4</sub>H}<sub>3</sub>(TEMPO)] reported by Evans,<sup>11</sup> and Cloke's account of CO<sub>2</sub> activation with a postulated Th(III) complex generated *in situ* using Na/K alloy to form carbonates and oxalates.<sup>12</sup> Germane to these studies, the

small molecule activation chemistry of "highly reduced" thorium synthons, which formally contain Th(IV) centres and reduced ligands, has been investigated more extensively.<sup>13</sup>

There are currently only several examples of U(III)-mediated white phosphorus activation,<sup>14</sup> with Cloke's account of a "slipped" cyclo-P<sub>4</sub> dianion in [U(C<sub>5</sub>Me<sub>5</sub>)[C<sub>8</sub>H<sub>4</sub>(SiPr<sup>i</sup>)<sub>2</sub>-1,4]]<sub>2</sub>(μ-η<sup>2</sup>:η<sup>2</sup>-P<sub>4</sub>) of most relevance here.<sup>14b</sup> For Th(IV) chemistry, Scherer has reported the reaction of [Th(C<sub>5</sub>H<sub>3</sub>Bu<sup>t</sup><sub>2</sub>-1,3)<sub>2</sub>(η<sup>4</sup>-C<sub>4</sub>H<sub>6</sub>)] with white phosphorus in refluxing toluene to give [Th(C<sub>5</sub>H<sub>3</sub>Bu<sup>t</sup><sub>2</sub>-1,3)<sub>2</sub>]<sub>2</sub>(μ-η<sup>3</sup>:η<sup>3</sup>-P<sub>6</sub>), with [Th(C<sub>5</sub>H<sub>3</sub>Bu<sup>t</sup><sub>2</sub>-1,3)<sub>2</sub>](μ-η<sup>3</sup>:η<sup>3</sup>-P<sub>3</sub>)[Th(C<sub>5</sub>H<sub>3</sub>Bu<sup>t</sup><sub>2</sub>-1,3)(Cl)] isolated when half an equivalent of MgCl<sub>2</sub> was added to the reaction mixture.<sup>15</sup>

The reaction of [Th(Cp'')<sub>3</sub>]<sup>6a</sup> with P<sub>4</sub> gives [Th(Cp'')<sub>3</sub>]<sub>2</sub>(μ-η<sup>1</sup>:η<sup>1</sup>-P<sub>4</sub>) (1) in poor yield as the only isolable product in either a 1:0.5 or 1:0.25 stoichiometry, with the yield approximately halved when P<sub>4</sub> is deficient (Scheme 1). The <sup>31</sup>P{<sup>1</sup>H} NMR spectra of the reaction mixtures showed that all almost P<sub>4</sub> was consumed, whereas we found that [U(Cp'')<sub>3</sub>]<sup>16</sup> does not react with P<sub>4</sub>, even under forcing conditions (16 hours reflux). To the best of our knowledge this is the first example of divergent reactivity profiles of isolated, structurally analogous Th(III) and U(III) complexes. This phenomenon can be attributed to differences in reduction potentials and electronic configurations of Th(III) and U(III) in [An(Cp'')<sub>3</sub>] (see below). It is noteworthy that a U(III) complex with a similar π-donor Cp ligand environment, [U(C<sub>5</sub>Me<sub>5</sub>)[C<sub>8</sub>H<sub>4</sub>(SiPr<sup>i</sup>)<sub>2</sub>-1,4]], has previously been shown to activate P<sub>4</sub>,<sup>14b</sup> and the reduction potential of P<sub>4</sub> is reported at −1.53 and −1.98 V, depending on the conditions employed.<sup>17</sup>

The <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of 1 in d<sub>6</sub>-benzene at 298 K exhibits two triplet signals [δ<sub>p</sub>: 227.59 (t, <sup>1</sup>J<sub>pp</sub> ≈ 400 Hz) and 328.86 ppm (t, <sup>1</sup>J<sub>pp</sub> ≈ 400 Hz)] and two broad unresolved multiplets [δ<sub>p</sub>: −246.55 (br m, Δ<sub>v</sub> ≈ 2000 Hz) and 10.35 ppm (br



Scheme 1. Synthesis of complex 1.

<sup>a</sup> School of Chemistry, The University of Manchester, Oxford Road, Manchester, M13 9PL, UK; E-mail: david.mills@manchester.ac.uk.

<sup>b</sup> Department of Chemistry, University College London, 20 Gordon Street, London, WC1H 0AJ, UK.

<sup>c</sup> Department of Chemistry, Lancaster University, Lancaster, LA1 4YB, UK.

† Electronic supplementary information (ESI) available: Full synthetic details, crystallography and DFT studies. CCDC 1426081. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/C5DT04528B. Additional research data supporting this publication are available from The University of Manchester eScholar repository at [insert DOI here when proofs available].


## COMMUNICATION

## Journal Name

m,  $\Delta_\nu \approx 2000$  Hz)]. The multiplicity for the two low field signals correlates with the solid state structure of **1**, by comparison with the chemical shift of the signal in the  $^{31}\text{P}$  NMR spectrum of  $\text{Cs}_2\text{P}_4$  [ $\delta_\text{P}$ : 348 ppm (s)].<sup>18</sup> The two high field signals are likely due to processes involving changes in hapticity of the cyclo- $\text{P}_4$  ring, but could not be confidently assigned. These were reproducibly observed in separate samples of **1** but elemental analysis values obtained were in close agreement with the solid state structure (see below). To probe the identity of the unassigned signals, VT  $^{31}\text{P}\{^1\text{H}\}$  NMR spectroscopy of **1** in toluene/ $d_8$ -toluene was performed from 193–353 K in 10 K increments (see ESI Figures S1–S2). At high temperatures (> 313 K) the two low field signals were not distinguishable from the baseline and the two high field multiplets simplified to broad signals. At low temperatures (< 233 K) the two low field signals resolved to two pairs of triplets [Data at 193 K,  $\delta_\text{P}$  = 216 (t,  $^1J_{\text{PP}}$  = 411 Hz), 220 (t,  $^1J_{\text{PP}}$  = 411 Hz), 317 (t,  $^1J_{\text{PP}}$  = 411 Hz), and 323 (t,  $^1J_{\text{PP}}$  = 411 Hz)]. We tentatively attribute this to restricted rotation about the Th...Th axis; the coalescence temperature for these signals is around 250 K (see ESI Figure S3). The two high field signals did not resolve sufficiently at this temperature for coupling constants to be extracted.  $^{31}\text{P}\{^1\text{H}\}$  NMR COSY experiments were performed in  $d_6$ -benzene at 298 K to determine if all four signals derived from the same molecule, but this was inconclusive despite extended acquisition times.

Only one singlet is observed in the  $^{29}\text{Si}\{^1\text{H}\}$  NMR spectrum of **1** at 298 K [ $\delta_\text{Si}$ : -8.02 ppm], showing that a pseudo-high symmetry environment of the silyl groups is adopted in solution at room temperature. The  $^1\text{H}$  and  $^{13}\text{C}\{^1\text{H}\}$  NMR spectra of **1** were unremarkable and two signals were observed for the silyl groups in each spectrum [ $\delta_\text{H}$ : 0.58 and 0.60 ppm;  $\delta_\text{C}$ : 2.66 and 2.86 ppm]. The UV/Vis spectrum of **1** exhibits three intense absorptions (see ESI Figure S6). The most intense absorption at 285 nm ( $\epsilon$  = 30,400  $\text{M}^{-1} \text{cm}^{-1}$ ) is assigned as the cyclo- $\text{P}_4$   $\pi \rightarrow \pi^*$  transition and the broad absorptions at 425 nm (9,000  $\text{M}^{-1} \text{cm}^{-1}$ ) and 477 nm (11,200  $\text{M}^{-1} \text{cm}^{-1}$ ), assigned as LMCT bands based on the similarity of this spectrum with closely related “slipped” cyclo- $\text{P}_4$  transition metal systems.<sup>1,19</sup> The spectral data for **1** are typical of a diamagnetic formulation, with two Th(IV) centres and a cyclo- $\text{P}_4^{2-}$   $6\pi$ -electron aromatic bridging dianion.

The solid state structure of **1** ( $\text{C}_7\text{H}_8$ ) was determined by single crystal XRD (Figure 1). Complex **1** adopts the cubic space group  $P\bar{a}3$ , with a crystallographic threefold rotation axis through the Th(1)–P(1) bond and the centre of the cyclo- $\text{P}_4$  ring. The two phosphorus atoms that are not bonded to thorium are each located over three symmetry-equivalent positions set at 1/3 occupancy, therefore only one conformation is shown for clarity. Complex **1** exhibits a  $\mu\text{-}\eta^1\text{:}\eta^1\text{-P}_4$  binding mode, which we attribute to the presence of three bulky Cp\* groups at each thorium centre. Steric effects have previously accounted for the “slipped” cyclo- $\text{P}_4$  binding mode in  $[\{\text{U}(\text{C}_5\text{Me}_5)[\text{C}_8\text{H}_4(\text{SiPr}^t_3)_2\text{-1,4}\}]_2(\mu\text{-}\eta^2\text{:}\eta^2\text{-P}_4)]$ .<sup>14b</sup> The Th–P distances in **1** [2.919(4) Å] are unremarkable, being similar to those observed in  $[\{\text{Th}(\text{C}_5\text{H}_3\text{Bu}^t\text{-1,3})_2(\mu\text{-}\eta^3\text{:}\eta^3\text{-P}_3)\{\text{Th}(\text{C}_5\text{H}_3\text{Bu}^t\text{-1,3})_2(\mu\text{-}\eta^3\text{:}\eta^3\text{-P}_3)\}]_2$  [2.913(7) Å mean].<sup>15</sup> The Th...Cp<sub>centroid</sub> distances in **1** [2.543(5) Å] are longer than those in  $[\text{Th}(\text{Cp}^*)_3]$  [2.519(2) Å mean]<sup>6a,7</sup> and are more similar to those reported for  $[\text{Th}(\text{Cp}^*)_3(\text{Cl})]$  [2.565 Å mean],<sup>20</sup> in agreement with a formal Th(IV) assignment in **1**.



**Figure 1.** Molecular structure of **1** ( $\text{C}_7\text{H}_8$ ) with selected atom labelling and displacement ellipsoids set at 30 % probability level. Th(1) and P(1) are on special positions, hence no equator quadratics could be generated. Hydrogen atoms, phosphorous and silyl group disorder components and lattice toluene have been removed for clarity. Symmetry operation to generate equivalent atoms:  $i = -x, -y, -z$ . Selected bond distances [Å] and angles [°]: Th(1)–P(1) 2.919(4), P(1)–P(2) 2.051(9), Th(1)–Cp<sub>centroid</sub> 2.543(5), Th(1)–P(1)–P(2) 133.0(2), P(1)–P(2)–P(1) 85.9(4), P(2)–P(1)–P(2) 94.1(4), Cp<sub>centroid</sub>...Th(1)–Cp<sub>centroid</sub> 119.1(2).

The Th...Cp<sub>centroid</sub> distances in **1** [2.543(5) Å] are longer than those in  $[\text{Th}(\text{Cp}^*)_3]$  [2.519(2) Å mean]<sup>6a,7</sup> and are more similar to those reported for  $[\text{Th}(\text{Cp}^*)_3(\text{Cl})]$  [2.565 Å mean],<sup>20</sup> in agreement with a formal Th(IV) assignment in **1**.

The cyclo- $\text{P}_4$  rhombus in **1** is planar by definition, with the endocyclic P–P angles of 86.4(5) and 93.6(5)° skewed significantly from the approximate square arrangement observed in  $[\{\text{U}(\text{C}_5\text{Me}_5)[\text{C}_8\text{H}_4(\text{SiPr}^t_3)_2\text{-1,4}\}]_2(\mu\text{-}\eta^2\text{:}\eta^2\text{-P}_4)]$ .<sup>14b</sup> The P–P distances in **1** [2.051(9) Å mean] are remarkably short compared with those in  $[\{\text{U}(\text{C}_5\text{Me}_5)[\text{C}_8\text{H}_4(\text{SiPr}^t_3)_2\text{-1,4}\}]_2(\mu\text{-}\eta^2\text{:}\eta^2\text{-P}_4)]$  [2.150(2) Å mean]<sup>14b</sup> and  $[\{\text{Th}(\text{C}_5\text{H}_3\text{Bu}^t\text{-1,3})_2(\mu\text{-}\eta^3\text{:}\eta^3\text{-P}_3)\{\text{Th}(\text{C}_5\text{H}_3\text{Bu}^t\text{-1,3})_2(\mu\text{-}\eta^3\text{:}\eta^3\text{-P}_3)\}]_2$  [2.185(9) Å mean],<sup>15</sup> being more similar to a P=P double bond length (2.04 Å).<sup>21</sup> This observation can be attributed to the unusual  $\mu\text{-}\eta^1\text{:}\eta^1$ -binding mode of the cyclo- $\text{P}_4$  unit in **1**. In this orientation  $\sigma$ -donation of phosphorus  $\text{sp}^2$ -hybridised lone pairs to vacant thorium orbitals should be the major component of the Th–P bonds, and due to the reduced  $\pi$ -contribution to these bonds most of the  $\pi$ -electron density is retained in the cyclo- $\text{P}_4$  ring.

A full model of **1** was studied by DFT using the TURBOMOLE (v6.6) quantum chemistry package with the hybrid-GGA exchange-correlation functional PBE0. A gas-phase geometry optimisation was followed by vibrational frequency analysis and a local energy minimum was confirmed. Calculated Th...Cp<sub>centroid</sub> distances (2.540 Å) were in excellent agreement with those calculated experimentally; Th–P bonds (2.964 Å) were in reasonable agreement, but P–P bonds (2.138 Å) were overestimated. This discrepancy was due to a small deviation from planarity in the Th– $\text{P}_4$ –Th unit, hence bonding analysis was performed on both PBE0-optimised and crystallographic structures. Interaction with the thorium centres manifests itself in splitting of the high-lying  $\sigma_1$ - and  $\pi_1$ -levels that are formally degenerate in the free  $\text{P}_4^{2-}$  dianion, with the  $\sigma_1$ -level split by 1.11 eV/0.52 eV at the experimental/PBE0-optimised geometry and the corresponding  $\pi_1$ -level split by 0.21 eV/0.14 eV, demonstrating the pronounced  $\sigma$ - and diminished  $\pi$ -contributions (Figure 2).




Figure 2.  $\sigma$ - and  $\pi$ -bonding orbitals of **1**, rendered at an isosurface of 0.015 a.u.

The Quantum Theory of Atoms in Molecules (QTAIM) was employed to further investigate the bonding in **1**. Parameters obtained at experimental and PBE0-optimised geometries were very similar (see ESI), so only the former are considered here. The calculated magnitude of the electron density at the bond critical point ( $\rho_{\text{BCP}}$ ) of the Th-P bond is 0.047 a.u., indicative of a predominantly ionic interaction with slightly greater covalent character than that found in Th-S and Th-Se single bonds.<sup>22</sup> Similar characteristics were found for the energy density,  $H$ , and the Th-P delocalisation index,  $\delta(\text{Th},\text{P})$ . Within the cyclo-P<sub>4</sub> ring, the P-P delocalisation index,  $\delta(\text{P},\text{P})$ , which in this situation can be regarded as a measure of bond order, was found to be 1.33, close to the formal value of 1.50 expected for an ideal aromatic system and slightly lower than the value of 1.45 calculated for free P<sub>4</sub><sup>2-</sup> at the same level of theory. This is indicative of a largely unperturbed aromatic  $6\pi$ -electron configuration. The calculated partial charge on the ring is -1.27 a.u., indicating a degree of charge transfer from the ring that is commensurate with the slight reduction in aromatic character.

We conclude that the enhanced reactivity of [Th(Cp'')<sub>3</sub>] over [U(Cp'')<sub>3</sub>] with P<sub>4</sub> necessitates future investigations into the reactivity of Th(III) complexes. We are especially interested in Th(III)-mediated small molecule activation chemistry, as these investigations could give novel reactivity profiles and bonding motifs. We thank The University of Manchester, the EPSRC (grant numbers EP/K039547/1, EP/L014416/1 and EP/J002208/1) and the Nuclear FIRST DTC for funding. AK thanks Lancaster University for access to the HEC high performance computing facility.

## Notes and references

- C. C. Cummins, *Prog. Inorg. Chem.*, 1998, **47**, 685.
- B. M. Cossairt, N. A. Piro and C. C. Cummins, *Chem. Rev.*, 2010, **110**, 4164.
- D. Corbridge, *Phosphorus: An Outline of its Chemistry, Biochemistry and Technology*, 5<sup>th</sup> ed., Elsevier, New York, 1994.
- R. Engel, *Synthesis of Carbon Phosphorus Bonds*, 2<sup>nd</sup> ed., CRC Press, Boca Raton, 2004.
- (a) S. T. Liddle, *Angew. Chem. Int. Ed.*, 2015, **54**, 8604; (b) H. S. La Pierre and K. Meyer, *Prog. Inorg. Chem.*, 2014, **58**, 303; (c) B. M. Gardner and S. T. Liddle, *Eur. J. Inorg. Chem.*, 2013, 3753; (d) O. P. Lam and K. Meyer, *Polyhedron*, 2012, **32**, 1; (e) P. L. Arnold, *Chem. Commun.*, 2011, **47**, 9005; (f) O. T. Summerscales and F. G. N. Cloke, *Struct. Bond.*, 2008, **127**, 87; (g) I. Castro-Rodríguez and K. Meyer, *Chem. Commun.*, 2006, 1353; (h) W. J. Evans and S. A. Kozimor, *Coord. Chem. Rev.*, 2006, **250**, 911; (i) M. Ephritikhine, *Dalton Trans.*, 2006, 2501; (j) I. Korobkov and S. Gambarotta, *Prog. Inorg. Chem.*, 2005, **54**, 321.
- (a) P. C. Blake, M. F. Lappert, J. L. Atwood and H. Zhang, *J. Chem. Soc., Chem. Commun.*, 1986, 1148; (b) J. S. Parry, F. G. N. Cloke, S. J. Coles and M. B. Hursthouse, *J. Am. Chem. Soc.*, 1999, **121**, 6867; (c) P. C. Blake, N. M. Edelstein, P. B. Hitchcock, W. K. Kot, M. F. Lappert, G. V. Shalimoff and S. Tian, *J. Organomet. Chem.*, 2001, **636**, 124; (d) J. R. Walensky, R. L. Martin, J. W. Ziller and W. J. Evans, *Inorg. Chem.*, 2010, **49**, 10007; (e) N. A. Siladke, C. L. Webster, J. R. Walensky, M. K. Takase, J. W. Ziller, D. J. Grant, L. Gagliardi and W. J. Evans, *Organometallics*, 2013, **32**, 6522.
- Th(II) complexes have recently been reported: R. R. Langeslay, M. E. Feiser, J. W. Ziller, F. Furche and W. J. Evans, *Chem. Sci.*, 2015, **6**, 517.
- L. J. Nugent, R. D. Baybarz, J. L. Burnett and J. L. Ryan, *J. Phys. Chem.*, 1973, **77**, 1528.
- R. J. Baker, *Coord. Chem. Rev.*, 2012, **256**, 2843.
- P. C. Blake, M. F. Lappert, R. G. Taylor, J. L. Atwood and H. Zhang, *Inorg. Chim. Acta*, 1987, **139**, 13.
- R. R. Langeslay, J. R. Walensky, J. W. Ziller and W. J. Evans, *Inorg. Chem.*, 2014, **53**, 8455.
- Z. E. Button, J. A. Higgins, M. Suvova, F. G. N. Cloke and S. M. Roe, *Dalton Trans.*, 2015, **44**, 2588.
- See, for example: (a) I. Korobkov, S. Gambarotta, G. P. A. Yap, *Angew. Chem. Int. Ed.*, 2003, **42**, 814; (b) I. Korobkov, S. Gambarotta, G. P. A. Yap, *Angew. Chem. Int. Ed.*, 2003, **42**, 4958; (c) A. Arunachalampillai, P. Crewdson, I. Korobkov and S. Gambarotta, *Organometallics*, 2006, **25**, 3856; (d) I. Korobkov, B. Vidjayacoumar, S. I. Gorelsky, P. Billone and S. Gambarotta, *Organometallics*, 2010, **29**, 692; (e) E. J. Schelter, R. Wu, B. L. Scott, J. D. Thompson, T. Cantat, K. D. John, E. R. Batista, D. E. Morris and J. L. Kiplinger, *Inorg. Chem.*, 2010, **49**, 924; (f) A. Mrutu, C. L. Barnes, S. C. Bart and J. R. Walensky, *Eur. J. Inorg. Chem.*, 2013, 4050; (g) W. Ren, H. Song, G. Zi and M. D. Walter, *Dalton Trans.*, 2012, **41**, 5965; (h) W. Ren, G. Zi and M. D. Walter, *Organometallics*, 2012, **31**, 672; (i) W. Ren, W. W. Lukens, G. Zi, L. Maron and M. D. Walter, *Chem. Sci.*, 2013, **4**, 1168; (j) B. Fang, W. Ren, G. Hou, G. Zi, D.-C. Fang, L. Maron and M. D. Walter, *J. Am. Chem. Soc.*, 2014, **136**, 17249; (k) B. Fang, G. Hou, G. Zi, D.-C. Fang and M. D. Walter, *Dalton Trans.*, 2015, **44**, 7927.
- (a) F. H. Stephens, Ph.D. Thesis, Massachusetts Institute of Technology, Cambridge, 2004; (b) A. S. P. Frey, F. G. N. Cloke, P. B. Hitchcock and J. C. Green, *New J. Chem.*, 2011, **35**, 2022; (c) D. Patel, F. Tuna, E. J. L. McInnes, W. Lewis, A. J. Blake and S. T. Liddle, *Angew. Chem. Int. Ed.*, 2013, **52**, 13334; (d) B. M. Gardner, F. Tuna, E. J. L. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Angew. Chem. Int. Ed.*, 2015, **54**, 7068.
- O. J. Scherer, B. Werner, G. Heckmann and G. Wolmershäuser, *Angew. Chem. Int. Ed.*, 1991, **30**, 553.
- M. del Mar Conejo, J. S. Parry, E. Carmona, M. Schultz, J. G. Brennan, S. M. Beshouri, R. A. Andersen, R. D. Rogers, S. Coles and M. B. Hursthouse, *Chem. Eur. J.*, 1999, **5**, 3000.
- I. M. Osadchenko and A. P. Tomilov, *Russ. J. Appl. Chem.*, 2006, **79**, 2033.
- F. Kraus, J. C. Aschenbrenner and N. Korber, *Angew. Chem. Int. Ed.*, 2003, **42**, 4030.
- C. Camp, L. Maron, R. G. Bergman and J. Arnold, *J. Am. Chem. Soc.*, 2014, **136**, 17652.
- P. C. Blake, M. A. Edelman, P. B. Hitchcock, J. Hu, M. F. Lappert, S. Tian, G. Müller, J. L. Atwood and H. Zhang, *J. Organomet. Chem.*, 1998, **551**, 261.
- P. Pyykkö and M. Atsumi, *Chem. Eur. J.*, 2009, **15**, 12770.

## COMMUNICATION

Journal Name

- 22 (a) M. B. Jones, A. J. Gaunt, J. C. Gordon, N. Kaltsoyannis, M. O. Neu and B. L. Scott, *Chem. Sci.* 2013, **4**, 1189; (b) A. C. Behrle, C. L. Barnes, N. Kaltsoyannis and J. R. Walensky, *Inorg. Chem.*, 2013, **52**, 10623; (c) A. C. Behrle, A. Kerridge and J. R. Walensky, *Inorg. Chem.*, 2015, **54**, 11625.

## Table of contents

Lappert's original Th(III) complex,  $[\text{Th}\{\text{C}_5\text{H}_3(\text{SiMe}_3)_{2-1,3}\}_3]$ , reduces white phosphorus to give a cyclo- $\text{P}_4$  dianion, which exhibits an unprecedented  $\mu\text{-}\eta^1:\eta^1\text{-}$  binding mode in the dinuclear Th(IV) product.

