

Green Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

ARTICLE

A new class of antimicrobial biosurfactants: quaternary ammonium sophorolipids

Cite this: DOI: 10.1039/x0xx00000x

E. I. P. Delbeke,^a B. I. Roman,^a G. B. Marin,^b K. M. Van Geem,^b and C. V. Stevens^aReceived 00th January 2012,
Accepted 00th January 2012


DOI: 10.1039/x0xx00000x

www.rsc.org/

New synthetic pathways are proposed for the production of a broad range of innovative sophorolipid amines and sophorolipid quaternary ammonium salts starting from microbially produced sophorolipids. The selective formation of an intermediate sophorolipid aldehyde proved to be a key synthetic step of the new derivatives. The sophorolipid quaternary ammonium salts were evaluated for their antimicrobial activity against Gram-negative and Gram-positive test strains. Minimum inhibitory concentration (MIC) values were determined for the active compounds. Derivatives with an octadecyl group on the nitrogen atom proved to be more active than the antibiotic gentamicin sulfate against all tested Gram-positive strains. The results show great promise for modified sophorolipids in the medical sector.

Introduction

In our modern society, the concept of sustainability gains more interest from day to day.¹ Indeed, over the last decades, the transition towards a bio-based economy has initiated. The significance of renewable resources for the chemical industry is gradually increasing because they are considered good alternatives for fossil resources, whose supplies are limited and which have a major environmental impact.² At present already more than 8% of all the chemicals produced in Europe are based on renewable resources.³ Most of these renewable resources are directly used as biomaterials such as paper, plastics, textile and fibres or as biochemicals in paint, detergents, and cosmetics. However, the incorporation of renewable resources in high value added products is mostly based on the same concept as the one adopted for fossil resources, i.e. starting from small hydrocarbon building blocks. Traditionally, fossil and renewable resources are broken down into base chemicals which are then used for the synthesis of the desired products. Using this methodology, the cost of renewable based pharmaceuticals outweighs the cost of their fossil counterparts, giving renewable resources a competitive disadvantage for the synthesis of these high value products. Obviously the business case would be far more interesting if renewable resources with a complex structure would be directly used as building blocks in a synthetic pathway. This will reduce the number of required steps to obtain the desired compounds and will additionally contribute to the sustainability of both process and end product. Target compounds for this 'green synthesis' approach should preferably possess the desired



Scheme 1. Microbial produced sophorolipid lactone **1** and sophorolipid acid **2**


properties or biological activities which the traditionally obtained competing molecules lack.

In this light, sophorolipids are very useful building blocks for chemical derivatization. They occur in a lactonic form or an open acid form (Scheme 1). These glycolipids are produced by micro-organisms from renewable resources through fermentation.⁴ The yeast *Starmerella bombicola* is the preferred producer, with output of around 400 g/L and a production price of 2 to 5 €/kg.^{4a} Due to the presence of a hydrophilic carbohydrate head and a hydrophobic lipid tail, sophorolipids possess surface-active properties classifying them as biosurfactants. They are low foaming surfactants which can be used for hard surface cleaning and automatic dishwashing rinse aid applications.⁵ Nevertheless, application of sophorolipids in the detergent sector is limited because they have a competitive disadvantage compared to synthetic surfactants in terms of production cost. Therefore, it is desirable to look for other application areas which are economically more attractive, e.g. in the medical sector. Sophorolipids feature beneficial biological activities such as anticancer, anti-HIV, sperm-immobilizing and antibacterial activity.⁶ Recently, the

self-assembly properties of sophorolipids have been described, demonstrating the formation of nanostructures with supramolecular chirality.⁷ Until now, sophorolipid production is restricted to only a few derivatives. Consequently, the variation in their surface-active properties is limited and their biological activities have not yet been optimized. Chemical modification offers the opportunity to extend the limited set of microbial derivatives. In this paper, chemical modification pathways are evaluated which increase the applicability of sophorolipids for medical applications.

At present, chemical and enzymatic modification of sophorolipids have been mostly limited to the sugar head or the lipid tail.⁸ Cleavage of the double bond, however, has only been described for the synthesis of short-chained sophorolipid acids or in ring-opening cross-metathesis reactions.⁹ It was never included in a synthetic pathway for the production of a functionalized building block towards modified derivatives. This is surprising because such modification would result in the formation of sophorolipid analogues with shorter chain lengths, which would be better soluble in water.

Therefore in this work, a series of chemical modification pathways is described for sophorolipids towards quaternary ammonium salt derivatives (Scheme 2). First, an ozonolysis reaction is performed on sophorolipids in order to transform the double bond into a reactive site. In a second key step, a nitrogen functionality is incorporated. Further modification leads to a library of sophorolipid quaternary ammonium salts, which are evaluated as antimicrobial compounds.




Scheme 2. Retrosynthetic scheme for sophorolipid quaternary ammonium salts **3**

Results and discussion

Synthesis of the sophorolipid aldehyde intermediate

A chemical modification pathway for sophorolipids was developed starting from the major microbial product, i.e. the diacetylated sophorolipid lactone **1**.¹⁰ This lactone was transformed into sophorolipid methyl ester **5** and, subsequently, into peracetylated analogue **6** according to literature procedures.^{8c, 8d} Cleavage of the double bond through ozonolysis, offering an aldehyde functionality, would create opportunities for the incorporation of nitrogen. These nitrogen containing derivatives can then easily be transformed into cationic surfactants.

The double bond was cleaved through ozonolysis in MeOH, and reductive work-up with NaBH(OAc)₃ furnished the peracetylated sophorolipid aldehyde **7** and methyl 9-oxononanoate side product **8** (Scheme 3).¹¹ At first, ozonolysis was attempted in CH₂Cl₂, followed by reductive work-up with dimethyl sulfide. Analysis of




Scheme 3. Chemical modification towards sophorolipid aldehyde **7**

the reaction mixture demonstrated the presence of the desired peracetylated sophorolipid aldehyde **7**, but methyl 9-oxononanoate **8** could not be detected. Instead, only the presence of a stable 1,2,4-trioxolane (ozonide) intermediate was observed. The presence of high concentrations of residual ozonides after work-up with dimethyl sulfide has already been described by Dussault *et al.*¹² Therefore, reaction conditions were changed towards the use of the more environmental friendly NaBH(OAc)₃. Besides, isolation of methyl 9-oxononanoate **8** is most desirable for a green synthetic pathway. This side product could be used as a valuable building block for other applications. Direct ozonolysis of the diacetylated sophorolipid lactone **1** was also evaluated. An intermediate dialdehyde was obtained after ozonolysis, but transesterification to the desired sophorolipid aldehyde was not possible. Analysis of the reaction mixture demonstrated the presence of aldol condensation product **9** formed from methyl 9-oxononanoate indicating the non-compatibility of the aldehyde function with alkaline reaction conditions (Scheme 4).

Care should be taken during the ozonolysis reaction due to the formation of unstable ozonides and peroxides, and the use of methanol as a solvent. The reaction is performed in a taped washing

Scheme 4. Observed aldol condensation product **9**



flask on a small scale of 10-15 g peracetylated sophorolipid methyl ester **6** at -78°C. Ozone is generated from dry air and is purged through the reaction mixture at a flow of 2.15 L/min and concentration of 6.2 mg/L. For large scale reactions, use of microreactor equipment should be considered.¹³


The purification of the sophorolipid aldehyde **7** proved to be a challenge due to the presence of saturated sophorolipids from the fermentation process and degradation products from the ozonolysis reaction. Multiple column chromatography purifications were evaluated, giving the highest yield and purity for automated flash chromatography with a gradient elution of

ethyl acetate and hexane. A sodium bisulfite addition reaction was performed after the chromatography purification to improve the purity but resulted in a very low yield.

Synthesis of a quaternary ammonium salt library

Next, a nitrogen functionality was introduced through reductive amination of the aldehyde function of **7** with a variety of secondary amines (Table 1). Complete conversion to the resulting sophorolipid tertiary amines **10** was obtained, encompassing methyl, butyl, benzyl and octadecyl groups. Purification of sophorolipid tertiary amines **10** was necessary to avoid purification of sophorolipid quaternary ammonium salts **11** in the next step, but resulted in low yields.


Table 1. Reductive amination towards sophorolipid amines **10**



Entry	R ¹ R ² NH	Yield (%)
1	dimethylamine	10a 38
2	<i>N</i> -methylbutylamine	10b 52
3	dibutylamine	10c 53
4	<i>N</i> -methylbenzylamine	10d 40
5	<i>N</i> -butylbenzylamine	10e 49
6	dibenzylamine	10f 48
7	<i>N</i> -methyloctadecylamine	10g 39


Subsequently, the sophorolipid tertiary amines **10** were quaternized with alkyl iodides in pressure vials to obtain the desired peracetylated sophorolipid quaternary ammonium salts **11** (Table 2). In a final step, the sugar head group of the quaternary ammonium salts **11** was deprotected to obtain the water soluble sophorolipid quaternary ammonium salts **3** (Table 3).

Table 2. Quaternization towards peracetylated sophorolipid quaternary ammonium salts **11**



Entry	10	R ³ I	Time (h)	Yield (%)
1	10a	methyl iodide	18	11a 91
2	10b	methyl iodide	18	11b 89
3	10c	methyl iodide	18	11c 96
4	10c	butyl iodide	48	11d 94
5	10d	methyl iodide	18	11e quant.
6	10e	methyl iodide	18	11f 89
7	10f	methyl iodide	18	11g quant.
8	10g	methyl iodide	18	11h 98
9	10g	butyl iodide	48	11i quant.

Table 3. Transesterification towards sophorolipid quaternary ammonium salts **10**



Entry	11	Yield (%)
1	11a	3a quant.
2	11b	3b 88
3	11c	3c quant.
4	11d	3d quant.
5	11e	3e quant.
6	11f	3f quant.
7	11g	3g 99
8	11h	3h 97
9	11i	3i 66

Evaluation of the antimicrobial activity

The antimicrobial activity of peracetylated sophorolipid aldehyde **7**, peracetylated sophorolipid amines **10a**, **10b**, **10c**, **10d**, **10f** and **10g**, peracetylated sophorolipid quaternary ammonium salts **11** and deprotected sophorolipid quaternary ammonium salts **3** was evaluated. The Gram-negative strains *Escherichia coli* LMG 8063 and *Klebsiella pneumoniae* LMG 2095, and the Gram-positive strains *Staphylococcus aureus* LMG 8064 and *Bacillus subtilis* LMG 13579 were chosen as test strains. The bioassay was carried out in 96-well microtiter plates at a concentration of approximately 0.5 mg/mL of the test compound and 10⁴ CFU/mL test bacteria. None of the test compounds showed significant growth inhibition of *Escherichia coli* LMG 8063 and *Klebsiella pneumoniae* LMG 2095. Compounds **11b**, **11c**, **11d**, **11e**, **11f**, **11g**, **11h**, **11i**, **3b**, **3h** and **3i** showed significant growth inhibition of *Staphylococcus aureus* LMG 8064 and *Bacillus subtilis* LMG 13579.

For the active compounds, the minimum inhibitory concentration (MIC) was determined against a test panel of four Gram-positive bacterial strains, namely *Staphylococcus aureus* LMG 8064, *Enterococcus faecium* LMG 11397, *Bacillus subtilis* LMG 13579 and *Streptococcus pneumoniae* LMG 16738. The MIC value is considered as the lowest concentration of the test compound for which a lack of visible bacterial growth is observed. This bioassay was carried out in 96-well microtiter plates at a concentration range between 100 and 2.5 µg/mL or 1000 and 5 µg/mL of the test compounds for respectively strong or weak inhibitors and 10⁴ CFU/mL test bacteria. MIC values for the active compounds are given in Table 4 together with MIC values for the antibiotic gentamicin sulfate. Microscopic analysis in addition to the determination of the MIC values revealed that lysis of the cells occurred at the active concentrations.

The lowest MIC value of 5 µg/mL was obtained with compounds **3h** and **3i** against all four Gram-positive test strains. Low MIC values of 10 µg/mL were obtained with compounds **11h** and **11i**, also against all four Gram-positive test strains. Interestingly, these activities lie in the same concentration

range as that of the antibiotic gentamicin sulfate. All four compounds perform as good or better as gentamicin sulfate against *E. faecium* and *S. pneumoniae*. Compounds **3h** and **3i** even perform as good as gentamicin sulfate against *S. aureus* and *B. subtilis*. For better comparison, the MIC values were converted on the basis of their molecular weight (Table 5). On this basis, we can conclude that compounds **3h**, **3i**, **11h** and **11i** are more active against all four Gram-positive test strains than the antibiotic gentamicin sulfate. These results show great

promise for further evaluation of the biological activities of the active derivatives such as synergistic effects of multiple compounds against specific test strains or activities of specific compounds against multiple test strains. The active derivatives also offer opportunities for their use in medical applications such as the inhibition of biofilm formation. These sophorolipid quaternary ammonium salts represent a new class of antimicrobial surfactants.

Table 4. MIC values ($\mu\text{g/mL}$) for the active compounds and the antibiotic gentamicin sulfate

	11b	11c	11d	11e	11f	11g	11h	11i	3b	3h	3i	gentamicin sulfate
<i>S. aureus</i>	>100	>100	>100	500	>100	50	10	10	>100	5	5	5
<i>E. faecium</i>	>100	>100	>100	>1000	>100	>100	10	10	>100	5	5	10
<i>B. subtilis</i>	>100	25	>100	1000	>100	50	10	10	>100	5	5	5
<i>S. pneumoniae</i>	>100	100	>100	1000	>100	100	10	10	>100	5	5	25

Table 5. MIC values (μM) for the active compounds and the antibiotic gentamicin sulfate

	11b	11c	11d	11e	11f	11g	11h	11i	3b	3h	3i	gentamicin sulfate
<i>S. aureus</i>	>101	>97	>93	489	>94	45	8	8	>144	6	5	10
<i>E. faecium</i>	>101	>97	>93	977	>94	>91	8	8	>144	6	5	21
<i>B. subtilis</i>	>101	24	>93	977	>94	45	8	8	>144	6	5	10
<i>S. pneumoniae</i>	>101	97	>93	>977	>94	91	8	8	>144	6	5	52

Conclusions

Microbial produced sophorolipids have been chemically modified into a series of novel sophorolipid amines and sophorolipid quaternary ammonium salts via an intermediate sophorolipid aldehyde. The sophorolipid amines and sophorolipid quaternary ammonium salts have been evaluated for their antimicrobial activity against the Gram-negative strains *Escherichia coli* LMG 8063 and *Klebsiella pneumoniae* LMG 2095, and the Gram-positive strains *Staphylococcus aureus* LMG 8064 and *Bacillus subtilis* LMG 13579. Eight of the peracetylated sophorolipid quaternary ammonium salts and three of the deprotected sophorolipid quaternary ammonium salts showed inhibition against the Gram-positive strains, but not against the Gram-negative strains. For the active compounds, their minimum inhibitory concentration (MIC) has been determined against the four Gram-positive strains *Staphylococcus aureus* LMG 8064, *Enterococcus faecium* LMG 11397, *Bacillus subtilis* LMG 13579 and *Streptococcus pneumoniae* LMG 16738. The best results are obtained with N,N-dimethyl,N-octadecyl-(8-L-[(2'',3'',3'',4'',4'',6'',6''-heptaacetoxy-2'-O- β -D-glucopyranosyl- β -D-glucopyranosyl)-oxy])nonan-1-ammonium iodide **11h**, N-benzyl,N-methyl,N-octadecyl-(8-L-[(2'',3'',3'',4'',4'',6'',6''-heptaacetoxy-2'-O- β -D-glucopyranosyl- β -D-glucopyranosyl)-oxy])nonan-1-ammonium iodide **11i**, N,N-dimethyl,N-octadecyl-(8-L-[(2 β -O- β -D-glucopyranosyl- β -D-glucopyranosyl)-oxy])nonan-1-ammonium iodide **3h** and with N-benzyl,N-methyl,N-octadecyl-(8-L-[(2 β -O- β -D-glucopyranosyl- β -D-glucopyranosyl)-oxy])nonan-1-ammonium

iodide **3i**. These four compounds are more active than the antibiotic gentamicin sulfate against all four Gram-positive test strains. These results show great promise for further evaluation of the biological activities of sophorolipid quaternary ammonium salts with an octadecyl group on the nitrogen atom and their use for medical applications. These sophorolipid quaternary ammonium salts represent a new class of antimicrobial surfactants.

Acknowledgements

The research leading to these results has received funding from the Long Term Structural Methusalem Funding by the Flemish Government (grant number BOF09/01M00409). The author gratefully acknowledge the company Ecover (Malle, Belgium) and the InBio research group (Department of Biochemical and Microbial Technology, Ghent University, Belgium) for the delivery of the sophorolipid starting compound, and the Laboratory for Microbiology (Ghent University, Belgium) for the evaluation of the antimicrobial activities.

Notes and references

^a SynBioC, Department of Sustainable Organic Chemistry and Technology, Ghent University, Coupure Links 653, 9000 Ghent, Belgium.

^b LCT, Department of Chemical Engineering and Technical Chemistry, Ghent University, Technologiepark 914, 9052 Ghent, Belgium.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/b000000x/

- (a) R. C. Brown, *Biorenewable resources. Engineering new products from agriculture*. Blackwell Publishing: USA, 2003; p 286; (b) A. A. Koutinas, C. Du, R. H. Wang and C. Webb, In *Introduction to chemicals from biomass*, ed. J. H. Clark and F. E. I. Deswarte, John Wiley & Sons, UK, 1 edn., 2008, pp 77-101; (c) J. O. Metzger and M. Eissen, *C. R. Chim.*, 2004, **7**, 569-581.
- (a) V. Vandermeulen, M. Van der Steen, C. V. Stevens and G. Van Huylenbroeck, *Biofuel Bioprod Bior*, 2012, **6**, 453-466; (b) H. L. Bos and J. P. M. Sanders, *Biofuel Bioprod Bior*, 2013, **7**, 246-259; (c) J. P. M. Sanders and H. L. Bos, *Biofuel Bioprod Bior*, 2013, **7**, 260-272; (d) C. V. Stevens, *Biofuel Bioprod Bior*, 2012, **6**, 495; (e) C. V. Stevens, *Biofuel Bioprod Bior*, 2011, **5**, 115.
- M. Chambers, M. Muecke, *J. Green Building*, 2010, **5**, 91.
- (a) I. N. A. Van Bogaert, K. Saerens, C. De Muynck, D. Develter, W. Soetaert and E. J. Vandamme, *Appl. Microbiol. Biotechnol.*, 2007, **76**, 23-34; (b) I. N. A. Van Bogaert, J. X. Zhang and W. Soetaert, *Process Biochem.*, 2011, **46**, 821-833.
- D. W. G. Develter and L. M. L. Lauryssen, *Eur. J. Lipid Sci. Technol.*, 2010, **112**, 628-638.
- (a) C. Jing, S. Xin, Z. Hui and Y. B. Qu, *Enzyme Microb. Technol.*, 2006, **39**, 501-506; (b) J. Chen, X. Song, H. Zhang, Y. B. Qu and J. Y. Miao, *Appl. Microbiol. Biotechnol.*, 2006, **72**, 52-59; (c) L. J. Shao, X. Song, X. J. Ma, H. Li and Y. B. Qu, *J. Surg. Res.*, 2012, **173**, 286-291; (d) S. L. Fu, S. R. Wallner, W. B. Bowne, M. D. Hagler, M. E. Zenilman, R. Gross and M. H. Bluth, *J. Surg. Res.*, 2008, **148**, 77-82; (e) V. Shah, G. F. Doncel, T. Seyoum, K. M. Eaton, I. Zalenskaya, R. Hagver, A. Azim and R. Gross, *Antimicrob. Agents Chemother.*, 2005, **49**, 4093-4100; (f) V. Shah, D. Badia and P. Ratsep, *Antimicrob. Agents Chemother.*, 2007, **51**, 397-400.
- (a) N. Baccile, N. Nassif, L. Malfatti, I. N. A. Van Bogaert, W. Soetaert, G. Pehau-Arnaudet and F. Babonneau, *Green Chem.*, 2010, **12**, 1564-1567; (b) A. S. Cuvier, J. Berton, C. V. Stevens, G. C. Fadda, F. Babonneau, I. N. A. Van Bogaert, W. Soetaert, G. Pehau-Arnaudet and N. Baccile, *Soft Matter*, 2014, **10**, 3950-3959.
- (a) S. K. Singh, A. P. Felse, A. Nunez, T. A. Foglia and R. A. Gross, *J. Org. Chem.*, 2003, **68**, 5466-5477; (b) A. Azim, V. Shah, G. F. Doncel, N. Peterson, W. Gao and R. Gross, *Bioconjugate Chem.*, 2006, **17**, 1523-1529; (c) K. S. Bisht, R. A. Gross and D. L. Kaplan, *J. Org. Chem.*, 1999, **64**, 780-789; (d) J. A. Carr and K. S. Bisht, *Tetrahedron*, 2003, **59**, 7713-7724; (e) J. A. Zerkowski, D. K. Y. Solaiman, R. D. Ashby and T. A. Foglia, *J. Surfactants Deterg.*, 2006, **9**, 57-62; (f) *US Pat.*, 4 195 177, 1980; (g) *US Pat.*, 4 305 931, 1981; (h) *US Pat.*, 4 305 929, 1981; (i) Y. F. Peng, D. J. Munoz-Pinto, M. T. Chen, J. Decatur, M. Hahn and R. A. Gross, *Biomacromolecules*, 2014, **15**, 4214-4227.
- (a) *US Pat.*, EP 1 953 237 A1, 2008; (b) Y. Peng, F. Totsingan, M. A. R. Meier, M. Steinmann, F. Wurm, A. Koh and R. A. Gross, *Eur. J. Lipid Sci. Technol.*, 2015, **117**, 217-228.
- S. L. K. W. Roelants, K. Ciesielska, S. L. De Maeseneire, B. Everaert, Q. Denon, H. Moens, B. Vanlerberghe, I. N. A. Van Bogaert, P. Van der Meer, B. De Vreese and W. Soetaert, *Biotechnol. Bioeng.*, submitted.
- S. Kyasa, T. J. Fisher and P. H. Dussault, *Synthesis-Stuttgart*, 2011, 3475-3481.
- C. Schwartz, J. Raible, K. Mott and P. H. Dussault, *Tetrahedron*, 2006, **62**, 10747-10752.
- (a) M. Nobis, D.M. Roberge, *Chemistry Today*, 2011, **29**, 56; (b) M. Irfan, T. N. Glasnov and C. O. Kappe, *Org Lett*, 2011, **13**, 984-987.