

Green Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/greenchem



Green Chemistry

COMMUNICATION

Sustainable oxidations with air mediated by gallic acid: Potential applicability in the reutilization of grape pomace†

Received 00th January 20xx,
Accepted 00th January 20xx

Jimena Scoccia,^a Marcelle D. Perretti,^b Diego M. Monzón,^b Fernando P. Crisóstomo,^b Víctor S. Martín^b Romen Carrillo*^b


DOI: 10.1039/x0xx00000x

www.rsc.org/

Gallic acid converts atmospheric oxygen into hydrogen peroxide, which is able to oxidize arylboronic acids as a proof of concept of sustainable oxidations. Moreover, tannic acid and grape pomace extract are also able to perform oxidations with air. Therefore this work unleashes an alternative method of reutilization and valorization of bio-waste rich in tannins.

Oxygen from air is certainly the most sustainable and affordable oxidant and oxygen atom source. However oxygen is in a triplet ground state which makes the reaction with singlet organic compounds spin forbidden.¹ Traditionally, such forbiddance is circumvented by using paramagnetic transition catalyst and/or high temperatures or irradiation.² In order to walk towards a higher sustainability, metal free methodologies for oxygen activation are highly desirable, particularly if such methodologies are based on non-toxic compounds from renewable sources at room temperature.^{3–9} Herein we report an environment friendly solution for oxygen activation by taking advantage of the known auto-oxidation of gallic acid in slightly basic media: Indeed gallic acid can be considered as the simplest prototype of tannin,¹⁰ and it can be extracted from vegetal sources such as oak bark or grape pomace,¹¹ or synthesized from glucose and microbial catalysis;¹² moreover, it has been reported that gallic acid is able to generate Reactive Oxygen Species (ROS) from oxygen.^{13–15} Thus the potential use of gallic acid for atmospheric oxygen activation towards sustainable oxidations has been herein explored and applied (Scheme 1). Additionally, there is an increasing social and legal pressure to develop sustainable processes for the use of natural 'feedstocks' in chemical synthesis as an alternative to hazardous organic reagents or scarce and expensive metal-based catalysts. Particularly attractive is the reutilization and valorisation of the

numerous residues generated by agricultural activities.^{16,17} Thus, the use of an abundant and underutilized bio-waste such as grape pomace, in sustainable oxidations with air has been also explored herein (Scheme 1).



Scheme 1 Sustainable oxidations with air by gallic acid or with grape pomace extract.

Oxidative hydroxylation of arylboronic acids to phenols (*ipso*-hydroxylation) was chosen as an optimal example to test the effective generation of hydrogen peroxide from air, and to optimize the oxidation conditions. Indeed, it is well known that arylboronic acids react efficiently with ROS. Thus, phenols are expected to be obtained in case that auto-oxidation of gallic acid efficiently generates reactive oxygen species, and no secondary/lateral reactions interfere in any way. Optimization was performed with 4-tertbutyl phenylboronic acid (**1a**, 0.5 mmol). It was dissolved in acetone (2 ml) together with gallic acid (0.5 mmol). Then a 1M solution of sodium bicarbonate (2 ml) was added and the reaction was stirred under air for 24h. Fortunately, 4-tertbutylphenol (**2a**) was obtained in excellent yield, as shown in Table 1 (entry 1).¹⁸ When aqueous phase was neutral or acidic, no oxidation was observed, even after longer reaction times (entries 2 and 3). It is important to highlight that bicarbonate is vital to our process. Using a 1M solution of K₂HPO₄ instead of bicarbonate, the oxidation was unsuccessful, although both salts provide comparable ionic strength and pH (entry 4). Therefore it is very likely that the role of bicarbonate is to activate the peroxide generated, which is proved below. Additionally, the amount of gallic acid could be decreased to 0.55 eq without any detriment on the yield nor in the reaction time (entry 5).¹⁵ Lesser amounts of gallic acid led to incomplete conversions and lower yields (entry 6). Finally, reactions on

^a Departamento de Química, Universidad Nacional del Sur, Av. Alem 1253, 8000 Bahía Blanca, Argentina. INQUISUR, CONICET, Av. Alem 1253, 8000 Bahía Blanca, Argentina.


^b Departamento de Química Orgánica, Instituto Universitario de Bio-Orgánica "Antonio González" (IUBO), Centro de Investigaciones Biomédicas de Canarias (CIBICAN), Universidad de La Laguna. Avda. Astrofísico Fco. Sánchez 2, Apdo. Correos 456, 38200 – La Laguna, S/C. de Tenerife (Spain). Phone: +34 922 318 583; e-mail: rocarril@ull.es

†Electronic Supplementary Information (ESI) available: Experimental details, characterization of the products and NMR spectra. See DOI: 10.1039/x0xx00000x

COMMUNICATION

aqueous phase exclusively, gave lower yields (entry 7). Ethanol was chosen as the organic co-solvent, not only because it gave a slightly better yield (entry 8), but also because ethanol is cheaper and has GRAS (generally recognized as safe according to U.S. Food and Drug Administration definition) status. Worth to mention, unchanged yields were obtained when reaction was carried out in the dark (entry 10), ruling out any role of light.¹⁹


Table 1 Optimization of the conditions for the ipso-hydroxylation of boronic acids^a



Entry	Gallic acid	Solvent Mixture	Isolated Yield
1	1 eq	Acetone/NaHCO ₃ (1M)	96
2	1 eq	Acetone/H ₂ O	Traces
3	1 eq	Acetone/TsOH (1M)	Traces
4	1 eq	Acetone/KHPO ₄ (1M)	16
5	0.55 eq	Acetone/NaHCO ₃ (1M)	96
6	0.4 eq	Acetone/NaHCO ₃ (1M)	68
7	0.55 eq	NaHCO ₃ (1M)	76
8	0.55 eq	Ethanol/NaHCO₃ (1M)	98
9	0.55 eq	Isopropanol/NaHCO ₃ (1M)	97
10 ^b	0.55 eq	Ethanol/NaHCO ₃ (1M)	98

^a 0.5 mmol of boronic acid used, 24h of reaction. ^b In the dark.


Once the optimal conditions were set up, the scope of the reaction was explored. As it can be seen in Scheme 2, yields were excellent regardless of the nature of the boronic acid employed. Only sterically crowded 2, 6-dimethylphenol (**2g**) and 2-cyanophenol (**2f**) were obtained in moderate yields. We should emphasize that *n*-decylboronic acid was also efficiently oxidized.



Scheme 2 Scope of ipso-hydroxylation of arylboronic acids, under optimized conditions. Conversion was complete in all cases.


On the other hand, arylboronic esters and potassium trifluoroborate salts were also efficiently oxidized with this methodology (Scheme 3).

Journal Name



Scheme 3 ipso-hydroxylation of boronic ester and trifluoroborate salt, under optimized conditions for 24h. Conversion was complete in both cases.

A few mechanistic experiments were performed. Firstly, the reaction of oxidation of phenylboronic acid was carried out under ¹⁸O₂ atmosphere, in order to prove that this methodology is indeed activating the atmospheric oxygen and incorporating it into the final products. As it can be seen in Scheme 4, incorporation of such heavier isotope of oxygen in the final compound is detected by MS.



Scheme 4 ¹⁸O labelling experiment.

In addition, ¹³C NMR confirmed the abovementioned activation of peroxide by bicarbonate. Such bicarbonate activated peroxide (BAP) is a known method for activating H₂O₂.²⁰ In such case, the actual oxidant is not hydrogen peroxide, but peroxymonocarbonate ion (HCO₄⁻) formed by the equilibrium reaction between bicarbonate ion and H₂O₂ (Figure 1). Peroxymonocarbonate ion is a reactive species but it can be detected by several techniques, such as ¹³C NMR. We prepared a 1:1 solution of methanol-*d*₄ and 99% ¹³C-enriched NaHCO₃ (1M) in deuterated water, analogous to the solution used in the reactions. Gallic acid was added and after 10 minutes stirring under air, the ¹³C NMR experiment at 25 °C was submitted. In the NMR spectrum it is clearly seen three peaks (Figure 1).




Figure 1 Detection of peroxymonocarbonate by ¹³C NMR.


The peak at 160.3 ppm is due to HCO₃⁻, while the peak at 158.9 ppm corresponds to peroxymonocarbonate HCO₄⁻. The third peak observed at 160.0 ppm is due to the known reaction between primary alcohols, such as methanol, with bicarbonate to form alkyl carbonate esters, in this case the methyl carbonate (MeCO₃⁻). The apparently small peak of peroxymonocarbonate detected is understandable not only due to its intrinsically instable nature, but also because in this case hydrogen peroxide

is generated progressively in situ in small amounts, compared to previously reported experiments,²⁰ where a high concentration of peroxide is employed. Worth to mention, the confirmation of BAP mechanism, paves the way for future combinations of this methodology with sustainable metals or, even better, phytoextracted metals.²¹

Finally, we pondered the application of this methodology as an alternative way for the reutilization and valorization of bio-waste,^{22,23} such as grape pomace, which is known to be rich in gallic acid and other tannins.¹¹ Grape pomace is an abundant agricultural residue: The wine industry generates a huge amount of grape by-products called pomaces, which account for about 20% of the weight of the grapes used to make wine.²³ Considering that grape is one of the world's largest fruit crops,²⁴ grape pomace represents a waste management issue both ecologically and economically. Thus, we decided to explore a grape pomace extract to carry out oxidations with air.²⁵

As a first approach, tannic acid was employed instead of gallic acid. Tannic acid is present in considerable amounts in wine byproducts.¹¹ The aim of this preliminary approach is to evaluate if complex derivatives of gallate behave similarly. Indeed, tannic acid displays a glucose bearing five digallates (Scheme 5). The optimal amount of tannic acid required for a complete oxidation in 24h was studied, being 0.2 eq. As it is seen in Scheme 5, oxidation of several boronic acids was successful.

Subsequently, we tested the oxidation of arylboronic acids directly with grape pomace extract. For this experiment, grape pomace from a local winery was employed (E. Páez winery). The red grapes *Vitis vinifera* (*Listan Negro*, a local cultivar) were cultivated in the Canary Islands, specifically in the north of Tenerife (Finca Los Angostos, La Cruz Santa, Los Realejos) and harvested at the beginning of October 2015. The grapes were pressed right after the harvest and left in contact with the grape juice for one week, while fermentation was going on. Grape seeds and skins were pressed and dried at air for two days. The extraction method chosen was analogous to other previously reported,²⁶ and it was not optimized. The grape pomace was left with ethanol/water/HCl (37%) (7:2.9:0.1) (3ml per g of pomace), for 3 days, and then filtered. To 2ml of such extract, boronic acid was added (0.1mmol), and finally 2ml of 1M solution of sodium bicarbonate (2 ml) was added and the reaction was stirred under air for 24h. These results are shown in Scheme 5 (purple). As it can be seen, yields were excellent considering the complex mixture present in the extract, and the non-optimized conditions employed. Further experiments on this and other reactions are being carried out in our lab.



Scheme 5 Yield of the oxidation of boronic acids by tannic acid (in green) and grape pomace extract (in purple). The latter oxidations were carried out in 0.1 mmol scale. Conversion was complete in all cases.

Conclusions

In summary, auto-oxidation of gallic acid in slightly basic media has proven to be an inexpensive and sustainable way to generate hydrogen peroxide in situ from air, which acts efficiently as an oxidant of arylboronic acids. The relevance of bicarbonate in the reaction media has proved to be related not to the pH, but to the activation of the peroxide, to produce the reactive peroxymonocarbonate. More importantly, other tannins such as tannic acid, and even grape pomace extract are able to carry out the oxidation reactions with air successfully. Therefore, these results unleash an alternative method for the reutilization of certain bio-wastes like grape pomace, which is rich in gallic acid and its derivatives: This work shows the potential utilization of the reactivity of compounds present in bio-waste, rather than the compounds themselves. Such an approach, therefore, could complement other traditional waste reutilization methods. Indeed, once the extraction of grape pomace has been performed, other traditional treatments of such waste can still be applied (biogas, fertilizer, etc.). Additionally, there are great environmental and economic advantages in substituting conventional chemical reagents for bio-waste, thus avoiding compounds which are very likely to be


more expensive or more toxic, or which simply have to be manufactured, causing additional costs and waste.

Acknowledgments

This work was supported by the European Commission (IMBRAIN project, FP7-REGPOT-2012-CT2012-31637-IMBRAIN) and the Spanish MINECO (CTQ2014-56362-C2-1-P), cofinanced by the European Regional Development Fund (ERDF). The authors are also very grateful to Esteban Páez for supplying the grape pomace.

Notes and references

- (a) B. F. Minaev, *Int. J. Quantum. Chem.*, 1980, **17**, 367-374; (b) A. L. Buchchenko and V. L. Berdinsky, *J. Phys. Chem.*, 1996, **100**, 18292-18299.
- (a) A. Bakac, *Inorg. Chem.*, 2010, **49**, 3584-3593. (b) R. Prabhakar, P. E. M. Siegbahn, B. F. Minaev and H. Ågren, *J. Phys Chem. B*, 2004, **108**, 13882-13892.
- G. Silveira-Dorta, D. M. Monzón, F. P. Crisóstomo, T. Martín, V. S. Martín and R. Carrillo, *Chem. Commun.*, 2015, **51**, 7027-7030.
- (a) H. Kawai, S. Okusu, Z. Yuan, E. Tokunaga, A. Yamano, M. Shiro and N. Shibata, *Angew. Chem. Int. Ed.*, 2013, **52**, 2221 – 2225.
- Y. Imada, H. Iida, S. Ono and S.-I. Murahashi, *J. Am. Chem. Soc.*, 2003, **125**, 2868-2869.
- F. Hollmann, A. Taglieber, F. Schulz and M. T. Reetz, *Angew. Chem. Int. Ed.*, 2007, **46**, 2903-2906.
- S. Chen and F. W. Foss Jr., *Org. Lett.*, 2012, **14**, 5150-5153.
- R. Liu, X. Liang, C. Dong and X. Hu, *J. Am. Chem. Soc.*, 2004, **126**, 4112-4113.
- R. Mu, Z. Liu, Z. Yang, Z. Liu, L. Wu and Z.-L. Liu, *Adv. Synth. Catal.*, 2005, **347**, 1333-1336.
- B. Badhani, N. Sharma and R. Kakkar, *RSC Adv.*, 2015, **5**, 27540-27557.
- (a) Y. Yilmaz and R. T. Toledo, *J. Agric. Food Chem.*, 2004, **52**, 255-260; (b) M. R. González-Centeno, M. Jourdes, A. Femenia, S. Simal, C. Rosselló and P.-L. Teissedre, *J. Agric. Food Chem.*, 2013, **61**, 11579-11587; (c) A. R. Fontana, A. Antoniolli and R. Bottini, *J. Agric. Food Chem.*, 2013, **61**, 8987-9003; (d) Y. Lu and L. Y. Foo, *Food Chem.*, 1999, **65**, 1-8; (e) T.-I. Lafka, V. Sinanoglou and E. S. Lazos, *Food Chem.*, 2007, **104**, 1206-1214.
- S. Kambourakis, K. M. Draths and J. W. Frost, *J. Am. Chem. Soc.*, 2000, **122**, 9042-9043.
- L. H. Russell Jr, E. Mazzi, R. B. Badisa, Z.-P. Zhu, M. Agharahimi, E. T. Oriaku and C. B. Goodman, *Anticancer Res.*, 2012, **32**, 1595-1602.
- P. A. Clapp, N. Du and D. F. Evans, *J. Chem. Soc. Faraday Trans.*, 1990, **86**, 2587-2592.
- V. Tulyathan, R. B. Boulton and V. L. Singleton, *J. Agric. Food Chem.* 1989, **37**, 844-849.
- (a) L. H. Keith, L. U. Gron and J. L. Young, *Chem. Rev.*, 2007, **107**, 2695-2708; (b) P. T. Anastas, *Crit. Rev. Anal. Chem.*, 1999, **29**, 167-175; (c) R. A. Sheldon, *Green Chem.*, 2014, **16**, 950-963; (d) R. A. Sheldon, *Green Chem.*, 2005, **7**, 267-278; (e) P. T. Anastas and J. C. Warner, *Green Chemistry: Theory and Practice*, Oxford University Press, New York, 1998.
- (a) P. R. Boruah, A. A. Ali, B. Saikia and D. Sarma, *Green Chem.*, 2015, **17**, 1442-1445; (b) B. Saikia, P. Borah and N. C. Barua, *Green Chem.*, 2015, **17**, 4533-4536; (c) P. R. Boruah, A. A. Ali, M. Chetia, B. Saikia and D. Sarma, *Green Chem.*, 2015, **17**, 11489-11492; (d) E. Saikia, S. J. Bora and B. Chetia, *RSC Adv.*, 2015, **5**, 102723-102726; (e) B. Saikia and P. Borah, *RSC Adv.*, DOI: 10.1039/C5RA20133K.
- The resulting compound of the oxidation of gallic acid in alkaline solutions is the highly coloured galloflavin. See: (a) R. Bohn and C. Graebe, *Chemische Berichte*, 1887, **20**, 2327-2331; (b) J. Grimshaw and R. D. Howarth, *J. Chem. Soc.*, 1956, 418-422.
- Reactions under pure O₂ led to similar yields. We chose open flask reactions because they are much more sustainable and affordable.
- (a) H. Yao and D. E. Richardson, *J. Am. Chem. Soc.*, 2000, **122**, 3220-3221; (b) D. E. Richardson, H. Yao, K. M. Frank and D. A. Bennett, *J. Am. Chem. Soc.*, 2000, **122**, 1729-1739; (c) B. Balagam and D. E. Richardson, *Inorg. Chem.*, 2008, **47**, 1173-1178; (d) E. V. Bakhmutova-Albert, H. Yao, D. E. Denevan and D. E. Richardson, *Inorg. Chem.*, 2010, **49**, 11287-11296.
- (a) V. Escande, E. Petit, L. Garoux, C. Boulanger, and C. Grison, *ACS Sustainable Chem. Eng.*, 2015, **3**, 2704-2715. (b) V. Escande, A. Velati, C. Garel, B.-L. Renard, E. Petit and C. Grison, *Green Chem.*, 2015, **17**, 2188-2199.
- R. A. D. Arancon, C. S. K. Lin, K. M. Chan, T. H. K. and R. Luque, *Energy Science and Engineering*, 2013, **1**, 53-71.
- G. Laufenberg, B. Kunz and M. Nystroem, *Bioresource Technology*, 2003, **87**, 167-198.
- More than 67 million tons in 2010 according FAOSTAT-FAO Statistical Database, 2010.
- Other polyphenolic compounds such as anthocyanins could have a beneficial or detrimental effect on the reactions. Further experiments are being carried out in our group in order to understand the effect of other compounds present in the extract.
- A. R. Fontana, A. Antoniolli and R. Bottini, *J. Agric. Food Chem.* 2013, **61**, 8987-9003.



Tannins like gallic acid and even grape pomace extract are able to perform oxidations with air thus unleashing an alternative method of reutilization and valorization of bio-wastes.