

Molecular BioSystems

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/molecularbiosystems

Computational predictions of corroles as a class of Hsp90 inhibitors†

Cite this: DOI: 10.1039/x0xx00000x

Ruijie D. Teo,^{a,b,†} Sijia S. Dong,^{a,b,†} Zeev Gross,^c Harry B. Gray,^b William A. Goddard^{*a,b}

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

Introduction

Corroles, and in particular the metal chelates of 5,10,15-tris(pentafluorophenyl)-2,17-bis(sulfonic acid)-corrole 1-H₃, 1-M (Scheme 1), have become increasingly important in recent years due to their utility in medical applications. The iron and manganese complexes display excellent catalytic activity for decomposition of the main reactive oxygen species (ROS) and reactive nitrogen species (RNS) that are implicated in numerous diseases. Many beneficial effects have been disclosed for 1-Fe in model systems of diabetes,¹ neurodegeneration,² optic neuropathy,³ and atherosclerosis.⁴ 1-Mn and similar manganese corroles are most effective for the attenuation of molecular and intracellular nitration. 1-Sb is able to photodynamically inactivate mold fungi spores.⁵ 1-Al and 1-Ga are highly

Corroles have been shown experimentally to cause cell cycle arrest, and there is some evidence that this might be attributed to an inhibitory effect of corroles on Heat shock protein 90 (Hsp90), which is known to play a vital role in cancer cell proliferation. In this study, we used molecular dynamics to examine the interaction of gallium corroles with Hsp90, and found that it can bind preferentially to the ATP-binding N-terminal site. We also found that structural variations of the corrole ring can influence the binding energies and affinities of the corrole to Hsp90. We predict that both the bis-carboxylated corrole (4-Ga) and a proposed 3,17-bis-sulfonated corrole (7-Ga) are promising alternatives to Ga(III) 5,10,15-tris(pentafluorophenyl)-2,17-bis(sulfonic acid)-corrole (1-Ga) as anti-cancer agents. fluorescent,⁶ and 1-Au and 1-Ga have been shown to exhibit cytotoxicity in a number of cancer cell lines.⁷

These amphiphilic bis-sulfonated metallo-corroles interact with a variety of proteins, including albumin⁸ and transferrin.⁹ Recently, metallo-corroles have been shown to bind tightly to low-density lipoprotein (LDL) and a sub-fraction of high-density lipoprotein (HDL) upon mixing with human serum to protect them from oxidative/nitrative damage.¹⁰


^a Materials and Process Simulation Center, California Institute of Technology, Pasadena, California 91125, USA. E-mail: wag@wag.caltech.edu

^b Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California 91125, USA.

^c Schulich Faculty of Chemistry, Technion—Israel Institute of Technology, Haifa 32000, Israel.

† These authors contributed equally to this work.

‡ Electronic supplementary information (ESI) available: For ESI see DOI:



Scheme 1. Synthesis of cytotoxic metallo-corroles 1-M.

1-Ga spontaneously assembles with a heregulin-modified protein directed at the human epidermal growth factor receptor (HER), leading to targeted tumor cell death in a mouse model. This can be observed by the intense fluorescence of the corrole rendering it suitable for both cellular and whole animal optical imaging.^{11,12} Although there have been large advances in elucidating the cancer cell-killing mechanism by 1-Ga,^{13,14} both free and protein-conjugated, many open questions about its intracellular behavior remain unanswered.

One such class of intracellular proteins that have been implicated in cancer is heat shock proteins (Hsp).¹⁸ Hsp are ubiquitous molecular chaperones distributed from bacteria to animal cells and essential mediators of signal-transduction and cell cycle progression in prokaryotes and eukaryotes.¹⁹ In particular, Hsp90, a subclass of Hsp that is often implicated in

This journal is © The Royal Society of Chemistry 2015

a) FeCl₂, pyridine, reflux
 b) Mn(OAc)₂, DMF, reflux
 c) GaCl₃, pyridine, reflux
 d) Au(OAc)₃, pyridine, 25°C
 e) Au(OAc)₃, pyridine, 25°C
 f) SbCl₃, pyridine, reflux

1-Fe 15
 1-Mn 16
 1-Ga 17
 1-Au 6
 1-Sb 5

Ref.

tumorigenesis and tumor progression, aids in a wide range of intracellular processes - protein assembly, trafficking, folding, degradation, and neurodegenerative diseases.^{20,21} Since Hsp90 has a concentration 2-10 times higher in tumors than normal cells, it has been recognized as a potential target for cancer.²² Indeed the development of Hsp90 inhibitors has been rapid – 17 agents had entered clinical trials in 2012.²³ Potent Hsp90 inhibitors such as geldanamycin and radicicol (Fig. 1) block the ATP binding/hydrolysis site at the N-terminal nucleotide-binding pocket, inhibiting its function, depleting oncogenic clients, and engendering antitumor activity.^{23,24,25} Studies in both human breast cancer cells and dopaminergic neural cells showed that 19-substituted benzoquinone ansamycin (BQA) macrocycles are effective Hsp90 inhibitors while possessing markedly reduced toxicity to normal endothelial and epithelial cells than their parent quinones.²⁶




Fig. 1 Structures of Geldanamycin **1** and Radicicol **2**.

Previous work showed that Tubocapsenolide A inhibits activity of the Hsp90-Hsp70 chaperone complex, leading to destabilization of Hsp90 client proteins and subsequent cell cycle arrest and apoptosis in MDA-MB-231 cells.²⁷ In 2012, Termini and coworkers found that 1-Ga causes mitotic cell cycle arrest in the same MDA-MB-231 cell line.²⁸ Most recently, protoporphyrin IX (PPIX) treatment was purported to inhibit the chaperone activity of Hsp90, accelerating protein degradation of HIF-1 α .²⁹ These developments, coupled with the structural similarity of PPIX and 1-Ga, led us to hypothesize that 1-Ga's tumor killing ability is from its inhibition of Hsp90. In this study, we predicted the binding site for 1-Ga and its derivatives (Fig. 2) to Hsp90 which we compared with the predicted binding sites of PPIX and ATP with Hsp90. Then we carried out molecular dynamics (MD) simulation of the docked complexes. These studies led to corrole-Hsp90 interactions that support our hypothesis suggesting how corroles might inhibit Hsp90. These findings also provide insight into designing potent corrole-based anti-cancer drugs.




Fig. 2 Compounds investigated in this study - Gallium corrole analogues **1-Ga** to **5-Ga**, and PPIX (with the exception of ATP).

Procedures

Preparation of the protein and ligands

1- to 5-Ga were constructed using the crystal structure of a structurally similar gold corrole.³⁰ Starting with the co-crystal structure of Hsp90 with geldanamycin (Protein Data Bank (PDB) code: 1YET) we removed the ligand and all crystallographic water molecules. The structure for PPIX was obtained from a co-crystal structure of human ferrochelatase with PPIX (PDB code: 2HRE) while the structure of ATP was obtained from the crystal structure of ATP-bound monomeric actin (PDB code: 2HF4). All ligands tested were then geometrically optimized with quantum mechanics (the B3LYP-D3 level of Density Functional Theory (DFT) that includes D3 corrections for van der Waals attraction) using the 6-31G**++ basis set for the non-metal atoms and the LACV3P^{31,32} basis set for the metal and including the Poisson-Boltzmann model for implicit solvation by water, as implemented in the Jaguar 8.0 package.³³ We obtained the Mulliken charges from these DFT calculations for use in docking the ligands to the protein structure of Hsp90.

Molecular docking

Molecular docking studies were performed using our GenDock^{34,35} procedure using the DREIDING force field (FF).³⁶ GenDock has been shown to be able to predict protein-ligand binding sites and energy consistent with experimental findings for a number of cases including G protein-coupled receptors^{34,37} and cyclin-dependent kinase 5.³⁸ We first generated all possible ligand-binding positions in the protein by filling the binding regions with spheres that are classified into boxes with 10 Å sides. Then we used PyMOL³⁹ to choose spheres in the N-terminal binding pocket of Hsp90⁴⁰ for docking. For each ligand, we screened 5 million poses without energy evaluation and kept the 98-109 poses that could fit into the chosen binding pocket. The poses were clustered into 3 families by RMSD and all poses were maintained. We then de-alanized these selected poses using SCREAM. Then we

neutralized all charged residues in the protein except for Lys58 and Lys112 before geometry optimization of the poses, as both Lys58 and Lys112 are important mediators in ligand-Hsp90 interactions.⁴¹

Molecular dynamics


Starting with the docked structures, we inserted the ligand-protein into an equilibrated water box and carried out Molecular Dynamics (MD) calculations to relax the ligand-protein complex while allowing water to diffuse throughout the complex. These calculations used Amber force field with the NAMD 2.9⁴² MD software. We carried out 10 ns of MD at 298K to select the best 1-Ga-Hsp90 complex from the docking studies. Our procedure was to use VMD⁴³ to build an equilibrated orthorhombic water box that included the ligand-protein complex at the center and extended 10 Å beyond the protein in all 6 directions. Then we eliminated all water molecules within 5 Å of the complex. Using tleap from AmberTools11,⁴⁴ we added sufficient Cl⁻ and Na⁺ ions to neutralize the complex and to give a physiological salt concentration of 0.9% w/v. The final system has 29996 atoms. Then we minimized the solvent for 5000 steps with the protein

and ligand fixed, followed by 1 ns of NPT dynamics of the solvent using a bath temperature of 310 K and a pressure of 1 atm. We then conjugate gradient minimized the whole resulting system for 5000 steps, followed by 10 ns NPT dynamics for the whole system at 310 K and 1 atm. We used the Nosé-Hoover Langevin piston pressure control, with a damping coefficient of 5 ps⁻¹. The Particle Mesh Ewald (PME) method was applied for electrostatic interactions. Finally, the analysis of MD trajectories were performed using VMD.

Corrole Amber Parameters for MD

Since Amber does not have a force field for corroles, we introduced new atom types and approximated the force constants from a published set of cobalt corrole parameters⁴⁵ as well as GAFF.⁴⁶ The well depth for Ga nonbonding interaction was taken from UFF.⁴⁷ For the equilibrium bond distances and angles, we used the average bond distances and angles from the structure optimized using DFT (B3LYP-D3/LAV3P for Ga and B3LYP-D3/6-31G**++ for all other atoms). These modified force field parameters are in tables S1-6 of the SI.

Fig. 3 Predicted ligand-Hsp90 structures at the N-terminal binding pocket of Hsp90 - a) 1-Ga, b) 2-Ga, c) 3-Ga, d) 4-Ga, e) 5-Ga,



f) PPIX, and g) ATP. Dotted lines represent possible interactions and binding modes, and atoms are classified by color according to type. The final pdb files for the coordinates of 1-Ga-Hsp90 and 4-Ga-Hsp90 are listed in the SI.

Results and Discussion

Of all the gallium corroles studied (Table 1), we find that 1-Ga has the strongest bonding to Hsp90 [based on the unified-cavity energy (ucav) and the snap binding energy (snapbe)]. 1-Ga also has the largest number of interactions of favourable interactions as indicated in Table 1. We find that the sulfonate groups of 1-Ga both make salt-bridges with the positively-charged amino group of Lys112, while Lys58 ‘hangs’ from the top to form a cation- π interaction with the conjugated corrole π -system (Fig. 3a). Most interesting is that we find an uncommon anion- π interaction between the negatively-charged carboxylate group of Asp102 and the electron-poor $-C_6F_5$ group. We also find interactions for the docking of other gallium corrole analogues, including 2-Ga, 3-Ga, and 5-Ga (Fig. 3). We predict that 4-Ga is bonded nearly as strongly as 1-Ga, but that 2-Ga and 3-Ga bind much more weakly (due to the lack of hydrophobic groups).

Also observed are axial interactions (Fig. 3a and 3e) as expected between the Lewis acidic Ga(III) and the amide group of Asn106.

DFT calculations were also performed to validate the FF calculations of these interactions (Fig. S2a-b). Comparing the predicted binding energies among 1- to 5-Ga, PPIX and ATP provides insight into designing of new Hsp90 inhibitors. We find that it is essential for the ligand to have three hydrophilic groups to achieve maximal binding at the N-terminal site, which is the ATP binding site and relatively polar.⁴⁰ For comparison we docked ATP to the N-terminal binding pocket, leading to interactions (Fig. 3g) consistent with those in the x-ray structure (PDB code: 1AM1). Since ATP is much smaller than the corroles and has more hydrophilic groups, we find that it binds more strongly than the other ligands investigated here.

Table 1. Docking results of the best poses of 1-5-Ga, PPIX, and ATP to Hsp90. Here Snapbe and Scored ucav are two measure of the bond strength. Note that negative numbers for the binding energy indicate stronger bonds. The atom labels of the ligands are shown in Fig. S1 of the SI.

Corrole-Hsp90 complex	Snapbe (kcal mol ⁻¹)	Scored ucav (kcal mol ⁻¹)	Type of interaction	Interaction	Bond length (Å)
1-Ga	-150.44	-131.54	Salt-bridge	(Lys112)NZ-O66	2.8
			Salt-bridge	(Lys112)NZ-O68	2.7
			Non-covalent	(Asn51)OD1-C57	4.0
			Cation- π	(Lys58)NZ-C18	4.1
			Anion- π	(Asp102)OD1-C37	3.2
			Non-covalent	(Asn106)OD1-Ga1	3.5
4-Ga	-149.27	-130.31	H-bond	(Asn51)HD21-O67	2.1
			Salt-bridge	(Lys112)NZ-O70	2.7
			Non-covalent	(Met98)SD-C55	3.2
5-Ga	-107.93	-89.39	Salt-bridge	(Lys58)NZ-O67	2.7
			Cation- π	(Lys58)NZ-C21	5.0
			Anion- π	(Asp102)OD1-C37	3.6
			Non-covalent	(Asn106)OD1-Ga1	3.1
3-Ga	-43.20	-33.53	H-bonding	(Asn51)HD21-O78	1.8
			Anion- π	(Asp102)OD2-C56	2.9
2-Ga	-31.93	-31.66	Anion- π	(Asp102)OD1-C60	3.1
PPIX	-141.55	-118.94	Cation- π	(Lys58)NZ-C29	3.4
			Salt-bridge	(Lys112)NZ-O27	2.7
ATP	-288.59	-261.30	Salt-bridge	(Lys112)NZ-O18	2.6
			H-bond	(Asn106)HD22-O14	1.9
			Salt-bridge	(Lys58)NZ-O16	2.5

Despite our prediction that ATP binds more strongly to Hsp90 than PPIX and 1- to 5-Ga, experiments show that PPIX and hemin directly impede the activation of hypoxia-inducible factor 1 α (HIF-1 α) by inhibiting Hsp90.^{29,48} The molecular details of the interactions between PPIX and Hsp90 have been addressed previously by computational docking studies.²⁹ It was predicted that the conjugated aromatic system of porphyrin ring could make cation- π interactions with Lys58, while one of the propionates form a hydrogen bond with Lys112. Indeed, our docking predictions confirm the purported cation- π interaction of Lys58 with the porphyrin, and the direct salt-bridge interaction of Lys112 with one of the carboxylate groups (Fig. 3f). We also predict that the Hsp90 binding energy of 1-Ga and 4-Ga is stronger than that of PPIX, suggesting that 1-Ga and 4-Ga might be better Hsp90 inhibitors than PPIX.

To confirm our docking results, we carried out 10 ns of MD simulation on the docked complex 1-Ga-Hsp90. We

consider that 10ns is sufficient for our analyses because the RMSD changes of the Hsp90 backbone change little during the last 5 ns (Fig. S3). The key interactions of 1-Ga with Hsp90 (Table 1) were analysed over the 10ns as shown in Fig. 4. We find that these important interactions are relatively stable during the MD simulation (Fig. 5), with expected changes as water diffuses into the binding site. At the beginning of the MD simulation, Lys112 seemed to be drawn slightly towards proximal groups (Gly135 and Asn106) probably because of the initial interactions with the water. However, after 5 ns, the salt-bridge interaction between the sulfonate group of 1-Ga and the protonated nitrogen of Lys112 eventually dominates and is maintained throughout (Fig. 4a). The non-covalent (Asn51 and -C₆F₅ ring) and cation- π interaction (Lys58 and corrole) were maintained throughout, while it is noted that the anion- π interaction between OD1 of Asp102 and the -C₆F₅ ring is present but weak. The axial interaction between the gallium center and the amide group of Asn106 was maintained throughout with only some aberrations at 9 ns (Fig. 4e). In addition, a sulfur- π interaction (-S and -C₆F₅) was observed (Fig. 4c). The pharmacophore of 1-Ga with Hsp90 is shown in Fig. 6.








Fig. 4 10 ns MD results of the interaction distances of 1-Ga to Hsp90 starting from the docking pose c4644. a) Lys112 NZ to O65, O68, b) Lys58 NZ to C16, c) Met98 SD to C50, d) Asn51 OD1 to C57, e) Asn106 OD1 to GA1, and f) Asp102 OD1 to C40.

Table 2. Docking results of 7-Ga to Hsp90.

Ligand-Hsp90 complex	Snapbe (kcal mol ⁻¹)	Scored ucav (kcal mol ⁻¹)	Type of interaction	Interaction	Bond length (Å)
7-Ga	-145.08	-124.50	Salt-bridge	(Lys112)NZ-O58	2.9
			Cation- π	(Lys58)NZ-C28	4.0
			Anion- π	(Asp102)OD2-C59	4.9

**Fig. 5** Final frame of the MD for 1-Ga/Hsp90 after 10 ns MD. The starting pose is shown in Fig. S3. Residues important to binding are highlighted (color code: red = O, blue = N, white = H, yellow = S, pink = Ga).**Fig. 6** Pharmacophore of 1-Ga to Hsp90. The Ga center interacts with Asn106 (not shown).**Fig. 8** Docking of 7-Ga to the N-terminal binding pocket of Hsp90. The final pdb files for the coordinates of 7-Ga-Hsp90 are listed in the SI.

We predict two interactions (salt-bridge and cation- π) between Lys58 and Lys 112 with 7-Ga (Fig. 8). The predicted binding energy (scored ucav energy) for 7-Ga is comparable to that of 4-Ga and 1-Ga (Table 2).

The results for the 10 ns of MD simulations are summarized in Fig. 9. We find that the docked binding mode is stable (Fig. S9), with all key interactions from the docking maintained during the MD (Fig. S10 and Fig. 10). Moreover we find the sulfur- π interaction as observed in 1-Ga (Fig. 9b), although both the sulfur- π and anion- π bond distances are about 1 Å longer than that observed in 6-Ga (Fig. S8).

Fig. 9 10ns MD interaction Ga to Hsp90 the docking Lys112 NZ to NZ to O56, b) C43, c) Lys58 Asp102 OD1 to





Fig. 10 Ending frame of 7-Ga to Hsp90 after 10ns MD. The starting pose is shown in Fig. S10. Relevant residues are highlighted.

results of the distances of 7-starting from pose c23708. a) O54, Lys112 Met98 SD to NZ to C11, d) C34.

Conclusions

Our docking and MD simulation results suggest that the experimentally observed cytotoxicity of 1-Ga might be the result of its inhibition of Hsp90. We also propose a new gallium corrole derivative, 7-Ga, as a potential Hsp90 inhibitor, suggesting it as a promising candidate for further experimental investigation. These studies suggest a new

understanding of the tumor-killing mechanism of metalloporphyrins, and we have provided a pharmacophore that can help develop this new class of Hsp90 inhibitors.

Acknowledgements

A Caltech/COH grant (H.B.G. and Z.G.) and the AACR-Thomas J. Bardos Science Education Award (R.D.T.) is gratefully acknowledged. We would also like to thank Y. C. Lam for helpful discussions.

Notes and references

- Z. Okun, L. Kupersmidt, T. Amit, S. Mandel, O. Bar-Am, M. B. H. Youdim and Z. Gross, *ACS Chem. Biol.*, 2009, **4**, 910.
- L. Kupersmidt, Z. Okun, T. Amit, S. Mandel, I. Saltsman, A. Mahammed, O. Bar-Am, Z. Gross and M. B. H. Youdim, *J. Neurochem.*, 2010, **113**, 363.
- A. Kanamori, M. M. Catrinescu, A. Mahammed, Z. Gross and L. A. Levin, *J. Neurochem.*, 2010, **114**, 488.
- A. Haber, A. Mahammed, B. Fuhrman, N. Volkova, R. Coleman, T. Hayek, M. Aviram and Z. Gross, *Angew. Chem. Int. Ed.*, 2008, **47**, 7896.
- A. Preuß, I. Saltsman, A. Mahammed, M. Pfitzner, I. Goldberg, Z. Gross and B. Röder, *J. Photochem. Photobiol. B*, 2014, **133**, 39.
- D. Kowalska, X. Liu, U. Tripathy, A. Mahammed, Z. Gross, S. Hirayama and R. P. Steer, *Inorg. Chem.*, 2009, **48**, 2670.
- R. D. Teo, H. B. Gray, P. Lim, J. Termini, E. Domeshek and Z. Gross, *Chem. Commun.*, 2014, **50**, 13789.
- A. Mahammed, H. B. Gray, J. J. Weaver, K. Sorasaene and Z. Gross, *Bioconjugate Chem.*, 2004, **15**, 738.
- A. Haber, H. Agadjanian, L. K. Medina-Kauwe and Z. Gross, *J. Inorg. Biochem.*, 2008, **102**, 446.
- A. Haber, M. Aviram and Z. Gross, *Chem. Sci.*, 2011, **2**, 295.
- H. Agadjanian, J. J. Weaver, A. Mahammed, A. Rentsendorj, S. Bass, J. Kim, I. J. Dmochowski, R. Margalit, H. B. Gray, Z. Gross and L. K. Medina-Kauwe, *Pharm. Res.*, 2006, **23**, 367.
- H. Agadjanian, J. Ma, A. Rentsendorj, V. Valluripalli, J. Y. Hwang, A. Mahammed, D. L. Farkas, H. B. Gray, Z. Gross and L. K. Medina-Kauwe, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 6105.
- J. Y. Hwang, J. Lubow, D. Chu, J. Ma, H. Agadjanian, J. Sims, H. B. Gray, Z. Gross, D. L. Farkas, and L. K. Medina-Kauwe, *Mol. Pharm.*, 2011, **8**, 2233.
- J. Y. Hwang, D. J. Lubow, D. Chu, J. Sims, F. Alonso-Valenteen, H. B. Gray, Z. Gross, D. L. Farkas and L. K. Medina-Kauwe, *J. Control. Release*, 2012, **163**, 368.
- A. Mahammed and Z. Gross, *J. Am. Chem. Soc.*, 2005, **127**, 2883.
- I. Saltsman, A. Mahammed, I. Goldberg, E. Tkachenko, M. Botoshansky and Z. Gross, *J. Am. Chem. Soc.*, 2002, **124**, 7411.
- A. Mahammed and Z. Gross, *J. Inorg. Biochem.*, 2002, **88**, 305.
- H. Charlotte, *Nat. Rev. Drug Discov.*, 2013, **12**, 346.
- Yamaki, H.; Nakajima, M.; Shimotohno, K. W.; Tanaka, *J. Antibiot.*, 2011, **64**, 635.
- R. Bhat, S. R. Tummalapalli and D. P. Rotella, *J. Med. Chem.*, 2014, **57**, 8718.
- D. S. Hong, U. Banerji, B. Tavana, G. C. George, J. Aaron and R. Kurzrock, *Cancer Treat. Rev.*, 2013, **39**, 375.
- S. Paul and S. Mahanta, *Mol. Cell. Biochem.*, 2014, **386**, 45.
- L. Neckers and P. Workman, *Clin. Cancer Res.*, 2012, **18**, 64.
- R. R. A. Kitson and C. J. Moody, *J. Org. Chem.* 2013, **78**, 5117.
- B. L. Dutton, R. R. A. Kitson, S. Parry-Morris, S. M. Roe, C. Prodromou and C. J. Moody, *Org. Biomol. Chem.*, 2014, **12**, 1328.
- R. R. A. Kitson, C.-H. Chang, R. Xiong, H. E. L. Williams, A. L. Davis, W. Lewis, D. L. Dehn, D. Siegel, S. M. Roe, C. Prodromou, D. Ross and C. J. Moody, *Nat. Chem.*, 2013, **5**, 307.
- W.-Y. Chen, F.-R. Chang, Z.-Y. Huang, J.-H. Chen, Y.-C. Wu and C.-C. Wu, *J. Biol. Chem.*, 2008, **283**, 17184.
- P. Lim, A. Mahammed, Z. Okun, I. Saltsman, Z. Gross, H. B. Gray and J. Termini, *Chem. Res. Toxicol.*, 2011, **25**, 400.
- W. H. Lee, J. M. Lee, C. Lim, S. Kim and S. G. Kim, *Chem. Biol. Interact.*, 2013, **204**, 49.
- K. E. Thomas, A. B. Alemayehu, J. Conradie, C. Beavers, and A. Ghosh, *Inorg. Chem.*, 2011, **50**, 12844.
- P. J. Hay and W. R. Wadt, *J. Chem. Phys.*, 1985, **82**, 270.
- P. J. Hay and W. R. Wadt, *J. Chem. Phys.*, 1985, **82**, 284.
- A. D. Bochevarov, E. Harder, T. F. Hughes, J. R. Greenwood, D. A. Braden, D. M. Philipp, D. Rinaldo, M. D. Halls, J. Zhang and R. A. Friesner, *Int. J. Quantum Chem.*, 2013, **113**, 2110.
- W. A. Goddard, S.-K. Kim, Y. Li, B. Trzaskowski, A. R. Griffith and R. Abrol, *J. Struct. Biol.*, 2010, **170**, 10.
- W. B. Floriano, N. Vaidehi, G. Zamanakos, and W. A. Goddard, *J. Med. Chem.*, 2004, **47**, 56.
- S. L. Mayo, B. D. Olafson and W. A. Goddard, *J. Phys. Chem.*, 1990, **94**, 8897.
- S. S. Dong, R. Abrol and W. A. Goddard, *ChemMedChem*, 2015, **10**, 650.
- N. Nair, W. Kudo, M. A. Smith, R. Abrol, W. A. Goddard and V. P. Reddy, *Bioorg. Med. Chem. Lett.*, 2011, **21**, 3957.
- The PyMOL Molecular Graphics System, Version 1.7.4 Schrödinger, LLC.
- C. E. Stebbins, A. A. Russo, C. Schneider, N. Rosen, N.; F. U. Hartl and N. P. Pavletich, *Cell*, 1997, **89**, 239.
- G. Banumathy, V. Singh, S. R. Pavithra and U. Tatu, *J. Biol. Chem.*, 2003, **278**, 18336.

Molecular Biosystems

- 42 J. C. Phillips, R. Braun, W. Wang, J. Gumbart, E. Tajkhorshid, E. Villa, C. Chipot, R. D. Skeel, L. Kale and K. Schulten, *J. Comp. Chem.*, 2005, **26**, 1781.
- 43 W. Humphrey, A. Dalke and K. Schulten, *J. Mol. Graphics*, 1996, **14**, 33.
- 44 D.A. Case, T.A. Darden, T.E. Cheatham, III, C.L. Simmerling, J. Wang, R.E. Duke, R. Luo, R.C. Walker, W. Zhang, K.M. Merz, B. Roberts, B. Wang, S. Hayik, A. Roitberg, G. Seabra, I. Kolossváry, K.F. Wong, F. Paesani, J. Vanicek, J. Liu, X. Wu, S.R. Brozell, T. Steinbrecher, H. Gohlke, Q. Cai, X. Ye, J. Wang, M.-J. Hsieh, G. Cui, D.R. Roe, D.H. Mathews, M.G. Seetin, C. Sagui, V. Babin, T. Luchko, S. Gusarov, A. Kovalenko, and P.A. Kollman, 2010, AMBER 11, University of California, San Francisco.
- 45 C. Zipp, PhD Thesis, 2014.
- 46 J. Wang, R. M. Wolf, J. W. Caldwell, P. A. Kollman and D. A. Case, *J. Comp. Chem.*, 2004, **25**, 1157.
- 47 A. K. Rappé, C. J. Casewit, K. Colwell, W. A. Goddard and W. Skiff, *J. Am. Chem. Soc.*, 1992, **114**, 10024.
- 48 J. M. Lee, W. H. Lee, H. Y. Kay, E.-S. Kim, A. Moon and S. G. Kim, *Int. J. Cancer*, 2012, **130**, 716.
- 49 A. Mahammed and Z. Gross, *J. Porphyrins Phthalocyanines*, 2010, **14**, 911.