

MedChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Identification of new hit scaffolds by INPHARMA-guided virtual screening

Cite this: DOI: 10.1039/x0xx00000x

Justyna Sikorska,[†] Luca Codutti,[†] Lars Skjærven,[†] Bettina Elshorst,[§] Rebeca Saez-Ameneiro,[†] Andrea Angelini,[†] Peter Monecke,[§] and Teresa Carlomagno^{*†§}

Received,
Accepted

DOI: 10.1039/x0xx00000x

www.rsc.org/

Structure-based drug design (SBDD) relies on the availability of high-quality structures that describe protein-ligand interactions. INPHARMA is an NMR-based method that allows the determination of ligand binding poses to accuracy higher than 2 Å. In this work, we demonstrate that INPHARMA can be used to find novel ligand scaffolds as inhibitors of a model system protein, the cyclin dependent kinase (Cdk-2). The workflow is as follows: first, we determine the binding poses for six low-affinity fragments of Cdk-2 and use them to derive a structure-based pharmacophore. Two of the ligands show an unexpected binding mode, which differs from the one observed in crystal structures of other kinases. Second, we use the INPHARMA-generated pharmacophore for virtual screening of the ZINC database; one of the hit compounds is found to bind Cdk-2 in the low μM range and shows selectivity for Cdk-2 against kinases of other families. Our results demonstrate that INPHARMA is an efficient structure-based tool in solution to evolve low affinity fragments into hit compounds.

Introduction

The quest for new drug leads comprises the *in silico* screening of ligands libraries with the help of an experimentally derived pharmacophore. In the absence of structural information for the target protein, “ligand based” pharmacophores are generated by aligning series of ligands with known biological activity. Conversely, “structure based” pharmacophores utilize knowledge of the protein structure to predict key protein-ligand interactions. The “structure-based” approach is powerful; however, its application is limited by the availability of structures of the protein and/or of protein-ligand complexes.¹

Next to X-ray crystallography, Nuclear Magnetic Resonance (NMR) spectroscopy has emerged as a potent technique for the determination of receptor-ligand interactions. NMR offers two strategies to detect ligand binding interfaces: (1) ligand spectra in the presence of the target protein report on the ligand pharmacophore (ligand-observed experiments); (2) protein spectra in the presence of the small molecule binder provide information on the protein interacting residues (protein-observed experiments). The ligand-observed approach is broadly applicable to low-affinity ligands in complex with proteins of any size and has the advantage not to require expensive isotope labeling of the target. Among ligand-observed experiments, the STD² and waterLOGSY³ permit the identification of ligand binding epitopes, *tr*-NOEs^{4, 5} determine the bioactive conformation of the ligand, while INPHARMA⁶⁻⁸ provides access to the relative binding mode of two low-affinity ligands interacting competitively within the same cavity of the target protein.

The INPHARMA method relies on the interligand, spin-diffusion mediated, transfer of magnetization (NOE) between the two ligands L1 and L2. As the ligands are competitive

binders, such NOEs do not originate from direct transfer of magnetization between L1 and L2, but rather from a spin-diffusion process, mediated by the protons of the receptor binding pocket. The intensity of the signals depends on the specific interactions of each of the two ligands with the protein; a number of such intermolecular NOEs describe the relative orientation of the two ligands in the receptor-binding pocket (Fig. S1A). The quantitative analysis of the INPHARMA NOEs enables ranking pairs of binding orientations obtained by docking L1 and L2 to the receptor.⁶ By using a combination of INPHARMA datasets acquired for multiple ligand pairs, molecular dynamic simulations and ensemble docking, we showed that INPHARMA reliably selects the correct ligand binding modes for a set of four to five ligands. For this selection process the INPHARMA information is measured for all possible pairwise combinations and the data are fitted simultaneously (INPHARMA-STRING).⁹

This work describes the implementation of INPHARMA-STRING in a classical structure-based drug-design process, aimed at evolving low-affinity fragment ligands into a hit compound. The pipeline that we develop is applied to the cyclin dependent kinases Cdk-2. First, we determine the binding orientations of six low-affinity fragments by INPHARMA-STRING and use them to build a three-dimensional, structure-based pharmacophore model. Second, we implement the 3D-pharmacophore in virtual screening of the compounds contained in the ZINC database. The procedure enables the discovery of a new scaffold, which binds Cdk-2 with improved affinity (up to two order of magnitude with respect to the low-affinity fragments) and shows selectivity for Cdk-2 against kinases of other families. The results demonstrate for the first time the applicability of INPHARMA-STRING to the evolution

of low-affinity fragments with low specificity into promising hit compounds.

Results and discussion

NMR studies of Cdk-2 ligands.

Six weakly binding ligands were used to explore the features of the Cdk-2 binding pocket (Fig. 1). LL4, ZIP and ADO are purine derivatives; M77 and LL6 display different substitutions of an isoquinoline ring while RSA is a quinazoline derivative. The pairwise combination of the six ligands results in 15 INPHARMA pairs; however, in the analysis we used INPHARMA experiments from only 11 pairs. The combinations LL6-M77, RSA-LL6 and RSA-M77 were excluded due to extreme overlap of the resonances of the two ligands; the ADO-LL4 pair was eliminated because of the low number of INPHARMA peaks. For each pair, three NOESY spectra were acquired with different mixing times (300ms, 500ms and 800ms), resulting in a total of 33 experiments and 144 correlation peaks (Table S1).




Fig. 1 Cdk-2 inhibitors used for the INPHARMA analysis. K_d values were measured by isothermal titration calorimetry and are: LL4, 62 μM ; ZIP, 35 μM ; RSA, 208 μM ; ADO, 184 μM ; M77, 460 μM ; LL6, 286 μM .

The INPHARMA information content was distributed non-homogeneously, as readily observed for other protein-ligand systems⁹. For the purine analogues LL4 and ZIP the key structural information was contained in INPHARMA-NOEs to purine hydrogens, while the signals from the benzyl and the 3-methyl-2-butenyl fragments were not used due to the extensive degeneracy of the chemical shifts. For LL6 and M77 (Fasudil) the majority of correlations resulted from the 8-isoquinolinesulfonamide fragment; the *N*-ethyl-guanidine and hexahydro-1,4-diazepine moieties overlapped with protein

resonances between 0-4 ppm. The most challenging ligand was adenosine (ADO), with INPHARMA peaks stemming exclusively from the H-2 and H-8 of adenine. Finally, for RSA all protons delivered peaks that could be used for the analysis.

Determination of Cdk-2-ligand interactions by INPHARMA-STRING.

According to the workflow proposed in⁹ the INPHARMA-STRING data were used to rank *in silico* generated models of the protein-ligand complexes (Fig. S1B). For the docking, we collected three representative structures of Cdk-2 (1HCK: ATP-bound; 1H1R: cyclin-bound; 2VTM: cyclin-free), and performed molecular dynamics (MD) simulations to explore the conformational space around the starting models. The resulting ensemble of MD-generated conformers was reduced by clustering analysis, providing 5 receptor structures from each simulation. The six ligands were subsequently docked using Glide¹⁰ and Surflex¹¹ to the 15 receptor structures and the best 10 scoring docking poses of each docking program were retained (generating a total of 150 docking poses per ligand per program = 300 docking poses per ligand). Each of the resulting structures was further subjected to a short MD simulation: five structures per trajectory were selected (Supplementary Information), resulting in 1500 binding modes per ligand. The binding free energy of each complex structure was estimated using MM/GBSA, and the 50% highest energy conformations were excluded from further analysis. The remaining binding poses (750 per ligand) were combined pairwise to calculate the theoretical INPHARMA-NOEs using the in-house program described before¹¹ and the K_d for each ligand measured by calorimetric analysis. The INPHARMA-NOEs predicted for each ligand pair were compared to the experimental ones by means of the Pearson correlation coefficient *R* (INPHARMA score); subsequently, the INPHARMA-STRING algorithm was implemented for selection of the binding modes that are uniquely in agreement with the experimental data.⁹

Briefly, the INPHARMA-STRING algorithm identifies consensus binding modes among poses with top-ranked INPHARMA scores. It returns a string of six docking modes (one per ligand) with high INPHARMA scores in all 11 combinations used for the selection. During the search, the number of high-ranking docking modes is gradually increased until the consensus string is found.

For our experimental system, the algorithm reached a consensus among the best-ranked 3.6% docking modes (Fig. 2 and S2). For LL4 and ZIP, the INPHARMA-STRING selected poses are close to those observed by crystallography (all atoms root mean square deviation, RMSD, of 0.8-1.4 Å and 1.4 Å, respectively; Fig. 2), thus providing a validation of the algorithm. To verify the stability of the selection, we analyzed the poses selected by INPHARMA-STRING up to 5% of the best-ranked docking modes and found that the poses were similar to each other, with an average RMSD <1.5 Å (Fig. S3). LL4 and ZIP showed hydrogen bonding to Glu-81 and Leu-83 in the Cdk-2 hinge region, as well as hydrophobic interaction

with Val-18 (Fig. 3). For LL4 an additional interaction to Lys-33 was occasionally observed (Fig. 3).




Fig. 2 Binding modes of Cdk-2 ligands identified by INPHARMA-STRING (consensus string in the 3.6% best-ranked poses). The INPHARMA-selected binding poses of ZIP, LL4 and ADO are represented as pink, blue and yellow structures (tubes) overlaid to their respective X-ray structures (wires); PDB codes; ZIP, 2EXM; ADO, 1HCK. The M77 and LL6 INPHARMA-selected binding poses are shown in orange and purple, and overlaid to the orientation of M77 bound to PKA (1Q8W.pdb). The INPHARMA-selected binding pose of RSA, shown in black, is compared to that of the 4-[3-hydroxyanilino]-6,7-dimethoxyquinazoline in complex with Cdk-2 (1D18.pdb).

The selection of the M77 and LL6 binding poses was also consistent among the best-ranked 3.5-5% docking modes (Fig. 2 and S3). For these ligands, no crystallographic structure is available in complex with Cdk-2, probably due to their low-affinity. When we compare the INPHARMA-STRING selected poses to the X-ray structure of M77 in complex with protein kinase A (PKA),¹² we observe clear differences (Fig. 2).¹³ Here, M77 and LL6 take advantage of the larger Cdk-2 binding site and occupy the frontal cavity (Fig. S4), as also confirmed by STD data (Fig. S5). The 8-isoquinolinesulfonamide fragment of both M77 and LL6 forms a hydrogen bond to Leu-83 as well as hydrophobic interactions with Ile-10 and Phe-82 (Fig. 3). In

support of this unexpected interaction mode, binding to the frontal cavity of Cdk-2 had been previously utilized for the design of compounds that, like flavopiridol,¹⁴ have higher selectivity for Cdk-2 in comparison to PKA.

For adenosine (ADO) two binding poses were selected by INPHARMA among the 5% best-ranked docking modes (Fig. 2 and S3). The first binding pose differs by 180° from the binding pose of ATP to Cdk-2 (X-ray structure 1HCK.pdb, Fig. 2), while the second one corresponds to the ATP binding mode. The INPHARMA score did not allow discrimination of the two orientations of ADO; this can be explained by the fact that the INPHARMA NOEs of ADO stem primarily from protons H2 and H8 of the aromatic ring and are therefore insensitive to 180° rotations around the H2-H8 axis.




Fig. 3 Schematic representation of the protein-ligand interactions for all analyzed compounds. We show all binding interactions observed in the binding poses selected by INPHARMA-STRING in the 5% top-ranked binding modes. The binding site residues within 4 Å from the ligand are indicated as spheres, while hydrogen bonds are represented by red arrows. Blue, hydrogen bond donor; red, hydrogen bond acceptor; green, hydrophobic interaction.

The pose of RSA selected by INPHARMA-STRING has a prominent hydrogen bond interaction between the N-3 of the quinazoline ring and Lys-33, as well as the methylamine of

RSA and Asp-145 (Fig. 3). An additional π - π interaction can be formed between the pyrimidine ring of RSA and Phe-80. However, when a larger population of binding poses was examined (5%), only the hydrogen bond between the N-3 of the quinazoline ring to Lys-33 was consistent through all binding modes. Noticeably all selected poses are similar to those of the N-phenylquinazolin-4-amine analogues (the closest analogue of RSA that has been co-crystalized with Cdk-2, 1DI8.pdb); however these compounds interact with the Cdk-2 binding site through an hydrogen bond between the N-1 of the quinazoline ring and Leu-83.¹³

Overall, the INPHARMA-selected poses show receptor–ligand interactions that are in accord with previously published structures of known Cdk-2 inhibitors, thus validating the robustness of the selection.¹⁵

Epitope mapping of selected Cdk-2 ligands with Saturation Transfer Difference (STD).

To independently validate the INPHARMA-selected binding modes, we applied saturation transfer difference (STD)² experiments to ZIP, LL4, LL6, M77 and RSA in the presence of Cdk-2 (Fig. S5). For LL4, ZIP and RSA the strongest saturation effects were observed for purine H2 and quinazoline H2, H7, H8, in good agreement with the INPHARMA-selected poses, which locate these protons in close proximity to Ile-10 or deep inside the binding cavity. For LL6 and M77, the STD data exclude an orientation of the ligand similar to that of M77 in PKA, for which the STD intensities of the buried protons H6, H7 should be higher than those of the solvent exposed protons H3, H4. Instead, we observe STD signals of medium intensity for H3, H4 and of weak intensity for H6, H7, which is in agreement with the INPHARMA-selected pose.

Structure-based pharmacophore generation, validation and virtual screening.

With the binding modes of these fragments in our hands, we set out to generate a common structure-based pharmacophore, to be used for virtual screening of the ZINC database.

First we generated the individual pharmacophores of ZIP, LL4, M77, LL6 and RSA using the following protocol. We analyzed all intermolecular interactions observed in the binding poses selected by INPHARMA-STRING in the top-ranking 5% structural pairs with LigandScout3.12¹⁶. Of these interactions we retained only those that were consistently present in all binding modes of the ligand under consideration. For ZIP and LL4 we decreased the size of the spheres describing the hydrogen bond interaction to Glu-81 and Leu-83 by 0.3 Å to increase pharmacophore specificity. For LL6, we excluded all interactions originating from the guanidinium group, due to the lack of experimental INPHARMA data stemming from this part of the molecule, which contains only exchangeable hydrogens. Furthermore, for M77, which is the weakest binder, we retained only the pharmacophoric features common to LL6.

The common-structure-based pharmacophore (CSB) of all five ligands was generated by the alignment of reference points (defined as amino acids residues of the binding site) followed by a merge of all pharmacophoric features. The final pharmacophore consisted of six interactions: two H-bond acceptors (HBA), one H-bond donor (HBD), three hydrophobic contacts (HY) as well as excluded volume spheres representing the surface of the Cdk-2 binding pocket (Fig. 4, S6 and S7). The HBA1 and HBD represent two known key interactions of Cdk-2 inhibitors with residues of the hinge region: the backbone NH of Leu-83 and the backbone oxygen of Glu-81. The HBA2 feature results from the interaction between the quinazoline nitrogen of RSA and the side chain amino group of Lys-33. Because of the high flexibility of Lys-33 in the Cdk-2 binding site, we also constructed a pharmacophore model where we excluded HBA2 and named it reduced-structure-based pharmacophore (RSB). The HY1 and HY2 contacts result from the interaction of the isoquinoline ring of LL6 and M77 with Ile-10 and Phe-82, while HY3 originates from the interaction of the aliphatic and aromatic fragments of ZIP and LL4, respectively, with Val-18.




Fig. 4 The common structure-based (CSB) pharmacophore used in virtual screening. H-bond acceptors (HBA) red arrows and spheres; H-bond donors (HBD) green arrows and spheres; hydrophobic features (HY), yellow spheres. The radius of the sphere represents the space where the feature can be localized. The excluded volumes are not indicated for clarity. Protein residues 80-82 (hinge region) are shown in cyan sticks.

The pharmacophoric features identified in this study are part of the comprehensive pharmacophore map derived for Cdk-2 from all Cdk-2 binders.¹⁵ This confirms the ability of INPHARMA to generate high quality binding modes for fragment ligands, which can be further used to generate pharmacophore models. Among all six features, HBA1, HBD, HY2 and HY3 are present in > 25% of the crystal structures of Cdk-2 in complex with ligands¹⁵, while HBA2 and HY1 are found with 22% occurrence.¹⁵

The ability of the CSB and RSB pharmacophore models to identify active compounds was estimated for a collection of entries composed of active and decoy molecules. The pool of active compounds consisted of 115 entries from the ChEMBL database with IC_{50} below 500 nM, while the 2438 decoys were generated with the program DUD-E (<http://dude.docking.org/>)¹⁷ from 33 active compounds. The performance of the pharmacophores was quantified by the enrichment factor (EF) and the area under the receiver operating characteristic curve (ROC) obtained for the top 1.5% of the scored database.¹⁸ The EF represents the ability of the pharmacophore to accurately identify active hits with respect to a random selection.¹⁸ The area under the ROC curve (AUC) is a function of both specificity and sensitivity and indicates the probability of ranking a randomly chosen active compound higher than a randomly chosen decoy. For the CSB pharmacophore LigandScout 3.12¹⁶ identified 2 compounds from the 115 active ligands and 2 from 2438 decoys, resulting in an $EF_{1.5\%} = 11.1$ and $AUC_{1.5\%} = 0.92$. With the RSB we selected 12 out of the 115 active compounds and 5 out of the decoys pool yielding $EF_{1.5\%} = 14$ and $AUC_{1.5\%} = 0.95$. Both pharmacophores were used to screen the 3D ZINC database,¹⁹ resulting in 1343 hits for CSB and 1206 hits for RSB (Fig. S8). The hits were docked to two Cdk-2 structures and the binding poses were ranked with both GlideScore²⁰ and X-score²¹. The two Cdk-2 structures used for the docking had been selected by INPHARMA-STRING in complex with ligands LL6 and RSA and represent two very different orientations of Lys-33 in the binding site. Those compounds were retained which ranked among the best 100 of both scores and fulfilled at least one H-bond and two hydrophobic interactions of the CSB or RSB pharmacophores (Fig. S9). This selection procedure resulted in 28 and 16 ligands from the CSB and the RSB pharmacophore, respectively (Table S2).

Evaluation of identified hit compounds by NMR.

Of the 44 compounds selected by *in silico* screening, we purchased 11 commercially available ones, representing 5 distinct chemical skeletons (Table S2, Fig. S8 and S10). Among them, ZINC20085956, and ZINC06781696 were excluded from further evaluation due to poor solubility. The remaining compounds were subjected to trNOE and STD experiments in the presence of Cdk-2 (Fig. S10). The 1,2,4-triazole-3-thiones derivatives did not show trNOE. Weak saturation transfer effects were observed only for ZINC02808739.

Five compounds exhibited solubility < 10 μ M; for these ligands, we could not record direct trNOE or STD experiments. Instead, we measured competition STD against M77 ($K_d = 460$ μ M) at the highest possible ligand concentration. A 40% decrease of the intensity of M77 H-8 was observed in the presence of 2.5 μ M of HDS029 (ZINC13682420). Interestingly, HDS029 had been previously identified as a potent inhibitor of the proteins ErbB1, ErbB2 and ErbB4 belonging to the receptor tyrosine kinase family.

Analogously, among the derivatives of 1H-pyrazolo[3,4-B]pyridin-3(2H)-one and benzimidazole, ZINC20085954 caused 45% decrease of the M77 H-8 peak at 2.5 μ M.

Overall, from 11 purchased compounds only 9 were sufficiently soluble for further analysis. Among those 3 were identified as Cdk-2 binders though STD experiments, resulting in a hit rate of 33% (defined as a percentage of Cdk-2 binding compounds among all purchased compounds). To independently confirm that ZINC20085954 and ZINC02808739 are kinase inhibitors with a novel chemical skeleton, both compounds were subjected to the kinase phosphorylation assay.




Fig. 5. (A) Chemical structure of ZINC20085954 and a representative of ALK inhibitors. (B) The proposed binding of ZINC20085954 with interacting protein residues shown in grey. The hydrophobic surface is represented in a brown/blue scale and the hydrogen bond to the hinge region is shown as red dashed line.

Kinase phosphorylation assay.

We tested ZINC20085954 and ZINC02808739 at Reaction Biology Corporation, using their kinase assay platform panel²². The ability of the ligand to inhibit the catalytic activity of Cdk-2/cyclin A, Cdk-2/cyclin E, PKA, and ALK kinases was evaluated at the highest soluble concentration of the ligands (Table 1). In addition, the binding affinity of ZINC20085954 and ZINC02808739 to Cdk-2/cyclin A was evaluated measuring the IC_{50} values. ZINC20085954 showed the strongest inhibition of Cdk-2 activity with IC_{50} of 2.6 μ M. In

addition, the compound displayed selectivity for Cdk-2 against PKA, albeit the fact that the fragments (Fig. 1) were not chosen as selective Cdk-2 binders (indeed M77 binds to PKA stronger than to Cdk-2). ZINC20085954 belongs to the family of the pyrazolo[3,4-B] pyridine derivatives, which were published as inhibitors of the ALK²³ during the course of this work (Fig. 5A); however, ALK preserved 87% of activity in the presence of 5 μ M of ZINC20085954, confirming its selectivity for Cdk-2/cyclin A (Table 1).

Binding mode of ZINC20085954 to Cdk-2.

To better understand the selectivity of ZINC20085954 towards Cdk-2, we generated models of the Cdk-2–ZINC20085954 complex by ensemble docking. The top 10 docking results converged to an orientation that fulfils four of the five features of the RSB pharmacophore (Fig. 5B and S9). ZINC20085954 entertains three hydrophobic interactions with Phe-82/Ile-10, Ala-31 and Val-18, and a hydrogen bond with the backbone nitrogen of the hinge region residue Leu-83. The ligand occupies both the frontal and the back cavities of Cdk-2, which explains its selectivity against PKA, lacking the frontal cavity (Fig. S4).

In addition to the interactions described by the pharmacophore, the bromine atom of ZINC20085954 makes contacts with the phenyl ring of the gatekeeper Phe-80, while the C5-methyl forms a CH- π interaction with Phe-82. These non-covalent interactions can contribute significantly to the affinity of ZINC20085954 for Cdk-2, and can rationalize the much lower activity of the compound against ALK, where Phe-80 and Phe-82 are substituted by leucines (Table 1).

Conclusions

In this work we propose an alternative route for structure based drug design in the absence of a high-resolution x-ray structures of the receptor-ligand complexes. The INPHARMA approach is especially relevant to the design of new leads for proteins that are challenging to crystallize, either by themselves or in complex with the ligands of interest; it also offers an attractive alternative for obtaining high-resolution structural information on protein-ligand interactions without extensive optimization of crystallographic conditions or assignment of NMR protein resonances. In this study, the method allowed us to identify significant differences in the binding mode of purine and isoquinoline derivatives to Cdk-2: while the hydrogen bond between one heterocyclic nitrogen and the NH of Leu-83 is formed by both class of ligands, the isoquinoline derivatives lack the possibility to form an additional hydrogen bond to the carbonyl of Glu-81. Consequently, the heterocyclic ring flips and occupies the frontal open cavity of Cdk-2, rather than the back cavity. The combination of the pharmacophoric features of both the purine and isoquinoline derivatives allows for the identification of a new hit molecule, which is selective for Cdk-

2 against PKA, lacking the frontal cavity. The novel scaffold can be used for further optimization into a high-activity ligand.

Acknowledgements

The authors acknowledge support from the BMBF (grant 0315870B) and the DFG (grant CA 294/6-1). We thank Vladimir Rybin and the EMBL Protein Expression and Purification Core Facility for performing ITC experiments and data analysis.

Notes and references

[†] EMBL, Structural and Computational Biology Unit, Meyerhofstraße 1, D-69117 Heidelberg, Germany

[§] Helmholtz Zentrum für Infektionsforschung, Inhoffenstrasse 7 Braunschweig, Germany

[§] Sanofi-Aventis Deutschland GmbH R&D LGCR/Structure, Design & Informatics, Industriepark Höchst, Bldg. G877, D-65926 Frankfurt am Main, Germany

Electronic Supplementary Information (ESI) available: computational and experimental procedures, additional figures and tables. See DOI: 10.1039/b000000x/

- G. L. Wilson and M. A. Lill, *Future Medicinal Chemistry*, 2011, **3**, 735-750.
- M. Mayer and B. Meyer, *Angewandte Chemie International Edition*, 1999, **38**, 1784-1788.
- C. Dalvit, P. Pevarello, M. Tatò, M. Veronesi, A. Vulpetti and M. Sundström, *J Biomol NMR*, 2000, **18**, 65-68.
- P. Balaram, A. A. Bothner-By and J. Dadok, *Journal of the American Chemical Society*, 1972, **94**, 4015-4017.
- G. M. Clore and A. M. Gronenborn, *Journal of Magnetic Resonance (1969)*, 1983, **53**, 423-442.
- J. Orts, C. Griesinger and T. Carlomagno, *Journal of Magnetic Resonance*, 2009, **200**, 64-73.
- J. Orts, J. Tuma, M. Reese, S. K. Grimm, P. Monecke, S. Bartoschek, A. Schiffer, K. U. Wendt, C. Griesinger and T. Carlomagno, *Angewandte Chemie International Edition*, 2008, **47**, 7736-7740.
- V. M. Sánchez-Pedregal, M. Reese, J. Meiler, M. J. J. Blommers, C. Griesinger and T. Carlomagno, *Angewandte Chemie International Edition*, 2005, **44**, 4172-4175.
- L. Skjærven, L. Codutti, A. Angelini, M. Grimaldi, D. Latek, P. Monecke, M. K. Dreyer and T. Carlomagno, *Journal of the American Chemical Society*, 2013, **135**, 5819-5827.
- R. A. Friesner, R. B. Murphy, M. P. Repasky, L. L. Frye, J. R. Greenwood, T. A. Halgren, P. C. Sanschagrin and D. T. Mainz, *Journal of Medicinal Chemistry*, 2006, **49**, 6177-6196.
- E. Kellenberger, J. Rodrigo, P. Muller and D. Rognan, *Proteins: Structure, Function, and Bioinformatics*, 2004, **57**, 225-242.
- C. Breitenlechner, M. Gaßel, H. Hidaka, V. Kinzel, R. Huber, R. A. Engh and D. Bossemeyer, *Structure*, 2003, **11**, 1595-1607.

13. A. K. Ghose, T. Herbertz, D. A. Pippin, J. M. Salvino and J. P. Mallamo, *Journal of Medicinal Chemistry*, 2008, **51**, 5149-5171.
14. A. Huwe, R. Mazitschek and A. Giannis, *Angewandte Chemie International Edition*, 2003, **42**, 2122-2138.
15. J. Zou, H.-Z. Xie, S.-Y. Yang, J.-J. Chen, J.-X. Ren and Y.-Q. Wei, *Journal of Molecular Graphics and Modelling*, 2008, **27**, 430-438.
16. G. Wolber and T. Langer, *Journal of Chemical Information and Modeling*, 2004, **45**, 160-169.
17. M. M. Mysinger, M. Carchia, J. J. Irwin and B. K. Shoichet, *Journal of Medicinal Chemistry*, 2012, **55**, 6582-6594.
18. *Pharmacophores and Pharmacophore Searches*, WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim, 2006.
19. J. J. Irwin, T. Sterling, M. M. Mysinger, E. S. Bolstad and R. G. Coleman, *Journal of Chemical Information and Modeling*, 2012, **52**, 1757-1768.
20. T. A. Halgren, R. B. Murphy, R. A. Friesner, H. S. Beard, L. L. Frye, W. T. Pollard and J. L. Banks, *Journal of Medicinal Chemistry*, 2004, **47**, 1750-1759.
21. R. Wang, L. Lai and S. Wang, *J Comput Aided Mol Des*, 2002, **16**, 11-26.
22. T. Anastassiadis, S. W. Deacon, K. Devarajan, H. Ma and J. R. Peterson, *Nat Biotech*, 2011, **29**, 1039-1045.
23. R. Rabot, K. Bedjeguelal, E. B. Kaloun, P. Schmitt, N. Rahier, P. Mayer and E. Fournier, Google Patents, 2014.

Ligand	Single dose concentration	Enzyme activity				IC50
		ALK	Cdk-2 /cyclin A	Cdk-2/cyclin E	PKA	Cdk-2/cyclin A
ZINC20085954	5 μ M	87 %	35 %	95 %	115 %	2.6 μ M
ZINC02808739	100 μ M	-	69 %	96 %	99 %	1000 μ M

Table 1. Inhibitory activity of virtual screening hits against various kinases, measured with ^{33}P – radiolabeled assay.