

# MedChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

# Discovery of high affinity inhibitors of *Leishmania donovani* N-myristoyltransferase

Mark D. Rackham,<sup>1, 4, §</sup> Zhiyong Yu,<sup>1, 5, §</sup> James A. Brannigan,<sup>2</sup> William P. Heal,<sup>1, 6</sup> Daniel Paape,<sup>3, 7</sup> K. Victoria Barker,<sup>1, 8</sup> Anthony J. Wilkinson,<sup>2</sup> Deborah F. Smith,<sup>3</sup> Robin J. Leatherbarrow<sup>1, 9</sup> and Edward W. Tate<sup>1\*</sup>

<sup>1</sup> Department of Chemistry, Imperial College London, South Kensington Campus, London, SW7 2AZ, U.K.

<sup>2</sup> Structural Biology Laboratory, Department of Chemistry, University of York, York, YO10 5DD, U.K.

<sup>3</sup> Department of Biology, University of York, York, YO10 5DD, U.K.

<sup>4</sup> Current address: GlaxoSmithKline, Gunnels Wood Road, Stevenage, SG1 2NY, U.K.

<sup>5</sup> Current address: International Discovery Service Unit, WuXi AppTec, Shanghai, 200131, China.

<sup>6</sup> Current address: Department of Chemistry, Kings College London, London SE1 1UL, U.K.

<sup>7</sup> Current address: Wellcome Trust Centre for Molecular Parasitology, Institute of Infection, Immunity and Inflammation, University of Glasgow, Glasgow, U.K.

<sup>8</sup> Current address: Dehns, Aspect House, 84-87 Queens Road, Brighton, BN1 3XE, U.K.

<sup>9</sup> Current address: Liverpool John Moores University, Egerton Court, 2 Rodney Street, Liverpool, L1 2UA, U.K.

§MDR and ZY contributed equally to the preparation of this manuscript

## ABSTRACT

*N*-Myristoyltransferase (NMT) is a potential drug target in *Leishmania* parasites. Scaffold-hopping from published inhibitors yielded the serendipitous discovery of a chemotype selective for *Leishmania donovani* NMT; development led to high affinity inhibitors with excellent ligand efficiency. The binding mode was characterised by crystallography and provides a structural rationale for selectivity.

## INTRODUCTION


Leishmaniasis are a spectrum of neglected tropical diseases resulting from the infection with parasites of the genus *Leishmania* spp., transmitted by the bite of the female *Phlebotomine* sandfly to the human host. The disease manifests in one of three forms (cutaneous, mucocutaneous and visceral)<sup>1</sup> and is responsible for an estimated 60,000 deaths and in excess of 500,000 new cases per annum.<sup>2</sup> The majority of fatalities stem from visceral leishmaniasis, which is predominately the result of infection by *Leishmania donovani* (Ld).<sup>3</sup> As with many neglected diseases, the leishmaniasis are most prevalent in developing and newly industrialized nations<sup>4</sup> and although treatments exist, the majority suffer from significant levels of resistance and toxicity.<sup>5</sup> *Leishmania* spp. infections remain challenging to treat chemotherapeutically; many currently marketed drugs display EC<sub>50</sub> values in the low  $\mu$ M range in *in vitro* models,<sup>5,6</sup> orders of magnitude higher than the potency typically targeted in an anti-infective drug discovery project. In order to combat the continued evolution of the parasite in developing resistance, new chemotherapies functioning by novel mechanisms of action are essential.

*N*-Myristoyltransferase (NMT) is a ubiquitous eukaryotic enzyme responsible for the attachment of the long chain fatty acid myristate to the N-terminus of substrate proteins. This

modification is crucial in many biological processes,<sup>7-10</sup> and NMT has been implicated as a highly promising drug target in fungal<sup>11</sup> and parasitic infections,<sup>12-14</sup> as recently demonstrated in human malaria parasites.<sup>15</sup> In the specific context of leishmaniasis, NMT has been validated genetically in *Leishmania major* (Lm),<sup>16</sup> and through chemical proteomic approaches in Ld,<sup>17</sup> and selective small molecule inhibitors of this drug target are highly desirable.

## RESULTS AND DISCUSSION

We have previously reported the discovery of LdNMT inhibitors by high-throughput screening,<sup>18-21</sup> as well as the discovery of *Plasmodium falciparum* (Pf) and *Plasmodium vivax* (Pv) NMT inhibitors by both high-throughput screening<sup>22</sup> and piggy-back/lead-hopping approaches.<sup>23-25</sup> Throughout this work, all compounds were routinely screened against all three parasitic enzymes (LdNMT, PfNMT and PvNMT) and *Homo sapiens* (Hs) NMT, where selectivity over the human orthologue was desired to minimise any potential toxicity from modulating endogenous myristoylation. 2,3-Substituted benzo[*b*]thiophenes were discovered as ligand-efficient inhibitors of PfNMT, which incidentally displayed selectivity over LdNMT (Figure 1).<sup>25</sup>




$K_i$  ( $\mu\text{M}$ ):<sup>a</sup> 0.83 (PfNMT)  
 0.08 (PvNMT)  
 3.20 (HsNMT)<sup>b</sup>  
 5.82 (LdNMT)

**Figure 1.** NMT affinity profile for 2,3-Substituted Benzo[*b*]thiophene **1**: <sup>a</sup> Enzyme apparent  $K_i$  values are calculated from the  $\text{IC}_{50}$  values using the Cheng–Prusoff equation (see Supporting Information) to enable cross-comparisons.  $\text{IC}_{50}$  values were the mean of two or more independent determinations. Standard deviation is within 20% of reported  $\text{IC}_{50}$ . <sup>b</sup>HsNMT affinities reported in this work refer to HsNMT1; no significant difference in inhibition was observed between HsNMT1 and HsNMT2 isoforms.<sup>18</sup>

During studies to reduce the molecular weight and lipophilicity of **1** in the further development of the series against plasmodial NMT, the bicyclic benzo[*b*]thiophene of **1** was truncated to a monocyclic scaffold (Scheme 1).




### Scheme 1: Synthesis of Truncated Inhibitor **5**<sup>a</sup>



<sup>a</sup>Reagents and Conditions: (a) 1-Boc-4-piperidinol, DIAD,  $\text{PPh}_3$ , THF, rt, 4 h, 94%; (b) NaOH, MeOH/ $\text{H}_2\text{O}$ , 50°C, 2 h, 90%; (c) (3-methoxyphenyl)methanol, EDCI, HOBt, DIPEA, MeCN, rt, 18 h; (d) 10% TFA in DCM ( $v/v$ ), rt, 2 h, 9% over 2 steps.

The affinity spectrum of compound **5** is strikingly distinct from that observed with benzo[*b*]thiophene **1** (Table 1). The activity against human and both *Plasmodium* enzymes drops by almost two orders of magnitude, whereas the affinity against LdNMT improves 8-fold, a significant improvement when considering the loss of 4 heavy atoms and attendant reduction in lipophilicity.

**Table 1.** Enzyme affinity and parasitic LE for mono- and bicyclic analogues **1**, **5** and **6**

no.	Structure	cLogP <sup>a</sup>	PfNMT		PvNMT		LdNMT		HsNMT K <sub>i</sub> (μM)
			K <sub>i</sub> (μM)	LE <sup>b</sup>	K <sub>i</sub> (μM)	LE	K <sub>i</sub> (μM)	LE	
<b>1</b>		4.0	0.83	0.30	0.08	0.35	5.82	0.28	3.20
<b>5</b>		2.9	>100	-	6.50	0.30	0.73	0.35	>100
<b>6</b>		3.0	6.00	0.29	2.42	0.31	0.28	0.36	>100

<sup>a</sup>cLogP values calculated with ChemAxon; <sup>b</sup>LE =  $[-\log(K_i)](1.374)/(\text{no. of heavy atoms})$ .

Based on this affinity profile and structural analysis of the binding mode of **1** in PvNMT,<sup>25</sup> it became clear that the monocyclic thiophene scaffold was unlikely to be a suitable chemotype for *Plasmodium* NMT inhibitors. However, **5** displays a very promising profile as a LdNMT inhibitor with excellent selectivity and LE, as well as sub-μM enzyme affinity. **5** was selected for further development, with the aim of discovering a new lead series for LdNMT inhibitors.


Although heterocycles are often preferred as bioisosteres for a phenyl ring, thiophene has been associated with cytochrome P450 inhibition and generally low “developability” during optimisation.<sup>26</sup> Replacing the thiophene with a phenyl was deemed preferable, and provided a

cost-effective synthetic route and versatility to variations around the scaffold. Pleasingly, this modification from thiophene to phenyl produced a further 3-fold improvement in enzyme affinity, whilst maintaining physicochemical properties and selectivity (Table 1).

Interestingly, compounds with a phenyl scaffold proved amenable for development into high affinity *Plasmodium* NMT inhibitors – this work shall be reported elsewhere.<sup>27</sup>


Compound **6** has a promising affinity profile, but further development of the series would require removal of the potentially labile ester moiety. We have previously reported the successful replacement of the ester moiety with an oxadiazole in related series,<sup>24,28</sup> the same bioisosteric replacement was implemented in the present series, with the aim of producing a drug-like series for further development. The relatively straightforward synthetic scheme (Scheme 2) enabled facile synthesis of a variety of analogues, thereby allowing investigation of the preferred substituents on this ring system.

**Scheme 2.** Synthesis of Oxadiazole Derivatives **10a-k**<sup>a</sup>




<sup>a</sup>Reagents: (a) BnBr, K<sub>2</sub>CO<sub>3</sub>, DMF, rt, 2 h, 72-89%; (b) DIAD, PPh<sub>3</sub>, *N*-Boc-4-OH piperidine, THF, rt, 4 h; (c) NaOH, MeOH/H<sub>2</sub>O, 50°C, 2 h, 80-95% over two steps; (d) NH<sub>2</sub>OH, EtOH, 80°C, 6 h, 98%; (e) i). EDCI, HOBT, **9**, DIPEA, CH<sub>3</sub>CN, rt, 4 h; ii). 0.5N NaOH, rt, 0.5 h; (f) 10% TFA in DCM, rt, 2 h, 20-60% over 2 steps.

**Table 2.** Enzyme affinity and cellular potency for inhibitors based on a phenyl scaffold



No.	R	cLogP <sup>a</sup>	LdNMT		HsNMT K <sub>i</sub> (μM)	Ld EC <sub>50</sub> (μM) <sup>c</sup>	Macrophage LD <sub>50</sub> (μM) <sup>d</sup>
			K <sub>i</sub> (μM)	LE <sup>b</sup>			
<b>10a<sup>e</sup></b>	H	3.4	0.06	0.37	1.24	> 30	-
<b>10b</b>	3-OMe	3.2	4.02	0.26	7.23	-	-
<b>10c</b>	3-Me	3.9	1.43	0.29	7.44	-	-
<b>10d</b>	4-Me	3.9	5.18	0.26	4.64	-	-
<b>10e<sup>e</sup></b>	4-OMe	3.2	3.04	0.26	0.86	-	-
<b>10f<sup>e</sup></b>	4-Cl	4.0	0.36	0.32	0.45	-	-
<b>10g<sup>e</sup></b>	4-F	3.5	0.03	0.37	0.59	10.2	29.5



<b>10h</b>	4-Br	4.2	0.68	0.30	0.26	-	-
<b>10i</b>	5-Me	3.9	0.24	0.32	0.76	-	-
<b>10j</b>	5-Cl	4.0	0.02	0.38	0.07	6.7	18.0
<b>10k</b>	5-OMe	3.2	0.05	0.35	0.76	13.5	27.9
<b>No.</b>	<b>Structure</b>	<b>cLogP</b>	<b>LdNMT <math>K_i</math> (<math>\mu</math>M)</b>	<b>LE</b>	<b>HsNMT <math>K_i</math> (<math>\mu</math>M)</b>	<b>Ld <math>EC_{50}</math> (<math>\mu</math>M)</b>	<b>Macrophage <math>LD_{50}</math> (<math>\mu</math>M)</b>
<b>13</b>		1.8	0.01	0.39	0.02	> 50	>90

<sup>a</sup>cLogP values calculated with ChemAxon; <sup>b</sup>LE =  $[-\log(K_i)] * 1.374 / (\text{no. of heavy atoms})$ ; <sup>c</sup>

*Ex vivo* *L.donovani* amastigotes; <sup>19 d</sup> LD<sub>50</sub> is an indication of general toxicity of a compound;

Moving from the ester linker in **6** to oxadiazole in **10a** produced a ~4-fold improvement in enzyme affinity alongside removal of the ester moiety. This compound displays measurable affinity against HsNMT but possesses selectivity of 20-fold in favour of the parasite enzyme, constituting a highly promising scaffold from which to investigate substituent effects on the phenyl ring (<sup>a</sup>Reagents: (a) BnBr, K<sub>2</sub>CO<sub>3</sub>, DMF, rt, 2 h, 72-89%; (b) DIAD, PPh<sub>3</sub>, *N*-Boc-4-OH piperidine, THF, rt, 4 h; (c) NaOH, MeOH/H<sub>2</sub>O, 50°C, 2 h, 80-95% over two steps; (d) NH<sub>2</sub>OH, EtOH, 80°C, 6 h, 98%; (e) i. EDCI, HOBt, 9, DIPEA, CH<sub>3</sub>CN, rt, 4 h; ii. 0.5N NaOH, rt, 0.5 h; (f) 10% TFA in DCM, rt, 2 h, 20-60% over 2 steps. ).

Comparison of **10c**, **10d** and **10i** shows the effect of scanning a methyl substituent around the 3-, 4- and 5- positions of the scaffold. In all cases this variation is less active than the unsubstituted parent **10a**, although among these three analogues the 5-substituted **10i** displays the highest affinity. This pattern repeats when comparing the methoxy-substituted **10b**, **10e**


and **10k**, indicating that the 5-position is preferred for substitution. 5-OMe (**10k**) provides improved potency over 5-Me (**10i**), and this trend continues on moving to the 5-Cl substituent, identifying **10j** as a high affinity LdNMT inhibitor with a  $K_i$  of 0.02  $\mu\text{M}$ .

The binding mode of **10j** to *Leishmania major* NMT (98% sequence identity to LdNMT with closely superimposable ligand binding sites)<sup>14,29</sup> was confirmed by determination of the crystal structure of the ternary complex of LmNMT, myristoyl-CoA and **10j** (PDB ID: 5a27). The inhibitor binds in the expected site adjacent to the enzyme's C-terminal carboxylate to which its piperidine forms an ion pairing interaction. This and other key interactions are conserved in a related series in PvNMT.<sup>24</sup> Surprisingly, in addition to the observed interactions between the protein and the ligand, the crystal structure revealed a ring-opened derivative of the 1,2,4-oxadiazole, rather than the desired heterocycle (see Figure S1, S2). However, proton NMR spectroscopy and high resolution mass spectrometry confirmed that the desired oxadiazole as the synthetic product, furthermore no degradation of **10j** was observed when this molecule was incubated at 37 °C over a wide pH range (pH 1-10, data not shown) over a period of 24 hours. This 'ring opening' phenomenon has been observed in related compounds within the program,<sup>27</sup> although to our knowledge, it has not been reported elsewhere. The origin of the ring-opened species observed in the crystals remains unclear, and further investigation is underway to determine the cause of this discrepancy (such as potential radiation damage). As characterisation of the compounds was consistent with the proposed oxadiazoles, the development of the series was continued.

A selection of compounds with LdNMT  $K_i < 0.1 \mu\text{M}$  (**10a**, **10g**, **10j**, **10k**) were tested against *ex-vivo* amastigotes, the clinically-relevant life cycle stage of the parasite, to determine the ability of these molecules to inhibit parasite growth. These molecules displayed a narrow (within 3-fold) therapeutic window against the macrophage host *in vitro*, and it was unclear whether the observed anti-parasitic activity is a result of NMT inhibition or non-


specific toxicity. Given the relatively high cLogP of the tested compounds (>3.5), it was hypothesised that reducing lipophilicity would lead to reduced macrophage toxicity.<sup>30</sup> We therefore elected to build on our previous success in targeting the S319 pocket through polar heterocyclic replacement of the 3-methoxyphenyl, with the aim of producing a high affinity LdNMT inhibitor with reduced lipophilicity.<sup>28</sup>

**Scheme 3.** Synthesis of Pyrazole Derivatives **13**<sup>a</sup>



<sup>a</sup> Reagents and Conditions: (a) 1-Boc-4-piperidinol, DIAD, PPh<sub>3</sub>. THF, rt, 4 h, 46%; (b) Hydrazine monohydrate, EtOH, reflux, 18 h, quantitative; (c) EDCI, HOBt, 2-(trimethyl-1H-pyrazol-4-yl)acetic acid,<sup>28</sup> DIPEA, THF:DMF (4:1, v/v), rt, 18 h; (d) TsCl, 1,2,2,6,6-pentamethylpiperidine, DCM, rt, 3 h; (e) 10% TFA in DCM (v/v), rt, 2h, 40% over three steps.


Replacing the 1,2,4-oxadiazole with the more polar 1,3,4-regioisomer<sup>31</sup> and utilising a 1,3,5-trimethylpyrazole motif to replace the hydrophobic methoxyphenyl moiety rendered compound **13** (Scheme 3). Encouragingly, **13** is roughly equipotent to **10j** yet is predicted to be two orders of magnitude less lipophilic, resulting in a highly promising compound. It should be noted that neither **10j** nor **13** display significant selectivity for LdNMT over HsNMT, highlighting a target for further development of the series. The crystal structure of **13** bound to LmNMT was solved (PDB ID: 5a28, Figure 2), and this alternative oxadiazole isomer did not display the ring-opened analogue (Figure S2).



**Figure 2: Crystal structure of 13 (blue) bound to LmNMT (grey).** (A) The chloro-substituent is buried within a hydrophobic pocket, and appears to have good shape complementarity with the enzyme active site. (B) In addition to the shape-complementarity of the scaffold, the compound appears to form a similar binding mode to that previously observed. (C) The trimethylpyrazole motif forms both a hydrogen bonding interaction with Ser330 and a hydrophobic stacking interaction with Phe90.

The basic piperidine moiety forms a polar interaction with the carboxylate of the C-terminal residue, Leu421 and a water-mediated interaction with Tyr92, mimicking the N-terminus of substrate peptides (**Error! Reference source not found.B**). In addition, the trimethyl pyrazole substituent forms  $\pi$ - $\pi$  and polar interactions with Phe90 and Ser330 respectively (**Error! Reference source not found.C**), completing the interactions observed with previous NMT inhibitors. Of particular interest are the interactions around the inhibitor core; a qualitative assessment shows the 5-chlorine inserting into a hydrophobic pocket within the active site, and a lack of space around the 4- position of the scaffold reinforces the lack of activity of the larger substitutions in this position, in the absence of evidence for plasticity in the binding site (Figure 2A). The rationale for the relative lack of activity of the 3-position substituents is less clear, but may be a result of placing a hydrophobic substituent in a water-filled cavity, or restriction of conformation of the piperidine moiety.

Interestingly, these conserved contacts of **1** and **13** are achieved with remarkably distinct trajectories through the binding site, and a comparison of the binding modes suggests a rationale for the large activity difference between PvNMT and LdNMT for **1** (Figure 3).



**Figure 3.** Comparison of the crystal structure of **1** (pink) bound to PvNMT (yellow) and **13** (blue) bound to LmNMT (grey). (A) Overlay of **1** in PvNMT and **13** in LdNMT, with the enzymes removed for clarity; the phenyl ring of the benzo[*b*]thiophene scaffold protrudes in the direction of potential 3- and 4- substitutions of the phenyl series. (B) The surface of the active site of LmNMT (grey) shows a constrained, water filled back pocket in the same position as that occupied by the benzo[*b*]thiophene. (C) Alanine to methionine substitution in LmNMT results in a more constrained pocket and a potential clash with the bulkier benzothiophene scaffold (red dash).

The phenyl ring of the benzothiophene scaffold in **1** occupies a region of the PvNMT binding site that is not occupied by **13** in LmNMT (Figure 3A), filling a volume occupied by two water molecules in LmNMT (Figure 3B). Comparison of the residues around this pocket shows some significant differences, the largest of which is Ala366 in PvNMT substituted for Met377 in LmNMT (Figure 3C). This shifts the protein backbone, and likely results in a more restricted pocket in LmNMT relative to PvNMT. The potential clash between the LmNMT backbone and the benzothiophene core of compound **1** provides a rationale for the

reduced affinity of this molecule for LdNMT compared to PvNMT, and also an explanation for the activity loss in molecules carrying 4-substituents larger than fluorine (**10d-f**, **10h** vs. **10a**, Table 2).

Compound **13** was tested against both macrophages and Ld amastigotes. Pleasingly, **13** did not display any macrophage toxicity at the top concentration tested (90  $\mu\text{M}$ ), reinforcing the earlier hypothesis that reducing macrophage toxicity can be achieved by reducing lipophilicity. However, this molecule showed no observed effect against *ex-vivo* amastigotes up to 50  $\mu\text{M}$ .

## CONCLUSION

In summary, commencing from a member of a previously described series of PfNMT inhibitors, compound **5** was discovered as a highly ligand efficient inhibitor of LdNMT, and subsequent modification of the scaffold yielded the 5-chlorophenyl moiety as the optimum core. However, the compounds showed weak activity when tested against *ex-vivo* amastigotes, and this appeared to correlate with macrophage toxicity; reduction of lipophilicity led to the discovery of compound **13**, a high affinity LdNMT inhibitor. The binding mode and selectivity could be rationalised by crystallography, and comparisons to previous structures. Whilst this molecule displayed no macrophage toxicity at all concentrations tested, it also failed to inhibit *L. donovani* amastigotes; the reasons for this lack of cellular activity remain unclear, but it is possible that this may be resolved through further optimisation of the amine basicity and lipophilicity balance in this series. Although NMT has great potential as a drug target in *Leishmania* parasites, these organisms are notoriously robust,<sup>6</sup> and the physicochemical properties required for optimal engagement of intracellular parasite drug targets in the intra-macrophage amastigote stage remain poorly defined. Further investigation of the influences of the physicochemical properties of this

series is currently underway, alongside the introduction of new NMT inhibitor chemotypes<sup>18</sup> with the aim of chemically validating NMT as a drug target in *Leishmania donovani*.

## AUTHOR INFORMATION

M. Rackham and Z. Yu contributed equally to the preparation of this manuscript.

\*EWT Contact Phone: +44 (0) 2075 943752 E-mail: [e.tate@imperial.ac.uk](mailto:e.tate@imperial.ac.uk)

## NOTES

The authors declare no competing financial interest.

## ACKNOWLEDGEMENTS

This work was supported by the Wellcome Trust (grant number: 087792), the Medical Research Council (grant numbers: 0900278 and U117532067) and the EU F97 project MALSIG (HEALTH-F3-2009-223044). The authors are grateful to Andrew Bell and Jennie Hutton for valuable discussions, Shirley Roberts for crystal handling and Johan Turkenburg for X-ray data processing. We also acknowledge the Diamond Light Source for Synchrotron facilities.

## ABBREVIATIONS USED

*N*-Myristoyltransferase (NMT), *Leishmania donovani* (Ld), *Leishmania major* (Lm), Ligand efficiency (LE), *Plasmodium falciparum* (Pf), *Plasmodium vivax* (Pv), 1-ethyl-3-(3-dimethylaminopropyl) carbodiimidehydrochloride (EDCI), hydroxybenzotriazole (HOBT), diisopropylazodicarboxylate (DIAD).

## REFERENCES

- 1 R. D. Pearson and A. de Queiroz Sousa, *Clin. Infect. Dis.*, 1996, **22**, 1–11.
- 2 P. Desjeux, *Comp. Immunol. Microbiol. Infect. Dis.*, 2004, **27**, 305–18.
- 3 F. Chappuis, S. Sundar, A. Hailu, H. Ghalib, S. Rijal, R. W. Peeling, J. Alvar and M. Boelaert, *Nat. Rev. Microbiol.*, 2007, **5**, 873–82.
- 4 WHO Leishmaniasis Programme, <http://www.who.int/leishmaniasis/en/itle>, 2010.
- 5 S. L. Croft, S. Sundar and A. H. Fairlamb, *Clin. Microbiol. Rev.*, 2006, **19**, 111–26.
- 6 P. Escobar, S. Matu, C. Marques and S. L. Croft, *Acta Trop.*, 2002, **81**, 151–7.
- 7 M. D. Resh, *Trends Mol. Med.*, 2012, **18**, 206–14.
- 8 E. W. Tate, K. A. Kalesh, T. Lanyon-Hogg, E. M. Storck and E. Thinon, *Curr. Opin. Chem. Biol.*, 2014, **24C**, 48–57.
- 9 M. H. Wright, W. P. Heal, D. J. Mann and E. W. Tate, *J. Chem. Biol.*, 2010, **3**, 19–35.
- 10 E. Thinon, R. A. Serwa, M. Broncel, J. A. Brannigan, U. Brassat, M. H. Wright, W. P. Heal, A. J. Wilkinson, D. J. Mann and E. W. Tate, *Nat. Commun.*, 2014, **5**, 4919.
- 11 K. Kawasaki, M. Masubuchi, K. Morikami, S. Sogabe, T. Aoyama, H. Ebiike, S. Niizuma, M. Hayase, T. Fujii, K. Sakata, H. Shindoh, Y. Shiratori, Y. Aoki, T. Ohtsuka and N. Shimma, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 87–91.
- 12 P. Pino, S. Sebastian, E. A. Kim, E. Bush, M. Brochet, K. Volkmann, E. Kozlowski, M. Llinás, O. Billker and D. Soldati-Favre, *Cell Host Microbe*, 2012, **12**, 824–34.
- 13 E. W. Tate, A. S. Bell, M. D. Rackham and M. H. Wright, *Parasitology*, 2014, **141**, 37–49.
- 14 J. A. Frearson, S. Brand, S. P. McElroy, L. A. T. Cleghorn, O. Smid, L. Stojanovski, H. P. Price, M. L. S. Guther, L. S. Torrie, D. A. Robinson, I. Hallyburton, C. P. Mpamhanga, J. A. Brannigan, A. J. Wilkinson, M. Hodgkinson, R. Hui, W. Qiu, O. G. Raimi, D. M. F. van Aalten, R. Brenk, I. H. Gilbert, K. D. Read, A. H. Fairlamb, M. A. J. Ferguson, D. F. Smith and P. G. Wyatt, *Nature*, 2010, **464**, 728–32.
- 15 M. H. Wright, B. Clough, M. D. Rackham, K. Rangachari, J. a Brannigan, M. Grainger, D. K. Moss, A. R. Bottrill, W. P. Heal, M. Broncel, R. a Serwa, D. Brady, D. J. Mann, R. J. Leatherbarrow, R. Tewari, A. J. Wilkinson, A. a Holder and E. W. Tate, *Nat. Chem.*, 2014, **6**, 112–21.
- 16 H. P. Price, M. R. Menon, C. Panethymitaki, D. Goulding, P. G. McKean and D. F. Smith, *J. Biol. Chem.*, 2003, **278**, 7206–14.



- 17 M. H. Wright, D. Paape, E. M. Storck, R. A. Serwa, D. F. Smith and E. W. Tate, *Chem. Biol.*, 2015, **22**, 342–54.
- 18 A. S. Bell, J. E. Mills, G. P. Williams, J. A. Brannigan, A. J. Wilkinson, T. Parkinson, R. J. Leatherbarrow, E. W. Tate, A. A. Holder and D. F. Smith, *PLoS Negl. Trop. Dis.*, 2012, **6**, e1625.
- 19 D. Paape, A. S. Bell, W. P. Heal, J. A. Hutton, R. J. Leatherbarrow, E. W. Tate and D. F. Smith, *PLoS Negl. Trop. Dis.*, 2014, **8**, e3363.
- 20 J. A. Hutton, V. Goncalves, J. A. Brannigan, D. Paape, M. H. Wright, T. M. Waugh, S. M. Roberts, A. S. Bell, A. J. Wilkinson, D. F. Smith, R. J. Leatherbarrow and E. W. Tate, *J. Med. Chem.*, 2014, **57**, 8664–70.
- 21 J. A. Brannigan, S. M. Roberts, A. S. Bell, J. A. Hutton, M. R. Hodgkinson, E. W. Tate, R. J. Leatherbarrow, D. F. Smith and A. J. Wilkinson, *IUCrJ*, 2014, **1**, 250–60.
- 22 V. Goncalves, J. A. Brannigan, D. Whalley, K. H. Ansell, B. Saxty, A. A. Holder, A. J. Wilkinson, E. W. Tate and R. J. Leatherbarrow, *J. Med. Chem.*, 2012, **55**, 3578–82.
- 23 T. O. Olaleye, J. A. Brannigan, S. M. Roberts, R. J. Leatherbarrow, A. J. Wilkinson and E. W. Tate, *Org. Biomol. Chem.*, 2014, **12**, 8132–7.
- 24 Z. Yu, J. A. Brannigan, D. K. Moss, A. M. Brzozowski, A. J. Wilkinson, A. A. Holder, E. W. Tate and R. J. Leatherbarrow, *J. Med. Chem.*, 2012, **55**, 8879–90.
- 25 M. D. Rackham, J. A. Brannigan, D. K. Moss, Z. Yu, A. J. Wilkinson, A. A. Holder, E. W. Tate and R. J. Leatherbarrow, *J. Med. Chem.*, 2013, **56**, 371–375.
- 26 T. J. Ritchie, S. J. F. Macdonald, S. Peace, S. D. Pickett and C. N. Luscombe, *Med. Chem. Commun.*, 2012, **3**, 1062.
- 27 Z. Yu, J. A. Brannigan, K. Rangachari, W. P. Heal, A. J. Wilkinson, A. A. Holder, R. J. Leatherbarrow and E. W. Tate, *Med. Chem. Commun.*, **Submitted**.
- 28 M. D. Rackham, J. A. Brannigan, K. Rangachari, S. Meister, A. J. Wilkinson, A. A. Holder, R. J. Leatherbarrow and E. W. Tate, *J. Med. Chem.*, 2014, **57**, 2773–2788.
- 29 J. A. Brannigan, B. A. Smith, Z. Yu, A. M. Brzozowski, M. R. Hodgkinson, A. Maroof, H. P. Price, F. Meier, R. J. Leatherbarrow, E. W. Tate, D. F. Smith and A. J. Wilkinson, *J. Mol. Biol.*, 2010, **396**, 985–99.
- 30 P. D. Leeson and B. Springthorpe, *Nat. Rev. Drug Discov.*, 2007, **6**, 881–90.
- 31 J. Boström, A. Hogner, A. Llinàs, E. Wellner and A. T. Plowright, *J. Med. Chem.*, 2012, **55**, 1817–30.

## TOC Graphic

