

# NJC

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



[www.rsc.org/njc](http://www.rsc.org/njc)

## Ecofriendly Synthesis of Halogenated Flavonoids and Evaluation of their Antifungal Activity

Roberta Bernini,<sup>a\*</sup> Marcella Pasqualetti,<sup>b</sup> Gianfranco Provenzano<sup>a</sup> and Sabrina Tempesta<sup>b</sup>

Received,  
Accepted

DOI:


www.rsc.org/

Brominated and chlorinated flavonoids belonging to different classes (flavanones, flavones and catechins) were prepared from the corresponding flavonoids by a simple and ecofriendly procedure based on the use of sodium halides, aqueous hydrogen peroxide and acetic acid. Pure samples of substrates and products were tested against *Trichoderma koningii*, *Fusarium oxysporum* and *Cladosporium cladosporioides*, common saprotrophic soil and seeds fungi, potentially pathogens for humans and their activity was expressed as linear mycelial growth inhibition (%). Among them, 8-chloro-5,7,3',4'-tetramethoxyepicatechin **29**, a novel catechin derivative, exhibited a remarkable effect against all tested fungi also at low concentrations.

### Introduction

Flavonoids are more than 5000 plant secondary metabolites involved in important biological processes such as nitrogen fixation, photosensitization, energy transfer, plant growth, control of respiration, and photosynthesis [1, 2]. In flowers they provide colors attractive to plant pollinators; in leaves they promote physiological survival of the plant, protecting it from pathogens and UV-B radiation. Flavonoids are found in fruits, vegetables, propolis, honey, tea, and wine and are of interest to human health as a result of beneficial biological effects such as anti-inflammatory, antioxidant, antiviral, antibacterial, anticancer, and antimicrobial activities [3-5].

Structurally, these compounds possess a common phenylbenzopyrone nucleus (C6-C3-C6) consisting of two aromatic rings (A and B) linked by an oxygenated central pyranic ring (C) and are categorized according to the saturation level of the central ring and the presence of carbonylic or alcoholic functionality. Among them, flavanones, flavones, and flavan-3-ols (catechins) are widely diffused in food and beverages (Figure 1).



**Fig. 1** Basic structure of (a) flavonoids; (b) flavanones; (c) flavones and (d) flavan-3-ols.

The introduction of one or more halogen atoms into A or B-ring of flavonoids produced the corresponding halogenated derivatives useful as synthetic intermediates [6-9]. In addition, halogen atoms confer to flavonoids additional biological properties [10, 11]. As example, a series of mono- and di-halogenated flavonoids have showed a remarkable antifungal activity against some human pathogens *e.g.* *Trichophyton longifusus*, *Candida albicans*, *Aspergillus flavus*, *Microsporium canis*, *Fusarium solanii* and *Candida glaberata* even if a structure-activity relationship was not described [12-14].

Generally, the halogenation of aromatic compounds was carried out using molecular halogens in chlorinated solvents. Even if efficient, this procedure has several environmental drawbacks due to the toxic and hazardous nature of reagents (chlorine gas and liquid bromine), and solvents. In addition, pollutant wastes were produced as by-products.

In order to overcome these problems, in the last few years several papers reported green procedures based on the generation of the electrophilic reagent  $X^+$  (*e.g.*  $Br^+$ ,  $Cl^+$ ) by an ecofriendly oxidation of halide ions. The most used benign oxidants are dimethyldioxirane (DMD), potassium peroxymonosulfate (Oxone®) and hydrogen peroxide ( $H_2O_2$ ). Although DMD is an efficient oxidizing agent and its preparation is an easy and safe operation, Oxone® is more readily accessible being commercially available and  $H_2O_2$  appears as the best candidate producing water as the only by-product [15, 16]. In combination with hydrochloric acid or sodium halides, DMD and Oxone® were very efficient for the halogenation of phenols, methoxybenzenes [17-19] and flavonoids [20, 21]; similarly,  $H_2O_2$  carried out the selective halogenation of phenols and methoxybenzene derivatives in good yields in the presence of catalysts [19] or in acetic acid [22, 23]. To the best of our knowledge, no papers about the halogenation of flavonoids under similar conditions are published.

As part of our research on the chemical modifications of naturally phenols to obtain bioactive compounds by ecofriendly procedures [24, 25], we prepared several halogenated flavonoids using a simple and environmentally benign way based on the use of sodium halides, aqueous solution of hydrogen peroxide and acetic acid as solvent. Final compounds were tested against common saprotrophic soil and seed fungi, potentially pathogens for humans e.g. *Trichoderma koningii*, *Fusarium oxysporum*, and *Cladosporium cladosporioides*, [26-28].

## Results and discussion

**Synthesis of halogenated flavonoids.** We firstly investigated the efficiency of the selected halogenating system on flavanone **1**, 5-methoxyflavanone **4**, 7-methoxyflavanone **8**, 5,7,4'-trimethoxyflavanone (methylated naringenin) **12** and 5,7,3',4'-tetramethoxyflavanone (methylated hesperetin) **15** (Scheme 1). All experimental data were reported in Table 1. In acetic acid, in the presence of NaX (X=Br, Cl) and H<sub>2</sub>O<sub>2</sub> (30% water solution), flavanone **1** was converted into the corresponding 6-halogenated derivatives **2** and **3** in 70% and 62% yield (Table 1, entries 1 and 3). According to our previous results [20], flavanone **1** was regioselectively brominated and chlorinated at position C-6 activated by the *para* ethereal oxygen of the heterocyclic C-ring. In our hands, the bromination and chlorination of flavanone **1** failed in ethanol (Table 1, entries 2 and 4). These results confirmed the crucial role of acetic acid in promoting the formation of the electrophilic specie X<sup>+</sup> (Br<sup>+</sup> and Cl<sup>+</sup>) responsible for the attack of the aromatic A-ring of flavanone and the success of the halogenation [22]. In the presence of H<sub>2</sub>O<sub>2</sub>/NaX/CH<sub>3</sub>COOH, the reaction proceeded with good regioselectivity also with activated substrates; the combination of the *ortho* and *para* effects both of the ethereal oxygen of the heterocyclic C-ring and methoxyl groups present into A-ring was responsible for the formation of the

exclusive (or main) halogenated regioisomer. The bromination of 5-methoxyflavanone **4** produced the corresponding 8-bromoderivative **5** in quantitative yield (Table 1, entry 5); the chlorination gave the 8-chloroderivative **6** as main product (72%) and the 6-chloroderivative **7** as secondary product (Table 1, entry 6). Similarly, 7-methoxyflavanone **8** afforded the corresponding 6-bromoderivative **9** in quantitative yield (Table 1, entry 7); 6-chloroderivative **10** and 8-chloroderivatives **11**, respectively in 70 and 18% yields (Table 1, entry 8). Finally, methylated naringenin **12** and methylated hesperetin **15** gave the corresponding 8-bromo and 8-chloroderivatives **13** and **14**, respectively in 95 and 85% yields; **16** and **17** in 88 and 76% yields (Table 1, entries 9-12). Compared to DMD/HCl and Oxone®/NaX systems, halogenated flavanones were obtained in better yields with the only exception of 6-bromo and 6-chloroflavanone **2** and **3**. Worth noting is the absence of 6,8-dihalogenated derivatives evidencing a better control of the regioselectivity of reaction.


On the basis of these experimental results, we extended the use of the H<sub>2</sub>O<sub>2</sub>/NaX/CH<sub>3</sub>COOH system to biologically relevant flavonoids such as 5,7-dimethoxyflavone (chrysin) **18** and catechin derivatives **21**, **26**. As depicted in Scheme 2, compound **18** was converted into the corresponding 8-bromo and 8-chloroderivatives **19** and **20** in 92 and 74% yields (Table 1, entries 13-14) whereas 5,7,3',4'-methylated catechin **21** and 5,7,3',4'-tetramethylepicatechin **26** resulted more reactive producing, respectively, the corresponding 8-halogenated compound **22**, **24**, **27**, **29** as main products and 6,8-derivatives **23**, **25**, **28**, **30** as secondary products (Table 1, entries 15-18).

These data were appealing because the H<sub>2</sub>O<sub>2</sub>/NaX/CH<sub>3</sub>COOH system resulted efficient with different classes of flavonoids producing the corresponding halogenated derivatives in good to excellent yields. In addition, to the best of our knowledge, catechin derivatives **22-25** and **27-30** are novel compounds.

Table 1 Halogenation of flavonoids depicted in Schemes 1-2.


Entry	Substrate <sup>a</sup>	Experimental conditions	Product (yield %) <sup>a</sup>
1	<b>1</b>	H <sub>2</sub> O <sub>2</sub> (3.0 eq.), AcOH (2.5 mL), NaBr (1.0 eq.), 25 °C, 24 h	<b>2</b> : 70
2	<b>1</b>	H <sub>2</sub> O <sub>2</sub> (3.0 eq.), EtOH (2.5 mL), NaBr (1.0 eq.), 25 °C, 24 h	<b>2</b> : --
3	<b>1</b>	H <sub>2</sub> O <sub>2</sub> (6.0 eq.), AcOH (2.5 mL), NaCl (1.0 eq.), 40 °C, 48 h	<b>3</b> : 62
4	<b>1</b>	H <sub>2</sub> O <sub>2</sub> (6.0 eq.), EtOH (2.5 mL), NaCl (1.0 eq.), 40 °C, 48 h	<b>3</b> :--
5	<b>4</b>	H <sub>2</sub> O <sub>2</sub> (3.0 eq.), AcOH (2.5 mL), NaBr (1.0 eq.), 25 °C, 4 h	<b>5</b> : 98
6	<b>4</b>	H <sub>2</sub> O <sub>2</sub> (3.0 eq.), AcOH (2.5 mL), NaCl (3.0 eq.), 25 °C, 24 h	<b>6</b> : 72; <b>7</b> : 20
7	<b>8</b>	H <sub>2</sub> O <sub>2</sub> (3.0 eq.), AcOH (2.5 mL), NaBr (1.0 eq.), 25 °C, 4 h	<b>9</b> : 95
8	<b>8</b>	H <sub>2</sub> O <sub>2</sub> (6.0 eq.), AcOH(2.5 mL, NaCl (6.0 eq.), 40 °C, 48 h	<b>10</b> : 70; <b>11</b> : 18
9	<b>12</b>	H <sub>2</sub> O <sub>2</sub> (3.0 eq.), AcOH (2.5 mL), NaBr (1.0 eq.), 25 °C, 4 h	<b>13</b> : 95
10	<b>12</b>	H <sub>2</sub> O <sub>2</sub> (6.0 eq.), AcOH (2.5 mL), NaCl (6.0 eq.), 25 °C, 8 h	<b>14</b> : 85
11	<b>15</b>	H <sub>2</sub> O <sub>2</sub> (1.0 eq.), AcOH (2.5 mL), NaBr (3.0 eq.), 25 °C, 4 h	<b>16</b> : 88
12	<b>15</b>	H <sub>2</sub> O <sub>2</sub> (6.0 eq.), AcOH (2.5 mL), NaCl (6.0 eq.), 25 °C, 24 h	<b>17</b> : 76
13	<b>18</b>	H <sub>2</sub> O <sub>2</sub> (3.0 eq.), AcOH (2.5 mL), NaBr (1.0 eq.), 25 °C, 24 h	<b>19</b> : 92
14	<b>18</b>	H <sub>2</sub> O <sub>2</sub> (6.0 eq.), AcOH (2.5 mL), NaCl (6.0 eq.), 40 °C, 24 h	<b>20</b> : 74
15	<b>21</b>	H <sub>2</sub> O <sub>2</sub> (3.0 eq.), AcOH (2.5 mL), NaBr (1.0 eq.), 25 °C, 4 h	<b>22</b> : 70; <b>23</b> : 22
16	<b>21</b>	H <sub>2</sub> O <sub>2</sub> (6.0 eq.), AcOH (2.5 mL), NaCl (6.0 eq.), 40 °C, 4 h	<b>24</b> : 60 <b>25</b> : 18
17	<b>26</b>	H <sub>2</sub> O <sub>2</sub> (3.0 eq.), AcOH (2.5 mL), NaBr (1.0 eq.), 25 °C, 4 h	<b>27</b> : 78; <b>28</b> : 20
18	<b>26</b>	H <sub>2</sub> O <sub>2</sub> (6.0 eq.), AcOH (2.5 mL), AcOH, NaCl (10.0 eq.), 25 °C, 4 h	<b>29</b> : 62; <b>30</b> : 10

<sup>a</sup> Calculated after chromatographic purification



a) AcOH or EtOH, H<sub>2</sub>O<sub>2</sub>, NaBr; b) AcOH or EtOH, H<sub>2</sub>O<sub>2</sub>, NaCl

Scheme 1 Halogenation of flavanones **1**, **4**, **8**, **12** and **15**.

Scheme 2 Halogenation of flavonoids **18**, **21** and **26**.

#### Evaluation of the antifungal activity.


Pure samples of flavonoids **4**, **8**, **12**, **15**, **18**, **21**, **26** and the corresponding halogenated derivatives **5**, **6**, **9**, **10**, **11**, **13**, **14**, **16**, **17**, **19**, **20**, **22**, **23**, **24**, **25**, **27**, **28** and **29** were screened in vitro at three different concentrations 0.5, 2.0 and  $8.0 \times 10^{-4}$  M [13-25] about their growth inhibitory activity against *Trichoderma koningii*, *Fusarium oxysporum* and *Cladosporium cladosporioides*, common saprotrophic soil and seed fungi, potentially pathogens for humans [26-28]. All experimental results, expressed as linear mycelial growth inhibition (%), were reported in Table 2 and grouped in the MDS plot depicted in Figure 2. The MDS approach showed a main distribution trend of the flavonoids along the first axis related to their antifungal activity that increases from right to left. 7-Methoxyflavanone **8** and 8-chloro-5,7,3',4'-tetramethoxyepicatechin **29**, located on the right of the plot (group A), were the most active against all species tested and the percent of radial growth inhibition was high also at lower tested concentration ( $0.5 \times 10^{-4}$  M) against *Trichoderma koningii* (respectively, 49.4% and 51.5%, Table 2). 8-Bromo-5-methoxyflavanone **5**, 6-bromo-7-methoxyflavanone **9**, 8-chloro methylated naringenin **14**, methylated hesperetin **15**, 8-bromo methylated hesperetin **16**, 8-chloro methylated hesperetin **17**, 8-bromo methylated chrysin **19**, 8-bromo-5,7,3',4'-tetramethoxycatechin **22**, 6,8-

dibromo-5,7,3',4'-tetramethoxycatechin **23**, 8-chloro-5,7,3',4'-tetramethoxycatechin **24**, 6,8-dichloro-5,7,3',4'-tetramethoxycatechin **25** and 6,8-dibromo-5,7,3',4'-tetramethoxyepicatechin **28** (group D) showed low antifungal activity against all tested fungi (mean growth reduction <25%, Table 2). Flavonoids distributed in the central part of the plot showed a relatively high activity and could be separated in two main groups: group B including 8-chloro-5-methoxyflavanone **6**, 8-chloro-7-methoxyflavanone **11** and 5,7,3',4'-tetramethoxycatechin **21**, active mainly against *Trichoderma koningii*; group C including 5-methoxyflavanone **4**, 6-chloro-7-methoxyflavanone **10**, methylated chrysin **18**, 8-chloro-5,7-methylated chrysin **20** and 8-bromo-5,7,3',4'-tetramethoxyepicatechin **27**, active against all fungal species. Finally, methylated naringenin **12** resulted active only against *Trichoderma koningii* and *Fusarium oxysporum*; 8-bromo methylated naringenin **13** and 5,7,3',4'-tetramethoxyepicatechin **26** were active, respectively, against *Trichoderma koningii* and all fungi but only at higher concentrations.

Table 2 Antifungal activity of tested flavonoids expressed as inhibition (%) of linear mycelial growth; significant differences at  $P < 0.05$ ; no significant values are unreported.

Flavonoid	<i>Trichoderma koningii</i>			<i>Fusarium oxysporum</i>			<i>Cladosporium cladosporioides</i>		
	$8 \times 10^{-4}$ M	$2 \times 10^{-4}$ M	$0.5 \times 10^{-4}$ M	$8 \times 10^{-4}$ M	$2 \times 10^{-4}$ M	$0.5 \times 10^{-4}$ M	$8 \times 10^{-4}$ M	$2 \times 10^{-4}$ M	$0.5 \times 10^{-4}$ M
4	45.3	43.0	41.1	29.6	28.4	21.4	47.1	36.8	25.6
5	15.1	16.3	21.3	19.2	15.1	16.3	21.2	15.0	13.2
6	48.4	25.0	23.2	22.3	20.4	12.5	26.5	23.1	17.4
8	82.2	70.7	49.4	55.8	49.7	27.1	70.3	49.6	26.4
9	21.9	13.1	5.0	11.1	6.9	6.7	7.8	14.8	10.5
10	48.9	34.6	31.3	28.6	35.4	24.4	38.3	32.0	20.7
11	43.3	30.2	28.3	24.4	18.9	12.3	23.9	25.1	5.8
12	74.6	61.5	18.4	60.0	52.8	14.1	17.4	18.6	10.2
13	70.4	20.8	10.3	27.5	18.7	14.9	24.1	9.0	---
14	22.5	15.6	---	20.4	20.7	15.5	7.8	---	---
15	8.9	10.7	6.9	15.2	12.5	9.8	6.7	7.3	---
16	19.5	---	4.9	12.3	21.7	10.8	16.9	8.6	4.7
17	15.9	---	5.9	13.9	17.4	8.3	8.1	---	---
18	52.9	59.5	34.5	27.0	23.7	18.2	28.2	21.5	16.5
19	23.6	21.1	---	17.1	10.2	-	12.6	---	---
20	80.2	48.1	32.4	41.8	26.5	14.9	47.7	18.9	13.8
21	38.2	48.4	22.8	16.4	18.2	9.6	14.1	17.1	11.3
22	23.7	5.3	21.0	22.8	7.1	6.8	16.5	4.5	---
23	20.0	17.4	6.7	40.6	26.0	5.7	10.8	8.6	5.1
24	32.8	17.5	6.5	---	4.2	---	13.3	5.6	-
25	29.3	23.8	6.8	9.3	4.5	6.8	---	---	---
26	70.2	12.5	---	36.3	10.2	---	48.9	21.5	5.9
27	68.7	66.9	22.4	32.5	22.3	6.5	40.2	27.3	17.6
28	25.4	29.0	22.8	6.2	5.9	---	---	---	---
29	73.2	72.2	51.5	43.7	39.3	19.7	55.2	47.4	19.3

Fig. 2 MDS analysis on fungal activity of tested compounds.



## Conclusions

A simple and environmentally benign bromination and chlorination procedure based on the use of aqueous solution of hydrogen peroxide, sodium halides (NaBr and NaCl) and acetic acid was applied for the first time on flavonoids to obtain the corresponding halogenated derivatives, useful synthetic intermediates and potentially bioactive compounds. The procedure was efficient with different representative classes of flavonoids; the halogenation proceeded with good regioselectivity induced by a combination of the effects both of the ethereal oxygen of the heterocyclic C-ring and methoxyl groups present into aromatic A-ring. To the best of our knowledge, catechin derivatives **22-25** and **27-30** appears as novel compounds. Substrates and the corresponding halogenated derivatives were tested against *Trichoderma koningii*, *Fusarium oxysporum*, and *Cladosporium cladosporioides*, common saprotrophic soil and seed fungi, potentially pathogens for humans. Although experimental data did not evidence a structure-activity relationship, almost all flavonoids showed an inhibitory activity; in addition it is remarkable the effect of 8-chloro-5,7,3',4'-tetramethoxyepicatechin **29** which resulted active against all tested fungi also at low concentrations.

## Experimental section

### Materials and methods

All reagents and flavonoids were of the highest grade available and used as such (Sigma-Aldrich, Milan, Italy). Chromatographic purifications were performed on columns packed with Merck silica gel 60, 230-400 mesh. Thin layer chromatography was carried out using Merck platen Kieselgel 60 F254.

$^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were recorded on a Bruker 200 and 400 MHz. Mass spectra were recorded with a gas chromatograph Shimadzu GC-17A equipped with an electron beam of 70 eV, a SPB column (25 mx0.30 mm and 0.25 mm film thickness) and a mass-selective detector QP 6000. The injector temperature was 280 °C. An isothermal temperature profile of 60 °C for 5 min, followed by a 10 °C/min temperature gradient to 250 °C for 10 min was used. Chromatographic grade helium was used as the carrier gas. High Resolution Mass Spectrometry (HRMS) analyses were recorded with Micromass Q-TOF micro Mass Spectrometer (Waters).

**Methylation reactions.** 5,7,4'-Trimethoxyflavanone (methylated naringenin) **12**; 5,7,3',4'-tetramethoxyflavanone (methylated hesperetin) **15**; 5,7,3',4'-tetramethylcatechin **21** and 5,7,3',4'-tetramethylepicatechin **26** were prepared as already reported by us [29, 30].

**5,7,4'-Trimethoxyflavanone (Methylated naringenin) (12).** Yield: 90%; colorless oil; spectroscopic data are according to those reported in the literature [29].

**5,7,3',4'-Tetramethoxyflavanone (Methylated hesperetin) (15).** Yield: 88%; colorless oil; spectroscopic data are according to those reported in the literature [29].

**5,7,3',4'-Tetramethoxycatechin (21).** Yield: 92%; yellow oil; spectroscopic data are according to those reported in the literature [30].

**5,7,3',4'-Tetramethoxyepicatechin (26).** Yield: 90%; yellow oil; spectroscopic data are according to those reported in the literature [30].

**Halogenation reactions.** A 30% aqueous solution of hydrogen peroxide (3.0-6.0 eq.) was added to solution of flavonoid (0.5 mmol) and sodium halide (1.0-3.0 eq.) in acetic acid (2.5 ml); then, the mixture was stirred at room temperature or 40 °C for 4-48 h depending on the substrate. The reaction was monitored by thin layer chromatography (TLC) and gas-mass

chromatography (GC-MS). At the end, the crude was treated with sodium thiosulfate and extracted with ethyl acetate (3x10 mL). The reunited organic fractions were dried over anhydrous sodium sulfate; after filtration, the solvent was evaporated under reduced pressure. Final products were isolated and purified by chromatographic column using CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH=9.5/05 as eluent and characterized by <sup>1</sup>H NMR, <sup>13</sup>C NMR, and HR-MS analysis.

**6-Bromoflavanone (2).** Yield: 70%; yellow oil; spectroscopic data are according to those reported in the literature [20].

**6-Chloroflavanone (3).** Yield: 62%; colorless oil; spectroscopic data are according to those reported in the literature [20].

**8-Bromo-5-methoxyflavanone (5).** Yield: 98%; Yellow oil; spectroscopic data are according to those reported in the literature [20].

**8-Chloro-5-methoxyflavanone (6).** Yield: 72%; yellow oil; spectroscopic data are according to those reported in the literature [20].

**6-Chloro-5-methoxyflavanone (7).** Yield: 20%; colorless oil; spectroscopic data are according to those reported in the literature [20].

**6-Bromo-7-methoxyflavanone (9).** Yield: 95%; colorless oil; spectroscopic data are according to those reported in the literature [20].

**6-Chloro-7-methoxyflavanone (10).** Yield: 70%; yellow oil; spectroscopic data are according to those reported in the literature [20].

**8-Chloro-7-methoxyflavanone (11).** Yield: 18%; colorless oil; spectroscopic data are according to those reported in the literature [20].

**8-Bromo-5,7,4'-trimethoxyflavanone (13).** Yield: 95%; colorless oil; spectroscopic data are according to those reported in the literature [20].

**8-Chloro-5,7,4'-trimethoxyflavanone (14).** Yield: 85%; colorless oil; spectroscopic data are according to those reported in the literature [20].

**8-Bromo-5,7,3',4'-tetramethoxyflavanone (16).** Yield: 88%; yellow oil; spectroscopic data are according to those reported in the literature [31].

**8-Chloro-5,7,3',4'-tetramethoxyflavanone (17).** Yield: 76%; colorless oil; spectroscopic data are according to those reported in the literature [31].

**8-Bromo-5,7-dimethoxyflavone (19).** Yield: 92%; colorless oil; spectroscopic data are according to those reported in the literature [32].

**8-Chloro-5,7-dimethoxyflavone (20).** Yield: 74%; colorless oil; spectroscopic data are according to those reported in the literature [32].

**8-Bromo-5,7,3',4'-tetramethoxycatechin (22).** Yield: 70%; yellow oil; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>), δ: 2.41-2.54 (dd, *J*<sub>1</sub> = 8.1 Hz, *J*<sub>2</sub> = 18.3 Hz, CH), 2.82-2.94 (dd, *J*<sub>1</sub> = 5.3 Hz, *J*<sub>2</sub> = 18.4 Hz, 1H, CH), 3.70 (s, 3H, OCH<sub>3</sub>), 3.86 (s, 9H, 3xOCH<sub>3</sub>), 4.01-4.11 (m, 1H, CH), 4.75 (d, *J* = 7.7 Hz, 1H, CH), 6.13 (s, 1H, Ph-H), 6.83-6.96 (m, 3H, Ph-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>), δ: 161.8, 148.5, 148.1, 134.5, 134.0, 120.5, 113.9, 110.9, 110.6, 93.6, 80.2, 79.5, 73.1, 57.0, 56.1, 55.8, 46.9, 35.2. C<sub>19</sub>H<sub>21</sub>BrO<sub>6</sub> requires C, 53.66; H, 4.98; O, 22.57. Found: C, 53.02; H, 5.01; O, 22.67.

**6,8-Dibromo-5,7,3',4'-tetramethoxycatechin (23).** Yield: 22%; yellow oil; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>), δ: 2.80-2.92 (dd, *J*<sub>1</sub> = 8.5 Hz, *J*<sub>2</sub> = 16.6 Hz, CH), 3.07-3.19 (dd, *J*<sub>1</sub> = 5.2 Hz, *J*<sub>2</sub> = 16.6 Hz, 1H, CH), 3.72 (s, 3H, OCH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 3.86 (s, 6H, 2xOCH<sub>3</sub>), 4.10-4.18 (m, 1H, CH), 4.75 (d, *J* = 7.7 Hz, 1H, CH), 6.83-6.86 (m, 3H, Ph-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>), δ: 162.7, 157.6, 149.7, 149.5, 134.1, 120.1, 112.6, 111.3, 107.6, 100.3, 99.4, 83.1, 67.1, 60.5, 59.9, 56.1, 56.0, 55.9, 27.5. C<sub>19</sub>H<sub>20</sub>Br<sub>2</sub>O<sub>6</sub> requires C, 45.26; H, 4.00; O, 19.04. Found: C, 45.89; H, 4.09; O, 18.87.

**8-Chloro-5,7,3',4'-tetramethoxycatechin (24).** Yield: 60%; yellow oil; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>), δ: 2.42-2.56 (dd, *J*<sub>1</sub> = 8.1 Hz, *J*<sub>2</sub> = 18.3 Hz, CH), 2.80-2.94 (dd, *J*<sub>1</sub> = 5.3 Hz, *J*<sub>2</sub> = 18.4 Hz, CH), 3.72 (s, 3H, OCH<sub>3</sub>), 3.86 (s, 9H, 3xOCH<sub>3</sub>), 4.00-4.12 (m, 1H, CH), 4.72 (d, *J* = 7.7 Hz, 1H, CH), 5.83 (s, 1H, CH), 6.85-6.98 (m, 3H, Ph-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>), δ: 161.8, 148.5, 148.1, 134.5, 134.0, 120.5, 113.9, 110.9, 110.6, 93.6, 80.2, 79.5, 73.1, 57.0, 56.1,

55.8, 46.9, 35.2. C<sub>19</sub>H<sub>21</sub>ClO<sub>6</sub> requires C, 59.92; H, 5.56; O, 25.21. Found: C, 60.02; H, 5.26; O, 24.99.

**6,8-Dichloro-5,7,3',4'-tetramethoxycatechin (25).** Yield: 18%; yellow oil; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>), δ: 2.78-2.90 (dd, *J*<sub>1</sub> = 8.5 Hz, *J*<sub>2</sub> = 16.6 Hz, 1H, CH), 3.06-3.20 (dd, *J*<sub>1</sub> = 5.2 Hz, *J*<sub>2</sub> = 16.6 Hz, 1H, CH), 3.74 (s, 3H, OCH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 3.86 (s, 6H, 2xOCH<sub>3</sub>), 4.10-4.20 (m, 1H, CH), 4.76 (d, *J* = 7.7 Hz, 1H, CH), 6.82-6.88 (m, 3H, Ph-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>), δ: 150.7, 148.8, 148.2, 134.9, 134.0, 120.6, 113.9, 110.8, 105.5, 97.8, 79.8, 79.6, 73.2, 59.5, 60.0, 55.9, 55.8, 55.7, 37.0. C<sub>19</sub>H<sub>20</sub>Cl<sub>2</sub>O<sub>6</sub> requires C, 54.95; H, 4.85; O, 23.12. Found: C, 55.25; H, 4.98; O, 23.42.

**8-Bromo-5,7,3',4'-tetramethoxyepicatechin (27).** Yield: 78%; yellow oil; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>), δ: 2.85-2.88 (m, 2H, CH<sub>2</sub>), 3.71 (s, 3H, OCH<sub>3</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 3.85 (s, 9H, 3xOCH<sub>3</sub>), 3.88 (s, 9H, 3xOCH<sub>3</sub>), 4.21-4.24 (m, 1H, CH), 4.91 (d, *J* = 7.7 Hz, 1H, CH), 6.13 (s, 1H, Ph-H), 6.85-7.05 (m, 3H, Ph-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>), δ: 161.9, 148.5, 148.2, 134.7, 134.2, 120.8, 113.9, 110.9, 110.8, 93.6, 80.4, 79.6, 73.1, 57.2, 56.0, 55.9, 46.8, 35.4. C<sub>19</sub>H<sub>21</sub>BrO<sub>6</sub> requires C, 53.66; H, 4.98; O, 22.57. Found: C, 54.02; H, 5.04; O, 22.87.

**6,8-Dibromo-5,7,3',4'-tetramethoxyepicatechin (28).** Yield: 20%; yellow oil; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>), δ: 2.85-2.88 (m, 2H, CH<sub>2</sub>), 3.70 (s, 3H, OCH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 3.86 (s, 6H, 2xOCH<sub>3</sub>), 4.10-4.18 (m, 1H, CH), 4.75 (d, *J* = 7.7 Hz, 1H, CH), 6.83-6.86 (m, 3H, Ph-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>), δ: 162.8, 157.6, 149.7, 149.5, 134.1, 120.1, 112.6, 111.3, 107.6, 100.4, 99.4, 83.2, 67.2, 60.4, 59.9, 56.2, 56.1, 55.9, 27.4. C<sub>19</sub>H<sub>20</sub>Br<sub>2</sub>O<sub>6</sub> requires C, 45.26; H, 4.00; O, 19.04. Found: C, 45.79; H, 4.12; O, 18.87.

**8-Chloro-5,7,3',4'-tetramethoxyepicatechin (29).** Yield: 62%; yellow oil; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>), δ: 2.85-2.88 (m, 2H, CH<sub>2</sub>), 3.72 (s, 3H, OCH<sub>3</sub>), 3.86 (s, 9H, 3xOCH<sub>3</sub>), 4.02-4.12 (m, 1H, CH), 4.74 (d, *J* = 7.7 Hz, 1H, CH), 5.88 (s, 1H, Ph-H), 6.86-7.00 (m, 3H, Ph-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>), δ: 157.8, 148.5, 148.1, 137.9, 134.2, 120.6, 113.9, 110.8, 102.5, 92.8, 80.4, 79.8, 73.0, 57.0, 56.2, 55.9, 55.8, 54.2, 34.8. C<sub>19</sub>H<sub>21</sub>ClO<sub>6</sub> requires C, 59.92; H, 5.56; O, 25.21. Found: C, 60.02; H, 5.26; O, 24.99.

**6,8-Dichloro-5,7,3',4'-tetramethoxyepicatechin (30).** Yield: 10%; yellow oil; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>), δ: 2.85-2.88 (m, 2H, CH<sub>2</sub>), 3.72 (s, 3H, OCH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 3.86 (s, 6H, 2xOCH<sub>3</sub>), 4.12-4.18 (m, 1H, CH), 4.76 (d, *J* = 7.7 Hz, 1H, CH), 6.80-6.86 (m, 3H, Ph-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>), δ: 150.6, 148.6, 148.0, 134.8, 134.5, 120.7, 113.9, 110.8, 105.5, 97.8, 79.9, 79.8, 73.0, 59.9, 59.7, 56.2, 56.0, 55.9, 37.3. C<sub>19</sub>H<sub>20</sub>Cl<sub>2</sub>O<sub>6</sub> requires C, 54.95; H, 4.85; O, 23.12. Found: C, 55.01; H, 4.75; O, 23.25.

#### Assay of antifungal activity.

Flavone and flavanone exhibit fungicidal activity at 10<sup>-3</sup>-10<sup>-5</sup> M [33, 34, 35]. Therefore, in this study flavonoid derivatives were tested at 0.5, 2.0 and 8.0 x 10<sup>-4</sup> M [13, 14, 25]. Each flavonoid was dissolved in acetone so that the final concentration of solvent in the test medium did not exceed 1% of the total solution composition; three solutions were prepared (0.5, 2.0 and 8.0 x 10<sup>-4</sup> M). *T. koningii*, *F. oxysporum* and *C. cladosporioides* were used for the antifungal tests. Before testing, each isolate was subcultured on MEA (DIFCO) to ensure optimal growth characteristics and purity. The isolates had been grown for 4-14 days on MEA at 25°C. Conidia suspensions were prepared in sterile water supplemented with 0.01% of Tween 80. Each suspension was diluted to obtain the final inoculum, which ranged from 0.5 x 10<sup>4</sup>-1.0 x 10<sup>4</sup> CFU/ml. The inoculum size was determined microscopically using Bürker's chamber and verified by plating 100 µL of serial dilutions of each inoculum onto an MEA plate and incubation until growth became visible. Each Petri dishes (90 mm) containing 12 mL of the medium



including the products in required concentrations (added to the agar at temperature below 50°C) was inoculated with 2 µl of the inoculum suspensions; 5-6 replications for each concentration and fungus were made. The Petri dishes were incubated at 25 °C in the dark to a clearly visible growth. Evaluation of linear growth was conducted by measuring mycelial diameters of each inoculated plate at broadest, medium and smallest diameter, and compared with the corresponding control. The inhibition (%) of linear mycelial mean growth of *T. koningii*, *C. cladosporioides* and *F. oxysporum* was calculated after incubation for 3, 5 and 6 days respectively. The data were evaluated by analysis of variance, probability of single differences was calculated at the 5% level. The data were also ordered by Multi-Dimensional Scaling (MDS) analysis using SYSTAT 10.0 Statistics [36, 37] to group the antifungal activity of all tested compounds; an input matrix of Euclidean distance (calculated on linear mycelial growth) and Kruskal loss function were utilized [38].

### Notes and references

<sup>a</sup> Dipartimento di Scienze e Tecnologie per l'Agricoltura, le Foreste, la Natura e l'Energia (DAFNE), Università degli Studi della Tuscia, Via S. Camillo De Lellis, 01100 Viterbo, Italy. Fax: +39 0761 357242; Tel: +39 0761 357452; E-mail: [berninir@unitus.it](mailto:berninir@unitus.it)

<sup>b</sup> Dipartimento di Scienze Ecologiche e Biologiche (DEB), Università degli Studi della Tuscia, Largo dell'Università, 01100 Viterbo, Italy

- 1 J. B. Harborne, H. Baxter. The Handbook of Natural Flavonoids, Vol. 1 and 2. Chichester, UK: Wiley and Sons; 1999.
- 2 J. B. Harborne, Williams, H. Advances in Research since 1992. *Phytochemistry* 2000, **55**, 481.
- 3 S. Kumar, A. K. Pandey. *Sci. World J.* 2013, **1**.
- 4 A. D. Agrawal. *Int. J. Pharm. Sci. Nanotechnol.* 2011, **4**, 1394.
- 5 T. P. Cushnie, *Int. J. Antimicrob. Ag.*, 2005, **26**, 343.
- 6 M. A. Selepe, F. A. Van Heerden, *Molecules*, 2013, **18**, 4739 and references therein.
- 7 R. Bernini, S. Cacchi, I. De Salve, G. Fabrizi, *Tetrahedron Lett.*, 2007, **48**, 4973.
- 8 D. Muller, J.-P. Fleury, *Tetrahedron Lett.*, 1991, **32**, 2229-2232.
- 9 X. Zheng, W.-D. Meng, F.-L. Qing, *Tetrahedron Lett.*, 2004, **45**, 8083-8085.
- 10 M. Marder, J. Zinzuk, M. I. Colombo, C. Wasowski, H. Viola, C. Wolfman, J. H. Medina, E. A. Ruveda, A. C. Paladini, *Bioorg. Med. Chem. Lett.*, 1997, **7**, 2003.
- 11 E. Ognibene, P. Bovicelli, W. Adriani, L. Saso, G. Laviola, *Prog. Neuro-Psycopa.*, 2008, **32**, 128.
- 12 A. Hasan, L. Rasheed. A. Malik, *Asian J. Chem.*, 2007, **19**, 937.
- 13 M. Weidenborner, H. C. Jha, *Mycol. Res.*, 1997, **101**, 733.
- 14 A. M. Silva, M. Weidenborner, J. A. S. Cavaliero, *Mycol. Res.*, 1998, **102**, 638.
- 15 P. B. De la Mare. Electrophilic Halogenation, Cambridge University Press: Cambridge, 1976, Chapter 5.
- 16 A. Podgorsek, M. Zupan, J. Iskra, *Angew. Chem, Int. Ed.*, 2009, **48**, 8424 and references therein.
- 17 Y.-J. Lee, H.-T. Wu, C.-N. Lin, *Lett. Org. Chem.*, 2009, **6**, 585-587.
- 18 V. V. Krishna Mohan Kandepi, N. Narender, *Synthesis*, 2012, **44**, 15-26 and references therein.
- 19 P. Bovicelli, E. Mincione, R. Antonioletti, R. Bernini, M. Colombari, *Synth. Commun.* 2001, **31**, 2955.
- 20 P. Bovicelli, R. Bernini, A. Antonioletti, E. Mincione, *Tetrahedron Lett.*, 2002, **43**, 5563.
- 21 A. Arnone, G. Nasini, W. Panzeri, O. Vajna de Pava, C. Zucca, *J. Chem. Res. (S)*, 2003, **11**, 683.
- 22 N. Narender, K. Reddy Suresh Kumar, K.V.V. Mohan Krishna, S. J. Kulkarni, *Tetrahedron Lett.*, 2007, **48**, 6124.
- 23 D. N. D'Aleo, S. R. Allard, C. C. Foglia, S. L. M. Parent, D. J. Rohr, C. Gottardo, C. D. MacKinnon, *Can. J. Chem.*, 2013, **91**, 679.
- 24 R. Bernini, E. Mincione, M. Barontini, G. Fabrizi, M. Pasqualetti, S. Tempesta, *Tetrahedron*, 2006, **62**, 7733.
- 25 R. Bernini, E. Mincione, G. Provenzano, G. Fabrizi, S. Tempesta, M. Pasqualetti, *Tetrahedron*, 2008, **64**, 7561.
- 26 T. Watanabe. Pictorial atlas of soil and seed fungi: morphologies of cultured fungi and key to species. **2002**. CRC Press.
- 27 G. S. De Hoog, J. Guarro, J. Gené, M. J. Figueras. Atlas of clinical fungi. **2000**. CBS, Utrecht, Netherlands.
- 28 K. H. Domsch, W. Gams, T.-H. Anderson. Compendium of soil fungi. **2007**. IHW-Verlag, Eching.
- 29 R. Bernini, E. Mincione, M. Cortese, R. Saladino, G. Gualandi, M. C. Belfiore, *Tetrahedron Lett.*, 2003, **44**, 4823.
- 30 R. Bernini, F. Crisante, P. Gentili, F. Morana, M. Pierini, M. Piras, *J. Org. Chem.*, 2011, **76**, 820.
- 31 P. Yaipakdee, L. W. Robertson, *Phytochemistry*, **2001**, **57**, 341.
- 32 G. Righi, R. Antonioletti, I. Proietti Silvestri, N. D'Antona, D. Lambusta, P. Bovicelli, *Tetrahedron*, 2010, **66**, 1294.
- 33 T. M. O'Neill, J. W. Mansfield, *Trans. Br. Mycol. Soc.*, 1982, **79**, 229.
- 34 L. Carrillo, S. E. Gomez Molina, M. Benitez Ahrendts, *Rev. Cient. Agraria*, 1999, **3**, 11.
- 35 M. P. Salas, G. Céliz, H. Geronazzo, M. Daz, S. L. Resnik, *Food Chem.*, 2011, **124**, 1411.
- 36 SPSS: Statistical package of the social sciences. 10.0. **1999**. SPSS Inc., Chicago, Illinois.
- 37 L. Wilkinson, M. Hill, E. Vang. SYSTAT: Statistics, Version 5.2 Edition. **1992** SYSTAT Inc. Evanston. Illinois.
- 38 J. Podani. Multivariate data analysis in ecology and systematics. **1994**. SPB Academic Publishing, The Hague, The Netherlands.

## TABLE OF CONTENTS

## Ecofriendly synthesis of halogenated flavonoids and evaluation of their antifungal activity

Roberta Bernini,<sup>\*</sup> Marcella Pasqualetti, Gianfranco Provenzano and Sabrina Tempesta

Chlorinated catechin derivative **29** resulted the most active compound against *Trichoderma koningii* and *Cladosporium cladosporioides*, common saprotrophic soil and seed fungi.

