

**NJC**

**Multicomponent pathway inspired regioselective synthesis
of 2,3,4-Trisubstituted 1H-Pyrroles via [3+2] Cycloaddition
Reaction**

Journal:	<i>New Journal of Chemistry</i>
Manuscript ID:	NJ-ART-02-2015-000302.R1
Article Type:	Paper
Date Submitted by the Author:	20-Mar-2015
Complete List of Authors:	Chaskar, Atul; National Taiwan University, Department of Chemistry Kalmode, Hanuman; Institute of Chemical Technology, Department of Dyestuff technology Vadagaonkar, Kamlesh; Institute of Chemical Technology, Department of Dyestuff technology Murugan, Kaliyappan; National Taiwan University, Department of Chemistry

SCHOLARONE™
Manuscripts

Multicomponent pathway inspired regioselective synthesis of 2,3,4-trisubstituted 1*H*-pyrroles via [3+2] cycloaddition reaction

Hanuman P. Kalmode,^a Kamlesh S. Vadagaonkar,^a Kaliyappan Murugan*^b and Atul C. Chaskar*^{a,b,c}

^a*Department of Dyestuff Technology, Institute of Chemical Technology, Mumbai 400019, India*

^b*National Taiwan University, Taipei 10617, Taiwan*

^c*National Centre for Nanosciences and Nanotechnology, University of Mumbai, Mumbai 400098, India*

Tel.: +91-22-3361 1111/ 2706; Fax: +91-22-3361 1020; E-mail: achaskar25@gmail.com.

Abstract: A copper catalyzed regioselective synthesis of 2,3,4-trisubstituted 1*H*-pyrroles by means of a novel three-component coupled domino reaction of aldehydes, ketones and alkyl isocyanoacetates is reported. This transformation proceeds through the formation of chalcone followed by [3+2] cycloaddition reaction to give the α -cuprioisocyanide, a cyclic organocopper intermediate, which on copper-hydrogen exchange followed by oxidation offers 2,3,4-trisubstituted 1*H*-pyrrole exclusively.

Keywords: Multicomponent reaction, Tandem [3+2] cycloaddition, 2,3,4-trisubstituted 1*H*-pyrrole, Regioselectivity, BODIPY dyes.

Introduction:

Pyrrole in general and 2,3,4-trisubstituted 1*H*-pyrroles in particular are the privileged scaffolds as they are the significant structural constituents of various natural products, pharmaceuticals and agrochemicals.^{1,2} 2,3,4-Trisubstituted 1*H*-pyrroles that possess coordinate bond forming ability are used in such applications as molecular sensing, semiconducting materials, lasers, image diagnosis, materials science, etc.³ Noteworthy is the strong absorption and very intense fluorescence of BODIPY (boron-dipyrromethene) dyes derived from pyrroles.⁴ The development of various convenient methods for the synthesis of pyrroles stems in these potential applications.

Several synthetic methodologies have been reported since Hantzsch⁵ first prepared pyrroles by the reaction of β -enaminones with α -haloketones. Notable amongst these are the Paal-Knorr synthesis⁶ using 1,4-diketones with amines, the Knorr synthesis using α -amino ketones and β -dicarbonyl compounds, the Barton-Zard synthesis using isocyanoacetates and nitroalkenes, van Leusen synthesis.⁷ In addition to these some other syntheses have also been reported.⁸⁻¹² In spite of their effectiveness these classical methods suffer from several shortcomings such as functional group non-compatibility, prolonged reaction time, formation of byproducts, less regiospecificity, harsh reaction conditions, multistep synthetic operation and tedious workup procedure.


In view of the aforementioned difficulties the development of a cost-effective, mild, convenient, straightforward and regioselective method for the synthesis of pyrroles from easily available chemical entities is highly desirable. Multi-component coupled domino reactions (MCDRs), with at least three different substrates reacting in a sequential manner to form a single compound, have provided a most powerful platform to access diversity as well as complexity in a limited number of reaction steps.¹³ These reactions can avoid timeconsuming and costly

processes for purification of various precursors. In addition, these reactions satisfy the objectives of an ideal synthesis and the principles of green chemistry.¹⁴ Therefore, the design of new MCDRs is a continuing challenge at the forefront of organic chemistry. In this context, in recent years atom-economical pyrrole synthesis has been accomplished by using isocyanides.¹⁵⁻²⁰ Their unusual reactivity and ability to form multiple bonds through a one-pot manner offered by tandem or multicomponent reactions, has been used to produce an array of heterocycles.²¹⁻²² Ley *et al.* reported the preparation of nitrosubstituted functionalized pyrroles *via* a one-pot three-component coupling of tosyl isocyanide, ethyl chloroformate and nitrostyrene.²³ Uno and co-workers used vinyl sulfones with ethyl isocyanoacetate for the synthesis of 3,4-disubstituted pyrrole-2-carboxylates.²⁴ 2,3,4-Trisubstituted 1*H*-pyrroles with a tosyl substituent at the C-2 position were synthesized from tosylmethyl isocyanide (TOSMIC) and vinyl azides.²⁵ Nevertheless, transition-metal catalyzed cycloaddition reactions of isocyanides with double or triple bonds have received enormous attention. Various groups have reported the synthesis of substituted pyrroles using copper, phosphine and silver catalyzed [3+2] cycloaddition of isocyanides and alkynes.²⁶⁻³⁴ Noteworthy is the formation of 2,4-di-EWG-substituted and 2,3-di-EWG-substituted pyrroles in presence of copper and phosphine catalyst, respectively. This implies that regiodifferentiated addition of isocyanides to alkynes is controlled by the catalytic species. However, some of these methods are associated with an inherent drawback of formation of mixture of regioisomers (Scheme 1).



Scheme 1. Formation of mixture of regioisomers


Considering the literature reports³⁵ and as a part of our ongoing research on the development of multicomponent approaches for the synthesis of heterocycles³⁶, we herein wish to report simple, efficient and regioselective synthesis of novel 2,3,4-trisubstituted 1*H*-pyrroles from easily available starting materials. Hitherto, ours is the first attempt in direct synthesis of 2,3,4-trisubstituted 1*H*-pyrroles *via* a three-component coupled domino reaction of aldehydes, ketones and alkyl isocyanoacetates (3) by using metal catalyst, to the best of our knowledge (Scheme 2).



Scheme 2. General regioselective one-pot synthesis of 2,3,4-trisubstituted 1*H*-pyrrole

Results and discussion

In order to establish the optimized reaction condition, we first investigated the reaction between acetophenone (**1a**), benzaldehyde (**2a**) and methyl 2-isocyanoacetate (**3a**) using different catalysts, bases and solvents. To our surprise, the combination of 10 mol % CuI and Cs₂CO₃ in methanol at room temperature in open air solely produced methyl 4-benzoyl-3-phenyl-1*H*-pyrrole-2-carboxylate (**4a**) regioselectively with excellent yield (Table 1, entry 11). Moderate yields of **4a** (53%, 45% and 48%) were obtained in presence of K₂CO₃, K^tBuO and NaOEt (Table 1, entries 6, 7 and 8) respectively. Use of DBU and 2,6-lutidine (Table 1, entries 9 and 10) also resulted in low yields (10% and 13%) of the product **4a**. We next carried out this domino reaction by adding all the reagents at the same time but could not detect the formation of product **4a** (Table 1, entry 12). This clearly highlights the significance of step-wise addition. The superior catalyst activity of CuI is observed in open air as compare to N₂ (Table 1, entries 13-14). Trace amount of product formation was observed in N₂ atmosphere, as oxygen present in the solvent drives oxidation of pyrroline to pyrrole. Various copper salts were tested under this condition for their catalytic activity but none of them were found to be as effective as CuI (Table 1, entries 15-19). Thus, firstly the Aldol condensation reaction involving the acetophenone (**1a**) and benzaldehyde (**2a**) took place in the presence of the base. When the reaction was finished and the corresponding α,β -unsaturated ketone was generated in the reaction media, the second reaction step ([3+2] cycloaddition) was achieved by adding catalyst and the methyl 2-isocyanoacetate (**3a**) to the reaction media.

Table 1. Optimization of reaction conditions^a

Entry	Catalyst (mol %)	base	Solvent	Yield (%) ^d
1	CuI	Cs ₂ CO ₃	DCE	62
2	CuI	Cs ₂ CO ₃	DME	68
3	CuI	Cs ₂ CO ₃	Dioxane	57
4	CuI	Cs ₂ CO ₃	THF	51
5	CuI	Cs ₂ CO ₃	CH ₃ CN	27
6	CuI	K ₂ CO ₃	MeOH	53
7	CuI	K ^t BuO	MeOH	45
8	CuI	NaOEt	MeOH	48
9	CuI	DBU	MeOH	10
10	CuI	2,6-Lutidine	MeOH	13
11	CuI	Cs ₂ CO ₃	MeOH	83
12 ^b	CuI	Cs ₂ CO ₃	MeOH	N.D.
13 ^c	CuI	Cs ₂ CO ₃	DMF	Trace
14 ^c	CuI	Cs ₂ CO ₃	MeOH	Trace
15	CuCl	Cs ₂ CO ₃	MeOH	44
16	CuBr	Cs ₂ CO ₃	MeOH	34
17	CuCl ₂	Cs ₂ CO ₃	MeOH	24
18	CuBr ₂	Cs ₂ CO ₃	MeOH	27
19	Cu(OAc) ₂	Cs ₂ CO ₃	MeOH	30


^aReaction conditions: acetophenone **1a** (1.0 mmol), benzaldehyde **2a** (1.3 mmol) and base (2.0 mmol) in 3 mL of solvent was stirred at R. T. for 3.5 h, catalyst (10 mol %), methyl 2-isocyanoacetate **3a** (1.0 mmol) in 2 mL solvent was added and stirring continued for 0.5 h, open air. ^bReaction conditions: acetophenone **1a** (1.0 mmol), benzaldehyde **2a** (1.3 mmol), CuI (10 mol %), methyl 2-isocyanoacetate **3a** (1.0 mmol) and Cs₂CO₃ (2.0 mmol) in 5 mL of MeOH was stirred at R. T. for 4.0 h, open air. ^cUnder N₂. ^dIsolated yields. N. D. = Not Detected.


In order to investigate the scope and limitations of this novel three-component coupled domino reaction of various aromatic or aliphatic ketones (**1**), aromatic or aliphatic aldehydes (**2**) and alkyl isocyanoacetates (**3**) were examined (Table 2). The reaction worked well and the corresponding pyrroles (**4a-4w**) were obtained in 48-90% yield within 2.5-6.0 hours. Electron neutral or weakly deactivated acetophenones (4-CH₃ and 3-Cl), substituted benzaldehydes bearing electron withdrawing (4-NO₂, 4-CN) substituents in the phenyl ring and methyl 2-isocyanoacetate (**3a**) offered excellent yields of the product (Table 2, entries **4b** and **4c**). The rate of chalcone formation followed by 1,3-dipolar cycloaddition reaction was highest (2.5 h) for the

above substrates. Benzaldehydes (**2**) and acetophenones (**1**) bearing halogens, electron neutral substituents on the phenyl ring and methyl 2-isocyanoacetate (**3a**) gave the corresponding 2,3,4-trisubstituted 1*H*-pyrroles in good to excellent yield (Table 2, entries **4d-4k**). Then the substituted benzaldehydes, acetone and methyl 2-isocyanoacetate (**3a**) were studied for the three-component coupled domino reaction. Benzaldehydes bearing halogens, electron neutral substituents in the phenyl ring, acetone and methyl 2-isocyanoacetate (**3a**) offered moderate to good yields of the products (Table 2, entries **4l-4q**). The steric hindrance near the double bond of chalcones generated *in-situ* from benzaldehydes (4-CH₃, 3-CH₃, 2-CH₃) and acetone, followed by [3+2] cycloaddition reaction with methyl 2-isocyanoacetate (**3a**) hampered the yield of products as well as rate of reactions, and the reactivity order was found to be *p* > *m* > *o*-isomer (Table 2, entries **4m**, **4n** and **4o**). As expected, benzaldehyde with an electron withdrawing 4-NO₂ group, acetone and methyl 2-isocyanoacetate (**3a**) took significantly shorter reaction time (3.5 h) and gave a higher yield of the product (Table 2, entry **4s**, 76%) than that offered by its methoxy counterpart (Table 2, entry **4r**, 55%). Interestingly acetaldehyde, acetone and methyl 2-isocyanoacetate (**3a**) also underwent the three-component coupled domino reaction to afford methyl 4-acetyl-3-methyl-1*H*-pyrrole-2-carboxylate (**4t**) in 48% yield.

The scope of this domino reaction was extended to ethyl 2-isocyanoacetate (**3b**). The acetophenone (**1a**), 4-fluorobenzaldehyde and ethyl 2-isocyanoacetate (**3b**) under optimal condition gave pyrrole (Table 2, entry **4u**) in 87% yield. To further extend the scope of this protocol we have carried out the reaction between acetophenone (**1a**), 4-bromobenzaldehyde and *tert*-butyl 2-isocyanoacetate (**3c**) to furnish the corresponding product (Table 2, entry **4v**) in 82% yield. Similarly, the reaction of acetone, acetaldehyde and *tert*-butyl 2-isocyanoacetate (**3c**) gave *tert*-butyl 4-acetyl-3-methyl-1*H*-pyrrole-2-carboxylate (Table 2, entry **4w**) in 53% yield.

Table 2. Scope of the construction of 2,3,4-trisubstituted 1*H*-pyrroles^{a,b}






^aReaction conditions: ketones **1** (1.0 mmol), aldehydes **2** (1.3 mmol) and Cs₂CO₃ (2.0 mmol) in methanol (3 mL) was stirred at R. T. for 2.0-5.5 h, CuI (10 mol %), alkyl isocynoacetate **3** (1.0 mmol) in methanol (2 mL) was added and stirring continued for 0.5 h, open air. ^bIsolated yields.


A possible mechanistic pathway of multicomponent coupled domino reaction is depicted in Scheme 3. First chalcone (**A**) is formed by the cross Aldol condensation of acetophenone (**1a**) and benzaldehyde (**2a**), followed by the reaction of methyl 2-isocynoacetate (**3a**) with copper iodide to give α -Cuprioisocyanide (**A'**). Then, [3+2] dipolar cycloaddition reaction of α -

Cuprioisocyanide (**A'**) and chalcone (**A**) give a cyclic organocopper intermediate (**B**) which on copper hydrogen exchange affords 3,4-dihydro-2*H*-pyrrole (**C**). Oxidation of the latter gives the methyl 4-benzoyl-3-phenyl-1*H*-pyrrole-2-carboxylate (**4a**).³⁷




Scheme 3. Proposed mechanism

Since the core objective of our research is to prepare 2,3,4-trisubstituted and 2,3,4,5-tetrasubstituted 1*H*-pyrroles for various promising applications, we explored the possibility of carrying out electrophilic substitution of 2,3,4-trisubstituted 1*H*-pyrrole (**4g**) at the C-5 position. We were delighted when the 5-bromo product (**5**) was obtained on treating pyrrole (**4g**) with *N*-bromosuccinimide in DMF at room temperature (Scheme 4). Thus, pyrrole (**4g**) is a possible intermediate for BODIPY dyes.



Scheme 4. Electrophilic substitution at C-5 position

Furthermore, the ester function in **4e** can be easily hydrolysed to the carboxylic acid (**6**), thereby offering a possible conjugation site for linking fluorophores available from such 2,3,4-trisubstituted 1*H*-pyrroles with biomolecules (Scheme 5).



Scheme 5: Hydrolysis of ester group

Conclusions

In conclusion, we have developed an operationally simple and highly efficient methodology for the synthesis of 2,3,4-trisubstituted 1*H*-pyrroles *via* a novel three-component coupled domino reaction. The striking feature of this protocol is the use of wide range of ketones, aldehydes and alkyl isocynoacetates to give the structural diversity of pyrroles at room temperature. This tandem process provides the moderate to excellent yields of the products (48-90%). Further use of these pyrroles in the synthesis of BODIPY dyes and establishment of conjugation with biomolecules is currently underway in our laboratory.

Experimental Section

General methods

Chemical reagents were obtained from commercial companies. All reactions were performed in round bottom flask and monitored by TLC performed on aluminium plates (0.25 mm, E. Merck) precoated with silica gel Merck 60 F-254. Developed TLC plates were visualized under a short-wavelength UV lamp. Reactions were conducted under open air, N₂, and O₂ atmosphere in solvents such as Acetonitrile, DCE, DME, dioxane, MeOH, THF, and DMF. Yields refer to spectroscopically (¹H, ¹³C NMR) homogeneous material obtained after column chromatography. Column chromatography was performed on silica gel (100-200 mesh size) supplied by S. D. Fine Chemicals Limited, India. IR spectra were recorded on a JASCO-FT/IR-4100 LE with attenuated total reflection (ATR) method. ¹H and ¹³C NMR were recorded in CDCl₃, CD₃OD and DMSO-*d*₆ solution with a Agilent 300, 500 and Brüker 400 MHz spectrometers. Chemical shifts (δ) are reported relative to SiMe₄ (δ = 0.0) as an internal standard. The number of protons (*n*) for a given resonance is indicated by *n*H. Peak multiplicities are designated by the following abbreviations: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; dd, doublet of doublet; br, broad; *J*, coupling constant in Hz. High-resolution mass spectra were obtained by using positive as well as negative electrospray ionization (ESI) by Time of Flight (TOF) method.

General experimental procedure for three-component coupled domino reaction:

A round bottom flask was charged with ketone **1** (1.0 mmol), aldehyde **2** (1.3 mmol) and Cs₂CO₃ (2.0 mmol) in methanol (3 mL) at R. T. TLC indicated the quantitative formation of chalcone after 2.0-5.5 h, to that, alkyl isocynoacetate **3** (1.0 mmol) and CuI (0.1 mmol) in

methanol (2 mL) was added. The reaction mixture was stirred at R.T. under open air atmosphere for 0.5 h. The solvent was removed under reduced pressure, the obtained residue was dissolved in ethyl acetate (30 mL) and washed with water. The above organic layer was dried over Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel (100-200 mesh size) with *n*-hexane : ethyl acetate (75 : 25) as eluent to afford desired product **4a-4w**.

Methyl 4-benzoyl-3-phenyl-1H-pyrrole-2-carboxylate 4a.

Off white solid, mp 140-142 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3256, 1724, 1694, 1632, 1508, 1264 and 761; ¹H NMR (400 MHz, CDCl₃) δ 9.82 (bs, 1H), 7.72 (dd, *J* = 7.2, 1.2 Hz, 2H), 7.46 (t, *J* = 7.4 Hz, 1H), 7.36-7.32 (m, 5H), 7.29-7.23 (m, 3H), 3.74 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 191.20, 161.40, 138.97, 133.12, 132.49, 132.00, 130.32, 129.41, 128.02, 127.74, 127.30, 127.21, 125.15, 120.44, 51.65; HRMS (ESI): calc. for [(C₁₉H₁₅NO₃)H] [M+H]⁻ 306.1131, found 306.1130.

Methyl 4-(4-methylbenzoyl)-3-(4-nitrophenyl)-1H-pyrrole-2-carboxylate 4b.

Pale yellow solid, mp 162-164 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3421, 1720, 1695, 1627, 1601, 1509, 1437, 1341 and 855; ¹H NMR (400 MHz, CDCl₃) δ 9.81 (bs, 1H), 8.15 (d, *J* = 8.4 Hz, 2H), 7.65 (d, *J* = 7.6 Hz, 2H), 7.50 (d, *J* = 8.8 Hz, 2H), 7.37 (d, *J* = 3.2 Hz, 1H), 7.19 (d, *J* = 8.0 Hz, 2H), 3.73 (s, 3H), 2.38 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 190.19, 160.86, 146.96, 143.42, 140.89, 136.07, 131.32, 130.08, 129.56, 129.10, 127.90, 125.09, 122.67, 121.03, 51.99, 21.66; HRMS (ESI): calc. for [(C₂₀H₁₆N₂O₅)H] [M+H]⁺ 365.1137, found 365.1111.

Methyl 4-(3-chlorobenzoyl)-3-(4-cyanophenyl)-1H-pyrrole-2-carboxylate 4c.

Off white solid, mp 182-184 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3273, 2234, 1727, 1707, 1644, 1517 and 746; ¹H NMR (400 MHz, CDCl₃) δ 9.57 (bs, 1H), 7.59 (distorted t, *J* = 7.2 Hz, 4H), 7.42 (distorted d, *J* = 7.6 Hz, 4H), 7.31 (t, *J* = 8.0, 7.2 Hz, 1H), 3.75 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 189.12, 160.77, 140.31, 138.41, 134.46, 132.33, 131.21, 130.35, 129.78, 129.41, 128.17, 127.28, 124.48, 121.21, 119.12, 111.04, 52.06; HRMS (ESI): calc. for [(C₂₀H₁₃ClN₂O₃)H] [M+H]⁺ 365.0693, found 365.0665.

Methyl 3-(4-fluorophenyl)-4-(4-methylbenzoyl)-1H-pyrrole-2-carboxylate 4d.

White solid, mp 128-130 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3316, 1728, 1692, 1634, 1517, 1263 and 841; ¹H NMR (400 MHz, CDCl₃) δ 9.56 (bs, 1H), 7.62 (d, *J* = 7.2 Hz, 2H), 7.34 (s, 1H), 7.28 (distorted t, *J* = 6.4, 6.0 Hz, 2H), 7.15 (d, *J* = 7.2 Hz, 2H), 6.95 (t, *J* = 8.4, 8.0 Hz, 2H), 3.74 (s, 3H), 2.37 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 190.74, 162.26 (d, *J*_{C-F} = 248.3 Hz), 161.25, 143.01, 136.26, 132.04 (d, *J*_{C-F} = 8.0 Hz), 131.99, 131.45, 129.65, 129.03, 128.88, 127.36, 125.34, 120.52, 114.40 (d, *J*_{C-F} = 21.4 Hz), 51.78, 21.64; HRMS (ESI): calc. for [(C₂₀H₁₆FNO₃)H] [M+H]⁺ 338.1192, found 338.1163.

Methyl 3-(4-bromophenyl)-4-(4-methylbenzoyl)-1H-pyrrole-2-carboxylate 4e.

Pale yellow solid, mp 166-168 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3262, 1730, 1692, 1636, 1504, 1261 and 754; ¹H NMR (400 MHz, CDCl₃) δ 9.57 (bs, 1H), 7.62 (d, *J* = 8.0 Hz, 2H), 7.39 (d, *J* = 8.0 Hz, 2H), 7.33 (d, *J* = 2.4 Hz, 1H), 7.20-7.15 (dd, *J* = 8.4, 8.0 Hz, 4H), 3.73 (s, 3H), 2.38 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 190.57, 161.21, 143.10, 136.23, 132.21, 132.04, 131.21, 130.57,

129.64, 128.94, 127.49, 125.19, 121.59, 120.54, 51.84, 21.66; HRMS (ESI): calc. for [(C₂₀H₁₆BrNO₃)H] [M+H]⁺ 398.0392, found 400.0354.

Methyl 4-benzoyl-3-(4-bromophenyl)-1H-pyrrole-2-carboxylate 4f.

Off white solid, mp 162-164 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3275, 1723, 1695, 1632, 1505, 1262 and 732; ¹H NMR (400 MHz, CDCl₃) δ 9.63 (bs, 1H), 7.71 (d, *J* = 7.6 Hz, 2H), 7.48 (t, *J* = 7.2 Hz, 1H), 7.40-7.34 (m, 5H), 7.19 (d, *J* = 8.0 Hz, 2H), 3.73 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 190.88, 161.20, 138.92, 132.28, 132.14, 132.03, 131.25, 130.57, 129.42, 128.25, 127.87, 124.99, 121.63, 120.69, 51.87; HRMS (ESI): calc. for [(C₁₉H₁₄BrNO₃)H] [M+H]⁺ 384.0235, found 386.0194.

Methyl 4-benzoyl-3-(4-chlorophenyl)-1H-pyrrole-2-carboxylate 4g.

Off white solid, mp 148-150 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3276, 1722, 1689, 1631, 1505, 1263 and 728; ¹H NMR (400 MHz, CDCl₃) δ 9.45 (bs, 1H), 7.71 (d, *J* = 7.2 Hz, 2H), 7.48 (distorted t, *J* = 7.2, 6.8 Hz, 1H), 7.36 (distorted t, *J* = 7.6, 7.2 Hz, 3H), 7.25 (s, 4H), 3.75 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 190.89, 161.19, 138.91, 133.33, 132.27, 131.71, 131.63, 131.24, 129.42, 128.23, 127.78, 127.64, 125.08, 120.72, 51.86; HRMS (ESI): calc. for [(C₁₉H₁₄ClNO₃)H] [M+H]⁺ 340.0740, found 340.0714.

Methyl 4-(3-chlorobenzoyl)-3-(4-chlorophenyl)-1H-pyrrole-2-carboxylate 4h.

White solid, mp 110-112 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3282, 1724, 1691, 1637, 1507, 1258 and 742; ¹H NMR (400 MHz, CDCl₃) δ 9.66 (bs, 1H), 7.62 (s, 1H), 7.57 (d, *J* = 7.6 Hz, 1H), 7.43 (d, *J* = 8.0 Hz, 2H), 7.30-7.23 (m, 5H), 3.74 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 189.52, 161.11, 140.42, 134.30, 133.48, 132.10, 131.68, 131.42, 131.09, 129.61, 129.49, 127.83, 127.69, 127.34, 124.72, 120.88, 51.92; HRMS (ESI): calc. for [(C₁₉H₁₃Cl₂NO₃)H] [M+H]⁺ 374.0350, found 374.0321.

Methyl 4-benzoyl-3-(4-fluorophenyl)-1H-pyrrole-2-carboxylate 4i.

White solid, mp 128-130 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3275, 1718, 1686, 1631, 1510, 1264 and 734; ¹H NMR (400 MHz, CDCl₃) δ 9.46 (bs, 1H), 7.69 (d, *J* = 7.2 Hz, 2H), 7.45 (d, *J* = 6.8 Hz, 1H), 7.37-7.25 (m, 5H), 6.94 (t, *J* = 8.0 Hz, 2H), 3.74 (s, 3H); ¹³C NMR (100 MHz, CD₃OD) δ 192.40, 162.10 (d, *J*_{C-F} = 243.7 Hz), 161.06, 139.18, 132.11 (d, *J*_{C-F} = 8.1 Hz), 131.86, 131.41, 129.92, 128.99, 128.86, 127.90, 124.13, 120.83, 113.53 (d, *J*_{C-F} = 21.5 Hz), 50.47; HRMS (ESI): calc. for [(C₁₉H₁₄FNO₃)H] [M+H]⁺ 324.1036, found 324.1005.

Methyl 4-(3-chlorobenzoyl)-3-(4-fluorophenyl)-1H-pyrrole-2-carboxylate 4j.

Off white solid mp 104-106 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3296, 1715, 1686, 1637, 1513, 1260 and 739; ¹H NMR (400 MHz, CDCl₃) δ 9.60 (bs, 1H), 7.61 (s, 1H), 7.55 (d, *J* = 7.6 Hz, 1H), 7.41 (d, *J* = 3.2 Hz, 2H), 7.28 (s, 3H), 6.95 (t, *J* = 8.8, 8.4 Hz, 2H), 3.74 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 189.74, 162.26 (d, *J*_{C-F} = 244.8 Hz), 161.18, 140.42, 134.24, 132.08, 132.05 (d, *J*_{C-F} = 7.9 Hz), 131.31, 129.55, 129.51, 128.79, 127.79, 127.34, 124.83, 120.79, 114.46 (d, *J*_{C-F} = 21.3 Hz), 51.88; HRMS (ESI): calc. for [(C₁₉H₁₃ClFNO₃)H] [M+H]⁺ 358.0656, found 358.0623.

Methyl 3-(4-bromophenyl)-4-(3-chlorobenzoyl)-1H-pyrrole-2-carboxylate 4k.

White solid, mp 144-146 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3298, 1730, 1698, 1630, 1505, 1255 and 744; ¹H NMR (400 MHz, CDCl₃) δ 9.69 (bs, 1H), 7.62 (s, 1H), 7.56 (d, *J* = 7.6 Hz, 1H), 7.44-7.38 (m, 4H), 7.28 (t, *J* = 7.6 Hz, 1H), 7.16 (d, *J* = 8.4 Hz, 2H), 3.74 (s, 3H); ¹³C NMR (100 MHz,

CDCl₃) δ 189.48, 161.11, 140.43, 134.31, 132.10, 131.98, 131.93, 131.09, 130.63, 129.63, 129.49, 127.89, 127.34, 124.65, 121.76, 120.85, 51.94; HRMS (ESI): calc. for [(C₁₉H₁₃BrClNO₃)H] [M+H]⁺ 417.9845, found 417.9822.

Methyl 4-acetyl-3-phenyl-1H-pyrrole-2-carboxylate 4l.

White solid, mp 170-172 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3250, 1728, 1697, 1638, 1512, 1270 and 768; ¹H NMR (400 MHz, CDCl₃) δ 9.81 (bs, 1H), 7.54 (d, *J* = 3.5 Hz, 1H), 7.37-7.34 (m, 3H), 7.30-7.28 (m, 2H), 3.63 (s, 3H), 2.02 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 194.43, 161.27, 134.02, 131.33, 129.83, 127.56, 127.47, 126.80, 126.37, 120.79, 51.49, 29.06; HRMS (ESI): calc. for [(C₁₄H₁₃NO₃)] [M]⁻ 243.0895 found 243.0817.

Methyl 4-acetyl-3-(*p*-tolyl)-1H-pyrrole-2-carboxylate 4m.

White solid, mp 184-186 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3253, 1726, 1694, 1642, 1516, 1268 and 765; ¹H NMR (400 MHz, CDCl₃) δ 9.85 (bs, 1H), 7.52 (d, *J* = 3.4 Hz, 1H), 7.24-7.15 (m, 4H), 3.64 (s, 3H), 2.37 (s, 3H), 2.03 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 194.61, 161.24, 137.11, 131.55, 130.83, 129.70, 128.33, 126.81, 126.41, 120.69, 51.44, 29.10, 21.24; HRMS (ESI): calc. for [(C₁₅H₁₅NO₃)-H] [M-H]⁻ 256.0973, found 256.0977.

Methyl 4-acetyl-3-(*m*-tolyl)-1H-pyrrole-2-carboxylate 4n.

White solid, mp 170-172 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3259, 1732, 1696, 1635, 1508, 1260 and 772; ¹H NMR (400 MHz, CDCl₃) δ 9.54 (bs, 1H), 7.54 (d, *J* = 3.5 Hz, 1H), 7.27 (d, *J* = 7.5 Hz, 1H), 7.15 (d, *J* = 7.6 Hz, 1H), 7.10-7.08 (m, 2H), 3.65 (s, 3H), 2.36 (s, 3H), 1.99 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 194.51, 161.13, 137.05, 133.79, 131.49, 130.48, 128.28, 127.47, 127.05, 126.90, 126.01, 120.69, 51.46, 29.10, 21.32; HRMS (ESI): calc. for [(C₁₅H₁₅NO₃)-H] [M-H]⁻ 256.0973, found 256.0974.

Methyl 4-acetyl-3-(*o*-tolyl)-1H-pyrrole-2-carboxylate 4o.

White solid, mp 162-164 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3248, 1731, 1687, 1629, 1502, 1255 and 774; ¹H NMR (400 MHz, CDCl₃) δ 9.77 (bs, 1H), 7.61 (d, *J* = 3.5 Hz, 1H), 7.27-7.16 (m, 3H), 7.11 (d, *J* = 7.9 Hz, 1H), 3.61 (s, 3H), 2.07 (s, 3H), 1.92 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 194.36, 161.19, 136.39, 134.05, 130.53, 129.53, 129.49, 127.72, 126.72, 126.49, 125.26, 120.70, 51.52, 28.54, 19.94; HRMS (ESI): calc. for [(C₁₅H₁₅NO₃)-H] [M-H]⁻ 256.0973, found 256.0972.

Methyl 4-acetyl-3-(4-bromophenyl)-1H-pyrrole-2-carboxylate 4p.

Off white solid, mp 150-152 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3275, 1734, 1690, 1640, 1513, 1250 and 765; ¹H NMR (400 MHz, CDCl₃) δ 9.89 (bs, 1H), 7.51 (d, *J* = 3.3 Hz, 1H), 7.47 (d, *J* = 8.5 Hz, 2H), 7.17 (d, *J* = 8.5 Hz, 2H), 3.65 (s, 3H), 2.13 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 193.76, 161.09, 132.88, 131.61, 130.63, 129.90, 126.78, 126.19, 121.65, 120.96, 51.59, 28.92; HRMS (ESI): calc. for [(C₁₄H₁₂BrNO₃)] [M]⁻ 321.0001 found 320.9922.

Methyl 4-acetyl-3-(4-chlorophenyl)-1H-pyrrole-2-carboxylate 4q.

Off white solid, mp 134-136 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3278, 1728, 1689, 1637, 1510, 1253 and 773; ¹H NMR (400 MHz, CDCl₃) δ 9.67 (bs, 1H), 7.44 (d, *J* = 3.3 Hz, 1H), 7.25 (d, *J* = 8.3 Hz, 2H), 7.15 (d, *J* = 8.2 Hz, 2H), 3.57 (s, 3H), 2.04 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 193.71, 161.06, 133.45, 132.35, 131.28, 129.90, 127.72, 126.62, 126.35, 121.01, 51.59, 28.93; HRMS (ESI): calc. for [(C₁₄H₁₂ClNO₃)H] [M+H]⁻ 278.0585, found 278.0584.

Methyl 4-acetyl-3-(4-methoxyphenyl)-1H-pyrrole-2-carboxylate 4r.

White solid, mp 132-134 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3251, 1720, 1694, 1625, 1501, 1254 and 763; ¹H NMR (400 MHz, CDCl₃) δ 9.73 (bs, 1H), 7.52 (d, *J* = 3.5 Hz, 1H), 7.21 (d, *J* = 8.8 Hz, 2H), 6.90 (d, *J* = 8.7 Hz, 2H), 3.82 (s, 3H), 3.65 (s, 3H), 2.04 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 194.53, 161.24, 159.00, 131.19, 131.05, 126.85, 126.34, 125.89, 120.75, 113.05, 55.09, 51.47, 29.09; HRMS (ESI): calc. for [(C₁₅H₁₄NO₄)-H] [M-H]⁻ 272.0922 found 272.0923.

Methyl 4-acetyl-3-(4-nitrophenyl)-1H-pyrrole-2-carboxylate 4s.

Pale yellow solid, mp 144-146 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3415, 1724, 1698, 1632, 1605, 1513, 1443, 1345 and 858; ¹H NMR (400 MHz, CDCl₃) δ 9.78 (bs, 1H), 8.20 (d, *J* = 8.6 Hz, 2H), 7.56 (d, *J* = 3.3 Hz, 1H), 7.45 (d, *J* = 8.6 Hz, 2H), 3.65 (s, 3H), 2.25 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 192.72, 160.67, 147.08, 141.28, 130.99, 128.68, 126.98, 125.72, 122.54, 121.36, 51.77, 28.47; HRMS (ESI): calc. for [(C₁₄H₁₂N₂O₅)H] [M+H]⁻ 289.0825, found 289.0824.

Methyl 4-acetyl-3-methyl-1H-pyrrole-2-carboxylate 4t.

White solid, mp 130-132 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3254, 1732, 1690, 1633, 1505, 1261 and 763; ¹H NMR (400 MHz, CDCl₃) δ 9.47(bs, 1H), 7.43 (d, *J* = 3.4 Hz, 1H), 3.85 (s, 3H), 2.59 (s, 3H), 2.39 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 194.04, 162.02, 129.17, 127.20, 125.54, 121.16, 51.46, 28.25, 11.58; HRMS (ESI): calc. for [(C₉H₁₁NO₃)H] [M+H]⁻ 182.0818, found 182.0817.

Ethyl 4-benzoyl-3-(4-fluorophenyl)-1H-pyrrole-2-carboxylate 4u.

White solid, mp 132-134 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3278, 1716, 1685, 1629, 1512, 1266 and 736; ¹H NMR (500 MHz, CDCl₃) δ 9.49 (bs, 1H), 7.70 (distorted dd, *J* = 8.0, 1.5 Hz, 2H), 7.47 (tt, *J* = 7.5, 1.5 Hz, 1H), 7.38 (d, *J* = 3.0 Hz, 1H), 7.34 (t, *J* = 8.0, 7.5 Hz, 2H), 7.31-7.27 (m, 2H), 6.95 (tt, *J* = 9.0, 3.0, 2.5, 2.0 Hz, 2H), 4.21 (q, *J* = 7.0 Hz, 2H), 1.16 (t, *J* = 7.0 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 190.77, 162.30 (d, *J*_{C-F} = 248.0 Hz), 161.28, 143.04, 136.30, 132.07 (d, *J*_{C-F} = 8.0 Hz), 131.48, 129.69, 129.07, 128.92, 127.40, 125.38, 120.55, 114.42 (d, *J*_{C-F} = 21.0 Hz), 61.24, 13.85; HRMS (ESI): calc. for [(C₂₀H₁₆FNO₃)H] [M+H]⁺ 338.1192, found 338.1197.

tert-Butyl 3-(4-bromophenyl)-4-(3-chlorobenzoyl)-1H-pyrrole-2-carboxylate 4v.

Light yellow solid, mp 158-160 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3296, 1726, 1696, 1632, 1501, 1252 and 739; ¹H NMR (400 MHz, CDCl₃) δ 9.68 (bs, 1H), 7.61 (s, 1H), 7.55 (d, *J* = 7.6 Hz, 1H), 7.43-7.37 (m, 4H), 7.27 (t, *J* = 7.6 Hz, 1H), 7.15 (d, *J* = 8.4 Hz, 2H), 1.54 (s, 9H); ¹³C NMR (100 MHz, CDCl₃) δ 189.47, 161.10, 140.42, 134.30, 132.09, 131.97, 131.92, 131.09, 130.62, 129.62, 129.48, 127.88, 127.33, 124.64, 121.75, 120.84, 81.82, 28.83; HRMS (ESI): calc. for [(C₂₂H₁₉BrClNO₃)H] [M+H]⁺ 460.0315, found 460.0317.

tert-butyl 4-acetyl-3-methyl-1H-pyrrole-2-carboxylate 4w.

White solid, mp 138-140 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3251, 1730, 1694, 1629, 1501, 1265 and 760; ¹H NMR (400 MHz, CDCl₃) δ 9.49(bs, 1H), 7.45 (d, *J* = 3.2 Hz, 1H), 2.61 (s, 3H), 2.41 (s, 3H), 1.54 (s, 9H); ¹³C NMR (100 MHz, CDCl₃) δ 194.06, 162.04, 129.18, 127.23, 125.55, 121.18, 81.72, 29.02, 28.26, 11.60; HRMS (ESI): calc. for [(C₁₂H₁₇NO₃)H] [M+H]⁺ 224.1286, found 224.1289.

Experimental procedure for bromination of 4g:

A round bottom flask was charged with methyl 4-benzoyl-3-(4-chlorophenyl)-1*H*-pyrrole-2-carboxylate **4g** (1.0 mmol), *N*-bromosuccinimide (1.2 mmol) and 5 mL DMF. The reaction mixture was stirred at 0 °C-R.T. for 2 h under nitrogen atmosphere. The reaction mixture was quenched with water (10 mL), extracted using ethyl acetate (3×10 mL). The combined organic layer was washed with brine solution and dried over Na₂SO₄ to afford the crude product. The crude product was purified by column chromatography on silica gel (100-200 mesh size) with *n*-hexane:ethyl acetate (80:20) as eluent to afford desired product **5**.

Methyl 4-benzoyl-5-bromo-3-(4-chlorophenyl)-1*H*-pyrrole-2-carboxylate 5.

Yield 88%, white solid, mp 172-174 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3287, 1711, 1515, 1439, 1282, and 783; ¹H NMR (300 MHz, CDCl₃) δ 10.16 (bs, 1H), 7.68-7.65 (m, 2H), 7.47-7.41 (m, 1H), 7.31-7.26 (m, 2H), 7.19-7.12 (m, 4H) 3.76 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 191.47, 160.72, 137.57, 133.76, 133.14, 131.78, 131.70, 130.88, 129.93, 128.32, 127.83, 125.28, 120.41, 107.81, 52.13; calc. for [(C₁₉H₁₃BrClNO₃)H] [M+H]⁺ 417.9845, found 417.9875.

Experimental procedure for hydrolysis of 4e:

A round bottom flask was charged with methyl 3-(4-bromophenyl)-4-(4-methylbenzoyl)-1*H*-pyrrole-2-carboxylate **4e** (1.0 mmol), LiOH.H₂O (10.0 mmol) and 5 mL THF:H₂O [1:1]. The reaction mixture was heated at 80 °C for 8 h. The reaction mixture was cooled to R.T., diluted with water (5 mL) and extracted using ethyl acetate (3×10 mL). The dried organic layer was concentrated under reduced pressure to afford the desired product **6**.

3-(4-bromophenyl)-4-(4-methylbenzoyl)-1*H*-pyrrole-2-carboxylic acid 6.

Yield 85%, Brown solid, mp 214-216 °C; IR (ATR) $\tilde{\nu}$ (cm⁻¹): 3214, 1690, 1629, 1603, 1500, 1402, and 754; ¹H NMR (300 MHz, DMSO-*d*₆) δ 12.04 (bs, 1H), 7.55 (d, *J* = 7.5 Hz, 2H), 7.25-7.17 (m, 6H), 7.03 (s, 1H), 2.35 (s, 3H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 190.00, 165.29, 141.71, 137.23, 134.92, 133.06, 129.36, 129.07, 128.78, 127.58, 126.83, 125.78, 122.55, 118.86, 21.15; HRMS (ESI): calc. for [(C₁₉H₁₄BrNO₃)H] [M+H]⁺ 384.0235, found 386.0249.

Acknowledgement

The financial support from the Council of Scientific and Industrial Research, New Delhi (sanction no. 01(2427)/10/EMR-II dated 28/12/2010 and pool scheme no. 8644-A), India, to carry out this work is gratefully acknowledged. H.P.K. and K.S.V. thank Council of Scientific and Industrial Research and University Grants Commission-Special Assistance Programme, New Delhi respectively for award of Senior Research Fellowships. A.C.C. thank DST for offering Fast Track Project SB/FT/CS-147/2013. The authors thank Prof. Prakash M. Bhate for his generous support.

Author Contributions

^aThese authors contributed equally to this work.

References

- 1 (a) D. E. N. Jacquot, M. Zöllinger and T. Lindel, *Angew. Chem.*, 2005, **117**, 2336–2338; (b) D. E. N. Jacquot, M. Zöllinger and T. Lindel, *Angew. Chem., Int. Ed.*, 2005, **44**, 2295–2298; (c) H. Garrido-Hernandez, M. Nakadai, M. Vimolratana, Q. Li, T. Doundoulakis and P. G. Harran, *Angew. Chem.*, 2005, **117**, 775–779; (d) H. Garrido-Hernandez, M. Nakadai, M. Vimolratana, Q. Li, T. Doundoulakis and P. G. Harran, *Angew. Chem., Int. Ed.*, 2005, **44**, 765–769; (e) H. Hoffmann and T. Lindel, *Synthesis*, 2003, 1753–1783; (f) D. Seidel, V. Lynch and J. L. Sessler, *Angew. Chem.*, 2002, **114**, 1480–1483; (g) D. Seidel, V. Lynch and J. L. Sessler, *Angew. Chem., Int. Ed.*, 2002, **41**, 1422–1425; (h) J. A. Johnson, N. Li and D. Sames, *J. Am. Chem. Soc.*, 2002, **124**, 6900–6903; (i) G. Dannhardt and W. Kiefer, *Arch. Pharm.*, 2001, **334**, 183–188; (j) T. Lindel, M. Hochgürtel, M. Assmann and M. Köck, *J. Nat. Prod.*, 2000, **63**, 1566–1569; (k) M. Adamczyk, D. D. Johnson and R. E. Reddy, *Angew. Chem.*, 1999, **111**, 3751–3753; (l) M. Adamczyk, D. Johnson and R. Reddy, *Angew. Chem., Int. Ed.*, 1999, **38**, 3537–3539; (m) S. Depraetere, M. Smet and W. Dehaen, *Angew. Chem.*, 1999, **111**, 3556–3558; (n) S. Depraetere, M. Smet and W. Dehaen, *Angew. Chem., Int. Ed.*, 1999, **38**, 3359–3361; (o) J. M. Gottesfeld, L. Neely, J. W. Trauger, E. E. Baird and P. B. Dervan, *Nature*, 1997, **387**, 202–205; (p) A. V. Gulevich, A. S. Dudnik, N. Chernyak and V. Gevorgyan, *Chem. Rev.*, 2013, **113**, 3084–3213; (q) M. A. Yurovskaya and R. S. Alekseyev, *Chem. Heterocycl. Compd.*, 2014, **49**, 1400–1425; (r) S. D. Joshi, U. A. More, V. H. Kulkarni and T. M. Aminabhavi, *Curr. Org. Chem.*, 2013, **17**, 2279–2304.
- 2 (a) K. Yamaji, M. Masubuchi, F. Kawahara, Y. Nakamura, A. Nishio, S. Matsukuma, M. Fujimori, N. Nakada, J. Watanabe and T. Kamiyama, *J. Antibiot.*, 1997, **50**, 402–411; (b) B. Fournier and D. C. Hooper, *Antimicrob. Agents Chemother.*, 1998, **42**, 121–128; (c) D. Perrin, B. van Hille, J.-M. Barret, A. Kruczynski, C. Etiévant, T. Imbert and B. T. Hill, *Biochem. Pharmacol.*, 2000, **59**, 807–819; (d) F. Micheli, R. Di Fabio, R. Benedetti, A. Maria Capelli, P. Cavallini, P. Cavanni, S. Davalli, D. Donati, A. Feriani, S. Gehanne, M. Hamdan, M. Maffeis, F. Maria Sabbatini, M. Elvira Tranquillini and M. V. Angela Viziano, *Farm.*, 2004, **59**, 175–183; (e) B. D. Roth, C. J. Blankley, A. W. Chucholowski, E. Ferguson, M. L. Hoefle, D. F. Ortwine, R. S. Newton, C. S. Sekerke, D. R. Sliskovic and M. Wilson, *J. Med. Chem.*, 1991, **34**, 357–366.
- 3 (a) H. Miyaji, W. Sato and J. L. Sessler, *Angew. Chem.*, 2000, **112**, 1847–1850; (b) H. Miyaji, W. Sato and J. Sessler, *Angew. Chem., Int. Ed.*, 2000, **39**, 1777–1780; (c) F.-P. Montforts and O. Kutzki, *Angew. Chem.*, 2000, **112**, 612–614; (d) F. Montforts and O. Kutzki, *Angew. Chem., Int. Ed.*, 2000, **39**, 599–601; (e) D.-W. Yoon, H. Hwang and C.-H. Lee, *Angew. Chem.*, 2002, **114**, 1835–1837; (f) D.-W. Yoon, H. Hwang and C.-H. Lee,

- Angew. Chem., Int. Ed.*, 2002, **41**, 1757–1759; (g) J. O. Jeppesen and J. Becher, *Eur. J. Org. Chem.*, 2003, **2003**, 3245–3266.
- 4 (a) M. Lazerges, K. I. Chane-Ching, S. Aeiyaich, S. Chelli, B. Peppin-Donnat, M. Billon, C. Lombard, F. Maurel and M. Jouini, *J. Solid State Electrochem.*, 2008, **13**, 231–238; (b) A. Kros, S. W. F. M. van Hövel, R. J. M. Nolte and N. A. J. M. Sommerdijk, *Sensors Actuators B Chem.*, 2001, **80**, 229–233; (c) M. C. Moreno-Bondi, G. Orellana, S. Hamilton, M. J. Hephher and J. Sommerville, *Sensors Actuators B Chem.*, 2005, **107**, 424–432; (d) A. Loudet and K. Burgess, *Chem. Rev.*, 2007, **107**, 4891–4932; (e) G. Ulrich, R. Ziessel and A. Harriman, *Angew. Chem., Int. Ed.*, 2008, **47**, 1184–1201; (f) N. Boens, V. Leen and W. Dehaen, *Chem. Soc. Rev.*, 2012, **41**, 1130–1172.
- 5 (a) A. Hantzsch, *Ber. Dtsch. Chem. Ges.*, 1890, **23**, 1474–1476; (b) A. Herath and N. D. P. Cosford, *Org. Lett.*, 2010, **12**, 5182–5185; (c) T. A. Moss and T. Nowak, *Tetrahedron Lett.*, 2012, **53**, 3056–3060; (d) V. Estévez, M. Villacampa and J. C. Menéndez, *Chem. Commun.*, 2013, **49**, 591–593; (e) F. Palacios, D. Aparicio, J. M. de los Santos and J. Vicario, *Tetrahedron*, 2001, **57**, 1961–1972.
- 6 (a) C. Paal, *Ber. Dtsch. Chem. Ges.*, 1884, **17**, 2756–2767; (b) L. Knorr, *Ber. Dtsch. Chem. Ges.*, 1884, **17**, 2863–2870; (c) A. R. Bharadwaj and K. A. Scheidt, *Org. Lett.*, 2004, **6**, 2465–2468.
- 7 (a) D. van Leusen, E. van Echten and A. M. van Leusen, *J. Org. Chem.*, 1992, **57**, 2245–2249; (b) A. M. van Leusen, H. Siderius, B. E. Hoogenboom and D. van Leusen, *Tetrahedron Lett.*, 1972, **13**, 5337–5340.
- 8 G. M. Robinson and R. Robinson, *J. Chem. Soc. Trans.*, 1918, **113**, 639–645.
- 9 V. Nair, A. U. Vinod and C. Rajesh, *J. Org. Chem.*, 2001, **66**, 4427–4429.
- 10 (a) S. Chiba, Y. Wang, G. Lapointe and K. Narasaka, *Org. Lett.*, 2008, **10**, 313–316; (b) Y.-F. Wang, K. K. Toh, S. Chiba and K. Narasaka, *Org. Lett.*, 2008, **10**, 5019–5022.
- 11 G. Minetto, L. F. Raveglia, A. Segal and M. Taddei, *Eur. J. Org. Chem.*, 2005, **2005**, 5277–5288.
- 12 (a) R. ten Have, F. R. Leusink and A. M. van Leusen, *Synthesis*, 1996, 871–876; (b) H. P. Dijkstra, R. ten Have and A. M. van Leusen, *J. Org. Chem.*, 1998, **63**, 5332–5338.
- 13 (a) L. F. Tietze, G. Brasche and K. M. Gericke, *Domino Reactions in Organic Synthesis*, Wiley-VCH, Weinheim, 2006; (b) A. S. Ivanov, *Chem. Soc. Rev.*, 2008, **37**, 789–811; (c) A. de Meijere, P. von Zezschwitz and S. Bräse, *Acc. Chem. Res.*, 2005, **38**, 413–422.
- 14 P. T. Anastas and J. C. Warner, *Green Chemistry: Theory and Practice*, Oxford University Press, New York, 1998, p.30.
- 15 A. Shaabani, M. B. Teimouri and S. Arab-Ameri, *Tetrahedron Lett.*, 2004, **45**, 8409–8413.
- 16 (a) T. Takizawz, N. Obata, Y. Suzuki and T. Yanagida, *Tetrahedron Lett.*, 1969, **10**, 3407–3410; (b) V. Nair, A. U. Vinod, N. Abhilash, R. S. Menon, V. Santhi, R. L. Varma, S. Viji, S. Mathew and R. Srinivas, *Tetrahedron*, 2003, **59**, 10279–10286.
- 17 C. Cao, Y. Shi and A. L. Odom, *J. Am. Chem. Soc.*, 2003, **125**, 2880–2881.
- 18 E. Barnea, S. Majumder, R. J. Staples and A. L. Odom, *Organometallics*, 2009, **28**, 3876–3881.

- 19 S. V. Ley, I. R. Baxendale, R. N. Bream, P. S. Jackson, A. G. Leach, D. A. Longbottom, M. Nesi, J. S. Scott, R. I. Storer and S. J. Taylor, *J. Chem. Soc., Perkin Trans. 1*, 2000, 3815–4195.
- 20 A. Solinas and M. Taddei, *Synthesis*, 2007, 2409–2453.
- 21 (a) T. Saegusa, Y. Ito, H. Kinoshita and S. Tomita, *J. Org. Chem.*, 1971, **36**, 3316–3323; (b) H. Takaya, S. Kojima and S.-I. Murahashi, *Org. Lett.*, 2001, **3**, 421–424; (c) Y. Motoyama, H. Kawakami, K. Shimozono, K. Aoki and H. Nishiyama, *Organometallics*, 2002, **21**, 3408–3416; (d) Y. Ito, M. Sawamura and T. Hayashi, *J. Am. Chem. Soc.*, 1986, **108**, 6405–6406; (e) A. Togni and S. D. Pastor, *J. Org. Chem.*, 1990, **55**, 1649–1664; (f) M. Sawamura, H. Hamashima and Y. Ito, *J. Org. Chem.*, 1990, **55**, 5935–5936; (g) T. Hayashi, E. Kishi, V. A. Soloshonok and Y. Uozumi, *Tetrahedron Lett.*, 1996, **37**, 4969–4972.
- 22 (a) F. Gerhart and U. Schöllkopf, *Tetrahedron Lett.*, 1968, **9**, 6231–6234; (b) A. M. van Leusen, J. Wildeman and O. H. Oldenzien, *J. Org. Chem.*, 1977, **42**, 1153–1159; (c) D. H. R. Barton and S. Z. Zard, *J. Chem. Soc., Chem. Commun.*, 1985, 1098–1100; (d) H. Saikachi, T. Kitagawa and H. Sasaki, *Chem. Pharm. Bull.*, 1979, **27**, 2857–2861; (e) T. Murakami, M. Otsuka and M. Ohno, *Tetrahedron Lett.*, 1982, **23**, 4729–4732.
- 23 I. Baxendale, C. Buckle, S. Ley and L. Tamborini, *Synthesis*, 2009, 1485–1493.
- 24 H. Uno, M. Tanaka, T. Inoue and N. Ono, *Synthesis*, 1999, 471–474.
- 25 W. Chen, J. Shao, Z. Li, M. A. Giulianotti and Y. Yu, *Can. J. Chem.*, 2012, **90**, 214–221.
- 26 S. Kamijo, C. Kanazawa and Y. Yamamoto, *J. Am. Chem. Soc.*, 2005, **127**, 9260–9266.
- 27 O. V. Larionov and A. de Meijere, *Angew. Chem., Int. Ed.*, 2005, **44**, 5664–5667.
- 28 J. Liu, Z. Fang, Q. Zhang, Q. Liu and X. Bi, *Angew. Chem., Int. Ed.*, 2013, **52**, 6953–6957.
- 29 M. Gao, C. He, H. Chen, R. Bai, B. Cheng and A. Lei, *Angew. Chem., Int. Ed.*, 2013, **52**, 6958–6961.
- 30 J. Song, C. Guo, P.-H. Chen, J. Yu, S.-W. Luo and L.-Z. Gong, *Chem. Eur. J.*, 2011, **17**, 7786–7790.
- 31 R. Grigg, M. I. Lansdell and M. Thornton-Pett, *Tetrahedron*, 1999, **55**, 2025–2044.
- 32 C. Arróniz, A. Gil-González, V. Semak, C. Escolano, J. Bosch and M. Amat, *Eur. J. Org. Chem.*, 2011, **2011**, 3755–3760.
- 33 (a) R. Wang, X.-P. Xu, H. Meng, S.-Y. Wang and S.-J. Ji, *Tetrahedron*, 2013, **69**, 1761–1766; (b) R. Wang, S.-Y. Wang and S.-J. Ji, *Tetrahedron*, 2013, **69**, 10836–10841; (c) X. Wang, X.-P. Xu, S.-Y. Wang, W. Zhou and S.-J. Ji, *Org. Lett.*, 2013, **15**, 4246–4249; (d) R. Wang, S.-Y. Wang and S.-J. Ji, *Org. Biomol. Chem.*, 2014, **12**, 1735–1740.
- 34 D. Zheng, S. Lia and J. Wu, *Chem. Commun.*, 2012, **48**, 8568–8570.
- 35 (a) V. Estévez, M. Villacampa and J. C. Menéndez, *Chem. Soc. Rev.*, 2014, **43**, 4633–4657; (b) V. Estévez, M. Villacampa and J. C. Menéndez, *Chem. Soc. Rev.*, 2010, **39**, 4402–4421; (c) N. A. Keiko, N. V. Vchislo, E. A. Verochkina and L. I. Larina, *Tetrahedron*, 2014, **70**, 8959–8970; (d) R. Pagadala, D. R. Kommidi, S. Kankala, S. Maddila, P. Singh, B. Moodley, N. A. Koorbanally and S. B. Jonnalagadda, *Org. & Biomol. Chem.*, 2015, **13**, 1800–1806; (e) G. Balme, *Angew. Chem., Int. ed.*, 2004, **43**,

- 6238–6241; (f) X. Lin, Z. Mao, X. Dai, P. Lu and Y. Wang, *Chem. comm.*, 2011, **47**, 6620–6622; (g) B. Khalili, P. Jajarmi, B. Eftekhari-sis and M. M. Hashemi, *J. Org. Chem.*, 2008, **73**, 2090–2095; (h) O. A. Attanasi, G. Favi, F. Mantellini, G. Moscatelli, S. Santeusano, C. Organica, U. Carlo and V. I. Maggetti, *J. Org. Chem.*, 2011, **76**, 2860–2866; (i) B. Jiang, Y. Li, M. Tu, S. Wang, S. Tu and G. Li, *J. Org. Chem.*, 2012, **77**, 7497–7505; (j) X. Liu, L. Hao, M. Lin, L. Chen and Z. Zhan, *Org. & Biomol. Chem.*, 2010, **8**, 3064–3072 (k) A. B. Atar, J. S. Kim, K. T. Lim and Y. T. Jeong, *New J. Chem.*, 2015, **39**, 396–402; (l) X. Feng, Q. Wang, W. Lin, G. Dou and Z. Huang, *Org. Lett.*, 2013, **15**, 2542–2545; (m) D. J. S. Cyr, N. Martin and B. A. Arndtsen, *Org. Lett.*, 2007, **9**, 449–452; (n) N. Bhunia and B. Das, *Synthesis*, 2013, **45**, 1045–1050; (o) P. Radha Krishna Murthi, D. Rambabu, M. V. Basaveswara Rao and M. Pal, *Tetrahedron Letters*, 2014, **55**, 507–509.
- 36 (a) K. S. Vadagaonkar, H. P. Kalmode, S. Prakash and A. C. Chaskar, *New J. Chem.*, DOI: 10.1039/c4nj02345e; (b) H. P. Kalmode, K. S. Vadagaonkar and A. C. Chaskar, *Synthesis*, 2015, **47**, 429–438; (c) K. S. Vadagaonkar, H. P. Kalmode, K. Murugan and A. C. Chaskar, *RSC Adv.*, 2015, **5**, 5580–5590; (d) H. P. Kalmode, K. S. Vadagaonkar and A. C. Chaskar, *RSC Adv.*, 2014, **4**, 60316–60326; (e) K. S. Vadagaonkar, K. Murugan, A. C. Chaskar and P. M. Bhate, *RSC Adv.*, 2014, **4**, 34056–34064; (f) B. Pawar, V. Padalkar, K. Phatangare, S. Nirmalkar and A. Chaskar, *Catal. Sci. Technol.*, 2011, **1**, 1641–1644; (g) S. Takale, S. Parab, K. Phatangare, R. Pisal and A. Chaskar, *Catal. Sci. Technol.*, 2011, **1**, 1128–1132; (h) A. Chaskar, V. Padalkar, K. Phatangare, B. Langi and C. Shah, *Synth. Commun.*, 2010, **40**, 2336–2340; (i) K. Phatangare, V. Padalkar, D. Mhatre, K. Patil and A. Chaskar, *Synth. Commun.*, 2009, **39**, 4117–4121; (j) A. Chaskar, V. Padalkar, K. Phatangare, K. Patil, A. Bodkhe and B. Langi, *Appl. Catal. A*, 2009, **359**, 84–87.
- 37 (a) Q. Cai, F. Zhou, T. Xu, L. Fu and K. Ding, *Org. Lett.*, 2011, **13**, 340–343; (b) F. Zhou, L. Fu, J. Wie, K. Ding, Q. Cai, *Synthesis*, 2011, 3037–3044.