



NPR

Calyculin: Nature's Way of Making the Sponge-derived Cytotoxin

Journal:	<i>Natural Product Reports</i>
Manuscript ID	NP-HIG-10-2015-000123.R2
Article Type:	Highlight
Date Submitted by the Author:	15-Feb-2016
Complete List of Authors:	Wakimoto, Toshiyuki; Hokkaido Univ, Egami, Yoko; Hokkaido Univ, Abe, Ikuro; The University of Tokyo, Graduate School of Pharmaceutical Sciences

SCHOLARONE™
Manuscripts

Calyculin: Nature's Way of Making the Sponge-derived Cytotoxin

Toshiyuki Wakimoto,[†] Yoko Egami,[†] Ikuro Abe[‡]

[†] Faculty of Pharmaceutical Sciences, Hokkaido University, Kita 12, Nishi 6, Kita-ku, Sapporo 060-0812, Japan

[‡] Graduate School of Pharmaceutical Sciences, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

Contents

- 1. Introduction**
- 2. Structure-activity relationship of calyculin A**
- 3. Biosynthetic gene cluster of calyculin A**
- 4. Phosphocalyculin A as a protoxin**
- 5. Calyculin A production in the sponge-microbe association**
- 6. Conclusions**

ABSTRACT: Calyculin A is a major cytotoxic compound isolated from the Japanese marine sponge *Discodermia calyx*. Its potent cytotoxicity is attributable to the specific inhibition of protein phosphatases 1 and 2A, as in the cases of okadaic acid and microcystins. Its chemical structure is well-designed for not only enzyme inhibition but also higher membrane permeability to impart its potent cytotoxicity. The biosynthetic gene cluster of this densely functionalized polyketide and nonribosomal peptide hybrid molecule was recently identified from the sponge-microbe association. The producer organism and the dynamic bioconversion process were also revealed. In this highlight, we focus on the recent studies addressing nature's design and biogenesis of the sponge-derived cytotoxin, calyculin A.

1. Introduction

The marine sponge *Discodermia calyx* inhabits relatively shallow water in the central to southern areas of Japan. Originally, this marine sponge was part of the vast collection of Japanese marine organisms amassed by the German biologist Ludwig Döderlein, who was a visiting Professor at the University of Tokyo from 1879-1881, and it was identified based on the specimen collected in the Gulf of Sagami, Japan.¹ Almost one hundred years later, Fusetani and co-workers reported that the marine sponge contains a major secondary metabolite, calyculin A (Fig. 1, **1**, 0.15% wet weight), showing potent activity in the starfish egg assay.² Subsequently, **1** was revealed to have pM range IC₅₀ values against several cancer cell lines and tumor promotion activity, ascribed to the specific and potent inhibition of protein phosphatases 1 and 2A (PP1 and 2A),³ thereby rendering it a valuable tool to evaluate intracellular signal transduction. Its densely functionalized structure is naturally well-designed for the specific inhibition of PPs as well as cytotoxicity, and it has also attracted considerable attention from synthetic chemists, which resulted in six total syntheses.⁴

The natural source of calyculins is not limited to *D. calyx*, as other sponge species also contain calyculin derivatives, such as calyculinamides from the New Zealand sponge *Lamellomorpha strongylata*,⁵ clavosines (**10**) from the Palauan sponge *Myriastra clavosa*,⁶ geometricin (**11**) from the Australian sponge *Lufferiella geometrica*,⁷ and swinhoeiamide (**12**) from the Papua New Guinean sponge *Theonella swinhoei* (Fig 2).⁸ However, no other organisms except for marine sponges have been reported to contain calyculin analogs. The cross-genera distribution of calyculin-related compounds suggests that they are produced by common or similar symbionts primarily associated with the marine sponges. Recently, the biosynthetic gene cluster and the producer organism were identified in the Japanese marine sponge *D. calyx*.⁹ This highlight focuses on the biological activity, the biosynthesis, and the producer organism of calyculins.

2. Structure-activity relationship of calyculin A

Phosphatase inhibitors, in addition to **1**, have been isolated from different biological sources, including cantharidin (**17**) from blister beetles,¹⁰ microcystins (**18**) from cyanobacteria,¹¹ and okadaic acid (**19**) from marine sponges¹² and dinoflagellates (Fig. 3).¹³ Actinomycetes are also important sources, producing tautomycin (**20**),¹⁴ fostriecin (**21**)¹⁵ and phoslactomycin (**22**) (Fig. 3).¹⁶ Notably, in addition to the distinct origins, the structural features of this class of molecules are also diverse, and include terpenoids such as cantharidin, mycrocystins with a peptidic nature, and polyketides including okadaic acid, tautomycin, fostriecin and phoslactomycin. In spite of the stark differences in their chemical structures, these natural molecules can inhibit the same enzyme in a competitive manner, although some inhibitors exhibited varied selectivities on PP1 and PP2A, c.f. moderate selectivity against PP2A by okadaic acid¹⁷ and fostriecin,¹⁵ and higher specificity of tautomycin¹⁴ towards PP1 than PP2A. This fact invokes the question of what common structural motifs are essential for the specific inhibition of protein phosphatases. Since all molecules possess a hydrophobic moiety, in addition to an acidic functionality as the phosphate mimic, these common structural features had been suggested to be indispensable for the enzyme inhibition.¹⁸⁻²⁰ In fact, the X-ray crystal structure of the complex between natural inhibitors and PPs has been found to firmly support the molecular basis for this proposal. Based on the crystal structure, the acidic functional groups of the PP inhibitors interact with the phosphate recognizing residue, Arg, of the enzyme. Furthermore, one of the Y-shaped grooves on the surface of the enzyme, which is occupied by hydrophobic amino acid residues (hydrophobic groove), indeed, interacts with the hydrophobic tails of the natural inhibitors.²¹⁻²⁴

Detailed structure-activity relationship studies have also been conducted on calyculins, with the natural variants isolated from the extract of the sponge *D. calyx* as well as semisynthetic analogs.²⁵ The tetraene moiety of **1** is photoisomerized into geometrical isomers, calyculins B (**2**), E (**6**), and F (**7**) (Fig. 1).^{26,27} Calyculin C (**3**) has an additional methyl group on C32 of **1**, and the geometrical isomers

of this series are calyculins D (**4**), G (**8**), and H (**9**) (Fig. 1).^{26,27} The tetraene group corresponds to the hydrophobic motif found in other phosphatase inhibitors, and therefore the IC_{50} values of the geometrical isomers decrease by up to 3 orders of magnitude. The semisynthetic 11,13-*O*-isopropylidene-calyculin A²⁵ (**13**) and the natural derivative, dephosphonocalyculin A²⁸ (**14**) (Fig. 2), completely lacked both the enzyme inhibition and cytotoxicity activities, demonstrating that the phosphate and the 1,3-diol are essential for the inhibitory activity against protein phosphatases. The considerable retention of the activity by calyculin J²⁹ (Fig. 2, **15**) indicated the importance of the 13-OH, rather than the 11-OH, for enzyme inhibition. On the other hand, hemicalyculin A²⁵ (Fig. 2, **16**), which is terminated by a nitrile in place of the oxazole ring, but retains the phosphate group, 13-OH, and tetraene, showed potent enzyme inhibition activities against both PP1 and PP2A comparable to those of **1**. Therefore, the peptide portion composed of two γ -amino acids does not seem to be involved in the binding to PPs. In fact, the corresponding portion could not be detected in the X-ray crystal structure of the complex between **1** and PP1 γ due to the indistinct electron densities, strongly supporting the idea that this motif does not function in the interaction.²³

These findings then engendered the next question about the role of the peptide portion. Remarkably, the cytotoxicity of **16** was significantly diminished, and was more than four orders of magnitude weaker than that of **1**,²⁵ which clearly indicated that the peptide portion is essential for the cytotoxicity of **1**. Again, the X-ray structure provided insight into the proposed role of this moiety. The free state of **1** forms the pseudocyclic conformation through an intramolecular salt bridge network between the phosphate group and the dimethylamino group, oxazole nitrogen, and 13-OH (Fig. 1).² This conformation masks the charges of the phosphate group and decreases the polarity of this molecule, and thus enhances the membrane permeability. However, upon binding to the PP1 γ , the pseudocyclic conformation is drastically changed into the extended conformation, to expose the

phosphate group to the amino acid residues in the phosphate recognition site.²³ Thus, the sophisticated natural design for the preservation of the cytotoxicity is embedded in the entire skeleton of **1**.

3. Biosynthetic gene cluster of calyculin A

Structurally, **1** has an oxazole ring that connects two biosynthetically dissimilar motifs: a peptide portion composed of two γ -amino acids and a polyketide portion containing a 5,6-spiroacetal, phosphate, tetraene, and terminal nitrile. Thus, these structural features suggest a non-ribosomal peptide and polyketide hybrid biosynthetic pathway, with some noncanonical modifications. One of these modifications is the β -branched methyl groups on C3 and C7, implying the *trans*-AT PKS origin.³⁰ The PCR analysis with degenerate primers for KS was performed on the *D. calyx* metagenome, to obtain the *trans*-AT type ketosynthase DNA fragments. The screening of a fosmid library with the DNA fragment encoding the *trans*-AT KS sequences successfully yielded the PKS-NRPS hybrid gene cluster, ranging over 150 kb. The PKS-NRPS modular assembly line is composed of nine ORFs encoding thirty-four modules aligned until the TE domain (Fig. 4), whereas the downstream ORFs, *calH* and *calI*, encode some extra modules, whose putative products are missing in the calyculin structure. The five A domains encoded in the *cal* gene are supposed to accept Ser, Gly, Ser, Ala, and Ser, according to the NRPS codes, and among them the latter two, the A4 and A5 domains, are encoded in the extra module (Fig. 4).³¹ The first two Ser residues recruited by the A1 and A3 domains are converted into trimethylserine and oxazole, respectively, which agree with the domain organization of the corresponding module (Fig. 4). The NRPS code of the A2 domain matched that accepting Gly, except for the replacement of Leu or Met by Thr, which presumably confers the promiscuity to this A domain and thus allows it to accept Ala as an additional substrate, to generate **3**.

Notably, the *cal* gene encodes 29 KS domains in total, and among them, nine are non-elongating KS⁰ domains (Fig. 4, KS1, 3, 6, 7, 10, 14, 17, 19, and 28). The fact that nearly one-third of

the total KS domains have non-elongating functions provides significant insights into the domain architecture exerting non-canonical modifications in the *trans*-AT PKS pathway. The *N*- and *O*-methylation proceeds on the module containing KS1 and KS19, respectively. The domain organization with KS28 and the succeeding DH catalyzes the β,γ -shift of the double bond to generate an enamide.^{32,33} The roles of KS6 and 7 are unknown, but they flank the NRPS module responsible for the oxazole formation. Aside from KS14, all of the remaining KS⁰ (Fig. 4, KS3, 10, and 17) domain-containing modules (Fig. 4, modules 4, 13, and 20) are involved in the α -hydroxylation. Modules 4 and 20 both have KS⁰ domains, and ACP could implement the α -hydroxylation by an unidentified oxygenase. Indeed, two monooxygenases are encoded upstream of the *cal* gene. While module 4 introduces the 34-OH, module 20 generates the 16-OH required for the 5,6-spiroacetal ring formation. Once the 16-OH is introduced, the tetrahydrofuran ring can be formed with the carbonyl group located at C19. The resulting lactol can generate the oxonium ion, which is captured by the 23-OH to afford the 5,6-spiroacetal. Since the natural C-19 *S*-isomer is thermodynamically more stable than the C-19 *R*-isomer,³⁴ the 5,6-spiroacetal could be formed in a non-enzymatic manner.

Above all, module 13, along with the stand-alone oxygenase (CalD), functions in the important process of the construction of the polyketide portion with an odd numbered carbon chain from C1 to C25. The exact mechanism of the chain shortening modification is still unknown, but a mechanism has been proposed, based on the domain organization. The architecture of Ox-KS⁰-DH suggests that α -hydroxylation by the Ox, followed by dehydration by the DH, generates the α -ketothioester intermediate (Fig. 4),³⁵ which in turn is taken up again by the Ox to undergo a Baeyer-Villiger type oxidation. If the next elongation reaction occurs on the ester carbonyl group of the mixed anhydride, then a carbon unit is eliminated as carbon dioxide. The other KS⁰ is KS14, and its role is still unknown. Thus, the *cal* gene is an intricate showcase of the module architectures potentiating the non-canonical modifications in the *trans*-AT PKS pathway.

By contrast, there are two major discrepancies between the *cal* gene organization and the calyculin structure. One is the stereochemistry of C17, which was predicted by the KR sequence (module 19) to be antipodal to that found in **1**. The stereochemical inversion may occur during the phosphorylation of the 17-OH at the post-PKS modification stage. The other discrepancy is the existence of the extra module encoded by *calH* and *call*. The nitrogen atom of Ala introduced by the A domain (A4) is obviously required for nitrile formation at the terminus. The succeeding domain architectures of modules 31 and 32 indicate the enamide formation at the corresponding position, as mentioned above. Considering that some natural calyculin derivatives have a terminal amide in place of the nitrile (Figs. 1 and 2), we envisioned that calyculin biosynthesis is terminated at the enamide functionality, to generate a terminal amide as nascent product. A phosphate-involved reaction was proposed³⁶ as the mechanism for the transformation of the amide into the nitrile, since three ORFs within the upstream region are annotated as phosphotransferases.

4. Phosphocalyculin A as a protoxin

The three phosphotransferases encoded in the upstream region of the gene cluster were subjected to heterologous expression and functional analysis. Unexpectedly, one of them, CalQ, appended a phosphate group onto **1** as well as calyculinamide A (**5**). Previously, **1** had been considered to be the final product of the biosynthesis, since it was the major metabolite in the sponge extract prepared in the conventional manner. However, the alcoholic extract of the sponge tissue, which was freeze-dried after flash-freezing in liquid nitrogen, afforded phosphocalyculin A (**30**) as the major metabolite, in place of **1**. This result suggested that an as yet unidentified phosphatase coexists along with **30**, and they are compartmentalized in the sponge tissue. The conventional extraction procedure induces decompartmentalization, which immediately converts **30** into **1**. Thus, **30** had been overlooked due to enzymatic dephosphorylation during the previous extraction procedure. Indeed, the induction of

the “activating” process was observed immediately after wounding the sponge tissue.^{25,37} As mentioned in the structure-activity relationships study, since the phosphate group is essential for phosphatase inhibition, **30** is 1,000 times less toxic than **1**. These findings provide not only the functional evidence of the gene cluster, but also the underlying mechanism of calyculin biogenesis.

Wound-activated chemical defense systems are prevalent among terrestrial plants (Fig. 5). Cyanogen glycoside is a representative example, in which tissue wounding induces a quick increase in hydrogen cyanide via the hydrolytic liberation of the aglycone.³⁸ The precursor molecule and the activating enzyme are compartmentalized in the plant tissue, which is readily disrupted by wounding to accomplish chemical defense. The activated defense strategy has also been found in a few marine sponges (Fig. 5).³⁹ The quick enzymatic transformation of the protoxin to the toxin often prevents the prevalence of this system from being recognized, since the homogenization of sponge tissue during the extraction of secondary metabolites results in decompartmentalization, thus giving rise to the same reaction as in the case of wound activation. The first example was found in the marine sponge *Aplysina aerophoba*, which accumulates brominated isoxazoline alkaloids. The isoxazoline alkaloids (**23**) are stored in spherulous cells and are enzymatically converted to the more active toxin, dienone amide (**25**), in response to tissue disruption.^{40,41} This bioconversion is a multistep process composed of the liberation of the isoxazoline moiety with a terminal nitrile, followed by demethylation, dehydration, and hydration of the nitrile. The enzymes catalyzing these steps are membrane-bound or membrane-associated enzymes except for that catalyzing the final step, nitrile hydration.⁴² Proksch and coworkers recently reported the isolation of an aeroplysinin-1 (**24**)-specific nitrile hydratase, which lacks sequence homology to known nitrile hydratases and has narrow substrate specificity.⁴² Similar bioconversion events have been found in the marine sponges *Luffariella variabilis* and *Aplysinella rhax*, and are relatively simpler processes involving deacetylation and desulfonation, respectively (Fig. 5).^{43,44} However, the enzymes responsible for the bioconversions have not been identified yet. If the

biosynthetic genes corresponding to the syntheses of these terpenoids and alkaloids are identified, it would be interesting to determine whether the enzymes involved in the deactivating or activating processes are clustered.

5. Calyculin A production in the sponge-microbe association

Faulkner and co-workers originally reported that the filamentous bacterium associated with Palauan *Theonella swinhoei* contained theopalauamide, by cell separation methodology.⁴⁵ This bacterium was subsequently identified as “*Candidatus Entotheonella palauensis*”.⁴⁶ Although the chemical localization study successfully illuminated the cellular distributions of the sponge-derived bioactive molecules, the site of biosynthesis might not always be the same as the site of storage. To determine the chemical localization in line with the production site, biosynthetic evidence is desirable. A few successful examples encompassing biosynthetic gene clusters have been reported. Several polychlorinated peptides isolated from the marine sponge *Dysidea herbacea* are closely related to barbamide, from the free-standing cyanobacterium *Lyngbya majuscula*.⁴⁷ Gerwick’s group applied the known sequence of the barbamide biosynthetic gene cluster to search for the biosynthetic gene responsible for the production of the polychlorinated peptides, which yielded the *bar* homolog from the sponge. A CARD-FISH analysis with the probe specific to the *bar* homolog revealed that the biosynthetic genes are housed within the cyanobacterial symbiont *Oscillatoria spongelliae*.⁴⁸ In 2014, Piel and co-workers reported that Entotheonella belongs to the candidate phylum Tectomicrobia, and the bacterium of the phylotype “*Candidatus Entotheonella factor*” is the producer of almost all metabolites isolated from the Japanese marine sponge *T. swinhoei* (yellow chemotype).⁴⁹ Quite recently, the white chemotype *T. swinhoei* was also found to harbor the similar filamentous bacterium “*Candidatus Entotheonella sarta*”, as the producer of misakinolide. The overlapped locations of the

16S rDNA and misakinolide were elucidated by CARD-FISH and MALDI imaging mass spectrometry, respectively, confirming that the biosynthesis and storage sites were identical.⁵⁰

A CARD-FISH analysis was also conducted with the calyculin biosynthetic gene cluster, in the sponge *D. calyx*.⁹ The gene cluster was localized within the filamentous bacterium “*Candidatus Entotheonella*” sp., as confirmed by the single cell analysis using laser microdissection.⁹ The data provided the first clues about the biosynthetic gene and the symbiotic producer involved in the wound-activated process to generate a cytotoxic compound derived from a marine sponge. Considering that calyculins have almost no antibacterial activity, the self-toxicity to *Entotheonella* is considered to be low. However, the potent cytotoxicity of calyculins would be harmful to the host sponge. The manner in which **1** is phosphorylated and deactivated by the bacterial symbiont suggests that the crosstalk between the sponge host and the symbiont plays an important role in not only the dynamic process for the on-demand generation of toxic **1**, but also the self-resistance systems of the host.

6. Conclusions

The overall picture of the biosynthetic mechanisms of sponge-derived bioactive molecules is still largely unclear. However, recent comprehensive studies have illuminated some intriguing pieces of the complicated processes, such as chemical localizations, real producers, and dynamic bioconversions. The discovery of the association of *Entotheonella* with the genera *Theonella* and *Discodermia* has given rise to more questions about the ubiquity of secondary metabolite production by *Entotheonella* in other sponges, and whether other bacterial taxa function as symbiotic producers. Future studies will provide answers to these questions.


Wound-activated bioconversion processes have been reported for brominated isoxazoline alkaloids and calyculins, which are both predominant metabolites in the sponges *A. aerophoba* and *D. calyx*, respectively. The spatial organization and the activation process might allow the sponges to

store large amounts of highly potent toxins. Recent advances in mass spectrometry-based imaging technology will clarify not only the detailed localization but also the dynamic transformations occurring in response to tissue disruption.

As potential applications, the enzymatic activation and deactivation processes of bioactive molecules have historically been utilized as biochemical tools, including certain antibiotics as expression markers. Thus, understanding nature's way of producing bioactive metabolites in the sponge-microbe association will provide significant opportunities, not only for addressing supply problems but also for further applications of these metabolites in biochemical research. The sponge factory may have many more treasures than the medicinally important bioactive molecules themselves.

ACKNOWLEDGMENTS

Our research on calyculin biosynthesis was supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, Japan. Support was also provided by the Mitsubishi Foundation, the Nagase Science Technology Foundation, Astellas Foundation for Research on Metabolic Disorders, the Sumitomo Foundation, the Takeda Science Foundation, and the Naito Foundation.

Figure 1. Structures of calyculin derivatives and crystal structure of *ent*-calyculin A.




Figure 2. Structures of calyculin derivatives.




Figure 3. Structures of natural product inhibitors against PP1 and PP2A.




Figure 4. Proposed biosynthetic pathway of calyculin A.





Aplysina sp.*Luffariella variabilis**Aplysinella rhax**Discodermia calyx*

Figure 5. Enzymatic wound-activated bioconversion in marine sponges.

REFERENCES

- (1) L. Döderlein, Studien an japanischen Lithistiden. *Zeitschr. f. wiss. Zool.* **40**, 62-104 (1883).
- (2) Y. Kato, N. Fusetani, S. Matsuanga, K. Hashimoto, S. Fujita, and T. Furuya. Calyculin A, a novel antitumor metabolites from the marine sponge *Discodermia calyx*, *J. Am. Chem. Soc.* 1986, **108**, 2780-2781.
- (3) M. Suganuma, H. Fujiki, H. Furuya-Suguri, S. Yoshizawa, S. Yamamoto, Y. Kato, N. Fusetani, and T. Sugimura, Calyculin A, and inhibitor of protein phosphatases, a potent tumor promoter on CD-1 mouse skin, *Cancer Res.* 1990, **50**, 3521-3525.
- (4) A. E. Fagerholm, D. Habrant, and A. M. P. Koskinen, Calyculins and related marine natural products as serine-threonine protein phosphatase PP1 and PP2A inhibitors and total syntheses of calyculin A, B, and C, *Mar. Drugs* 2010, **8**, 122-172.
- (5) E. J. Dumdei, J. W. Blunt, M. H. G. Munro, and L. K. Pannell, Isolation of calyculins, calyculinamides, and swinholide H from the New Zealand deep-water marine sponge *Lamellomorpha strongylata* *J. Org. Chem.* 1997, **62**, 2636-2639.
- (6) X. Fu, F. J. Schmitz, M. Kelly-Borges, T. L. McCready, and C. F. B. Holmes, Clavosines A-C from the marine sponge *Myriastrra clavosa*: Potent cytotoxins and inhibitors of protein phosphatases 1 and 2A, *J. Org. Chem.* 1998, **63**, 7957-7963.
- (7) S. Kehraus, G. M. König, and A. D. Wright, A new cytotoxic calyculinamide derivative, geometricin A, from the Australian sponge *Luffariella geometrica*. *J. Nat. Prod.* 2002, **65**, 1056-1058.

- (8) R. A. Edrada, R. Ebel, A. Supriyono, V. Wray, P. Schupp, K. Steube, R. van Soest, and P. Proksch, Swinhoeiamide A, a new highly active calyculin derivative from the marine sponge *Theonella swinhoei*. *J. Nat. Prod.* 2002, **65**, 1168-1172.
- (9) T. Wakimoto, Y. Egami, Y. Nakashima, Y. Wakimoto, T. Mori, T. Awakawa, T. Ito, H. Kenmoku, Y. Asakawa, J. Piel, and I. Abe, Calyculin biogenesis from a pyrophosphate protoxin produced by a sponge symbiont, *Nat. Chem. Biol.* 2014, **10**, 648-655.
- (10) R. E. Honkanen, Cantharidin, another natural toxin that inhibits the activity of serine/threonine protein phosphatases types 1 and 2A, *FEBS Lett.* 1993, **330**, 283-286.
- (11) C. MacKintosh, K. A. Beattie, S. Klumpp, P. Cohen, G. A. Codd, Cyanobacterial microcystin-LR is a potent and specific inhibitor of protein phosphatases 1 and 2A from both mammals and higher plants, *FEBS Lett.* 1990, **264**, 187-192.
- (12) K. Tachibana, P. J. Scheuer, Y. Tsukitani, H. Kikuchi, D. V. Engen, J. Clardy, Y. Gopichand, and F. J. Schmitz, Okadaic acid, a cytotoxic polyether from two marine sponges of the genus *Halichondria*, *J. Am. Chem. Soc.* 1981, **103**, 2469-2471.
- (13) Y. Murakami, Y. Osjima, and T. Yasumoto, Identification of okadaic acid as atoxic component of a marine dinoflagellate *Prorocentrum lima*, *Nihon Suisan Gakkaishi*, 2006, **72**, 4907-4916.
- (14) S. Mitsuhashi, N. Matsuura, M. Ubukata, H. Oikawa, H. Shima, and K. Kikuchi, Tautomycetin is a novel and specific inhibitor of serine/ threonine protein phosphatase type 1, PP1, *Biochem. Biophys. Res. Commun.* 2001, **287**, 328-331.
- (15) M. Roberge, C. Tudan, S. M. F. Hung, K. W. Harder, F. R. Jirik, and H. Anderson, Antitumor drug fostriecin inhibits the mitotic entry checkpoint and protein phosphatases 1 and 2A, *Cancer Res.* 1994, **54**, 6115-6121.

- (16) T. Usui, G. Marriott, M. Inagaki, G. Swarup, and H. Osada, Protein phosphatase 2A inhibitors, phoslactomycins. Effects on the cytoskeleton in NIH/3T3 cells. *J. Biochem.* 1999, **125**, 960-965.
- (17) A. Takai, and G. Miekes, Inhibitory effect of okadaic acid on the *p*-nitrophenyl phosphate phosphatase activity of protein phosphatases, *Biochem. J.* 1991, **275**, 233-239.
- (18) J. R. Bagu, B. D. Sykes, M. M. Craig, and C. F. B. Holmes, A molecular basis for different interactions of marine toxins with protein phosphatase-1. *J. Biol. Chem.* 1997 **272**, 5087–5097.
- (19) V. Gupta, A. K. Ogawa, X. Du, K. N. Houk, and R. W. Armstrong, A model for binding of structurally diverse natural product inhibitors of protein phosphatases PP1 and PP2A. *J. Med. Chem.* 1997, **40**, 3199–3206.
- (20) C. -M. Gauss, J. E. Sheppeck II, A. C. Nairn, and R. Chamberlin, A molecular modeling analysis of the binding interactions between the okadaic acid class of natural product inhibitors and the Ser-Thr phosphatases, PP1 and PP2A, *Bioorg. Med. Chem.* 1997, **5**, 1751–1773.
- (21) J. Goldberg, H. B. Huang, Y. G. Kwon, P. Greengard, A. C. Nairn, and J. Kuriyan, Three-dimensional structure of the catalytic subunit of protein serine/threonine phosphatase-1, *Nature* 1995, **376**, 745-753.
- (22) J. T. Maynes, K. S. Bateman, M. M. Cherney, A. K. Das, H. A. Luu, C. F. B. Holmes, and M. N. G. James, Crystal structure of the tumor-promoter okadaic acid bound to protein phosphatase-1, *J. Biol. Chem.* 2001, **276**, 44078-44082.
- (23) A. Kita, S. Matsunaga, A. Takai, H. Kataiwa, T. Wakimoto, N. Fusetani, M. Isobe, and K. Miki, Crystal structure of the complex between calyculin A and the catalytic subunit of protein phosphatase 1, *Structure* 2002, **10**, 715-724.

- (24) M. S. Keller, R. Page, and W. Peti, Crystal structures of protein phosphatase-1 bound to nodularin-R and tautomycin: A novel scaffold for structure-based drug design of serine/threonine phosphatase inhibitors, *J. Mol. Biol.* 2009, **385**, 11-21.
- (25) T. Wakimoto, S. Matsunaga, A. Takai, and N. Fusetani, Insight into binding of calyculin A to protein phosphatase 1: isolation of hemicalyculin A and chemical transformation of calyculin A, *Chem. Biol.* 2002, **9**, 309-319.
- (26) Y. Kato, N. Fusetani, S. Matsunaga, K. Hashimoto, and K. Koseki, Isolation and structure elucidation of calyculins B, C, and D, novel antitumor metabolites, from the marine sponge *Discodermia calyx*, *J. Org. Chem.* 1998, **53**, 3930-3932.
- (27) S. Matsunaga, H. Fujiki, D. Sakata, and N. Fusetani, Calyculins E, F, G, and H, additional inhibitors of protein phosphatases 1 and 2A, from the marine sponge *Discodermia calyx*, *Tetrahedron* 1991, **47**, 2999-3006.
- (28) S. Matsunaga, T. Wakimoto, N. Fusetani, and M. Sugauma, Isolation of dephosphonocalyculin A from the marine sponge *Discodermia calyx*, *Tetrahedron Lett.* 1997, **38**, 3763-3764.
- (29) S. Matsunaga, T. Wakimoto, N. Fusetani, Isolation of four new calyculins from the marine sponge *Discodermia calyx*, *J. Org. Chem.* 1997, **62**, 2640-2642.
- (30) E. J. N. Helfrich, and J. Piel, Biosynthesis of polyketides by *trans*-AT polyketide synthases, *Nat. Prod. Rep.* 2015, in press.
- (31) T. Stachelhaus, H. D. Mootz, and M. A. Marahiel, The specificity-conferring code of adenylation domains in nonribosomal peptide synthetases. *Chem. Biol.* 1999, **6**, 493-505.

- (32) B. Kusebauch, B. Busch, K. Scherlach, M. Roth, and C. Hertweck, Functionally distinct modules operate two consecutive $\alpha,\beta \rightarrow \beta,\gamma$ double-bond shifts in the rhizoxin polyketide assembly line. *Angew. Chem. Int. Ed.* 2010, **49**, 1460-1464.
- (33) J. Moldenhauer, D. C. G. Götz, C. R. Albert, S. K. Bischof, K. Schneider, R. D. Süßmuth, M. Engeser, H. Gross, G. Bringmann, and J. Piel, The final steps of bacillaene biosynthesis in *Bacillus amyloliquefaciens* FZB42: direct evidence for β,γ dehydration by a *trans*-acyltransferase polyketide synthase. *Angew. Chem. Int. Ed.* 2010, **49**, 1465-1467.
- (34) A. B. Smith, III, G. K. Friestad, J. Barbosa, E. Bertounesque, K. G. Hull, M. Iwashima, Y. Qiu, B. A. Salvatore, P. G. Spoor, and J. J.-W. Duan, Total synthesis of (+)-calyculin A and (-)-calyculin B: asymmetric synthesis of the C(9-25) spiroketal dipropionate subunit, *J. Am. Chem. Soc.* 1999, **121**, 10468-10477.
- (35) M.-C. Tang, H.-Y. He, F. Zhang, and G.-L. Tang, Baeyer–Villiger oxidation of acyl carrier protein-tethered thioester to acyl carrier protein-linked thiocarbonate catalyzed by a monooxygenase domain in FR901464 biosynthesis. *ACS Catal.* 2013, **3**, 444-447.
- (36) P. B. Gomes, M. Nett, H.-M. Sattler, K. Martin, and C. Hertweck, Bezerramycins A-C, antiproliferative phenoxazinones from *Streptomyces griseus* featuring carboxy, carboxamide or nitrile substituents, *Eur. J. Org. Chem.* 2010, **2**, 231-235.
- (37) Y. Egami, T. Wakimoto, and I. Abe, Phosphocalyculin C as apyrophosphate protoxin of calyculin C in the marine sponge *Discodermia calyx*, *Bioorg. Med. Chem. Lett.* 2014, **24**, 5150-5153.
- (38) A. Mithöfer, and W. Boland, Plant defense against herbivores: chemical aspects, *Annu. Rev. Plant. Biol.* 2012, **63**, 431-450.

- (39) S. Rohde, S. Nietzer, and P. J. Schupp, Prevalence and mechanisms of dynamic chemical defenses in tropical sponges, *PLoS ONE* 2015, **10**, e0132236.
- (40) R. Ebel, M. Brenzinger, A. Kunze, H. J. Gross, and P. Proksch, Wound activation of protoxins in marine sponge *Aplysina aerophoba*, *J. Chem. Ecol.* 1997, **23**, 1451-1462.
- (41) C. Thoms, R. Ebel, and P. Proksch, Activated chemical defense in *Aplysina* sponge revisited, *J. Chem. Ecol.* 2006, **32**, 97-123.
- (42) B. Lipowicz, N. Hanekop, L. Schmitt, and P. Proksch, An aeroplysinin-1 nitrile hydratase isolated from the marine sponge *Aplysina cavernicola*, *Mar. Drugs* 2013, **11**, 3046-3067.
- (43) P. Ettinger-Epstein, C. A. Motti, R. de Nys, A. D. Wright, C. N. Battershill, and D. M. Tapiolas, Acetylated sesterterpenes from the great barrier reef sponge *Luffariella variabilis*, *J. Nat. Prod.* 2007, **70**, 648-651.
- (44) C. Thoms, and P. J. Schupp, Activated chemical defense in marine sponges – a case study on *Aplysinella rhax*, *J. Chem. Ecol.* 2008, **34**, 1242-1252.
- (45) C. A. Bewley, N. D. Holland, and D. J. Faulkner, Two classes of metabolites from *Theonella swinhoei* are localized in distinct populations of bacterial symbionts, *Experientia* 1992, **52**, 716-722.
- (46) E. W. Schmidt, A. Y. Obraztsova, S. K. Davidson, D. J. Faulkner, and M. G. Haygood, Identification of the antifungal peptide-containing symbiont of the marine sponge *Theonella swinhoei* as a novel δ -proteobacterium, “*Candidatus Entotheonella palauensis*”, *Mar. Biol.* 2000, **136**, 969-977.

- (47) J. Orjala, and W. H. Gerwick, Barbamide, a chlorinated metabolite with molluscicidal activity from the Caribbean cyanobacterium *Lyngbya majuscula*, *J. Nat. Prod.* 1996, **59**, 427-430.
- (48) P. M. Flatt, J. T. Gautschi, R. W. Thacker, M. Musafija-Girt, P. Crews, and W. H. Gerwick, Identification of the cellular site of polychlorinated peptide biosynthesis in the marine sponge *Dysidea (Lamellodysidea) herbacea* and symbiotic cyanobacterium *Oscillatoria spongelliae* by CARD-FISH analysis, *Mar. Biol.* 2005, **147**, 761-774.
- (49) M. C. Wilson, T. Mori, C. Rückert, A. R. Uria, M. J. Helf, K. Takada, C. Gernert, U. Steffens, N. Heycke, S. Schmitt, C. Rinke, E. J. N. Helfrich, A. O. Brachmann, C. Gurgui, T. Wakimoto, M. Kracht, M. Crüsemann, U. Hentschel, I. Abe, S. Matsunaga, J. Kalinowski, H. Takeyama, and J. Piel, An environmental bacterial taxon with a large and distinct metabolic repertoire, *Nature* 2014, **506**, 58-62.
- (50) R. Ueoka, A. R. Uria, S. Reiter, T. Mori, P. Karbaum, E. E. Peters, E. J. N. Helfrich, B. I. Morinaka, M. Gugger, H. Takeyama, S. Matsunaga, and J. Piel, Metabolic and evolutionary origins of actin-binding polyketides from diverse organisms, *Nat. Chem. Biol.* 2015, **11**, 705-712.