

Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Visible-light-induced cleavage of 4- α -amino acid substituted naphthalimides and its application in DNA photocleavage

Cite this: DOI: 10.1039/x0xx00000x

Jin Zhou \ddagger , Canliang Fang \ddagger , Ying Liu, Yao Zhao, Nan Zhang, Xiangjun Liu, Fuyi Wang, Dihua Shangguan*Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

A new kind of visible-light photocleavable molecules, 4- α -amino acid substituted naphthalimides was reported. The cleavage occurred at the C-N bond between 4-amino and amino acid residue and released a 4-amino-naphthalimide. A lysine substituted naphthalimide exhibited strong DNA photocleavage activity irradiated under a blue light LED.

Organic molecules capable of releasing active moieties after light triggering, are termed photocages or photocleavable (photolabile, photoreleasable, photoremovable or photoactivatable) molecules. These molecules are extensively exploring in biological applications, organic synthesis, and photolithographic techniques as tools for spatial and temporal control¹. However, most of photocleavable molecules require irradiation with short wavelength light (< 400 nm)^{1b}, which limit their applications in living biosystems. Therefore, large efforts have been made to develop photocleavable chromophores that absorb visible light or near-IR light², especially the photocleavable fluorophores that have the ability to perform both as “phototrigger” for active molecule release and “fluorophore” for active molecule visualization. So far, only a few fluorophores have been modified as photocleavable groups triggered by visible light, such as coumarin³, xanthene⁴, perylene⁵ and organic-inorganic hybrid ruthenium compounds⁶.

Naphthalimide derivatives have been used in a large variety of fields. As a class of DNA-intercalating agents, they have been extensively explored as antitumor agents⁷. Due to the electronic push-pull structure, 3- and 4-aminonaphthalimides show strong absorption and emission with large Stokes' shift in the visible region; and have been widely developed as fluorescent and colorimetric probes for detection of ions and bioactive molecules⁸, as well as for cellular imaging⁹. These probes are considered to have high photostability irradiated at their absorption maxima^{9a, 10}. To the best of our knowledge, no photocleavable naphthalimide has been reported. Here, we

report the photocleavable property of 4- α -amino acid substituted naphthalimides after triggered by visible light. A designed lysine substituted naphthalimide (DNlys) was demonstrated to have strong DNA photocleavage activity when irradiated with a blue light LED (465-470 nm).

In our research, we observed that a lysine substituted naphthalimide, **1** could be gradually cleaved under the room light. Because 4-aminonaphthalimides are very stable under the room light, we suspected that the photocleavage of **1** might relate to the α -carboxyl group of lysine. Therefore we synthesized seven 4- α -amino acid substituted naphthalimides (compounds 1-7) and eight 4-amino substituted naphthalimides (without α -carboxyl group) (compounds 8-15) (Fig.1). These 4-aminonaphthalimides were dissolved in ethanol, ethanol containing 1% trifluoroacetic acid (TFA) or ethanol containing 1% ammonia, then put in glass bottles and exposed in daylight for 2 h. The thin layer chromatography (TLC) assay showed that all the seven α -amino acid substituted naphthalimides were photocleaved (compared with lane a) in all the three solutions, however other 4-amino substituted naphthalimides were not changed except for compound **13** in 1% TFA (Fig. 1). It worth to note that **13** (contain γ -carboxyl group) was photocleaved in 1% TFA solution, but not in neutral and basic condition (1% ammonia). These set of results suggest that 4- α -amino acid substituted naphthalimides are highly photolabile.

To understand the mechanism of the photocleavage of 4- α -amino acid substituted naphthalimides, we compared the photocleaved fluorescent product of **1** and **2** after irradiated in ethanol solution by HPLC with detection at 430 nm (Fig.S1). The HPLC results showed that the photocleaved fluorescent product of both compounds were identical. The ¹H NMR and ¹³C-NMR spectra of the photocleaved fluorescent product (Fig.S2) only showed ¹H signals higher than 6.5 ppm and ¹³C signals higher than 105 ppm, suggesting no saturated alkyl was

linked on the product. The EI-MS showed the molecular ion peak at m/z 212, which corresponded to compound **16** (Fig.1).




Fig.1 Photocleavage of 4-amino substituted naphthalimides. Top: structures of 4-amino substituted naphthalimides; 1-7: substituted by α -amino acids; 8-15: substituted by other amino compounds. Bottom: TLC analysis of 4-amino substituted naphthalimides (1-15) after exposed in daylight for 2 h; lane a: without irradiation; lane b: irradiated in 1% TFA; Lane c: irradiated in ethanol; lane d: irradiated in 1% ammonia.

The other part of the photocleaved products of 4- α -amino acid substituted naphthalimides was investigated with compound **6**, because the nitrobenzoyl moiety of **6** can be monitored by UV detector (270 nm). HPLC analysis showed that a peak (15 min) of photocleaved product only with the UV absorption at 270 nm (without absorption at 430 nm) was observed (Fig.S3). However, this product was unstable; and changed to many other compounds during purification (Fig.S4 and S5). In the ESI-MS analysis of the reaction mixture of photocleaved **6**, we found one group of molecules with molecular weights of 296, 280, 264, 250, 232; and the MS/MS analysis of these molecules (296, 280, 264 and 250) showed fragment ions of 233 and 150, which suggests a series of molecules containing nitrobenzoyl moiety (Fig. S6).

Based on the structure of 4- α -amino acid substituted naphthalimides, the formation of a five-membered-ring intramolecular hydrogen-bond between the α -carboxyl group and the amino group may be critical for the photocleavage of these 4- α -amino acid substituted naphthalimides, which can explain why other 4-amino-1,8-naphthalimides (without α -carboxyl group) can not be photocleaved (Fig. 2). 4-amino-1,8-naphthalimides are known to be the intramolecular charge

transfer (ICT) fluorophores with “push–pull” substituent pairs (electron donor/accepter)⁸. The photochemical excitation leads to the ICT from 4-amino group to the excited imide, resulting in the cleavage of the C-N bond of the α -amino acid and the intramolecular hydrogen transfer from α -carboxyl group to amino group to produce compound **16** and a radical of amino acid residue. Subsequently the radical may react with surrounding molecules and/or further decompose to form various products.




Fig. 2 The photocleavage of 4- α -amino acid substituted naphthalimides.

Naphthalimides represent an important class of DNA binders that have been extensively explored as antitumor agents, and a few of them have been shown to exhibit the activity of photoinduced DNA cleavage under UV light irradiation^{11,12}. It is well known that UV-light can cause DNA damage. Above results have shown the photocleavable property of 4- α -amino acid substituted naphthalimides, this property may endow them with the capacity of DNA photocleavage. To prove this hypothesis, we synthesized a lysine substituted naphthalimide, DNLys (N-dimethylaminopropyl-4-(1-carboxyl-5-amino-amyamino)-1,8-naphthalimide) and its photocleaved product, DNNH (N-dimethylaminopropyl-4-amino-1,8-naphthalimide) (Fig.3, and Fig.S7). The introduction of dimethylaminopropyl group and lysine is to increase the affinity of naphthalimide to DNA, because the amino groups bear positive charges under physiological conditions^{9d} and can enhance the DNA cleavage efficiency of light-activated DNA cleaver¹³. The absorption and emission spectra of DNLys, DNNH showed that the maximum absorption were in the range of 435-445 nm (Fig.S8), and the maximum emission were in the range of 545-550 nm (Fig.S9). The addition of DNA slightly decreased their absorption and slightly increased their emission.

The photocleavage experiment of DNLys was performed under a LED array (3W; 465-470 nm). 82% of DNLys was observed to be cleaved to DNNH after exposed to blue light in methanol in 1 h (Fig.S10). However, the photocleavage of DNLys in phosphate buffered saline (PBS, pH 7.4) was much slower than that in methanol (Fig.3), only 23% of DNLys was cleaved in 4 h. It is interesting that the addition of DNA could accelerate the photocleavage of DNLys in PBS, and 19%, 32% and 63% of it were cleaved in 1, 2 and 4 h (Fig.3). These results suggest that the interaction of DNA and DNLys could promote the photocleavage of DNLys and implied the feasibility for DNA cleavage. These results further confirm that 4- α -amino

acid substituted naphthalimides can be photocleaved by visible light and product a stable fluorescent 4-amino naphthalimides.




Fig.3. The structure of DNLys/DNNH and HPLC analysis of photocleaved DNLys (200 μM). (a) photocleaved in pH 7.4 PBS; (b) photocleaved in pH 7.4 PBS in the presence of CT-DNA (100 μM in base pairs). Samples were irradiated under 465-470 nm LED light for 1, 2, and 4 h at room temperature, and then applied for RP-HPLC assay, Mobile phase: methanol/ H_2O (0.1% TFA), 4:6; detected at 245 nm.

The DNA photocleavage experiments were performed using a closed supercoiled pBR322 DNA under irradiation with a blue LED at room temperature. The cleaved DNA was analyzed on a 1% agarose gel. After cleavage, supercoiled pBR322 DNA (form I) would convert to relaxed circular DNA (form II). As shown in Fig.4a, DNLys did not cleave DNA without irradiation. After irradiation for 0.5 h, DNA cleavage was observed in the presence of different concentrations of DNLys, the cleavage efficiency increased with the increase of DNLys concentration. 2-5 μM DNLys could cause most of DNA cleaved under irradiation. As expected, 30 μM DNNH (photocleavage product of DNLys) did not cause notable DNA cleavage under the same condition. The kinetics of DNA photocleavage showed that the intact DNA greatly decreased with the irradiation time prolonging, and most of plasmid DNA was consumed within 30 min of irradiation (Fig. 4b). The control experiments showed that very little of DNA damage was observed in the absence of DNLys even after irradiation for 60 min. Since DNNH also undergoes the photo-induced ICT process, these results suggest that the DNA cleavage was related to the photocleavage of DNLys, not to the ICT excited state of 4-aminonaphthalimides.

The photoinduced DNA damage by synthetic DNA cleavage agents usually involves the reactive oxygen species, such as hydroxyl radical, singlet oxygen and superoxide anion¹⁴. In order to whether the reactive oxygen species are responsible for DNA cleavage by light-activated DNLys, standard scavengers and inhibitors of reactive oxygen species were added respectively to the reaction system, i.e. singlet oxygen scavenger (sodium azide (NaN_3) and histidine (His)), superoxide radical scavenger (dithiothreitol (DTT)), and hydroxyl radical scavenger (salicylic acid (SA) and ethanol (EtOH)). As shown in Fig.4c, certain inhibition effect of all the test scavengers was observed. Among them, the singlet oxygen scavenger, NaN_3 and histidine showed the strongest and the

second strongest inhibition effect on the DNA cleavage. DNA photocleavage is usually caused by reactive oxygen species or the direct reaction with the light-activated DNA cleaver^{13b-d}. This set of results suggests that the DNA photocleavage in the presence of DNLys may involve the reactive oxygen species, especially the singlet oxygen¹⁵.




Fig.4 Cleavage of closed supercoiled pBR 322 DNA (0.5 $\mu\text{g}/20 \mu\text{L}$) irradiated under blue LED light at room temperature. a (left) agarose gel assay of DNA cleaved by different concentration (μM) of DNLys or DNNH after 0.5 h irradiation; (right) quantified plots of DNA as a function of DNLys concentration. b. (left) gel assay of DNA cleaved by DNLys (5 μM) after irradiation for different time (min); (right) quantified plots of DNA as a function of irradiation time. c gel (left) and quantified (right) assay of effects of additives on DNA cleavage by DNLys (5 μM) after 0.5 h irradiation, Ctr1: DNA only without irradiation; Ctr1 2: DNA only; Ctr1 3: DNA+DNLys; additives: DTT (50 mM); NaN_3 (100 mM); SA (salicylic acid, 20 mM); His (histidine, 5 mM); EtOH (ethanol, 2 M).

The preceding results have shown that the DNA cleavage was only caused by the photocleavage of DNLys, not caused by irradiated DNNH, suggesting that the singlet oxygen was not generated from photosensitized 4-aminonaphthalimide. The photocleavage of DNLys produced a radical of amino acid residues, thus the singlet oxygen may be generated by the reaction of dissolved oxygen with the excited radical states. Even so, we could not exclude the direct mechanism of DNA damage by the excited biradical states because of the high activity of the biradical. Further research will focus on the mechanistic details of photocleavage of 4- α -amino acid substituted naphthalimides, as well as of the DNA cleavage by irradiated DNLys.

Above results showed that 4- α -amino acid substituted naphthalimides could be photoactivated with blue light and released a fluorescent product, 4-amino naphthalimides. The 4- α -amino acid substituted naphthalimides can be easily synthesized by substitution reaction of 4-bromine-1,8-naphthalimides with different α -amino acids. Furthermore, through the reaction with the amino acid residues, the 4- α -amino acid substituted naphthalimides (e.g. lysine and glutamic acid substituted) could be further linked to peptides, proteins and other biologically interesting molecules. Additionally, the

substituent at the 9-position N-atom of naphthalimides (imide N-atom) can be easily changed to different molecules by reaction of 4-bromine-1, 8-naphthalanhydride with corresponding compounds containing NH₂ group. Therefore the 4- α -amino acid substituted naphthalimide fluorophore could act as a multifunctional platform for the construction of light-control systems by linking different functional molecules. The visible light activation makes the constructed systems hold the potential in biological applications, e.g. spatial and temporal control of drug delivery. The fluorescence emission of 4- α -amino acid substituted naphthalimides and their photocleaved products makes the constructed systems work on visualization. The photoactivation of DNLYs showed the photocleavage ability to DNA, suggesting the potential application in phototherapy after further modifying the 4- α -amino acid substituted naphthalimides to enhance the DNA binding ability.

Conclusions

In summary, we describe the photocleavable property of 4- α -amino acid substituted naphthalimides upon blue light triggering. The photocleavage of these molecules was occurred at the C-N bond between 4-amino and amino acid residue and released a fluorescent product, 4-amino-naphthalimide. DNA was found enhance the photocleavage of a lysine-substituted naphthalimides (DNLYs), which caused the cleavage of DNA. Because of the ease of the modification on the 4-position and 9-position of naphthalimide ring, this finding provided a multifunctional platform for the construction of light-control systems.

We gratefully acknowledge the financial support from Grant 973 Program (2011CB935800, 2011CB911000 and 2013CB933700) and NSF of China (21275149, 21375135, 21205124 and 21321003).

Notes and references

Beijing National Laboratory for Molecular Sciences, Key Laboratory of Analytical Chemistry for Living Biosystems, Institute of Chemistry, Chinese Academy of Sciences, Beijing, 100190, china
Tel/Fax:86-10-62528509; E-mail: sgdh@iccas.ac.cn.

‡ Primary authors contributed equally to this work.

Electronic Supplementary Information (ESI) available: [experimental information and figures]. See DOI:10.1039/c000000x/

- 1 (a) G. Mayer and A. Heckel, *Angew. Chem.-Int. Edit.*, 2006, **45**, 4900; (b) P. Klan, T. Solomek, C. G. Bochet, A. Blanc, R. Givens, M. Rubina, V. Popik, A. Kostikov and J. Wirz, *Chem. Rev.*, 2013, **113**, 119; (c) Q. Y. Liu and A. Deiters, *Accounts Chem. Res.*, 2014, **47**, 45.
- 2 A. P. Gorka, R. R. Nani, J. J. Zhu, S. Mackem and M. J. Schnermann, *J. Am. Chem. Soc.*, 2014, **136**, 14153.
- 3 (a) L. Fournier, I. Aujard, T. Le Saux, S. Maurin, S. Beaupierre, J. B. Baudin and L. Jullien, *Chemistry-a European Journal*, 2013, **19**, 17494; (b) J. P. Olson, H. B. Kwon, K. T. Takasaki, C. Y. Q. Chiu, M. J. Higley, B. L. Sabatini and G. C. R. Ellis-Davies, *J. Am. Chem. Soc.*, 2013, **135**, 5954.
- 4 P. Sebej, J. Wintner, P. Muller, T. Slanina, J. Al Anshori, L. A. P. Antony, P. Klan and J. Wirz, *Journal of Organic Chemistry*, 2013, **78**, 1833.
- 5 A. Jana, M. Iqbal and N. D. P. Singh, *Tetrahedron*, 2012, **68**, 1128.
- 6 (a) L. Zayat, C. Calero, P. Albores, L. Baraldo and R. Etchenique, *J. Am. Chem. Soc.*, 2003, **125**, 882; (b) M. Salierno, C. Fameli and R. Etchenique, *Eur. J. Inorg. Chem.*, 2008, 1125.
- 7 (a) M. Lv and H. Xu, *Curr. Med. Chem.*, 2009, **16**, 4797; (b) S. Banerjee, E. B. Veale, C. M. Phelan, S. A. Murphy, G. M. Tocci, L. J. Gillespie, D. O. Frimannsson, J. M. Kelly and T. Gunnlaugsson, *Chem. Soc. Rev.*, 2013, **42**, 1601.
- 8 (a) X. H. Qian, Y. Xiao, Y. F. Xu, X. F. Guo, J. H. Qian and W. P. Zhu, *Chem. Commun.*, 2010, **46**, 6418; (b) C. L. Fang, J. Zhou, X. J. Liu, Z. H. Cao and D. H. Shangguan, *Dalton Trans.*, 2011, **40**, 899; (c) J. Zhou, H. Y. Liu, B. Jin, X. J. Liu, H. B. Fu and D. H. Shangguan, *J. Mater Chem C*, 2013, **1**, 4427; (d) B. C. Zhu, X. L. Zhang, Y. M. Li, P. F. Wang, H. Y. Zhang and X. Q. Zhuang, *Chem. Commun.*, 2010, **46**, 5710; (e) B. C. Zhu, X. L. Zhang, H. Y. Jia, Y. M. Li, H. P. Liu and W. H. Tan, *Org. Biomol. Chem.*, 2010, **8**, 1650.
- 9 (a) J. Zhou, C. L. Fang, T. J. Chang, X. J. Liu and D. Shangguan, *J. Mater Chem B*, 2013, **1**, 661; (b) D. Srikun, E. W. Miller, D. W. Dornaille and C. J. Chang, *J. Am. Chem. Soc.*, 2008, **130**, 4596; (c) M. H. Lee, J. H. Han, P. S. Kwon, S. Bhuniya, J. Y. Kim, J. L. Sessler, C. Kang and J. S. Kim, *J. Am. Chem. Soc.*, 2012, **134**, 1316; (d) J. Zhou, A. Chang, L. Wang, Y. Liu, X. Liu and D. Shangguan, *Org. Biomol. Chem.*, 2014, **12**, 9207.
- 10 P. A. Panchenko, O. A. Fedorova and Y. V. Fedorov, *Russ. Chem. Rev.*, 2014, **83**, 155.
- 11 (a) Z. G. Li, Q. Yang and X. H. Qian, *Bioorg. Med. Chem. Lett.*, 2005, **15**, 3143; (b) *Bioorg. Med. Chem.*, 2005, **13**, 4864; (c) *Bioorg. Med. Chem.*, 2005, **13**, 3149; (d) I. Saito, M. Takayama and S. Kawanishi, *J. Am. Chem. Soc.*, 1995, **117**, 5590.
- 12 B. M. Aveline, S. Matsugo and R. W. Redmond, *J. Am. Chem. Soc.*, 1997, **119**, 11785.
- 13 (a) I. Hatial, P. S. Addy, A. K. Ghosh and A. Basak, *Tetrahedron Lett.*, 2013, **54**, 854; (b) B. Breiner, K. Kaya, S. Roy, W.-Y. Yang and I. V. Alabugin, *Org. Biomol. Chem.*, 2012, **10**, 3974; (c) W. Y. Yang, S. Roy, B. Phrathep, Z. Rengert, R. Kenworthy, D. A. R. Zorio and I. V. Alabugin, *J. Med. Chem.*, 2011, **54**, 8501; (d) W. Y. Yang, B. Breiner, S. V. Kovalenko, C. Ben, M. Singh, S. N. LeGrand, Q. X. A. Sang, G. F. Strouse, J. A. Copland and I. V. Alabugin, *J. Am. Chem. Soc.*, 2009, **131**, 11458; (e) S. V. Kovalenko and I. V. Alabugin, *Chem. Commun.*, 2005, 1444.
- 14 (a) B. Armitage, *Chem. Rev.*, 1998, **98**, 1171; (b) C. Bohne, K. Faulhaber, B. Giese, A. Hafner, A. Hofmann, H. Ihmels, A. K. Kohler, S. Pera, F. Schneider and M. A. L. Sheepwash, *J. Am. Chem. Soc.*, 2005, **127**, 76; (c) Y. F. Xu, X. Y. Huang, X. H. Qian and W. Yao, *Bioorg. Med. Chem.*, 2004, **12**, 2335; (d) N. Chowdhury, M. Gangopadhyay, S. Karthik, N. D. P. Singh, M. Baidya and S. K. Ghosh, *J. Photochem. Photobiol. B-Biol.*, 2014, **130**, 188; (e) W. Yao, X. H. Qian and Q. Y. Hu, *Tetrahedron Lett.*, 2000, **41**, 7711; (f) P. Yang, Q. Yang, X. H. Qian, L. P. Tong and X. L. Li, *J. Photochem. Photobiol. B-Biol.*, 2006, **84**, 221.
- 15 F. Xue, C. Z. Xie, Y. W. Zhang, Z. Qiao, X. Qiao, J. Y. Xu and S. P. Yan, *J. Inorg. Biochem.*, 2012, **115**, 78.