

Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Investigation of glycofullerene dynamics by NMR spectroscopy

*Olof Engström*¹, *Antonio Muñoz*², *Beatriz M. Illescas*², *Nazario Martín*², *Renato Ribeiro-Viana*³,
*Javier Rojo*³, and *Göran Widmalm*^{1*}

¹Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University, S-106 91 Stockholm, Sweden.

²Departamento de Química Orgánica, Facultad de Química, Universidad Complutense, E-28040 Madrid, Spain.

³Glycosystems Laboratory, Instituto de Investigaciones Químicas (IIQ), CSIC – Universidad de Sevilla, Av. Américo Vespucio 49, E-41092 Seville, Spain.

*Corresponding author. Email: goran.widmalm@su.se

Keywords: Glycofullerene, Carbohydrates, NMR relaxation, PFG Diffusion, Model-free, Flory exponent, Rebinding

ABSTRACT

Glycofullerenes, in which carbohydrate molecules are attached via a linker to a [60]fullerene core, facilitate spherical presentation of glyco-based epitopes. We herein investigate the dynamics of two glycofullerenes, having 12 and 36 mannose residues at their periphery, by NMR translational diffusion and quantitative ¹³C relaxation studies employing a model-free approach for their interpretation. The sugar residues are shown to be highly flexible entities with $S^2 < 0.2$ in both compounds. Notably, the larger glycofullerene with longer linkers show faster internal dynamics and higher flexibility than its smaller counterpart. The dynamics and flexibility as well as the slower translational diffusion of the larger glycofullerene, thereby favoring rebinding to a receptor, may together with its spatial extension explain why it is better than the smaller one at blocking the DC-SIGN receptor and inhibit the infection by pseudotyped Ebola virus particles.

INTRODUCTION

Carbohydrate-protein interactions are fundamental events in many physiological and pathological processes such as cell development and differentiation, embryogenesis, inflammation, pathogen infection, tumor progression and metastasis.^{1,2} These interactions are weak, but in nature this fact is compensated by a multivalent interaction between several copies of the carbohydrate ligands and the corresponding receptors (lectins) leading to strong and highly selective binding.^{3,4} Therefore, the development of carbohydrate multivalent tools is required to understand and explore these processes. Although there are some illustrative examples where a careful design to position ligands at a

precise distance on a specific scaffold have been achieved,^{5,6} these are scarce and most of the examples described in the literature try to explore different scaffolds in trial-and-error testing to look for the best presentation of carbohydrate ligands in order to facilitate efficient multivalent interactions with the corresponding lectins. Among these scaffolds, the most widespread ones are dendrons, dendrimers, polymers, cyclodextrins, calixarens, gold nanoparticles and liposomes.^{7,8} In this context, the possibility of using a fullerene molecule as a suitable spherical platform for a globular multivalent presentation of carbohydrates has been explored, namely, a [60]fullerene was used to achieve a globular presentation of carbohydrate ligands based on a Bingel hexakis-adduct functionalization strategy.^{9,10} Each malonate moiety is then endowed with two or six carbohydrate units, respectively, affording fullerenes bearing 12 or 36 carbohydrate entities resulting in a globular shape. Preliminary work using a cellular assay has shown that glycofullerenes of this type (compounds **1** and **2** in Figure 1, Figure S1) are able to block the DC-SIGN receptor and inhibit the infection by pseudotyped Ebola virus particles.¹¹ Somewhat surprisingly, the concentration for inhibition of infection differs by one order of magnitude between compounds **1** and **2** (IC₅₀ of **2** and 0.3 μM, respectively). These results highlight that, besides the importance of the number of ligands (valency), the accessibility of carbohydrate entities at the fullerene surface is of primary importance in order to interact in a suitable way with the corresponding receptors. In fact, to the best of our knowledge there has not been any study describing the disposition of ligands on the fullerene surface related to interactions with a receptor. With the aim to understand the different behavior and the influence of the flexibility of the carbohydrate groups at the surface of the fullerene, we have carried out different NMR experiments on two glycofullerenes. In particular, we have performed NMR translational diffusion measurements and quantitative ¹³C NMR relaxation studies. These studies provide experimental information related to structural aspects that can help in the design of multivalent systems relying on rigid scaffolds based on carbon structures and attached carbohydrate entities that are flexible.

RESULTS AND DISCUSSION

The translational diffusion of glycofullerenes **1** and **2**, which have M_w of 5.1 and 16.5 kDa respectively, was investigated by ¹H NMR pulsed-field-gradient (PFG) diffusion experiments performed on samples in D₂O under assumed dilute conditions (0.25 mg·ml⁻¹)¹² and at a higher concentration (25 mg·ml⁻¹). The fitted diffusion coefficients (D_0 and D_t , respectively) are presented in Table 1. D_0 and D_t were used to calculate radii of hydration ($R_{H,0}$ and $R_{H,t}$) employing the Stokes-Einstein relationship, results of which are also presented in Table 1. The change in concentration hardly affects **2**, whereas an effective increase for R_H by a factor of two takes place for **1** when the concentration is increased by two orders of magnitude, possibly caused by some aggregation at the higher concentration.

The change in translational diffusion upon generation growth in a dendrimer can yield insight into packing and solvation of the molecule. In the case of an ideal fractal dendrimer, D_0 has a power dependence of the molecular weight ($D_0 \propto M_w^{-\nu}$), with ν being the Flory exponent. Under poor solvent conditions, the growth of a dendrimer will be compact and homogenous thus yielding a Flory exponent of $\nu = 0.33$, which has been shown by computer simulations and scattering experiments.¹³ Under good solvent conditions, the growth of a dendrimer can be described as for a linear polymer ($\nu = 0.60$) if extended linkers at low generations are assumed, but at higher generations the

crowding of linkers will cause deviations.¹⁴ Analysis of D_0 as a function of M_w revealed the exponent ν to be 0.65, (Figure 2) i.e., in agreement with values of $\nu = 0.60$ and 0.67 that have been calculated for dendrimers in a good solvent utilizing Monte Carlo simulations.^{13,15} However, these results can be contrasted with our previous study of glycodendrimers where $\nu = 0.34$ was obtained.¹² The difference may be rationalized by the long linker present in glycofullerene **2**, thus allowing for semidilute condition and good solvation.¹⁵

¹³C NMR relaxation measurements were performed on the concentrated samples, for sensitivity reasons, at three different magnetic field strengths (11.75, 14.09 and 16.44 T corresponding to ¹H frequencies at 500, 600 and 700 MHz) to enable investigation of dynamic properties of glycofullerenes. The relaxation parameters R_1 , R_2 and (hetero)nuclear Overhauser effect (*NOE*) were measured for the endocyclic carbons of the mannosyl group, in consecutive order, inversion recovery, CPMG spin-echo and dynamic NOE experiments, and the results are presented in Figure 3 and Table S1. The slightly smaller *NOE* and the considerably higher R_2 relaxation rate of **1** in relation to **2** indicate that the former glycofullerene has a dynamic behavior that is more in the slow motion regime than the latter, in contrast to the results from the translation diffusion measurements. This difference may be caused by additional internal motions that influence the spectral density of the spins.

A model-free (MF) analysis was carried out to facilitate interpretation of the ¹³C NMR relaxation data. The computer program *Relax* was utilized,^{16,17} in which different MF approaches describing the spectral density function and different diffusion tensors are fitted to relaxation data. Solely isotropic diffusion tensors were considered due to the spherical geometry and symmetry of the glycofullerenes and τ_c were fixed at values calculated from D_t . Fitting of the NMR relaxation data using an extended MF equation¹⁸ resulted in lowest χ^2_{red} for the two glycofullerenes. In this approach the spectral density is described by three motions, namely, the global rotational diffusion and two local motions, a fast and a slow one:

$$J(\omega) = \frac{S^2\tau_c}{1+(\omega\tau_c)^2} + \frac{(1-S_f^2)\tau_f}{1+(\omega\tau_f)^2} + \frac{(S_f^2-S^2)\tau_s}{1+(\omega\tau_s)^2} \quad (1)$$

The global, the slow and the fast motions are described by correlation times, τ_c , τ_s and τ_f , respectively. The generalized order parameter (S^2) describes the amplitude of the local motions (where a value of unity corresponds to absence of local motion) and it can be bisected into S_f^2 and S_s^2 for the fast and the slow motions, respectively. S^2 can be calculated as the product of the two other order parameters, if one assumes that the fast internal motion is axially symmetric and independent of the slow motions. Equation 1 can be further simplified by assuming that τ_f is too short to have a substantial contribution to the spectral density:

$$J(\omega) = \frac{S^2\tau_c}{1+(\omega\tau_c)^2} + \frac{(S_f^2-S^2)\tau_s}{1+(\omega\tau_s)^2} \quad (2)$$

Correlation times and order parameters obtained from the extended MF fitting are presented in Table 2.

It has been suggested in the literature that in the case of dendrimers, the segmental reorientation that causes NMR relaxation is governed by three processes namely, (i) the overall global rotational diffusion, (ii) pulsating motions in the linker which become more prominent at the periphery and (iii) local reorientation of the segments.¹⁹ The

extended MF approach seems appropriate since the model includes three motions as expected of a dendrimer-like compound. Values of $S^2 < 0.2$ for the mannosyl groups in compounds **1** and **2** in this study indicate that the global rotational diffusion of the glycofullerene *per se* has a small contribution to the spectral density, which instead is governed by motions related to τ_s , possibly associated to the pulsating motion of the linker. The fitted τ_s are of the same magnitude as the principal correlation times as found for the glycodendrimers (~ 1 ns).¹² Interestingly, τ_s is shorter in **2** than in **1** (0.25 and 0.52 ns, respectively), likely a result of the extended linker and this difference may be the underlying reason for the observation of a spectral density more in the fast motion regime for the former compound in relation to the latter.

It is important to include parameters of different field strengths when interpreting NMR relaxation data, as has been highlighted in the literature.^{20,21} The exclusion of R_2 data, which had been acquired at 11.75 T, in the *Relax* analysis resulted in proposed models²² including R_{ex} parameters of large amplitude, describing motions on slower timescales. However, these results can appear from a forced fitting of parameters to a model, especially since neither R_1 nor *NOE* are dependent on R_{ex} . The inclusion of R_2 data acquired at two different fields confirmed that R_{ex} was an artifact of the fit.

The two mannofullerenes are both able to inhibit infection by Ebola virus in a model system by blocking binding to the DC-SIGN receptor. Their efficiency as measured by IC_{50} values differed by one order of magnitude even though the flexibility of both systems is very high with respect to the spatial presentation of the mannosyl groups, that on their own have well-defined 4C_1 chair conformations on the timescales investigated herein. The binding to the tetrameric DC-SIGN receptor takes place in a Ca^{2+} -dependent way through complexation with O3 and O4 of the mannose residue.^{23,24} The distance between two receptor binding sites is on the order of 40 Å.²⁵ Thus, in order for the multivalent construct to not only bind by statistical reassociation, where the high density of ligands facilitates consecutive rebinding, the geometrical restraint has to be fulfilled in order to facilitate chelate binding at the tetrameric receptor. Furthermore, additional geometric restrains have to be invoked if binding resulting in receptor clustering should occur.

Whereas both mannofullerenes are multivalent, each linker is monovalent in compound **1** but trivalent (branched) in **2**, which will result in a higher local density in the latter once binding has occurred and the process of statistical rebinding is in progress. The enthalpic term for the binding of a mannose residue in the two glycofullerenes to a receptor site should not differ and therefore the differences in the inhibition capability will be associated with other aspects. Besides the geometrical restraints essential for bridging different receptor sites the entropic costs for restraining the spacers upon binding will be different, being higher restricting the longer linker due to the larger degrees of freedom. The surface density of the ligands on the fullerenes is of importance as well as the kinetics of the rebinding process. Furthermore, once association of a ligand entity to the receptor has taken place both undesirable steric effects can take place due to incompatible ligand spacing but also beneficial steric interactions (steric stabilization) to competing virus-associated N-glycans carrying high-mannose structures may occur. The entropic loss due to long linkers can be circumvented by using a rigid core of defined length as recently reported by Ordanini

et al.,⁶ where the length of the central rod was investigated and optimal affinity toward DC-SIGN was obtained by a combination of a rod and a short flexible linker.

The approximate spatial extensions of the two mannofullerenes **1** and **2** were investigated by construction of molecular mechanics models. These revealed that, to a first approximation, the molecular size is comparable to that determined by translational diffusion measurements, namely, $R_{H,0} = 12.5 \text{ \AA}$ and $R_{H,0} = 26.7 \text{ \AA}$ for **1** and **2**, respectively. Thus, whereas compound **1** is able to reassociate, chelate binding to a second receptor site is severely limited, if possible at all, with a diameter of $< 30 \text{ \AA}$. Compound **2**, on the other hand, spanning a distance $> 50 \text{ \AA}$ across, will be able to engage in chelate binding to another site of the tetrameric DC-SIGN receptor as well as being involved in potential receptor clustering. Although these glycofullerenes do have a rigid core of a well-defined diameter they have not been optimized as described above for the rod-containing compounds. Instead the present study reveals the very high flexibility of both systems. The larger slower moving particle **2** with even more flexible linkers resulting in a very low S^2 value for the mannosyl residues arranged in a branched manner thereby facilitating rebinding²⁶ together with its spatial extension may be the reasons for its higher efficiency when binding to the DC-SIGN receptor inhibiting the Ebola virus.

CONCLUSIONS

The extended linker facilitates good solvation of segments as such and the glycofullerene as a whole, as indicated by the Flory exponent of $\nu = 0.65$. Also, extending the linker promotes the pulsating motion of the segment, reflected in the low S_s^2 and short τ_s parameters of **2**, consistent with relatively fast pulsating motions of large magnitude. These molecular properties together with the spatial extension of the molecule can shed light on why **2** is an efficient multivalent inhibitor. The enhanced effect seen in inhibition experiment of DC-SIGN may be explained by the kinetic concept of rebinding.²⁶ Accessible binding motifs attached to flexible linkers anchored on scaffolds of slow globular diffusion (as the case for glycofullerenes) will increase the binding due to a locally increased concentration of ligands. Furthermore, the extra branches in **2**, compared to **1**, increase the probability of interaction.²⁷ The rebinding mechanism is predicted to rely on low binding energy barriers of individual binding motifs (as for carbohydrates). Also, the rebinding benefits from slow translational diffusion as well as high flexibility in linkers, features that we herein show are characteristic for glycofullerene **2**.

EXPERIMENTAL

NMR samples of compounds **1** and **2** (synthesis described elsewhere^{9,11}) were prepared by dissolving the glycofullerenes in D₂O followed by treatment with Chelex 100 to remove divalent cations. Subsequent freeze-drying of the solution filtered from chelating resin was followed by re-dissolution in D₂O (pH = 5) to give samples of either high concentration (25 mg·mL⁻¹) for relaxation measurements or low concentration (0.25 mg·mL⁻¹) for translational diffusion measurements. The NMR experiments were performed at 25 °C, where the temperature had been calibrated²⁸ with a methanol-*d*₄ sample prior the experiments.

The translational diffusion coefficients of compounds **1** and **2** were measured with pulsed-field-gradient (PFG) ¹H NMR experiments on both concentrated and dilute samples. The experiments were performed on a 600 MHz Bruker

Avance III NMR spectrometer equipped with a 5 mm TXI ($^1\text{H}/^{13}\text{C}/^{31}\text{P}$) probe, where the Z-gradient had been calibrated to compensate for gradient inhomogeneities by using a Gadolinium-doped water sample (1% H_2O in D_2O + 1 $\text{mg}\cdot\text{mL}^{-1}$ GdCl_3) with a literature value of $D_t = 1.90 \times 10^{-9} \text{ m}^2\cdot\text{s}^{-1}$ for the HDO resonance.²⁹ The diffusion time delay (Δ) was set to 60 ms and the gradient pulse length (δ) was set to 4.0 ms; each experiment was acquired with 32k data points, with gradient strengths going from 5% to 95% of maximum (55.7 $\text{G}\cdot\text{cm}^{-1}$), using either 32 (concentrated sample) or 208 scans (dilute sample). The decay of the sugar bulk region integral (3 – 4 ppm) was used to calculate the translational diffusion coefficient by fitting a Stejskal–Tanner type equation³⁰ to the data. The experiments were repeated four to ten times in order to obtain statistical information.

The ^{13}C relaxation measurements were performed on three different NMR spectrometers operating at various field strengths, namely a Bruker AVANCE III 700 MHz spectrometer (16.44 T, 699.970 MHz) equipped with a 5 mm TCI Z-Gradient Cryoprobe, a Bruker AVANCE III 600 MHz spectrometer (14.09 T, 600.130 MHz) equipped with a 5 mm BBO probe and a Bruker AVANCE 500 MHz spectrometer (11.75 T, 500.130 MHz) equipped with a TCI Z-Gradient Cryoprobe. FIDs of 32k data points were recorded with either 1k or 2k scans with the carrier set in the region 70 – 80 ppm and spectral widths of 100 – 120 ppm. An exponential window function of 5 Hz and zero-filling to at least 64k data points were applied to all FIDs prior Fourier transformation. Fast inversion recovery experiments³¹ were used to measure R_1 relaxation rates and were performed at the three magnetic fields using an inter-scan delay of at least 3 s and six mixing times, ranging from 10 ms to 5 s, in a random order in each experiment. The relaxation rates were then fitted from the peak intensities using an in-house matlab (MathWorks, R2012a) script. The R_2 relaxation rates were measured by the use of Carr-Purcell-Meiboom-Gill (CPMG) spin echo experiments³² at two magnetic field strengths (11.75 T and 16.44 T), using interscan delays of 4 s and pulse train refocusing delays of 0.5 ms. Each experiment contained eight mixing times, ranging from 8 ms to 190 ms, which were applied in a random order, and R_2 relaxation rates were calculated using the fitting routine in the vendor-supplied software Topspin 3.1. The NOE was measured at the three field strengths using the dynamic NOE experiment in pairs, with either a long (2 s) or short (1 ms) irradiation and inter-scan delays of at least 8 s. Enhancement factors were calculated as the ratio of peak intensities between the two experiments. All experiments were repeated 2 – 5 times from which averages and standard deviations were calculated for the NMR relaxation observables.

The model-free analysis was performed using the program *Relax*.^{16,17} Averages and standard deviations of the NMR relaxation data were used as input and an isotropic diffusion tensor was assumed, using global rotational diffusion correlation times that had been calculated from the translational diffusion coefficients³³ measured by the NMR PFG diffusion experiments performed on the concentrated samples of compounds **1** and **2**.

ACKNOWLEDGEMENTS

This work was supported by grants from the Swedish Research Council and the Knut and Alice Wallenberg Foundation.

REFERENCES

- (1) Varki, A. *Glycobiology* **1993**, *3*, 97.
- (2) Lee, Y. C.; Lee, R. T. *Acc. Chem. Res.* **1995**, *28*, 321.
- (3) Mammen, M.; Choi, S.-K.; Whitesides, G. M. *Angew. Chem. Int. Ed.* **1998**, *37*, 2755.
- (4) Lundquist, J. J.; Toone, E. J. *Chem. Rev.* **2002**, *102*, 555.
- (5) Kitov, P. I.; Sadowska, J. M.; Mulvey, G.; Armstrong, G. D.; Ling, H.; Pannu, N. S.; Read, R. J.; Bundle, D. R. *Nature* **2000**, *403*, 669.
- (6) Ordanini, S.; Varga, N.; Porkolab, V.; Thépaut, M.; Belvisi, L.; Bertaglia, A.; Palmioli, A.; Berzi, A.; Trabattoni, D.; Clerici, M.; Fieschi, F.; Bernardi, A. *Chem. Commun.* **2015**, *51*, 3816.
- (7) Houseman, B. T.; Mrksich, M. *Host-Guest Chemistry* **2002**, *218*, 1.
- (8) Choi, S.-K. *Synthetic Multivalent Molecules: Concepts and Biomedical Applications*; John Wiley & Sons, Inc.: New Jersey, USA, 2004.
- (9) Nierengarten, J.-F.; Iehl, J.; Oerthel, V.; Holler, M.; Illescas, B. M.; Muñoz, A.; Martín, N.; Rojo, J.; Sánchez-Navarro, M.; Cecioni, S.; Vidal, S.; Buffet, K.; Durka, M.; Vincent, S. P. *Chem. Commun.* **2010**, *46*, 3860.
- (10) Sánchez-Navarro, M.; Muñoz, A.; Illescas, B. M.; Rojo, J.; Martín, N. *Chem. Eur. J.* **2011**, *17*, 766.
- (11) Luczkowiak, J.; Muñoz, A.; Sánchez-Navarro, M.; Ribeiro-Viana, R.; Ginieis, A.; Illescas, B. M.; Martín, N.; Delgado, R.; Rojo, J. *Biomacromolecules* **2013**, *14*, 431.
- (12) Novoa-Carballal, R.; Säwén, E.; Fernandez-Megia, E.; Correa, J.; Riguera, R.; Widmalm, G. *Phys. Chem. Chem. Phys.* **2010**, *12*, 6587.
- (13) Rathgeber, S.; Pakula, T.; Urban, V. *J. Chem. Phys.* **2004**, *121*, 3840.
- (14) Klos, J. S.; Sommer, J. U. *Macromolecules* **2009**, *42*, 4878.
- (15) Lue, L. *Macromolecules* **2000**, *33*, 2266.
- (16) d'Auvergne, E. J.; Gooley, P. R. *J. Biomol. NMR* **2008**, *40*, 107.
- (17) d'Auvergne, E. J.; Gooley, P. R. *J. Biomol. NMR* **2008**, *40*, 121.
- (18) Clore, G. M.; Szabo, A.; Bax, A.; Kay, L. E.; Driscoll, P. C.; Gronenborn, A. M. *J. Am. Chem. Soc.* **1990**, *112*, 4989.
- (19) Markelov, D. A.; Lyulin, S. V.; Gotlib, Y. Y.; Lyulin, A. V.; Matveev, V. V.; Lahderanta, E.; Darinskii, A. A. *J. Chem. Phys.* **2009**, *130*, 044907.
- (20) Meirovitch, E.; Shapiro, Y. E.; Polimeno, A.; Freed, J. H. *J. Phys. Chem. A* **2006**, *110*, 8366.
- (21) Pinto, L. F.; Correa, J.; Martin-Pastor, M.; Riguera, R.; Fernandez-Megia, E. *J. Am. Chem. Soc.* **2013**, *135*, 1972.
- (22) Mandel, A. M.; Akke, M.; Palmer III, A. G. *Biochemistry* **1996**, *35*, 16009.
- (23) Menon, S.; Rosenberg, K.; Graham, S. A.; Ward, E. M.; Taylor, M. E.; Drickamer, K.; Leckband, D. E. *Proc. Natl. Acad. Sci. U. S. A.* **2009**, *106*, 11524.
- (24) Cecioni, S.; Imberty, A.; Vidal, S. *Chem. Rev.* **2015**, *115*, 525.
- (25) Varga, N.; Sutkeviciute, I.; Ribeiro-Viana, R.; Berzi, A.; Ramdasi, R.; Daggetti, A.; Vettoretti, G.; Amara, A.; Clerici, M.; Rojo, J.; Fieschi, F.; Bernardi, A. *Biomaterials* **2014**, *35*, 4175.
- (26) Weber, M.; Bujotzek, A.; Haag, R. *J. Chem. Phys.* **2012**, *137*, 054111.
- (27) Kitov, P. I.; Bundle, D. R. *J. Am. Chem. Soc.* **2003**, *125*, 16271.
- (28) Findeisen, M.; Brand, T.; Berger, S. *Magn. Reson. Chem.* **2007**, *45*, 175.
- (29) Damberg, P.; Jarvet, J.; Gräslund, A. *J. Magn. Reson.* **2001**, *148*, 343.
- (30) Stejskal, E. O.; Tanner, J. E. *J. Chem. Phys.* **1965**, *42*, 288.
- (31) Canet, D.; Levy, G. C.; Peat, I. R. *J. Magn. Reson.* **1975**, *18*, 199.
- (32) Meiboom, S.; Gill, D. *Rev. Sci. Instrum.* **1958**, *29*, 688.
- (33) Price, W. S. *Concepts in Magnetic Resonance* **1997**, *9*, 299.

TABLES

Table 1. Translational diffusion coefficients measured under dilute (D_0) and concentrated (D_t) conditions and calculated $R_{H,0}$ and $R_{H,t}$.


Glycofullerene	$D_0 \times 10^8 / \text{cm}^2 \text{s}^{-1}$	$D_t \times 10^8 / \text{cm}^2 \text{s}^{-1}$	$R_{H,0} / \text{\AA}$	$R_{H,t} / \text{\AA}$
1	159(7)	90.7(5.0)	12.5	21.9
2	74.4(2.6)	83(0.5)	26.7	24.0

Table 2. Extended model-free (MF) parameters fitted from ^{13}C NMR relaxation data of glycofullerenes **1** and **2**.

	χ^2_{red}	S^2	S_f^2	S_s^2 ⁱ	τ_s (ns)	τ_c (ns) ⁱⁱ
1	7.4	0.17±0.02	0.81±0.02	0.217	0.52±0.04	11.74
2	10.7	0.04±0.002	0.87±0.04	0.051	0.25±0.05	15.32

ⁱCalculated as S_f^2/S^2 ⁱⁱCalculated from D_t employing the Debye-Stokes equation

FIGURES AND LEGENDS



TOC




Figure 1. Schematic representation of the mannofullerene compounds in this study. Green spheres = α -D-mannoses, pentagons = triazoles, wavy lines = polyethylene glycol linkers and cyan or magenta spheres = [60]fullerene of **1** and **2**, respectively. For a more detailed representation of the structures, see supplementary information.




Figure 2. Translational diffusion (D_0) as a function of molecular mass (M_w). The lines predict the ideal density growth of the zeroth generation glycofullerene **1** under poor ($\nu=0.33$, dashed) and good ($\nu=0.60$, solid) solvent conditions. The relatively lower D_0 value of **2** indicates a less compact molecular structure compared to **1** in D_2O solution.




Figure 3. Average *NOE*, R_1 and R_2 parameters of endocyclic carbon atoms measured of the mannosyl residues for glycofullerenes **1** (cyan) and **2** (magenta) at magnetic field strength of 11.75 T, 14.09 T and 16.44 T. Error bars represent one standard deviation.