

Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

Total Synthesis of Calothrixin B via Sequential Sonogashira Coupling/Copper-Catalyzed Oxidative Cyclization

Nagarajan Ramkumar and Rajagopal Nagarajan*

Received 00th January 20xx,
Accepted 00th January 20xx


DOI: 10.1039/x0xx00000x

www.rsc.org/

A total synthesis of antimalarial indolo[3,2-*f*]phenanthridine alkaloid, Calothrixin B is reported. The key intermediate, ketoester **11** was assembled using sequential Sonogashira coupling and intra/intermolecular fashioned copper-catalyzed oxidative cyclization reactions.

Introduction

Calothrixin B **1** is a pentacyclic quinone that possesses a indolo[3,2-*f*]phenanthridine or quinolino[4,3-*b*]carbazole skeleton and is unique among other naturally occurring alkaloids (Fig. 1). Since its first isolation from *Calothrix* cyanobacteria in 1999¹, several research groups have reported the synthesis and biological studies of Calothrixin B, principally because of its potent biological activities, such as anti-malarial and anti-cancer activities.^{2, 3} Various synthetic methods have been reported for the production of Calothrixin B, including a metallation strategy,⁴ Pd-catalyzed coupling,⁵ electrocyclization,⁶ hetero Diels-Alder reaction,⁷ Friedel-Crafts acylation/alkylation⁸ and a radical reaction.⁹ In addition, two biomimetic approaches for Calothrixin B synthesis have also been reported.¹⁰



Calothrixin B 1


Figure 1. Structure of Calothrixin B

The Sonogashira coupling reaction is widely used in the synthesis of 1,*n*-enynes system, which is an important unit in biologically active natural compounds.¹¹ Copper-catalyzed oxidative cyclization of 1,*n*-enynes is an elegant tool for the synthesis of 4-carbonyl quinoline system that is found in many natural compounds.¹² Herein, we report a novel total synthetic

route for Calothrixin B involving sequential Sonogashira coupling and copper-catalyzed oxidative cyclization reactions.

Results and discussion

The retrosynthetic disconnection pathway for the synthesis of Calothrixin B **1** is shown in Scheme 1. We envisaged that **1** could be derived from ketoester **11** through subsequent ester hydrolysis, cyclodehydration and debenzylation. The key intermediate **11** could be accessed from compounds **8** and **10** using intra/intermolecular copper-catalyzed oxidative cyclization. In turn, the compounds **8** and **10** could be obtained from the Sonogashira coupling of **7** with **3** and **9** respectively.




Scheme 1. Retrosynthetic approach to Calothrixin B (1)

School of Chemistry, University of Hyderabad, Hyderabad-500046, India.

E-mail: mnc@uohyd.ernet.in (Dr. R. Nagarajan); Fax: +91 40-23012460


Electronic Supplementary Information (ESI) available: Copies of ¹H, ¹³C and Mass spectra for all the compounds. See DOI: 10.1039/x0xx00000x

We embarked on the proposed synthetic route by conducting a Sonogashira coupling reaction of 2-ethynyl-1*H*-indole **2** and (*E*)-ethyl 3-((2-iodophenyl)amino)acrylate **3** produced a coupled product **4** in the presence of Pd(PPh₃)₂Cl₂ (5 mol%), CuI (10 mol%) and diisopropylamine (DIPA) (2 equiv.) in THF with 92% yield (Scheme 2). Our next step was to focus on copper-catalyzed oxidative cyclization reaction of enyne **4**. Most surprisingly, on carrying out this reaction using CuCl₂·2H₂O (10 mol%), Phen (20 mol%) and 1,4-diazabicyclo[2.2.2]octane (DABCO) (2 equiv.) in DMF at 100 °C under oxygen atmosphere (using balloon), we isolated benzo[*f*]indolo[1,2-*b*][2,7]naphthyridine-7,14-dione **6** in 78% yield instead of the expected ketoester **5**. From this result, we concluded that ketoester **5** was formed *in situ* and readily lactamized to a compound **6** due to the presence of free indole NH in the enyne **4**. The isolated compound **6** has been previously reported in literature.¹³ Data from spectroscopic analysis of the compound **6** matched literature report in all aspects (see Supporting information).




Scheme 2. Synthesis of benzo[*f*]indolo[1,2-*b*][2,7]naphthyridine-7,14-dione (**6**)

In order to achieve the desired outcome, we protected the free NH in indole with a benzyl group **7**. After this modification, 1-benzyl-2-ethynyl-1*H*-indole **7** was subjected to Sonogashira coupling reactions with (*E*)-ethyl 3-((2-iodophenyl)amino)acrylate **3** and 2-iodoaniline **9** by employing Pd(PPh₃)₂Cl₂ (5 mol%), CuI (10 mol%) and diisopropylamine (DIPA) (2 equiv.) in THF at room temperature. These reactions afforded **8** and **10** in 94% and 96% yield respectively (Scheme 3).



Scheme 3. Sonogashira coupling reaction of 1-benzyl-2-ethynyl-1*H*-indole (**7**)


With the coupled products **8** and **10** in hand, we were able to obtain the key intermediate **11** in two ways; intramolecular and intermolecular copper-catalyzed oxidative cyclization reactions (Scheme 4). Firstly, we carried out intramolecular copper-catalyzed oxidative cyclization of enyne **8**. After several screening conditions, we found that the conversion of **8** to **11** proceeded smoothly in the presence of CuCl₂·2H₂O (10 mol%), Phen (20 mol%) and 1,4-diazabicyclo[2.2.2]octane (DABCO) (2 equiv.) in DMF at 100 °C under oxygen atmosphere with 82% yield in 3 h. This reaction proceeded through a sequence of intramolecular carbocupration of the alkyne **8** followed by isomerization, elimination and further oxidation gave the desired product **11**.^{12e} Next, we carried out intermolecular copper-catalyzed oxidative cyclization reaction of **10** with ethyl acrylate **12**. The conversion of **10** to **11** was effected using 20 mol% CuCl and 1,4-diazabicyclo[2.2.2]octane (DABCO) (2 equiv.) in NMP at 100 °C under oxygen atmosphere for 6 h.



Scheme 4. Copper-catalyzed oxidative cyclization

At this juncture, having the key intermediate **11** in hand, we performed further functional group manipulations, consisting of ester hydrolysis, cyclodehydration and debenzoylation (Scheme 5). The ketoester **11** was hydrolyzed using NaOH/EtOH to generate ketoacid **13**, which was in turn

converted into a known *N*-benzyl calothrixin **14** by treatment with concd H₂SO₄ at 140 °C. The *N*-benzylated calothrixin **14** was readily deprotected by treatment with 10% Pd/C and HCOONH₄ in MeOH at reflux to finally obtain the target compound **1** in 71% yield. Spectroscopic data of **1** and **14** are fully consistent with those previously reported in literatures.^{1,7}



Scheme 5. Reactions to complete synthesis of Calothrixin B (**1**)

Conclusions

In summary, the total synthesis of Calothrixin B **1** has been accomplished using a novel route comprising of sequential Sonogashira coupling and copper-catalyzed oxidative cyclization reactions with an overall yield of 40.6% to 41.8% over 5 steps. This synthetic strategy provides a promising platform to synthesize novel indolonaphthyridine ring systems with ease.

Experimental Section

General Information and Materials

The NMR experiments were performed with 400 or 500 MHz spectrometer, and chemical shifts are expressed in ppm (δ) with TMS as an internal reference. Coupling constant *J* values are given in Hz. IR spectra were recorded by using KBr pellets or neat. ESI-TOF mass analyzer type was used for the HRMS measurements. Reactions were carried out under an inert atmosphere, referring to the use of nitrogen, and monitored by TLC. Column chromatography was performed on silica gel (100–200 mesh) in glass columns to purify the compounds. Solvents tetrahydrofuran (THF), *N,N*-dimethylformamide (DMF), methanol, ethyl acetate and dichloromethane were dried by using standard distillation methods. Commercially available reagents and solvents were used without further purification and were purchased. Melting points were determined using open capillary tubes and are uncorrected.

Preparation of (*E*)-ethyl 3-((2-iodophenyl)amino)acrylate (**3**)

Bis-dichloro(acetonitrile)palladium(II) (24 mg, 0.1 mmol), *p*-benzoquinone (98 mg, 1 mmol), lithium chloride (390 mg, 10 mmol) and ethyl acrylate **12** (100 mg, 1 mmol) were mixed in

THF (5 mL) at room temperature. After stirring for 15 min, 2-iodoaniline **9** (210 mg, 1 mmol) in THF (5 mL) was added. The resulting mixture was stirred for 12 h at room temperature and the solvent was removed under a reduced pressure. The residue was dissolved in ethyl acetate (75 mL) and repeatedly washed with 1 M NaOH solution and water. The organic layer was separated, dried over anhydrous Na₂SO₄ and the solvent was removed to give the crude product which was purified by silica gel column chromatography using petroleum ether.

R_f = 0.60 (petroleum ether: EtOAc, 95:5); colourless oil (240 mg, 80%); IR (neat): 3380, 1730 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz): δ 10.16 (s, br, 1H), 7.81 (d, *J* = 7.6 Hz, 1H), 7.32 (t, *J* = 7.2 Hz, 1H), 7.25–7.20 (m, 1H), 7.03 (d, *J* = 8.0 Hz, 1H), 6.75 (t, *J* = 7.2 Hz, 1H), 4.97 (d, *J* = 8.4 Hz, 1H), 4.28 (q, *J* = 6.8 Hz, 2H), 1.35 (t, *J* = 7.2 Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz): δ 169.7, 141.8, 141.6, 139.9, 129.4, 123.6, 113.6, 89.7, 87.3, 59.6, 14.5; HRMS (ESI): [M+H]⁺ calcd for C₁₁H₁₂INO₂ 317.9991, found 317.9987.

Preparation of 1-benzyl-2-ethynyl-1*H*-indole (**7**)

To a solution of 1-benzyl-2-ethynyl-1*H*-indole **2** (150 mg, 1 mmol) in DMF (10 mL), K₂CO₃ (293 mg, 2 mmol) was added and stirred for 30 min at room temperature. Benzyl bromide (0.25 mL, 2 mmol) was added and the mixture was stirred for 12 h under nitrogen atmosphere at room temperature. After completion of reaction (TLC), the reaction mixture was poured into water (100 mL) and extracted with ethyl acetate (75 mL). The organic layer was washed with brine, dried over anhydrous Na₂SO₄ and the solvent was removed to give the crude product which was purified by silica gel column chromatography using petroleum ether.

R_f = 0.64 (petroleum ether:EtOAc, 95:5); light yellow solid (235 mg, 96%); mp 112–114 °C; IR (KBr): 3320, 3095, 2260, 1445, 890 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz): δ 7.64 (d, *J* = 7.6 Hz, 1H), 7.33–7.14 (m, 8H), 6.94 (s, 1H), 5.49 (s, 2H), 3.48 (s, 1H); ¹³C NMR (CDCl₃, 100 MHz): δ 137.5, 136.6, 128.6 (2C), 127.4, 127.2, 126.7 (2C), 123.5, 121.2, 120.8, 120.4, 110.2, 109.0, 83.6, 75.6, 47.9; HRMS (ESI): [M+H]⁺ calcd for C₁₇H₁₃N 232.1126, found 232.1119.

General procedure for Sonogashira coupling reaction:

A mixture of (*E*)-ethyl 3-((2-iodophenyl)amino)acrylate **3** (150 mg, 1 mmol) or 2-iodoaniline **9** (150 mg, 1 mmol), Pd(PPh₃)₂Cl₂ (0.05 mmol) and CuI (0.1 mmol) were dissolved in THF (5 mL). Then DIPA (2 mmol) was added when the solution became clear. After stirring the mixture for 15 min at room temperature, corresponding indole **2** or **7** (1 mmol) was added. The reaction mixture was further allowed to stir for 2 h at room temperature. After completion of the reaction (TLC), the solvent was evaporated. The crude mixture was purified by silica gel column chromatography using petroleum ether/ethyl acetate.

(*E*)-Ethyl 3-((2-((1*H*-indol-2-yl)ethynyl)phenyl)amino)acrylate (**4**)

R_f = 0.44 (petroleum ether:EtOAc, 95:5); white solid (144 mg, 92%); mp 128–130 °C; IR (KBr): 3380, 3345, 2985, 1750, 1585, 1335, 725 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz): δ 11.04 (d, *J* = 12.4 Hz, 1H), 10.57 (s, 1H), 7.65 (d, *J* = 8.0 Hz, 1H), 7.50 (d, *J* = 7.6 Hz, 1H), 7.46–7.41 (m, 1H), 7.33–7.23 (m, 3H), 7.16–7.12 (m, 2H), 6.99 (t, *J* = 7.6 Hz, 1H), 6.86 (d, *J* = 0.8 Hz, 1H), 5.03 (d, *J* =

8.0 Hz, 1H), 4.44 (q, $J = 7.2$ Hz, 2H), 1.49 (t, $J = 6.8$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 170.8, 141.8, 141.3, 136.7, 130.9, 129.5, 127.8, 123.3, 121.7, 121.0, 120.1, 118.8, 110.9, 110.8, 110.6, 107.5, 90.5, 88.3, 87.8, 59.9, 14.5; HRMS (ESI): $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{21}\text{H}_{18}\text{N}_2\text{O}_2$ 331.1446, found 331.1441.

(E)-Ethyl 3-((2-((1-benzyl-1H-indol-2-yl)ethynyl)phenyl)amino)acrylate (8)

$R_f = 0.48$ (petroleum ether:EtOAc, 95:5); yellow solid (186 mg, 94%); mp 146–148 °C; IR (neat): 3350, 2985, 2310, 1755, 1430, 1235, 760 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz): δ 10.70 (d, $J = 8.0$ Hz, 1H), 7.65 (d, $J = 8.0$ Hz, 1H), 7.44 (dd, $J = 1.2, 7.6$ Hz, 1H), 7.35–7.20 (m, 10H), 7.14 (d, $J = 7.6$ Hz, 1H), 7.09 (d, $J = 8.4$ Hz, 1H), 6.96 (t, $J = 7.6$ Hz, 1H), 5.62 (s, 2H), 4.97 (d, $J = 8.4$ Hz, 1H), 4.22 (q, $J = 7.6$ Hz, 2H), 1.33 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 169.7, 141.7, 141.2, 137.8, 132.5, 129.8, 128.7, 128.6 (2C), 127.8, 127.3, 126.9 (2C), 123.2, 121.7, 121.5, 121.1, 120.2, 112.0, 110.8, 110.1, 109.3, 90.8, 89.4, 88.0, 59.4, 48.2, 14.5; HRMS (ESI): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{28}\text{H}_{24}\text{N}_2\text{O}_2$ 443.1736, found 443.1730.

2-((1-Benzyl-1H-indol-2-yl)ethynyl)aniline (10)

$R_f = 0.36$ (petroleum ether:EtOAc, 90:10); yellow solid (212 mg, 96%); mp 108–110 °C; IR (KBr): 3440, 3360, 3080, 2956, 1467, 1332, 1280, 764 cm^{-1} ; ^1H NMR (CDCl_3 , 500 MHz): δ 7.65 (d, $J = 9.5$ Hz, 1H), 7.33–7.28 (m, 3H), 7.26–7.21 (m, 3H), 7.16 (t, $J = 7.0$ Hz, 4H), 7.13 (s, 1H), 6.93–6.68 (m, 2H), 5.54 (s, 2H), 4.03 (s, br, 2H); ^{13}C NMR (CDCl_3 , 500 MHz): δ 147.9, 137.7, 136.9, 132.1, 130.1, 128.7 (2C), 127.5, 127.4, 126.4 (2C), 123.2, 121.9, 121.0, 120.3, 117.9, 114.3, 109.9, 107.9, 107.1, 92.0, 86.1, 47.8; HRMS (ESI): $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{23}\text{H}_{18}\text{N}_2$ 323.1548, found 323.1540.

General procedure for intramolecular copper-catalyzed oxidative cyclization reaction:

To a solution of appropriate enyne **4** or **8** (100 mg, 1 mmol) in DMF (2 mL) was added $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ (0.1 mmol), DABCO (2 mmol) and 1,10-Phen (0.2 mmol). The reaction mixture was then stirred for 3 h at 100 °C under O_2 atmosphere. The resulting mixture was quenched with water and extracted with EtOAc (2 × 30 mL). The combined organic extracts were washed with brine, dried over anhydrous Na_2SO_4 and concentrated. The crude product was purified by silica gel column chromatography using petroleum ether/ethyl acetate.

Benzo[*f*]indolo[1,2-*b*][2,7]naphthyridine-7,14-dione (6)¹³

$R_f = 0.57$ (petroleum ether:EtOAc, 80:20); yellow solid (70 mg, 78%); mp 244–246 °C (lit.¹³ mp 249 °C); IR (KBr): 1690, 1660, 1573, 1415, 1384, 1245, 780 cm^{-1} ; ^1H NMR (CDCl_3 , 500 MHz): δ 9.95 (s, 1H), 9.74 (dd, $J = 1.0, 8.5$ Hz, 1H), 8.60 (dd, $J = 1.0, 8.5$ Hz, 1H), 8.26 (dd, $J = 1.0, 8.5$ Hz, 1H), 7.94 (td, $J = 1.5, 8.5$ Hz, 1H), 7.85 (td, $J = 1.0, 8.5$ Hz, 1H), 7.77 (dt, $J = 1.0, 7.5$ Hz, 1H), 7.72 (d, $J = 0.5$ Hz, 1H), 7.63 (td, $J = 1.0, 8.5$ Hz, 1H), 7.41 (td, $J = 1.0, 8.0$ Hz, 1H); ^{13}C NMR (CDCl_3 , 125 MHz): δ 177.7, 158.3, 151.7, 149.5, 136.7, 134.0, 133.8, 132.4, 130.7, 130.5, 130.3, 128.7, 127.8, 125.7, 123.9, 123.3, 122.7, 117.9, 117.1; HRMS (ESI): $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{19}\text{H}_{10}\text{N}_2\text{O}_2$ 299.0820, found 299.0820.

Ethyl 4-(1-benzyl-1H-indole-2-carbonyl)quinoline-3-carboxylate (11)

$R_f = 0.42$ (petroleum ether:EtOAc, 80:20); yellow solid (84 mg, 82%); mp 148–150 °C; IR (KBr): 3120, 2850, 1765, 1650; 1430, 1245 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz): δ 9.55 (s, 1H), 8.22 (d, $J = 8.4$ Hz, 1H), 7.80 (t, $J = 8.4$ Hz, 1H), 7.64 (d, $J = 8.0$ Hz, 1H), 7.55 (d, $J = 8.4$ Hz, 1H), 7.48 (t, $J = 7.2$ Hz, 1H), 7.39–7.22 (m, 9H), 6.27 (d, $J = 15.6$ Hz, 1H), 5.95 (d, $J = 16.0$ Hz, 1H), 4.32 (m, 2H), 1.26 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 186.3, 164.6, 150.2, 149.7, 149.2, 138.8, 137.9, 131.8, 129.6, 128.7 (2C), 128.2, 127.9, 127.5, 126.9 (2C), 125.6, 125.0, 124.4, 121.8, 120.9, 119.9, 116.5, 111.1, 61.8, 49.2, 13.9; HRMS (ESI): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{28}\text{H}_{22}\text{N}_2\text{O}_3$ 457.1528, found 457.1520.

Procedure for intermolecular copper-catalyzed oxidative cyclization reaction:

To a solution of 2-((1-benzyl-1H-indol-2-yl)ethynyl)aniline **10** (100 mg, 1 mmol) and ethyl acrylate **12** (50 μL , 1.5 mmol) in NMP (2 mL), CuCl (0.2 mmol) and DABCO (2 mmol) were added. The reaction mixture was then stirred for 6 h at 100 °C under O_2 atmosphere. The resulting mixture was quenched with water and extracted with EtOAc (2 × 30 mL). The combined organic extracts were washed with brine, dried over anhydrous Na_2SO_4 and concentrated. The crude product was purified by silica gel column chromatography using petroleum ether/ethyl acetate (80:20) afforded **11** as yellow solid (105 mg, 78%).

Synthesis of 4-(1-benzyl-1H-indole-2-carbonyl)quinoline-3-carboxylic acid (13)

To a solution of ethyl 4-(1-benzyl-1H-indole-2-carbonyl)quinoline-3-carboxylate **11** (75 mg, 1 mmol) in ethanol (3 mL), NaOH (21 mg, 3 mmol) was added and refluxed for 1 h. After completion of the reaction (TLC), crushed ice was added to the mixture and neutralized with 10% dil HCl to pH 7. The solid formed was filtered off, washed with water and dried. The crude product was used in the next step without further purification.

Note: This compound **13** exists as 'rotamers'

$R_f = 0.38$ (DCM:MeOH, 90:10); yellow solid (84 mg, 98%); mp 218–220 °C; IR (neat): 3110, 2930, 1740, 1690, 1455, 1375, 890 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz): δ 10.05 (s, br, 2H), 9.70 (s, 1H), 9.69 (s, 1H), 8.22 (t, $J = 7.6$ Hz, 2H), 7.72–7.65 (m, 2H), 7.60 (t, $J = 7.6$ Hz, 2H), 7.53 (d, $J = 8.4$ Hz, 1H), 7.48–7.25 (m, 18H), 7.19 (t, $J = 7.2$ Hz, 2H), 7.11 (t, $J = 7.6$ Hz, 1H), 6.17 (d, $J = 16.0$ Hz, 2H), 5.97 (d, $J = 16.0$ Hz, 2H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 187.0, 186.0, 167.2, 167.0, 150.6, 150.1 (2C), 149.8, 148.2, 140.6, 138.8, 138.1, 137.8 (2C), 134.7, 132.4, 129.7, 128.69 (4C), 128.62 (2C), 128.3, 128.28, 128.23, 127.4, 127.3, 127.0, 126.9 (6C), 126.2, 125.8, 125.1, 125.0, 124.8, 123.4, 121.8, 121.3, 120.9 (2C), 120.8, 116.4, 116.0, 111.1 (2C), 49.2, 48.6; HRMS (ESI): $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{26}\text{H}_{18}\text{N}_2\text{O}_3$ 407.1395, found 407.1387.

Synthesis of 12-benzyl-7H-indolo[3,2-*j*]phenanthridine-7,13(12H)-dione (14)

4-(1-Benzyl-1H-indole-2-carbonyl)quinoline-3-carboxylic acid **13** (70 mg) was added to concd H_2SO_4 (2 mL) and heated to 140 °C for 2 h. After completion of the reaction (TLC), the

reaction mixture was poured into ice cold water (50 mL) and extracted with ethyl acetate (70 mL). The organic layer was separated, washed with 10% aq NaOH and dried over anhydrous Na₂SO₄. The solvent was removed to give the crude residue which was purified by silica gel column chromatography using petroleum ether/ethyl acetate.

R_f = 0.30 (petroleum ether:EtOAc, 80:20); red solid (84 mg, 78%); mp 258–260 °C (lit.⁷ mp 264 °C); IR (KBr): 3050, 1650, 1530, 1430, 1248 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz): δ 9.80 (s, 1H), 9.55 (d, *J* = 8.8 Hz, 1H), 8.46 (d, *J* = 8.0 Hz, 1H), 8.19 (d, *J* = 8.0 Hz, 1H), 7.83 (t, *J* = 7.2 Hz, 1H), 7.74 (t, *J* = 8.0 Hz, 1H), 7.46–7.42 (m, 3H), 7.35–7.29 (m, 3H), 7.23 (d, *J* = 7.2 Hz, 2H), 6.01 (s, 2H); ¹³C NMR (CDCl₃, 100 MHz): δ 182.0, 181.0, 152.1, 147.8, 140.1, 136.2, 135.1, 133.2, 131.3, 130.3, 130.1, 128.9 (2C), 128.0, 127.8, 127.7, 126.6 (2C), 125.1, 124.4, 123.9, 123.3, 123.1, 117.7, 111.5, 48.5; HRMS (ESI): [M+H]⁺ calcd for C₂₆H₁₆N₂O₂ 389.1290, found 389.1282.

Synthesis of 7*H*-indolo[3,2-*f*]phenanthridine-7,13(12*H*)-dione (1) (Calothrixin B)

To a solution of 12-benzyl-7*H*-indolo[3,2-*f*]phenanthridine-7,13(12*H*)-dione **14** (50 mg, 0.13 mmol) in MeOH (5 mL), 10% Pd/C (14 mg, 0.13 mmol) and ammonium formate (162 mg, 2.5 mmol) were added and refluxed for 3 h under nitrogen atmosphere. After completion of the reaction using TLC, the reaction mixture was filtered through celite pad and washed with ethyl acetate (50 mL). The solvent was evaporated to give crude oil which was purified by silica gel column chromatography gave the titled product.

R_f = 0.30 (petroleum ether:EtOAc, 80:20); red solid (27 mg, 71%); mp 296–298 °C (lit.¹ mp ≥ 300 °C); IR (neat): 3420, 1650 cm⁻¹; ¹H NMR (DMSO-*d*₆, 400 MHz): δ 13.11 (s, br, 1H), 9.56 (s, 1H), 9.52 (d, *J* = 8.0 Hz, 1H), 8.12 (d, *J* = 8.0 Hz, 2H), 7.91 (t, *J* = 8.0 Hz, 1H), 7.84 (t, *J* = 8.4 Hz, 1H), 7.57 (d, *J* = 8.4 Hz, 1H), 7.42 (t, *J* = 7.6 Hz, 1H), 7.34 (t, *J* = 7.6 Hz, 1H); ¹³C NMR (DMSO-*d*₆, 100 MHz): δ 181.2, 180.7, 151.6, 147.9, 138.8, 138.4, 132.9, 132.0, 130.6, 130.28, 130.21, 127.6, 125.2, 124.7, 123.7, 123.0, 122.7, 115.9, 114.3; HRMS (ESI): [M+H]⁺ calcd for C₁₉H₁₀N₂O₂ 299.0820, found 299.0822.

Acknowledgements

We gratefully acknowledge DST for financial support (Project No SR/S1/OC-70/2008). NR thanks CSIR, New Delhi for senior research fellowship and contingency grant.

Notes and reference

- 1 R. W. Rickards, J. M. Rothschild, A. C. Willis, N. M. de Chazal, J. Kirk, K. Kirk, K. J. Saliba and G. D. Smith, *Tetrahedron* 1999, **55**, 13513.
- 2 (a) A. W. Schmidt, K. R. Reddy and H.-J. Knölker, *Chem. Rev.*, 2012, **112**, 3193; (b) H.-J. Knölker and K. R. Reddy, in *The Alkaloids*, ed. G. A. Cordell, Elsevier Science, 2008, vol. 65, p 1–430; (c) H.-J. Knölker and K. R. Reddy, *Chem. Rev.*, 2002, **102**, 4303; (d) C. M. Miller and F. O. McCarthy, *RSC Adv.*, 2012, **2**, 8883; (e) T. Choshi and S. Hibino, *Heterocycles* 2009, **77**, 85; (f) S. Tohyama, T. Choshi, K. Matsumoto, A.

- Yamabuki, Y. Hieda, J. Nobuhiro and S. Hibino, *Heterocycles* 2010, **82**, 397.
- (a) N. T. Doan, R. W. Rickards, J. M. Rothschild and G. D. Smith, *J. Appl. Phycol.*, 2000, **12**, 409; (b) N. T. Doan, P. R. Stewart and G. D. Smith, *Microbiol. Lett.*, 2001, **196**, 135; (c) X. Chen, G. D. Smith and P. Waring, *J. Appl. Phycol.*, 2003, **15**, 269; (d) P. H. Bernardo, C. L. L. Chai, G. A. Heath, P. J. Mahon and G. D. Smith, *J. Med. Chem.*, 2004, **47**, 4958; (e) P. H. Bernardo, C. L. L. Chai, M. L. Guen, G. D. Smith and P. Waring, *Bioorg. Med. Chem. Lett.*, 2007, **17**, 82; (f) Q. A. Khan, J. Lu and S. M. Hecht, *J. Nat. Prod.*, 2009, **72**, 38; (g) N. Hatae, R. Satoh, H. Chiba, T. Osaki, T. Nishiyama, M. Ishikura, T. Abe, S. Hibino, T. Choshi, C. Okada and E. Yoyota, *Med. Chem. Res.*, 2014, **23**, 4956.
- (a) T. R. Kelly, Y. Zhao, M. Cavero, and M. Torneiro, *Org. Lett.*, 2000, **2**, 3735; (b) N. Ramkumar and R. Nagarajan, *J. Org. Chem.*, 2014, **79**, 736; (c) D. Mal, J. Roy and K. Biradha, *Org. Biomol. Chem.*, 2014, **12**, 8196.
- (a) P. H. Bernardo, W. Fitriyanto and C. L. L. Chai, *Synlett* 2007, 1935; (b) T. Abe, T. Ikeda, R. Yanada and M. Ishikura, *Org. Lett.*, 2011, **13**, 3356; (c) T. Abe, T. Ikeda, T. Choshi, S. Hibino, N. Hatae, E. Toyata, R. Yanada and M. Ishikura, *Eur. J. Org. Chem.*, 2012, 5018; (d) S. M. Bhosale, R. L. Gawade, V. G. Puranik and R. S. Kusrkar, *Tetrahedron Lett.*, 2012, **53**, 2894; (e) N. Ramkumar and R. Nagarajan, *J. Org. Chem.*, 2013, **78**, 2802; (f) S. A. Kaliyaperumal, S. Banerjee and U. K., S. Kumar, *Org. Biomol. Chem.*, 2014, **12**, 6105; (g) S. M. Bhosale, M. A. Momin, S. Kunjir, P. R. Rajamohan and R. S. Kusrkar, *Tetrahedron Lett.*, 2014, **55**, 155.
- (a) S. Tohyama, T. Choshi, K. Matsumoto, A. Yamabuki, K. Ikegata, J. Nobuhiro and S. Hibino, *Tetrahedron Lett.*, 2005, **46**, 5263; (b) B. M. Ramalingam, V. Saravanan and A. K. Mohanakrishnan, *Org. Lett.*, 2013, **15**, 3726; (c) V. Saravanan, B. M. Ramalingam and A. K. Mohanakrishnan, *Eur. J. Org. Chem.*, 2014, 1266.
- (a) D. Sissouma, S. C. Collet and A. Y. Guingant, *Synlett* 2004, 2612; (b) D. Sissouma, L. Maingot, S. Collet and A. Guingant, *J. Org. Chem.*, 2006, **71**, 8384; (c) L. Maingot, F. Thuaud, D. Sissouma, S. Collet, A. Guingant and M. Evain, *Synlett* 2008, 263.
- (a) P. H. Bernardo, C. L. L. Chai and J. A. Elix, *Tetrahedron Lett.*, 2002, **43**, 2939; (b) P. H. Bernardo and C. L. L. Chai, *J. Org. Chem.*, 2003, **68**, 8906.
- (a) M.-L. Bannasar, T. Roca and F. Ferrando, *Org. Lett.*, 2006, **8**, 561; (b) S. Xu, T. Nguyen, I. Pomilio, M. C. Vitale and S. E. Velu, *Tetrahedron* 2014, **70**, 5928.
- (a) A. Yamabuki, H. Fujinawa, T. Choshi, S. Tohyama, K. Matsumoto, K. Ohmura, J. Nobuhiro and S. Hibino, *Tetrahedron Lett.*, 2006, **47**, 5859; (b) K. Matsumoto, T. Choshi, M. Hourai, Y. Zamami, K. Sasaki, T. Abe, M. Ishikura, N. Hatae, T. Iwamura, S. Tohyama and J. Nobuhiro, *Bioorg. Med. Chem. Lett.*, 2012, **22**, 4762; (c) C. S. P. McErlean, J. Sperry, A. J. Blake and C. J. Moody, *Tetrahedron* 2007, **63**, 10963; (d) J. Sperry, C. S. P. McErlean, A. M. Z. Slawin and C. J. Moody, *Tetrahedron Lett.*, 2007, **48**, 231.
- (a) K. Sonogashira, Y. Tohda and N. Hagihara, *Tetrahedron Lett.*, 1975, **16**, 4467; (b) K. Sonogashira, *J. Organomet. Chem.*, 2002, **653**, 46; (c) J. M. Pedersen, W. R. Bowman, M. R. J. Elsegood, A. J. Fletcher and P. J. Lovell, *J. Org. Chem.*, 2005, **70**, 10615; (d) C. Thongsornklee and R. L. Danhauser, *J. Org. Chem.*, 2005, **70**, 2364; (e) R. Chinchilla and C. Nájera, *Chem. Rev.*, 2007, **107**, 874.
- (a) N. T. Patil, H. Wu and Y. Yamamoto, *J. Org. Chem.*, 2005, **70**, 4531; (b) Z.-Q. Wang, W.-W. Zhang, L.-B. Gong, R.-Y. Tang, X.-H. Yang, Y. Liu and J.-H. Li, *Angew. Chem., Int. Ed.*, 2011, **50**, 8968; (c) C. Fehr, I. Farris and H. Sommer, *Org. Lett.*, 2006, **8**, 1839; (d) K. K. Toh, S. Sanjaya, S. Sahnoun, S. Y. Choung and S. Chiba, *Org. Lett.*, 2012, **14**, 2290; (e) X.-F. Xia,

ARTICLE

Journal Name

- L.-L. Zhang, X.-R. Song, X.-Y. Liu and Y.-M. Liang, *Org. Lett.*, 2012, **10**, 2480; (f) S. E. Allen, R. R. Walvoord, R. Padilla-Salinas and M. C. Kozlowski, *Chem. Rev.*, 2013, **114**, 6234.
- 13 D. H. Dethe and G. M. Murhade, *Eur. J. Org. Chem.*, 2014, 6953.