

Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Organic & Biomolecular Chemistry

REVIEW

Fluorinated Porphyrinoids as Efficient Platforms for New Photonic Materials, Sensors, and Therapeutics

N. V. S. Dinesh K. Bhupathiraju,^{a,†} Waqar Rizvi,^{a,†} James D. Batteas,^b Charles Michael Drain^{a,c,*}

Received 00th August 20xx,
Accepted 00th 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Porphyrinoids are robust heterocyclic dyes studied extensively for applications in medicine and as photonic materials because of their tunable photophysical properties, diverse means of modifying the periphery, and the ability to chelate most transition metals. Commercial applications include phthalocyanine dyes in optical discs, porphyrins in photodynamic therapy, and as oxygen sensors. Most applications of these dyes require exocyclic moieties to improve solubility, target disease, modulate photophysical properties, or direct self-organization into architectures with desired photonic properties. The synthesis of the porphyrinoid depends on the desired application, but the de novo synthesis often involves several steps, is time consuming, and results in low isolated yields. Thus, applications of core porphyrinoid platforms that can be rapidly and efficiently modified to evaluate new molecular architectures allows researchers to focus on the design concepts rather than the synthetic methods, and opens porphyrinoid chemistry to a broader scientific community. We have focused on several widely available, commercially viable porphyrinoids as platforms: *meso*-perfluorophenylporphyrin, perfluorophthalocyanine, and *meso*-perfluorophenylcorrole. The perfluorophenylporphyrin is readily converted to the chlorin, bacteriochlorin, and isobacteriochlorin. Derivatives of all six of these core platforms can be efficiently and controllably made via mild nucleophilic aromatic substitution reactions using primary S, N, and O nucleophiles bearing a wide variety of functional groups. The remaining fluoro groups enhance the photo and oxidative stability of the dyes and can serve as spectroscopic signatures to characterize the compounds or in imaging applications using ¹⁹F NMR. This review provides an overview of the chemistry of fluorinated porphyrinoids that are being used as a platform to create libraries of photo-active compounds for applications in medicine and materials.

1. INTRODUCTION

Porphyrinoids e.g. porphyrins (Pors), phthalocyanines (Pcs), corroles (Cors), chlorins (Chls), bacteriochlorins (Bacs) and isobacteriochlorins (Ibacs) have applications in diverse fields such as medicine,¹ solar energy conversion,² optical sensors,³ electronic devices,⁴ supramolecular systems,⁵ photovoltaic

^a Department of Chemistry and Biochemistry, Hunter College and Graduate Center of the City University of New York (CUNY), 695 Park Avenue, New York, NY 10065.

^b Department of Chemistry, Texas A&M University, College Station, TX 77842.

^c The Rockefeller University, 1230 York Avenue, New York, NY 10065.

Email-cdrain@hunter.cuny.edu

† N V S Dinesh K Bhupathiraju and Waqar Rizvi contributed equally to this work.



N V S Dinesh K Bhupathiraju

N V S Dinesh K Bhupathiraju was born in India. He obtained his Master's degree in physical organic chemistry (2008) from Osmania University, India. He obtained his Ph.D. in organic chemistry (2013) from Louisiana State University, USA under the guidance of Prof. Maria Graça

H. Vicente. Presently, he is working as a visiting lecturer in the Department of Chemistry and Biochemistry and as a Research Associate in the laboratory of Prof. Charles M. Drain, Hunter College, CUNY. His research projects involve theranostic nanomedicines for cancer and development of new dyes for solar energy harvesting materials for coating on windows in urban applications.



Waqar Rizvi

Waqar Rizvi was born in Sialkot, Pakistan and moved to the U.S.A at the age of eleven. He obtained his Bachelor's degrees in Biochemistry and Bioinformatics (2012) from Hunter College of The City University of New York (CUNY). He is currently

pursuing his Ph.D. in nanotechnology and materials chemistry at the Graduate Center, CUNY under the supervision of Prof. Charles M. Drain. His research is focused on developing new supramolecular porphyrinoid based dyes for materials and medicinal applications such as solar energy harvesting, biosensors, catalysts, cancer imaging and therapeutics. He is currently an adjunct lecturer at Hunter College, CUNY and teaches at First Academy, Little Neck, NY.

Review

cells,⁶ and catalysis.^{7, 8} Porphyrinoids are large aromatic dyes that are 1 to 1.5 nm in diameter. The Por core has 11 π bonds (Figure 1) and the Pc has an additional 8 in the fused isoindole structure (Figure 1), Cor and Chl have 10 while Ibac and Bac have 9 each. These macromolecules are the focus of research because of their stability under a wide range of environmental conditions, absorption in the visible and near infra-red (NIR) range of the electromagnetic spectrum (400 nm to 1000 nm), many derivatives possess long lived triplet excited states that sensitize singlet oxygen ($^1\text{O}_2$) formation to potentiate phototoxicity as anticancer and antibiotic agents, and the redox properties of metalloporphyrinoids are used in catalysis.⁹⁻¹² These macrocycles can chelate with nearly every metal in periodic table which results in variable functional properties such as tuneable excited state lifetimes, redox potentials, catalytic activities, molar absorption coefficients, and molecular dynamics. Functional groups on the periphery of structurally rigid porphyrinoids provides topological diversity making these dyes well suited for the engineering of supramolecular materials.⁹

Fluorous porphyrinoids are usually more stable towards oxidative degradation,¹³⁻¹⁵ have different excited state dynamics because of changes on the HOMO-LUMO energy gap, and exhibit different pharmacokinetics.¹⁶ The perfluorophenylporphyrin (TPPF₂₀), fluorophthalocyanine (PcF₁₆) and perfluorophenylcorrole (CorF₁₅) are now commercially available. TPPF₂₀ is converted to the corresponding Chl (CF₂₀), Ibac (IF₂₀), and Bac (BF₂₀) compounds (Figure 1) using Diels-Alder and 1,3-dipolar cycloaddition

reactions.¹⁷⁻¹⁹ Compared to Por, Chl, Ibac, and Bac have NIR absorbance but still show high $^1\text{O}_2$ yield for PDT.¹⁷ These fluorinated porphyrinoids can be used as ^{19}F magnetic resonance imaging (MRI) agents for *in vivo* imaging in combination with fluorescence spectroscopy.²⁰⁻²² These molecules are strongly stabilized due to the high electronegativity of fluorines, (Pauli electronegativity²³ for F $\frac{1}{4}$ 3.98). For Pcs, in particular, the electron-withdrawing halogens tend to lower the LUMO which makes the molecule more stable to ambient oxidation.²⁴




Figure 1 Fluorinated porphyrinoid platforms.



Charles M Drain

Charles Michael Drain is Professor and Chair of Chemistry at Hunter College, CUNY and adjunct faculty at Rockefeller University. He started his career in chemistry at the University of Missouri in St. Louis. He received his Ph.D. from Tufts University in the laboratory of Barry B. Corden where he worked on porphyrin synthesis. He did postdoctoral work in the laboratory of

David Mauzerall at Rockefeller University where he examined self-organizing systems composed of porphyrins and lipid bilayers and developed one of the first examples of a purely organic, synthetic phototransistor. He also examined all-D amino acid-containing channel-forming antibiotic peptides with R. Bruce Merrifield. He was a guest researcher in the laboratory of Jean-Marie Lehn at the University Louis Pasteur in Strasbourg, France, where he developed methodologies to self-assemble porphyrins. He was a research fellow in the Holt Holten/Kirmaier laboratory at Washington University studying the complex dynamics of nickel porphyrins. Since joining Hunter College, 1996, his research continues to focus on the design, synthesis, and characterization of self-assembled and self-organized photonic systems.



James D Batteas

James D Batteas is a Professor of Chemistry and Materials Science and Engineering at Texas A&M University. He earned a B.S. in Chemistry at the University of Texas at Austin in 1990 and a Ph.D. in Chemistry from the University of California at Berkeley in 1995. He is an expert in materials chemistry of surfaces and interfaces with research activities covering a broad range of

fundamental surface and interfacial phenomena. These include investigations of charge transport in organic molecules on surfaces (measured by STM and modeled by density functional theory), nanoparticle catalysis, semiconducting nanomaterials, plasmonics, tribology, "smart" surfaces, and self-organizing nanoscale materials for device applications in optoelectronics and chemical sensing. His research in tribology includes AFM studies of atomic scale friction and wear of oxides and 2D materials such as graphene. His group also employs the use of large-scale molecular dynamics simulations to explore the assemblies of molecules on nanoparticles and their interactions. He is a Fellow of the Royal Society of Chemistry and on the Editorial Boards of ACS Central Science and RSC Advances.

Despite the aforementioned advantages of fluorinated porphyrinoids, many applications are hampered by poor solubility and/or processability. For example, insolubility in water and no selectivity towards disease hamper use as therapeutics, while amorphous aggregation limits photonics applications. *De novo* synthesis of all but the simplest porphyrinoids results in low yields which impedes facile assessments of derivatives in materials and medicine.

N,N-Dimethylformamide (DMF), acetonitrile, chloroform, and toluene are common solvents for metallation of free base Por.²⁵ In 1990 Kadish et al.,²⁶ in an attempt to metallate meso-TPPF₂₀ in DMF, obtained meso-tetrakis(2,3,5,6-tetrafluoro-4-(dimethylamino)phenyl)porphyrinato metal complexes. Refluxing TPPF₂₀ in DMF without the metal salt also resulted in the dimethylamino substituted Por. The trace dimethylamine formed at high temperatures is nucleophilic enough to replace the *p*-fluoro group via a nucleophilic aromatic substitution reaction. This opened a new avenue for research using fluorinated porphyrinoids as platforms to develop new dyes for diverse applications.

In 1991, Mansuy and co-workers²⁷ studied the nucleophilic substitution reaction on free base and metal analogues of TPPF₂₀ including one with bromine on β -pyrrolic positions. They used S, N and O nucleophiles with electron donating functionality and observed higher yields of tetra substituted products with selective substitution on all four *p*-fluoro positions. But in the case of nucleophiles with electron withdrawing functional groups they observed complex mixtures. Later, lithium organic compounds were used by Kalisch and Senge to successfully substitute the Por core using the same reaction.^{28, 29} This substitution usually takes place via a two-step addition-elimination reaction mechanism in aprotic solvent using amine,^{26, 30} thiol,³¹⁻³⁴ or hydroxy³⁵⁻³⁸ groups as nucleophiles. Boyle and co-workers reported the substitution chemistry on mono-(pentafluorophenyl)porphyrin containing different thiol functional groups. Both one step and two step reactions were reported. Thiols were added to mono-




Figure 2 Scheme of nucleophilic substitution on fluorinated porphyrinoids (substitution in the order of softness S>N>O).

(pentafluorophenyl)-porphyrin in DMF/Et₃N (one step reaction)³² or mono-(pentafluorophenyl)porphyrin was reacted with Na₂S first and then a range of electrophiles where transient thiolate intermediate was obtained in situ to give a thio tether link (two step reaction).³⁴ For TPPF₂₀ the *p*-fluorine atom is selectively substituted prior to the *o*- or the *m*-fluorines,²² but the *m*- position can be substituted with greater equivalents of the nucleophile and elevated temperatures.

Thiols result in high yields and take less time compared to amines and hydroxyls. The nucleophile softness dictates the efficiency of the substitution in the order S>N>O (Figure 2).^{39, 40} Both the free base and the metallo fluorinated porphyrinoids are suitable for substitution chemistry, or the dyes can be metallated post modification. By controlling stoichiometry and reaction conditions, the TPPF₂₀, CF₂₀, IF₂₀ and BF₂₀ platforms can be appended with 1-4 groups.^{33, 41} Note that asymmetric moieties appended onto the CF₂₀, IF₂₀ and BF₂₀ platforms result in diastereomers at the bridge head positions. Because of the orthogonal geometry of the aryl groups and the barrier to rotation, substitution of more than one *m*- position results in atropisomers. The PcF₁₆ can be appended with 1-16 groups.⁴²

For example, several biomolecules such as sugars, polyamines, polyethylene glycols (PEG), peptides, amino acids, and boron clusters have been appended to one or more of the fluorinated dyes for disease targeting.^{41, 43-47} There are few examples where more than one biomolecular motif (e.g. cell targeting, hydrophilic, or chemotherapeutic) can be appended to a dye but the molecular topology of the porphyrinoids allows a range of substitutions.^{41, 46} For biotargeting and biosensing, the multiple valence significantly increases binding to the target. Other strategies that exploit the topological richness of the porphyrinoids are as drug carriers, e.g. boron clusters for boron neutron capture therapy (BNCT).^{41, 46} A variety of alkanes can be appended to drive self-organization into nanoscale thin photonic films and/or to manipulate the photophysical properties of the dyes leading to a more effective application in materials such as solar energy harvesting.^{42, 48-51} Appending exocyclic motifs onto the redox active metalloporphyrinoids results in molecules and materials with improved catalytic properties for organic transformations.^{52, 53} This review will focus mainly on nucleophilic substitution of fluorine analogues belonging to Pors, Pcs, Chls, Cors, Bacs and Ibacs. We will discuss their physicochemical as well as biomedical and material properties and applications.

2. SUBSTITUTED FLUORINATED PORPHYRINOID FOR BIOLOGICAL APPLICATIONS

Porphyrinoid based photosensitizers (PS) have been used in medicine in treating various types of diseases (oncology and non-oncology) such as bladder, brain, breast, skin, lung, esophagus, and bronchial cancers,^{54, 55} age-related macular degeneration, cardiovascular disease, ophthalmic disease, hair growth, acne, and psoriasis using photodynamic therapy (PDT).⁵⁶⁻⁶² These PS are presently studied for their application in treating microbial infections (bacterial, fungal, parasitic, and viral infections) using photodynamic antimicrobial chemotherapy (PACT).^{55, 63-66} These PS have also been used to diagnose the above mentioned diseases and so are considered as effective theranostic (imaging and treatment) agents.

PDT treatment involves three important elements: PS, which should accumulate specifically in the targeted site, triplet oxygen, and light. For PDT, the patient is administered a non-toxic PS dye. After sufficient time for maximum drug uptake by the cancer cells, the tumor is selectively irradiated with light to result in cytotoxic reactive oxygen species, which results in tumor destruction.⁵⁴ Presently there are few approved PDT agents (Photofrin, protoporphyrin IX, meso-tetrahydroxyphenylchlorin (m-THPC), and verteporfin) to treat several diseases.^{54, 67} These porphyrinoid based PS have several disadvantages such as poor water solubility, poor light absorption in NIR, poor disease selectivity and light sensitivity after treatment. Other treatment methods such as BNCT involving carboranes and porphyrinoids were also studied, where porphyrinoids were used as imaging and also adjunct treatment (PDT) method.^{68, 69}

Important ongoing approaches for improving potency of PS are fluorinated derivatives that modulate pharmacokinetics, $^1\text{O}_2$ production, photostability, and lipophilicity. In addition to the intrinsic fluorescence of the porphyrinoids, magnetic resonance of fluorinated compounds (^{19}F -NMR),⁷⁰⁻⁷² ^{18}F positron emission tomography (PET) imaging,^{13, 73} carbon-fluorine spectroscopy (CFR),¹⁶ and fluoro-Raman spectroscopy (FRS)⁷⁴ are all enabled for sensitive and minimally invasive imaging of disease or as sensors.

Another advantage of these porphyrinoid platforms is the ease of substituting multiple copies of exocyclic biotargeting motifs to exploit multivalency effects, or several different motifs to direct the dyes to different targets.¹⁶ In this section we discuss the nucleophilic substitution of various bio-motifs on fluorinated porphyrinoids for medical applications.

2.1. Porphyrins

Marzilli and co-workers reported the synthesis of 5,10,15,20-tetrakis[2,3,5,6-tetrafluoro-4-(2-trimethylammoniummethyl-aminephenyl)]porphyrin via nucleophilic substitution of dimethyl-1,1-ethylenediamine for DNA binding studies.³⁰ Boyle




Figure 3 Thioglycosylated cationic porphyrin, see ref. 75

and co-workers reported a water soluble Por containing three glucose moieties substituted on para positions of perfluorophenyl group and a positively charged N-alkyl-pyridyl moiety via nucleophilic substitution (see Figure 3).⁷⁵ These compounds were then studied on human colorectal adenocarcinoma (HT29) cells for their PDT activity. Comparison of LD90 (concentration required to kill 90% of cells) showed that 5-(N-octadecyl-pyridyl)-10,15,20-tris(4-thioglycosyl-2,3,5,6-tetrafluorophenyl)porphyrin had the best phototoxicity (irradiated with red light) along with low dark toxicity (absence of light).

In 2001, we reported the synthesis of a symmetrical sugar substituted TPPF₂₀ via nucleophilic substitution of thioglucose and thiogalactose to obtain PGlc4 and PGal4 (Figure 4)⁷⁶ in high yield. These compounds were then tested on cancer cell line (human breast cancer cells (MDA-MB-231), and rat fibroblasts (normal (3Y1), partially transformed (3Y1c-Src), and fully transformed (3Y1v-Src))) for their uptake and PDT efficacy.³¹ Among the two conjugates PGlc₄ showed high cell uptake and better PDT efficacy compared to PGal₄. When PGlc₄ was studied for its uptake in normal cells (Rat fibroblast 3Y1) vs. cancer cells (rat fibroblast 3Y1c-Src and 3Y1v-Src) there was higher uptake in 3Y1v-Src cell line, intermediate uptake in 3Y1c-Src cell line and no uptake in 3Y1 cell line. These studies suggested that sugar appended porphyrinoids show selectivity towards cancer cells compared to normal cells.

Hirohara and co-workers extended the study comparing Zn, Pd and PtPGlc₄ with the free base PGlc₄ (Figure 4) for their phototoxicity.⁷⁷ Here, they started with the free base TPPF₂₀ and metallated with Zn, Pd and Pt before carrying out the nucleophilic substitution of acetyl protected thio-glucose in presence of diethylamine (DEA) as base and DMF as solvent at room temperature. The final step was deprotection of acetate groups. Human epithelial (HeLa) cell line was used to test phototoxicity of these compounds. Zn(II) and Pd(II)PGlc₄ showed high phototoxicity due to heavy-atom effect. Surprisingly, Pt(II)PGlc₄ showed a low phototoxicity. The same group also studied the effect of substitution patterns on the phototoxicity of glycosylated Pors. They similarly carried out




Figure 4 Metallated and non-metallated S-sugar-porphyrins, see ref. 33, 76, 77.




Figure 5 Mono substituted O-galactosyl porphyrin, see ref. 78

the synthesis of five possible *S*-glucose (mono-, 5,10-di-, 5,15-di-, tris- and tetrakis-) substituted Pors (Figure 4).³³ Cell uptake and phototoxicity studies of these conjugates were carried using HeLa cell line. Among these conjugates, the 5,15-di-substituted glucose Por showed the highest cell uptake and phototoxicity.

Cavaleiro and co-workers reported several glycosylated Pors and studied their significance of antiviral activity against herpes simplex virus types 1 and 2 in Vero cells. They synthesized mono substituted O-galactosylPor (Figure 5)⁷⁸ via nucleophilic substitution of isopropylidene protected O-galactopyranose on TPPF₂₀ using NaH base, followed by deprotection of isopropylidene. This compound showed very good antiviral activity even at lower than maximum noncytotoxic concentration.

In 2006, our group studied the substitution of aminoPEG, ethylenediamine, and lysine groups on to TPPF₂₀ (Figure 6) under microwave irradiation yielding the symmetric tetra substituted Por in quantitative yields.⁷⁹ In this study, we used *N*-methylpyrrolidine (NMP) as solvent without any base. Conventional heating at 60-70° C takes more than 20 hours with six different possible products while microwave reaction took less than 30 minutes with only tetra substituted product. Kral and co-workers reported the synthesis, cell uptake and phototoxicity studies of glycol Pors (Figure 7).⁴⁵ In their study they substituted O-glycols and aminoethanol on the TPPF₂₀. DMF solvent and NaH base were used to substitute the O-glycols. They obtained moderate yields of tetra substituted product as the reaction was carried at low temperatures and hard nucleophiles such as oxygen requires heating for better yields. Then for the substitution of aminoethanol they used dioxane as solvent with no base at 150° C to obtain the tetra amino substituted product with free -OH group in high yields. This shows that soft nucleophiles preferentially substitute the fluorine in presence of harder nucleophiles. The hydroxy terminated Pors exhibited a better intracellular uptake and phototoxicity compared to methoxy terminated derivatives when tested *in vitro* on human promyelocytic leukemia (HL60) and mouse mammary carcinoma (4T1) cell lines and *in vivo* on NuNu mice bearing human breast carcinoma MDA-MB-231.




Figure 6 Symmetric tetra amino substituted porphyrin, see ref. 79




Figure 7 Symmetricmetallated and metal free glycol porphyrins and free base chalcone porphyrin derivatives, see ref. 45, 80, 81.

Boyle and co-workers reported the synthesis of metallated (Ni and Ga) and free base tetra substituted thioPEG conjugates in high yields (Figure 7).⁸⁰ These conjugates were evaluated with a human Caucasian colon adenocarcinoma (Caco2) cell line for their phototoxicity efficacy and found that PEGs containing higher chain length showed high phototoxicity due to better water solubility compared to Por derivative containing shorter PEGs. For the Ga(III)Por derivatives, those with longer PEG chains bearing an -OPh axial ligand showed best phototoxicity results of the nine derivatives studied.

Cavaleiro and co-workers reported the synthesis, cellular uptake, subcellular localization and phototoxicity of mono and tetra substituted O-chalconePor derivatives (Figure 7).⁸¹ In this study, O-chalcone was reacted with TPPF₂₀ in presence of K₂CO₃ as base in DMSO as solvent at 100° C to obtain derivatives in moderate yields. These compounds were tested

Review

Organic & Biomolecular Chemistry

on monkey kidney cell line (COS-7). The tetra substituted derivative showed significant uptake compared to the mono substituted derivative, neither compound was phototoxic, and they localize in the nucleus and/or perinuclear loci. The same group reported the synthesis of porphyrin-quinolone conjugates where they synthesized five Zn(II) complexes (mono-, 5,10-di, 5,15-di, tri, and tetra). O-propargyl groups were appended on to TPPF₂₀ via nucleophilic substitution followed by metallation (Figure 8).³⁶ Then using 1,3-dipolar cycloaddition reaction, they conjugated 6-azidoquinoline on to preformed Zn(II) complexed O-propargylPors to yield five porphyrin-quinolone derivatives in quantitative yields.

Kral and co-workers reported the synthesis of porphyrin- β -cyclodextrin (β -CD) or γ -cyclodextrin (γ -CD) conjugates (Figure 9). They also conducted binding studies of several chemotherapy drugs with cyclodextrin (CD) for versatile drug delivery and *in vitro* and *in vivo* studies.⁸² Binding studies revealed that doxorubicin has good binding affinity towards




Figure 8 Zinc metallated porphyrin-quinolone conjugates, see ref. 36.




Figure 9: Porphyrin-CD conjugates, see ref. 82.

porphyrin- γ -CD **3** and paclitaxel has good binding affinity towards porphyrin- β -CD conjugates **1**, **2** and **4**. Supramolecular carrier-chemotherapy drug complexes were then tested *in vitro* using 4T1 and human chronic myelogenous leukemia K562 cell lines, and *in vivo* using BALB/c mice transplanted with 4T1 cells. These studies reveal that combination therapy (PDT and chemotherapy) can be used as an efficient method to treat cancers.

Cavaleiro and co-workers reported the synthesis and ¹O₂ studies of porphyrin-phosphoramidate conjugates (Figure 10).⁸³ They carried out nucleophilic aromatic substitution of aminoalkyl phosphoramidates (different alkyl chain lengths) onto TPPF₂₀ to obtain mono and tetra substituted porphyrin-phosphoramidate conjugates. Among these conjugates tetra substituted porphyrin-phosphoramidates showed higher ¹O₂ values and among the tetra substituted derivatives, the porphyrin-phosphoramidate with butyl chain (n = 4) showed the highest ¹O₂ value.

Tomé and co-workers reported the synthesis of tetra cationic Pors (Figure 11) and studied their activity against fungal conidia (antifungal activity).⁸⁴ In this study, 4-mercaptopyridine was substituted on TPPF₂₀ using DEA as base at room temperature to yield tetra substituted mercaptopyridylPor. Sulphur, being the softer nucleophile than nitrogen, undergoes nucleophilic substitution at these mild conditions. Additionally, iodomethane and 1-iodopentane were reacted respectively to obtain tetra cationic Por derivatives. Compounds were tested on *Penicilliumchrysogenum* conidia strain. Among these cationic derivatives tetramethylpyridyl cationic Por showed better photodynamic inactivation. Recently, they reported cationic Pors with inverted pyridinium groups (Figure 11)⁸⁵ where they reacted 4-hydroxypyridine with TPPF₂₀. Nitrogen, being a softer nucleophile than oxygen, is observed to undergo substitution and tautomerize to pyridone form. Further reaction with DMSO, cationic methoxypyridylporphyrin was obtained.




Figure 10 Porphyrin-phosphoramidateconjugates, see ref. 83.

Recently, Vicente and co-workers published the synthesis and *in vitro* studies of carboranePor derivatives containing three *p*-carborane and cancer targeting units such as polyamines, glucose, peptide and amino acid. In addition, symmetric Por containing four *p*-carborane units for applications in BNCT and/or PDT (Figure 12) were reported.^{41, 46} Symmetric tetra substituted porphyrin-*p*-carborane was obtained in quantitative yields. The PEG derivative was further conjugated to peptide (YRFA) and amino acid (lysine) via an amide coupling reaction. Phototoxicity, dark toxicity, cell uptake, localization and *in vitro* BBB studies were carried on all these conjugates. None of the conjugates showed dark toxicity when tested on human glioma T98G cell line and the polyamine, spermine, conjugates are most taken up by these cells and the




Figure 11 Tetra cationic pyridyl porphyrins, see ref. 84, 85.




Figure 12 Asymmetrical carborane-porphyrin conjugates for BNCT, see ref. 41, 46.

most phototoxic. Human carcinoma HEP2 cell line studies showed that polyamine conjugates were mainly localized in Golgi, endoplasmic reticulum, and mitochondria while the glucose conjugate was localized in mitochondria and lysosomes. Arginine and peptide (YRFA) localize in Golgi, lysosomes, and mitochondria. Moderate permeability of these compounds was observed when *in vitro* blood brain barrier (BBB) studies on human brain capillary endothelial hCMEC/D3 cells were conducted. The low dark toxicity and good phototoxicity for some of the polyamine conjugates reveal that these compounds could be promising agents for treating cancers with BNCT and PDT combined.

2.2. Chlorins, Isobacteriochlorins and Bacteriochlorins

A way to improve the cancer diagnostics potential and PDT efficacy of porphyrinoids (e.g. treating deep tumor tissues) is to enhance the NIR absorption bands.⁸⁶⁻⁸⁸ Chl, Ibac, and Bac are considered promising PS due to a strong NIR absorption and minimal dark toxicity. Chls are derivatives of Pors with one double bond on the pyrrole missing, Ibac are missing two double bonds on adjacent pyrroles and Bacs are missing two double bonds on opposite pyrroles. CF₂₀, IF₂₀ and BF₂₀ (Figure 1) synthesis from TPPF₂₀ was reported by Cavaleiro and co-workers.¹⁷

Vicente and co-workers utilized the CF₂₀ to substitute 1-mercapto-*o*-carborane (Figure 13).⁶⁸ This compound showed no dark toxicity and localized in cell lysosomes when tested using HEP2 cell line. Fluorescence microscopy studies revealed that this compound is mostly found in the cytoplasmic membrane when tested using B16F1 melanotic melanoma cells. *In vitro* and *in vivo* results suggest that this compound can be used as PDT agent in conjunction with BNCT.⁶⁹ Kalinin and co-workers reported the synthesis of boronated Chls and Pors via regioselective substitution of lithiocarboranes onto TPPF₂₀ and CF₂₀ (Figure 13).⁸⁹




Figure 13: Symmetrical tetra substituted carborane-chlorin conjugates, see ref. 68, 89.

We reported the synthesis of three nonhydrolysable tetrathioglycosylated porphyrinoids (CGlc₄, IGlc₄, and BGlc₄).⁹⁰ Hirohara and co-workers reported the synthesis and *in vivo* studies of S-glucose and S-galactose substituted Chl derivatives (Figure 14).⁹¹ These glucose and galactose chlorin derivatives were synthesized via nucleophilic aromatic substitution of acetyl protected S-glucose or acetyl protected S-galactose onto CF₂₀ followed by deprotection. *In vitro* studies of CGlc₄ and CGal₄ are compared with previously reported PGlc₄ and PGal₄³¹ using HeLa cell line. Both CGlc₄ and PGlc₄ showed continuous uptake up to 24 hours while other two compounds reached plateau after 12 hours.




Figure 14 Pd and free base CGlc₄, free base CGal₄, free base IGlc₄, and free base BGlc₄. See ref. 91, 92, 94.

We reported the photophysical properties and *in vitro* studies of CGlc₄, IGlc₄, and BGlc₄ (Figure 14).⁹² CGlc₄ and IGlc₄ have stronger and redder absorption bands compared to PGlc₄ and increased fluorescence quantum yield by 6- and 12-fold, respectively in phosphate buffered saline (PBS). BGlc₄ has the lowest energy Q-band and is considerably red shifted to 730 nm with nearly 50 fold greater absorbance but the fluorescence quantum yield is similar to PGlc₄. Fluorescence imaging studies of these conjugates using MDA-MB-231 and mouse embryonic fibroblast cells K: Molv NIH 3T3 reveal that these compounds are taken up by cancer cells. These compounds were also studied for two photon fluorescence imaging on Chinese hamster ovary cells and we found that IGlc₄ has good two photon absorption (TPA) between 760 – 880 nm, and the cross section is 24.5 GM at 860 nm.⁹³

Hirohara and co-workers then reported the synthesis of Pd(II) complexes of CGlc₄ and studied *in vitro* photocytotoxicity (Figure 14).⁹⁴ As expected from numerous prior studies,^{95, 96} the presence of the heavy atom Pd(II) enhances the singlet oxygen quantum yield. However, they found that free base CGlc₄ has higher *in vitro* photocytotoxicity compared to the Pd(II) complex using HeLa, metastatic B16-BL6, weakly metastatic B16F1, and metastatic 4T1 cell lines. Then, Morita and co-workers used xenograft tumor models to compare the PDT efficacy of CGlc₄, non-glycosylated conjugate (CF₂₀), and aspartyl chlorin (NPe₆) using several different human cancer cell lines.⁹⁷ CGlc₄ showed better PDT efficacy and suppressed tumor growth with no side effects.⁹⁸

Recently, Bruckner and co-workers reported the nucleophilic substitution reaction of thioPEG with different chain lengths onto meso-tetrakis(pentafluorophenyl) oxazolochlorins to obtain tetra substituted pegylated oxazoloChl.⁹⁹

2.3. Corroles

Cors are porphyrinoids with potential uses as PS. Cors are tetrapyrrolic macrocycles with 18 π conjugated electrons but with one less meso carbon and stabilize +3 metal ions.¹⁰⁰ Cors have larger optical cross sections in the red compared with Pors due to stronger Q absorption bands between 450 nm and 650 nm. Cors can have high ¹O₂ generation quantum efficacy,¹⁰¹ thermal and photochemical stability,¹⁰¹ and can be metabolized faster *in vivo*.^{101, 102} Cors are extensively studied as new generation PS for applications in PDT.^{39, 103}

The first synthesis of fluorinated Cor and water soluble pyridyl substituted fluorinated Cor derivatives was reported by Gross and co-workers.^{100, 104} In 2007, our group explored the substitution chemistry on fluorinated Cor (CorF₁₅) with acetate protected glucose, xylose and thiopyridine.³⁹ When thioglucose and thio xylose were mixed with CorF₁₅ a library of six different products were observed as expected. An attempt to make a library of 18 compounds by mixing CorF₁₅ with thioglucose, thioxylose and thiopyridine resulted in only 17 (Figure 15). From this study, we also observed that due to increase in acidity of Cor compared to Por, the substitution is somewhat slower and NMP solvent is best for substitution of nucleophiles on Cor. The higher acidity is because of three NH pyrroles compared to two in Por. Barata and co-workers reported substitution of protected O-glycosyl moiety on CoF₁₅,¹⁰⁵ resulting in mono substituted product with two positional isomers (Figure 15). *In vitro* uptake and phototoxicity studies of these compounds using human acute T cell leukaemia (Jurkat cells) were carried. There was good cell uptake but low phototoxicity.

Osuka and co-workers reported the synthesis of both mono and tri substituted derivatives of fluorinated Cor via nucleophilic substitution of several amines (primary and secondary) with moderate to high yields (Figure 16).¹⁰⁶ Recently, Schoefberger and co-workers reported both metallated and free base amino acid conjugates of CoF₁₅ (Figure 17).¹⁰⁷ Thiazine ring formation was observed when CoF₁₅ was reacted with cysteine. This is due to the presence of nucleophiles S and N on the amino acid. The reaction was carried using strong base (NaH) and heat in DMSO. Due to these harsh conditions the softer S nucleophile reacts first replacing *p*-fluoro group followed by N reacting at meta position forming a stable six membered thiazine ring.




Figure 15 Combinatorial library of sugars and pyridyl substituted corroles, see ref. 39, 105.




Figure 16 Mono and tri amino substituted corrole derivatives, see ref. 106.




Figure 17 Mono and di substituted amino acid-corrole derivatives, see ref. 107.

2.4. Phthalocyanines

Pcs are chemically stable, and have a strong absorption peak in the NIR region of the optical spectra ($\lambda_{\max} \approx 670 \text{ nm}$, $\epsilon \approx 10^5 \text{ L mol}^{-1} \text{ cm}^{-1}$) where light penetrates deeper into tissues and have good $^1\text{O}_2$ quantum yields.¹⁰⁸⁻¹¹⁰ Due to these properties, extensive research has been conducted on their synthesis and functionalization to tune their photochemical and photophysical properties.¹¹¹ Disadvantages of these porphyrinoids include difficulties in purification, formation of isomeric mixtures, and strong aggregation. To solve these issues, present research focuses on fluorinated analogues.¹⁶ Several bio-motifs containing S, N, and O groups can be easily substituted using click substitution chemistry to obtain pure Pcs with better water solubility. Our group was the first to report the synthesis of a zinc phthalocyanine (ZnPc) appended with eight nonhydrolysable thioglucose units obtained via nucleophilic substitution of acetyl protected thioglucose onto commercially available ZnPcF₁₆ (Figure 18).¹¹² Fluorescence microscopy of this compound on MDA-MB-231 cell line showed a very good uptake but the compound aggregated resulting in quenching of fluorescence. Slow disaggregation of the glycolPc inside the cell or tissue may allow these compounds to be used diagnostically or therapeutically.⁶⁷

Tomé and co-workers reported the synthesis and photophysical properties of glycodendritic conjugates of Pors and Pcs bearing eight and sixteen D-galactopyranose units on triazine linkers, respectively (Figure 18).¹¹³ When $^1\text{O}_2$ studies were conducted, high quantum yields were observed, suggesting the potential as PDT agents. Recently, *in vitro* studies of hexadeca-galactophthalocyanine dendrimer (Figure 18) were carried on two different human bladder cancer cell lines, UM-UC-3 and HT-1376, for their PDT efficacy.¹¹⁴ They observed no dark toxicity, high cell uptake, and high phototoxicity. These results were attributed due to the presence of galectin-1 and GLUT1 receptors on the two aforementioned bladder cell lines. The same group synthesized the cationic octa substituted thiopyridyl phthalocyanine (Figure 18) via nucleophilic substitution of

thiopyridine on ZnPcF₁₆ followed by methylation. The product was obtained in quantitative yields.¹¹⁵ Antiviral PDT activity was studied on this compound but did not show promising results.




Figure 18 Symmetrical octa and hexadecaglycosyl-ZnPc, see ref. 112, 113, 115.

Recently, Tomé and co-workers reported Pc-CD conjugates (Figure 19) for cancer PDT.¹¹⁶ Three such derivatives Pc- α -CD, Pc- β -CD and Pc- γ -CD were synthesized via nucleophilic substitution of cyclo-maltohexaose (α -CD), cyclo-maltoheptaose (β -CD) and cyclo-maltooctaose (γ -CD) on two fluorine atoms of PcF₁₆. Then, they tested these compounds on UM-UC-3 bladder cancer cell line for phototoxicity and results revealed that Pc- α -CD and Pc- γ -CD exhibited higher phototoxicity compared to Pc- β -CD.




Figure 19 Amphiphilic cyclodextrin-Pc conjugates, see ref. 116.

2.5. Triply bridged fused porphyrin

Current FDA approved PS have one-photon absorption peaks in the visible region (400nm to 650 nm).⁵⁴ Dyes that have stronger absorption cross section in the NIR region such as Chl, Bac, IBac, Pc, and Por dimers were also prepared^{90, 112, 117} for treatment of deeper seated tumors.¹¹⁸ Recently, research has been carried on two-photon absorption (TPA) dyes that absorb

two lower energy photons simultaneously to arrive at the same excited state as single photon absorptions. These TPA dyes have absorption bands in the NIR or infrared region (800-1100 nm) so scattering of these wavelengths in tissues are much less than higher energy photons.

Osuka and co-workers report that hexakis(pentafluorophenyl) substituted meso-meso-linked Zn(II) diporphyrin has little fluorescence, but a good TPA, and good ¹O₂ quantum yield.¹¹⁹ However, this compound is neither soluble in aqueous media nor specific towards diseased tissues for medical applications. Our group recently reported the synthesis and phototoxicity of the hexathioglucose derivative of the triply bridged fused Por dimer (Figure 20).⁴⁷ This compound showed good ¹O₂ quantum yield (0.78 \pm 0.03) in DMSO. *In vitro* phototoxicity studies carried on MDA-MB-231 breast cancer cell line showed significant cell death (IC₅₀ = 13 μ M) in agreement with ¹O₂ results when white light was exposed for 20 min, and after 24 hours 75% of cells were necrotic. No dark-toxicity was observed.




Figure 20 Hexathioglucose triply bridged fused porphyrin, see ref. 47.

3. SUBSTITUTED FLUORINATED PORPHYRINOIDS FOR APPLICATIONS IN MATERIALS

The tremendous potential applications of Por, Pc and Cor derivatives come from their photophysical and electrochemical properties, remarkable stability, and ability to chelate to nearly every metal to further diversify the chemical properties. These applications include catalysis,^{52, 120-125} solar energy harvesting dyes,^{10, 42, 48, 49, 51, 126} sensors,^{3, 35, 71, 127-130} and tracers.³⁸ The multifunctional properties are modulated by appending various chemical moieties on the periphery of the macrocycles using nucleophilic aromatic substitution chemistry and by varying core metal. As photonic materials, the exocyclic motifs can be used to dictate the supramolecular organization of the dyes, which determines the efficiency of energy or electron transfer, determines the hierarchical order of films, and affords opportunities for molecular recognition of

analytes. While metalloporphyrins can be good catalysts and excellent sensors, Cors have been employed as environmental tracers. Porphyrinoid based solar energy harvesting devices have also been extensively studied.

3.1. Catalysis

Over the last few decades, many research groups have reported that synthetic metalloporphyrins can be efficient and selective catalysts for various oxidation and reduction reactions.^{120, 131-134} Pors with Fe, Mn, Co, and other metal ions have been used as catalysts in laboratory-scale reactions using a variety of oxygen sources.¹³⁵ Such catalytic reactions can mimic the activity of cytochrome P450 monooxygenation reactions.⁵² The efficiency and catalytic activity of metalloporphyrins depends on a few variables; temperature, pressure, solvent, source of oxidant, axial ligand, and the structure of the substrate. These catalysts are observed to be effective in homogeneous systems. Hydrocarbon epoxidation and hydroxylation are typical products of these reactions.¹²⁰ A variety of oxygen donors from molecular oxygen to peroxides, peracids, and iodosylbenzene are used for these reactions.¹³⁶ Due to ease, availability, cost and greener approach, the molecular oxygen in air is preferable as oxygen donor because no energy is used to manufacture and store it.

However, there are a few drawbacks to these catalysts as secondary reactions deactivate the catalytic species, often leading to low turnover number (TON). Self-oxidative degradation during the catalytic process is a key issue that limits use of metalloporphyrin as catalysts. To overcome this, electron withdrawing groups such as fluorines or bulky substituents are often appended to the Por macrocycle.^{137, 138} The fluorinated porphyrinoid platforms allow facile substitution chemistry to append exocyclic moieties that modulate catalysis. Supramolecular Por catalysts are designed to prevent self-oxidative degradation and to enhance substrate selectivity for quicker catalysis and better TON.^{52, 53} To minimize oxidative damage and to develop metalloporphyrins¹³⁹⁻¹⁴³ with more robust structures, our group prepared aggregated metalloporphyrins into organic nanoparticles (ONP)^{52, 53}.

McKeown and co-workers¹²⁴ have reported that hydroquinone oxidation catalytic activity is enhanced in spiro-linked porphyrinoid polymer network. There was a significant difference seen in comparison to the small molecular mass derivative for H₂O₂ decomposition, cyclohexene oxidation and hydroquinone oxidation. These materials mimic the biological systems in which the rigid spiro-linked network creates space around the active sites which allows the substrate to gain access and reaction to take place. There is definitely a potential for heterogeneous catalysis of network polymers of intrinsic microporosity (PIMs) with catalytic active units incorporated within.

Breslow and co-workers¹²⁵ used the simple click-type chemistry to append β -CD onto the TPPF₂₀ macrocycle (Figure 21). In this case the β -CD behaves as an artificial binding pocket for catalysis while the natural Por core catalytic properties are retained. They report that fluorines on the phenyl rings stabilize the Por against autooxidative destruction. This stability results in a high TON and the appended β -CD provides substrate and regio/stereoselectivity in the hydroxylation of steroid substrates.

George and co-workers¹²² reported Por derivatized with fluorinated 'ponytails' as biphasic catalysts (Figure 21) to achieve photo-oxidations in supercritical carbon dioxide with ¹O₂. This study revealed that the fluorinated Por photocatalyst can be recycled 10 times. This is a dramatic increase in catalyst stability compared to previous studies.

Nakagaki and co-workers¹²¹ have reported ethylene glycol substitution onto the core TPPF₂₀ platform (Figure 21) and found that *p*-fluorines on the meso-aryl groups substitute preferentially. After metallation of the free-base macrocycles with iron and manganese, these materials were immobilized on layered double hydroxide (LDH) and silica supports. These materials are found to be catalytically active in the oxidation of (Z)-cyclooctene and cyclohexane by iodosylbenzene. The stability of the materials allowed their reuse for at least three additional cycles of (Z)-cyclooctene oxidation. Earlier this year, the same group¹²⁰ published substituting galactodendritic moieties onto the core TPPF₂₀ platform (Figure 21). The supported Cu(II) and Zn(II) complexes are catalytically active towards oxidation of alkenes, alcohols, and catechol and can be reused.

Our group^{52, 53} has published a green chemistry approach to develop ONP from substituted metalloporphyrins for oxidative catalysis of olefins and alkenes. These 10-30 nm ONP are easily formed by injecting the metalloporphyrin in a host solvent such as DMSO or DMF into water while vigorously mixing. Since water is the primary solvent (ca. 95%) and the ONP activate molecular oxygen in air, these catalysts avoid halogenated solvents or toxic oxygen sources such as iodosylbenzene. The energy requirements are minimal because the reactions are run at ambient temperature and pressure. We appended highly fluorinated alkyl groups on the para positions of the core TPPF₂₀ (Figure 21). Oxidative catalysis studies on cyclohexene with various Fe(III) and Mn(III) metalloporphyrins show that the rate of oxidation by the ONP is much slower than the solvated compounds, but the TONs are 10-50-fold greater. An onion peel mechanism was proposed to explain the slow reactivity and increased TON for these ONP, where only the outer layer of Por are available for the oxidative process, and when these are oxidatively compromised, they exfoliate to expose the next catalytically active layer.

Superoxide (SO) anion radical has dramatic damaging effects on living organisms. The main biological defence system to




Figure 21 Tetra substituted metallated and non-metallated porphyrin thio and oxy derivatives, see ref. 52, 53, 120, 121, 122, 124, 125.

mitigate SO are superoxide dismutase (SOD) enzymes.^{144, 145} SODs are enzymes that catalyse the dismutation of SO to hydrogen peroxide and oxygen very efficiently. SOD contains a redox-active metal with the capability of oxidizing and reducing SO. The endogenous SOD may not be sufficient to remove the highly produced SO under some therapeutic conditions, but direct injection of these enzymes as pharmaceutical agents have problems in terms of stability, immunogenicity and bioavailability.¹³⁴ These problems invite research to develop stable and non-toxic low molecular weight SOD mimics for therapeutic applications.

Gross and co-workers¹³⁴ reported that metal complexes such as Mn(III) or Fe(III) Cors^{146, 147} (Figure 22) catalyze the dismutation of SO. Fe(III) and Mn(III) complexes of the 3,17-bis-sulfonated Cor have been found to be excellent catalysts for the decomposition of peroxyntirite.¹³⁴ The *p*-fluorines have been substituted with methoxy groups to enhance the stability and solubility of these Cors.¹³⁴ The iron Cor derivative is found to be more reactive than the manganese complex. The reactions of metalloporphyrins start with reduction of SO rather than oxidation.

3.2. Sensors

The last 40 years witnessed an explosive growth in developing sensors and biosensors as portable and economical tools in analytical chemistry. The strong colours, fluorescence, electrochemical activity, stability, and ligation properties of fluorinated porphyrinoids enable them to be the functional heart of sensors.¹³⁰ The fluorinated porphyrinoid platforms facilitate substitution chemistry to append ligands of choice

onto the macrocycle to recognize and report on a diverse array of analytes.




Figure 22 Tri methoxy-substituted bis-sulfonated corroles. See ref 134.

Anions play crucial roles in a number of chemical, biological and environmental processes.^{148, 149} Diseases such as osteoporosis, Alzheimer's, and cystic fibrosis are caused by perturbations in anion fluxes across cell membranes.¹⁵⁰⁻¹⁵² Anions including cyanide and arsenate are considered a threat when present in the environment which requires a need for systems capable of controlling and monitoring these species.¹⁵³ MetalloPors with metals such as zinc, rhodium, silver, gold or copper can be used as cyanide sensors because of the essential affinity of cyanide to these metals.¹³⁰

The environmental and biological importance of anions accounts for a critical need for a sensor design to effectively detect these species. Amines and polyamines,¹⁵⁴ calixpyrroles,¹⁵⁵ subphthalocyanines,¹⁵⁶ Cors,¹⁵⁷ and Pors¹⁵⁸ are used to create anion receptors. Anion recognition involves the NH groups of the pyrrole units in Pors. However, with the discovery of substitution chemistry on these macrocycles, recognition can take place outside the macrocycle. Due to sterics, the inner NH groups are less effective hydrogen bond donors compared to the substituents on the periphery of the macrocycle. For instance, several Pors bearing NH groups at the ortho positions of meso-aryl substituents have been demonstrated as being effective anion sensors.¹⁵⁸ Pors bearing substituents at the para position of the meso-aryl groups can also be used for anion recognition. TPPF₂₀ can be used as a platform to build new polyamine anion hosts that are effective receptors for dihydrogen phosphate in organic media.¹⁵² Tomé and co-workers¹⁵² have substituted various diamines onto the core TPPF₂₀ platform for development of phosphate anion sensors (Figure 23). These sensors are reported to interact preferentially with phosphate anions both in organic solutions and in basic aqueous media when coated on gold piezoelectric quartz crystals.

Recently, Bruckner and co-workers¹³⁰ showed pyrrole-modified Pors bearing a lactone functional group at the macrocycle periphery as optical cyanide sensors. Nucleophilic attack of the cyanide onto the lactone moiety is proved to be

the sensing mechanism. A change in the hybridization of the carbonyl carbon couples electronically to the macrocycle and results in a 100 nm red-shifted absorbance band and the Soret band decreases in intensity upon cyanide addition. Substitution of the *p*-fluorine on the pentafluorophenyl moiety with PEG groups leads to water-soluble derivatives allowing for cyanide sensing in purely aqueous solutions.

Dissolved oxygen (DO) is related to water quality, medicine, food freshness, respiration and metabolism in living organisms, and many other biological applications.¹⁵⁹⁻¹⁶¹ Mechanism such as fluorescence quenching, oxygen pressure and electrochemistry are used to develop oxygen sensors.^{35, 160-164} Pt(II) and Pd(II) Pors^{165, 166} are typical chemical materials acting as fluorescence-based oxygen sensors because of their long triplet state, variable lifetimes from a few to tens of microseconds and respectable triplet-triplet energy transfer from these compounds to oxygen molecules. When this energy transfer takes place, the emission intensities and the lifetimes of these sensors is shortened. Pt(II)TPPF₂₀ is an excellent sensor because of its good response to oxygen concentrations and high photostability compared to other chromophores.^{167, 168} Meldrum and co-workers³⁵ reported the synthesis of a new sensor for DO. Simple click-type substitution chemistry was used on the Pt(II)TPPF₂₀ platform. The resulting sensor carries four cross-linkable methylacrylate functional moieties (Figure 23) which can be polymerized and cross-linked with other monomers for polymer sensing film. The sensor was then linked to hydrophilic poly(2-hydroxyethylmethacrylate)-copolyacrylamide (PHEMA) and hydrophobic polystyrene (PSt) films. The long fluorescence life time and sufficient triplet-triplet energy transfer between the porphyrinic based sensors and oxygen molecules results in a decrease of the oxygen sensors' emission intensities and a shortening of their lifetimes.




Figure 23 Symmetric metallated and free base oxy and amino substituted porphyrin for applications as sensors, see ref. 35, 38, 128, 152.

Recently, Pcs, coumarinderivatives,¹⁶⁹ and naphthalocyanines¹⁷⁰ have been used as fluorescent markers for petroleum fuel derivatives. These compounds are not visible with naked eye under visible light. However, because of the large optical cross sections and fluorescent properties, they can be easily detected by UV light. In this case, some characteristics of efficient fluorescent markers include 1) good fuel solubility; 2) ambient light invisibility; 3) prolonged shelf-life and stability; 4) fluorescence emissions in the regions other than those of the fuel itself; 5) cost effective and efficient synthesis at large scale.^{38, 171} Porphyrinoids, Pors in particular, are excellent sensor dyes that possess intrinsic photophysical properties that match the requirements for a fuel labelling marker. The desired characteristics for a good fuel labelling system are obtained by substituting non-polar moieties on the commercially available TPPF₂₀ platform resulting "invisible" fluorescent biodiesel markers.³⁸ Serra and co-workers³⁸ have reported the substitution of 1-hexadecyloxy- and ethoxy-substituents onto the *p*-fluoro positions of TPPF₂₀ (Figure 23). The developed sensor dyes are soluble in both hydrophobic and hydrophilic environments. The compounds were found to be stable in biodiesel environment for up to three months. Because of the high stability, solubility and ease of preparation, these markers are reported as efficient and cost effective.

Cosnier and co-workers¹²⁸ have shown the substitution of *p*-fluoro groups on zinc and manganese TPPF₂₀ macrocycles with electropolymerizable pyrrole groups (Figure 23). The electro-oxidation of these metalloporphyrins with functionalized pyrrole groups leads to electrogeneration of polypyrrole films and the electrochemical behaviour of the Pors is retained. The catalytic properties of these materials are associated with the advantages conferred by the electrochemically formed polymers and allow use as a new oxidase based biosensor involving a reductive detection of H₂O₂.

3.3. Solar

Porphyrinoid molecules have great potential applications because of their light absorbing properties. Their stability in harsh environments makes them suitable for applications such as solar energy harvesting. In addition, many of these dyes are low-cost, which makes them suitable for photovoltaic (PV) devices. Pc organic dyes have energy band gaps that are well-matched to the incident solar spectrum.⁹ However, use of Pcs in low-cost solar cells faces complications because of poor solubility in organic solvents and narrow absorption bandwidths in the red (Q-band) and ultraviolet (B-band) wavelengths.⁹ Pcs have been used in bulk heterojunction (BHJ) solar cells.^{49, 126, 172, 173} Some solar cell designs contain zinc and copper Pc complexes and also contain various C₆₀ derivatives, where vapour-deposition techniques are used to make specified layers.¹²⁶ Soluble dyes have also been used in layered devices or BHJ solar cells.^{126, 174} The commercially available fluorinated Pc platforms allow for click-type alkylation

chemistry and design of a series of robust, chemically compatible dyes with tuneable optical band gaps and energy levels.^{42, 48, 49, 126} The resulting dyes absorb a different portion of the solar spectrum in these devices. The semiconductor active layer is composed of a blend of Pc derivatives having different optical band gaps in order to capture a larger fraction of the solar spectrum. Substitution of peripheral fluoro groups on hexadecafluorophthalocyaninatozinc(II) (ZnPcF₁₆) by thioalkanes allows for better solubility. The frontier molecular orbitals (HOMO and LUMO) are primarily delocalized on the ring periphery. Substitution of electron-withdrawing groups with electron-donating groups outside the macrocycle affects the orbital energy levels.¹²⁶ The destabilization of the HOMO is more than the LUMO which results in decreased optical band gaps and the corresponding absorption wavelengths shifts toward longer wavelengths in the NIR.^{48, 49, 126} This substitution chemistry is used to cover a different and larger portion of the solar spectrum.

We have reported the substitution of thioalkyl substituents onto the ZnPcF₁₆ platform for fine tuning the absorbance of this dye (Figure 24).¹²⁶ The number of thioalkane versus fluorine substituents directly dictates the red-shift in the absorbance spectrum, especially of the Q-bands. There is a ca. 6.5 nm red shift for every fluorine substituted by thioalkane. Furthermore, solar cells made of blends of three dyes result in better efficiency than expected from individual dye performance because of broader coverage of the solar spectrum and synergistic charge migration. We also reported⁴⁹ substituting the core ZnPcF₁₆ platform with diverse thioalkanes to control the morphology and the packing of dyes in BHJ devices (Figure 24). Charge migration is not hindered because the Pc molecules have substantially overlapping HOMO-LUMO gaps. Similarly, from ZnPcF₁₆ we have synthesized⁴⁸ a dye bearing long alkane moieties on all 16 positions (Figure 24). This promotes self-organization into nanoparticles and nano films on indium tin oxide (ITO) and glass surfaces. The particles




Figure 24 Thio substituted phthalocyanines, see ref. 48, 49, 126.

are highly monodispersed and changing the concentration of the chromophores in solution can control their size upon surface deposition. We report substitution of some thioalkyl groups but any primary thiolated moiety can be appended to investigate the role of exocyclic motifs on the properties of nanomaterials.

Por is also robust enough to be incorporated in light harvesting devices. Plants use various chlorophylls to cover as much of the solar spectrum as possible.¹⁷⁵ Natural dyes are not robust enough to incorporate into devices because they are typically not stable for a prolonged times, such as the span of years for a solar energy device. A useful solar cell would be expected to have 15 to 20 years of life. The TPPF₂₀ and Pcf₁₆ may be able to withstand these conditions, so are considered good candidates for light harvesting. The maximum absorption of Por is a B (Soret) band at 390–430 nm with high molar coefficients and several Q-bands between 500 and 650 nm. Thus, TPPF₂₀ and its derivatives can be used to cover the blue end of the solar spectrum, while derivatives of Pcf₁₆ cover the red end. These macrocycles combined have good optical densities in the middle of solar spectrum and afford a range of dyes to incorporate into various devices.

We have done studies⁴² using the TPPF₂₀ platform to put fluoroalkanes on the macrocycle to investigate the self-organization of the chromophore with C₆₀ via a co-deposition process into thin-films (Figure 25). We observed a significant quenching of the Por fluorescence because of electron transfer to the C₆₀ electron acceptor. Kral and co-workers¹⁷⁶ have also reported functionalizing the Por macrocycle with highly fluoruous groups (Figure 25). The authors suggest that the compounds have potential in drug delivery, transport of gases, and solar cells if morphological packing is appropriate.

For a PV device to work efficiently and properly there has to be a continuous conductive medium between the light absorbing dye and the electron acceptor to transfer the electron and create a charge separation. Our group has recently developed a Por dimer linked by a flexible dithiol tether (Figure 26).⁵¹ Upon mixing the dimer with C₆₀ and C₇₀ we discovered that it preferentially binds with fullerene C₇₀ over C₆₀ in toluene solution. The dimer-fullerene complex also forms stable aggregates when casted on glass. This type of binding studies can open opportunities for future research where the electron acceptor is kept in close proximity to the dye for a more efficient charge transfer without modification of the C₆₀, which is usually detrimental to the system.

We have shown^{50, 177} that click chemistry can be used to append thioalkyl tethers which can be attached to Au surface for rapid assessment of chain length and dye association (Figure 27). Free base and zinc Por complexes were studied and both were found to bind near the edges of alkane thiol layer. Zn complexes were observed to form small aggregates held together via π - π stacking or pyridyl-Zn interactions.




Figure 25 Highly fluorinated symmetric porphyrin, see ref. 42, 176.

Among these two the free base acts as a stochastic switch and the Zn complex aggregates show evidence of quantum cascade. The effective overlap of frontier orbitals favours charge transfer in the Zn complex while minimal charge transfer was observed for the free base Por.




Figure 26 Zn porphyrin dimer connected via thio alkyl tether, see ref. 51.




Figure 27 A. Porphyrin molecule with thiol tether (optimized structure from DFT calculations) (Zn or H₂ is grey, carbon is aqua blue, hydrogen is white, nitrogen is blue, fluorine is pink, and sulphur is yellow). B. Scheme showing geometry of thioalkyl substituted porphyrin inserting between alkanethiols. see ref. 50, 177. Reproduced with permission from ref 50. Copyright 2015 American Chemical Society

4. CONCLUSIONS AND FUTURE PERSPECTIVE

One major issue with the synthesis of functional groups covalently bound to porphyrinoids for applications in medicine and materials is the cost and ease of production at multi-gram scales. To bring the synthetic costs in ranges for potential commercialization, straight forward and high yield synthetic organic strategies are needed for the construction of substituted porphyrinoid conjugates. Thus, the fluorinated porphyrinoids (TPPF₂₀, PcF₁₆, CF₂₀, IF₂₀, BF₂₀ and CorF₁₅) are used as platforms to obtain an array of porphyrinoid derivatives via straight forward nucleophilic aromatic substitution. Compared to non-fluorinated porphyrinoid derivatives, fluorinated ones show an equally diverse range of physical and chemical properties. The presence of fluorines also improves the stability of these molecules to oxidative damage and brings further functionality. Various nucleophiles such as glycol, sugars, polyamines, peptides, amino acids, and carboranes were substituted on fluorinated porphyrinoids in order to improve biological efficacy, solar energy absorbance, molecular packing in thin films, catalytic activity and stability. Another advantage of this strategy is that various motifs such as disease targeting and for therapy can be readily tagged onto a fluorinated porphyrinoid. Lately, fused fluorinated porphyrinoids with NIR absorption and/or TPA have been investigated and showed potential as PS. These fluorinated porphyrinoid platforms continue to be exploited in designing drugs that can effectively target and treat diseases such as cancers, be the signal transducer in sensors, as catalysts, and to develop photonic films for solar cells. For the ZnPcF₁₆, exchange of the fluorine group for an thioalkane leads to a

Review

systematic red-shift of the absorbance spectrum, thereby creating a family of closely related dyes with substantial alignment of the HOMO-LUMO gaps, that then facilitate charge transport in materials composed of these dyes. Since the aryl groups on TPPF₂₀ are orthogonal to the macrocycle, substitution of the *p*-fluorine has only small effects on the photophysics. The ability to chelate to most metals allows for creation of molecular catalysts and sensors.

Many biomolecules are robust under the conditions used for the nucleophilic substitution on these porphyrinoids, but some proteins and peptides may not. To append sensitive biomolecules, e.g. antibodies, a tether can be appended on to the preformed fluorinated porphyrinoid followed by conjugation of the sensitive biomolecule. The remaining fluorine groups present on these porphyrinoids after substitution (16 remaining on the TPPF₂₀, CF₂₀, IF₂₀, BF₂₀ platforms; 12 remaining on the CorF₂₀ platform; and 1-15 remaining on the P_{CF}₁₆ platform) can be used as adjunct imaging modalities, e.g. ¹⁹F NMR imaging, carbon-fluorine spectroscopy (CFR), and fluoro-Raman spectroscopy (FRS). Fluorines can be substituted with radio isotopes such as ¹⁸F (*t*_{1/2} = 110 mins), ¹²⁵I (*t*_{1/2} = 13.22 hours), ¹²³I (*t*_{1/2} = 8 days) via reactions of fluorinated porphyrinoids with K¹⁸F/Na¹³¹I/Na¹²³I and crown ethers. The K⁺ and Na⁺ chelator, kryptofix, may allow F-18, I-123 and I-131 labelling at elevated temperatures. ¹⁸F porphyrinoids can be used as PET imaging agents without altering the photophysics for fluorescence imaging, and ¹²³I porphyrinoids as single photon emission computed tomography (SPECT) imaging agents. ¹³¹I can be used as radiotherapy and imaging agent to treat cancers effectively as adjuvant to PDT. The biomedical applications of fluorinated porphyrinoids has been reviewed.¹⁶

Por molecules have a propensity to bind with electron acceptors such as C₆₀ and C₇₀. This opens a new window of research for further development of these materials where the electron acceptor is trapped in the complex allowing for a better connection and charge transfer. These complexes may result in more efficient solar cells and devices. Various thioalkane nucleophiles have been substituted onto these core platforms, wherein the alkanes dictate nanostructure morphologies.^{52, 181} The mechanism of nanoparticle (NP) formation, the size of the NPs, the stability, and the organization of the Pors in the NPs exquisitely depends on the exact structure of the Por and the methods to make the NPs.¹⁸¹

Recent advances in porphyrin-based metal-organic-framework (MOF) materials for catalysts, gas storage, and sensors builds upon earlier work in the self-assembly of porphyrins by metal ion coordination.^{10, 182, 183} Clicking metal ion coordination groups to these core platforms may facilitate the discovery and development of new MOF and hierarchically structured porphyrinoid materials. Hierarchical structure is of paramount importance in photonic materials.¹⁸⁴ As illustrated by the work described herein, these six and other fluorinated porphyrinoid

platforms enable the rapid, facile discovery of new theranostic, sensor, and photonic materials without complex, multistep reactions and tedious purifications.

Acknowledgements

Supported by the National Science Foundation – United States (NSF) through CHE-0847997 and CHE-1213962 to C.M.D. Hunter College science infrastructure is supported by the NSF, the City University of New York and the National Institute on Minority Health and Health Disparities of the National Institutes of Health under award number G12MD007599.

Notes and references

1. in *The Porphyrin Handbook-Applications: Past, Present and Future*, eds. K. Kadish, K. M. Smith and R. Guilard, Academic Press, New York, 2000, vol. 6.
2. D. M. Guldi, *Chem. Soc. Rev.*, 2002, **31**, 22-36.
3. M. I. J. Stich, L. H. Fischer and O. S. Wolfbeis, *Chem. Soc. Rev.*, 2010, **39**, 3102-3114.
4. M. O. Senge, M. Fazekas, E. G. A. Notaras, W. J. Blau, M. Zawadzka, O. B. Locos and E. M. Ni Mhuirheartaigh, *Adv. Mater.*, 2007, **19**, 2737-2774.
5. C. M. Drain, K. C. Russell and J.-M. Lehn, *Chem. Commun.*, 1996, 337-338.
6. M. G. Walter, A. B. Rudine and C. C. Wamser, *J. Porphyrins Phthalocyanines*, 2010, **14**, 759-792.
7. C.-M. Che, V. K.-Y. Lo, C.-Y. Zhou and J.-S. Huang, *Chem. Soc. Rev.*, 2011, **40**, 1950-1975.
8. S. Nia, X. Gong, C. M. Drain, M. Jurow, W. Rizvi and M. Qureshy, *J. Porphyrins Phthalocyanines*, 2010, **14**, 621-629.
9. I. Radivojevic, A. Varotto, C. Farley and C. M. Drain, *Energy Environ. Sci.*, 2010, **3**, 1897-1909.
10. C. M. Drain, A. Varotto and I. Radivojevic, *Chem. Rev.*, 2009, **109**, 1630-1658.
11. K. Davia, D. King, Y. Hong and S. Swavey, *Inorg. Chem. Commun.*, 2008, **11**, 584-586.
12. Y.-J. Ko, K.-J. Yun, M.-S. Kang, J. Park, K.-T. Lee, S. B. Park and J.-H. Shin, *Bioorg. Med. Chem. Lett.*, 2007, **17**, 2789-2794.
13. F. M. D. Ismail, *J. Fluorine Chem.*, 2002, **118**, 27-33.
14. D. O'Hagan and H. S. Rzepa, *Chem. Commun.*, 1997, 645-652.
15. D. Seebach, *Angew. Chem., Int. Ed. Engl.*, 1990, **29**, 1320-1367.
16. T. Goslinski and J. Piskorz, *J. Photochem. Photobiol., C*, 2011, **12**, 304-321.
17. A. M. G. Silva, A. C. Tomé, M. G. P. M. S. Neves, A. M. S. Silva and J. A. S. Cavaleiro, *J. Org. Chem.*, 2005, **70**, 2306-2314.
18. A. M. G. Silva, A. C. Tome, M. G. P. M. S. Neves, A. M. S. Silva and J. A. S. Cavaleiro, *Chem. Commun.*, 1999, 1767-1768.
19. A. M. G. Silva, A. C. Tomé, M. G. P. M. S. Neves and J. A. S. Cavaleiro, *Tetrahedron Lett.*, 2000, **41**, 3065-3068.

20. K. C. Partlow, J. Chen, J. A. Brant, A. M. Neubauer, T. E. Meyerrose, M. H. Creer, J. A. Nolta, S. D. Caruthers, G. M. Lanza and S. A. Wickline, *FASEB J.*, 2007, **21**, 1647-1654.
21. R. Murugesan, S. English, K. Reijnders, K.-i. Yamada, J. A. Cook, J. B. Mitchell, S. Subramanian and M. C. Krishna, *Magn. Reson. Med.*, 2002, **48**, 523-529.
22. H. R. A. Golf, H.-U. Reissig and A. Wiehe, *Eur. J. Org. Chem.*, 2015, **2015**, 1548-1568.
23. A. L. Allred, *J. Inorg. Nucl. Chem.*, 1961, **17**, 215-221.
24. Q. Tang, H. Li, Y. Liu and W. Hu, *J. Am. Chem. Soc.*, 2006, **128**, 14634-14639.
25. J. W. Buchler, in *The Porphyrins*, ed. D. Dolphin, Academic: New York, 1978, vol. **1**, pp. 389-483.
26. K. M. Kadish, B. C. Han, M. M. Franzen and C. Araullo-McAdams, *J. Am. Chem. Soc.*, 1990, **112**, 8364-8368.
27. P. Battioni, O. Brigaud, H. Desvaux, D. Mansuy and T. G. Traylor, *Tetrahedron Lett.*, 1991, **32**, 2893-2896.
28. W. W. Kalisch and M. O. Senge, *Angew. Chem.*, 1998, **110**, 1156-1159.
29. M. O. Senge, *Acc. Chem. Res.*, 2005, **38**, 733-743.
30. J. E. McClure, L. Baudouin, D. Mansuy and L. G. Marzilli, *Biopolymers*, 1997, **42**, 203-217.
31. X. Chen, L. Hui, D. A. Foster and C. M. Drain, *Biochemistry*, 2004, **43**, 10918-10929.
32. S. J. Shaw, K. J. Elgie, C. Edwards and R. W. Boyle, *Tetrahedron Lett.*, 1999, **40**, 1595-1596.
33. S. Hirohara, M. Nishida, K. Sharyo, M. Obata, T. Ando and M. Tanihara, *Bioorg. Med. Chem.*, 2010, **18**, 1526-1535.
34. S. J. Shaw, C. Edwards and R. W. Boyle, *Tetrahedron Lett.*, 1999, **40**, 7585-7586.
35. Y. Tian, B. R. Shumway and D. R. Meldrum, *Chem. Mater.*, 2010, **22**, 2069-2078.
36. F. d. C. Santos, A. C. Cunha, M. C. B. V. de Souza, A. C. Tomé, M. G. P. M. S. Neves, V. F. Ferreira and J. A. S. Cavaleiro, *Tetrahedron Lett.*, 2008, **49**, 7268-7270.
37. N. B. McKeown, S. Hanif, K. Msayib, C. E. Tattershall and P. M. Budd, *Chem. Commun.*, 2002, 2782-2783.
38. A. C. B. Figueira, K. T. de Oliveira and O. A. Serra, *Dyes Pigm.*, 2011, **91**, 383-388.
39. D. Samaroo, M. Vinodu, X. Chen and C. M. Drain, *J. Comb. Chem.*, 2007, **9**, 998-1011.
40. J. Kvičala, M. Beneš, O. Paleta and V. Král, *J. Fluorine Chem.*, 2010, **131**, 1327-1337.
41. N. V. S. D. K. Bhupathiraju, X. Hu, Z. Zhou, F. R. Fronczek, P.-O. Couraud, I. A. Romero, B. Weksler and M. G. H. Vicente, *J. Med. Chem.*, 2014, **57**, 6718-6728.
42. A. Varotto, L. Todaro, M. Vinodu, J. Koehne, G.-y. Liu and C. M. Drain, *Chem. Commun.*, 2008, 4921-4923.
43. C. M. Drain and S. Singh, in *The Handbook of Porphyrin Science with Applications to Chemistry, Physics, Materials Science, Engineering, Biology and Medicine* eds. K. Kadish, K. M. Smith and R. Guilard, World Scientific Publisher, Singapore, 2010, vol. **3**, pp. 485-537.
44. A. F. Mironov and M. A. Grin, *J. Porphyrins Phthalocyanines*, 2008, **12**, 1163-1172.
45. J. Králová, T. Bříza, I. Moserová, B. Dolenský, P. Vašek, P. Poučková, Z. Kejík, R. Kaplánek, P. Martásek, M. Dvořák and V. Král, *J. Med. Chem.*, 2008, **51**, 5964-5973.
46. N. V. S. D. K. Bhupathiraju and M. G. H. Vicente, *Bioorg. Med. Chem.*, 2013, **21**, 485-495.
47. S. Singh, A. Aggarwal, N. V. S. D. K. Bhupathiraju, B. Newton, A. Nafees, R. Gao and C. M. Drain, *Tetrahedron Lett.*, 2014, **55**, 6311-6314.
48. M. Jurow, A. Varotto, V. Manichev, N. A. Travlou, D. A. Giannakoudakis and C. M. Drain, *RSC Adv.*, 2013, **3**, 21360-21364.
49. M. J. Jurow, B. A. Hageman, E. DiMasi, C.-Y. Nam, C. Pabon, C. T. Black and C. M. Drain, *J. Mater. Chem. A*, 2013, **1**, 1557-1565.
50. A. E. Schuckman, B. W. Ewers, L. H. Yu, J. P. C. Tomé, L. M. Pérez, C. M. Drain, J. G. Kushmerick and J. D. Batteas, *J. Phys. Chem. C*, 2015, **119**, 13569-13579.
51. M. Jurow, C. Farley, C. Pabon, B. Hageman, A. Dolor and C. M. Drain, *Chem. Commun.*, 2012, **48**, 4731-4733.
52. A. Aggarwal, S. Singh, J. Samson and C. M. Drain, *Macromol. Rapid Commun.*, 2012, **33**, 1220-1226.
53. A. Aggarwal, S. Singh and C. M. Drain, *J. Porphyrins Phthalocyanines*, 2011, **15**, 1258-1264.
54. M. Ethirajan, Y. Chen, P. Joshi and R. K. Pandey, *Chem. Soc. Rev.*, 2011, **40**, 340-362.
55. T. J. Dougherty, C. J. Gomer, B. W. Henderson, G. Jori, D. Kessel, M. Korbelik, J. Moan and Q. Peng, *J. Natl. Cancer Inst.*, 1998, **90**, 889-905.
56. P. Jichlinski and H.-J. Leisinger, *Urol. Res.*, 2001, **29**, 396-405.
57. D. E. J. G. J. Dolmans, D. Fukumura and R. K. Jain, *Nat. Rev. Cancer*, 2003, **3**, 380-387.
58. M. R. Detty, S. L. Gibson and S. J. Wagner, *Journal of medicinal chemistry*, 2004, **47**, 3897-3915.
59. C. Fritsch, G. Goerz and T. Ruzicka, *Arch. Dermatol.*, 1998, **134**, 207-214.
60. R.-M. Szeimies, M. Landthaler and S. Karrer, *J. Dermatol. Treat.*, 2002, **13**, s13-s18.
61. S. Bonneau and C. Vever-Bizet, *Expert Opin. Ther. Pat.*, 2008, **18**, 1011-1025.
62. D. Phillips, *Photochem. Photobiol. Sci.*, 2010, **9**, 1589-1596.
63. K. Konopka and T. Goslinski, *J. Dent. Res.*, 2007, **86**, 694-707.
64. A. Kubin, F. Wierrani, U. Burner, G. Alth and W. Grunberger, *Curr. Pharm. Des.*, 2005, **11**, 233-253.
65. L. L. Trannoy, J. W. M. Lagerberg, T. M. A. R. Dubbelman, H. J. Schuitmaker and A. Brand, *Transfusion*, 2004, **44**, 1186-1196.
66. J. P. C. Tomé, M. G. P. M. S. Neves, A. C. Tomé, J. A. S. Cavaleiro, M. Soncin, M. Magaraggia, S. Ferro and G. Jori, *J. Med. Chem.*, 2004, **47**, 6649-6652.
67. S. Singh, A. Aggarwal, N. V. S. D. K. Bhupathiraju, G. Arianna, K. Tiwari and C. M. Drain, *Chemical Reviews*, 2015, **115**, 10261-10306.
68. E. Hao, E. Friso, G. Miotto, G. Jori, M. Soncin, C. Fabris, M. Sibrian-Vazquez and M. G. H. Vicente, *Org. Biomol. Chem.*, 2008, **6**, 3732-3740.
69. R. Hiramatsu, S. Kawabata, H. Tanaka, Y. Sakurai, M. Suzuki, K. Ono, S.-I. Miyatake, T. Kuroiwa, E. Hao and M. G. H. Vicente, *J. Pharm. Sci.*, 2015, **104**, 962-970.
70. C. Belle, C. Béguin, S. Hamman and J.-L. Pierre, *Coord. Chem. Rev.*, 2009, **253**, 963-976.
71. M. Malet-Martino, V. Gilard, F. Desmoulin and R. Martino, *Clin. Chim. Acta*, 2006, **366**, 61-73.
72. R. Martino, V. Gilard, F. Desmoulin and M. Malet-Martino, *J. Pharm. Biomed. Anal.*, 2005, **38**, 871-891.

73. O. Couturier, A. Luxen, J.-F. Chatal, J.-P. Vuillez, P. Rigo and R. Hustinx, *Eur. J. Nucl. Med. Mol. Imaging*, 2004, **31**, 1182-1206.
74. F. Mena, B. Mena and O. Sharts, *Faraday Discuss.*, 2011, **149**, 269-278.
75. S. Ahmed, E. Davoust, H. Savoie, A. N. Boa and R. W. Boyle, *Tetrahedron Lett.*, 2004, **45**, 6045-6047.
76. P. Pasetto, X. Chen, C. M. Drain and R. W. Franck, *Chem. Commun.*, 2001, 81-82.
77. S. Hirohara, M. Obata, H. Alitomo, K. Sharyo, T. Ando, S. Yano and M. Tanihara, *Bioconjugate Chem.*, 2009, **20**, 944-952.
78. J. P. C. Tomé, M. G. P. M. S. Neves, A. C. Tomé, J. A. S. Cavaleiro, A. F. Mendonça, I. N. Pegado, R. Duarte and M. L. Valdeira, *Bioorg. Med. Chem.*, 2005, **13**, 3878-3888.
79. D. Samaroo, C. E. Soll, L. J. Todaro and C. M. Drain, *Org. Lett.*, 2006, **8**, 4985-4988.
80. R. E. Mewis, H. Savoie, S. J. Archibald and R. W. Boyle, *Photodiagn. Photodyn. Ther.*, 2009, **6**, 200-206.
81. V. V. Serra, F. Camoes, S. I. Vieira, M. A. F. Faustino, J. P. C. Tomé, D. C. G. A. Pinto, M. G. P. M. S. Neves, A. C. Tomé, A. M. S. Silva, E. F. da Cruz e Silva and J. A. S. Cavaleiro, *Acta Chim. Slov.*, 2009, **56**, 603-611.
82. J. Králová, Z. Kejík, T. Bříza, P. Poučková, A. Král, P. Martásek and V. Král, *J. Med. Chem.*, 2010, **53**, 128-138.
83. L. F. Pedrosa, M. C. de Souza, M. A. F. Faustino, M. G. P. M. S. Neves, A. M. S. Silva, A. C. Tomé, V. F. Ferreira and J. A. S. Cavaleiro, *Aust. J. Chem.*, 2011, **64**, 939-944.
84. M. C. Gomes, S. M. Woranovicz-Barreira, M. A. F. Faustino, R. Fernandes, M. G. P. M. S. Neves, A. C. Tome, N. C. M. Gomes, A. Almeida, J. A. S. Cavaleiro, A. Cunha and J. P. C. Tome, *Photochem. Photobiol. Sci.*, 2011, **10**, 1735-1743.
85. D. C. S. Costa, V. F. Pais, A. M. S. Silva, J. A. S. Cavaleiro, U. Pischel and J. P. C. Tomé, *Tetrahedron Lett.*, 2014, **55**, 4156-4159.
86. E. D. Sternberg, D. Dolphin and C. Brückner, *Tetrahedron*, 1998, **54**, 4151-4202.
87. R. Bonnett, *Chem. Soc. Rev.*, 1995, **24**, 19-33.
88. R. K. Pandey, *J. Porphyrins Phthalocyanines*, 2000, **4**, 368-373.
89. V. Olshevskaya, A. V. Zaitsev, A. L. Sigan, E. G. Kononova, P. V. Petrovskii, N. D. Chkanikov and V. N. Kalinin, *Dokl. Chem.*, 2010, **435**, 334-338.
90. C. M. Drain, S. Singh, D. Samaroo, S. Thompson, M. Vinodu and J. P. C. Tome, *Proc. SPIE*, 2009, **7380**, 73902K-73901-73909.
91. S. Hirohara, M. Obata, H. Alitomo, K. Sharyo, T. Ando, M. Tanihara and S. Yano, *J. Photochem. Photobiol., B*, 2009, **97**, 22-33.
92. S. Singh, A. Aggarwal, S. Thompson, J. P. C. Tomé, X. Zhu, D. Samaroo, M. Vinodu, R. Gao and C. M. Drain, *Bioconjugate Chem.*, 2010, **21**, 2136-2146.
93. A. Aggarwal, S. Thompson, S. Singh, B. Newton, A. Moore, R. Gao, X. Gu, S. Mukherjee and C. M. Drain, *Photochem. Photobiol.*, 2014, **90**, 419-430.
94. S. Hirohara, Y. Kawasaki, R. Funasako, N. Yasui, M. Totani, H. Alitomo, J. Yuasa, T. Kawai, C. Oka, M. Kawaichi, M. Obata and M. Tanihara, *Bioconjugate Chem.*, 2012, **23**, 1881-1890.
95. E. G. Azenha, A. C. Serra, M. Pineiro, M. M. Pereira, J. Seixas de Melo, L. G. Arnaut, S. J. Formosinho and A. M. d. A. Rocha Gonsalves, *Chem. Phys.*, 2002, **280**, 177-190.
96. A. Gorman, J. Killoran, C. O'Shea, T. Kenna, W. M. Gallagher and D. F. O'Shea, *J. Am. Chem. Soc.*, 2004, **126**, 10619-10631.
97. S. Sakuma, E. Otake, K. Torii, M. Nakamura, A. Maeda, R. Tujii, H. Akashi, H. Ohi, S. Yano and A. Morita, *J. Porphyrins Phthalocyanines*, 2013, **17**, 331-342.
98. M. Tanaka, H. kataoka, M. Mabuchi, S. Sakuma, S. Takahashi, R. Tujii, H. Akashi, H. Ohi, S. Yano, A. Morita and T. Joh, *Anticancer Res.*, 2011, **31**, 763-770.
99. J. Ogikubo, J. L. Worlinsky, Y.-J. Fu and C. Brückner, *Tetrahedron Lett.*, 2013, **54**, 1707-1710.
100. Z. Gross, N. Galili and I. Saltsman, *Angew. Chem., Int. Ed. Engl.*, 1999, **38**, 1427-1429.
101. B. Ventura, A. Degli Esposti, B. Koszarna, D. T. Gryko and L. Flamigni, *New J. Chem.*, 2005, **29**, 1559-1566.
102. Y. Lili, S. Han, S. Lei, Z. GuoLiang, L. HaiYang, W. Hui and J. LiangNian, *Sci. China: Phys., Mech. Astron.*, 2010, **53**, 1491-1496.
103. D. T. Gryko and B. Koszarna, *Org. Biomol. Chem.*, 2003, **1**, 350-357.
104. Z. Gross and N. Galili, *Angew. Chem., Int. Ed. Engl.*, 1999, **38**, 2366-2369.
105. T. A. F. Cardote, J. F. B. Barata, M. A. F. Faustino, A. Preuß, M. G. P. M. S. Neves, J. A. S. Cavaleiro, C. I. V. Ramos, M. G. O. Santana-Marques and B. Röder, *Tetrahedron Lett.*, 2012, **53**, 6388-6393.
106. T. Hori and A. Osuka, *Eur. J. Org. Chem.*, 2010, **2010**, 2379-2386.
107. M. Schmidlehner, F. Faschinger, L. M. Reith, M. Ertl and W. Schoefberger, *Appl. Organomet. Chem.*, 2013, **27**, 395-405.
108. L. M. O. Lourenc, M. G. P. M. S. Neves, J. A. S. Cavaleiro and J. P. C. Tome, *Tetrahedron*, 2014, **70**, 2681-2698.
109. B. Roeder, D. Naether, T. Lewald, M. Braune, C. Nowak and W. Freyer, *Biophys. Chem.*, 1990, **35**, 303-312.
110. N. Sekkat, H. v. d. Bergh, T. Nyokong and N. Lange, *Molecules*, 2011, **17**, 98-144.
111. C. C. Leznoff and A. B. P. Lever, eds., *Phthalocyanines, Properties and Applications*, Wiley, V.C.H., New York, 1989.
112. A. Aggarwal, S. Singh, Y. Zhang, M. Anthes, D. Samaroo, R. Gao and C. M. Drain, *Tetrahedron Lett.*, 2011, **52**, 5456-5459.
113. S. Silva, P. M. R. Pereira, P. Silva, F. A. Almeida Paz, M. A. F. Faustino, J. A. S. Cavaleiro and J. P. C. Tome, *Chem. Commun.*, 2012, **48**, 3608-3610.
114. P. M. Pereira, S. Silva, J. A. Cavaleiro, C. A. Ribeiro, J. P. Tome and R. Fernandes, *PLoS one*, 2014, **9**, e95529-e95541.
115. J. B. Pereira, E. F. A. Carvalho, M. A. F. Faustino, R. Fernandes, M. G. P. M. S. Neves, J. A. S. Cavaleiro, N. C. M. Gomes, A. Cunha, A. Almeida and J. P. C. Tomé, *Photochem. Photobiol.*, 2012, **88**, 537-547.
116. L. M. O. Lourenco, P. M. R. Pereira, E. Maciel, M. Valega, F. M. J. Domingues, M. R. M. Domingues, M. G. P. M. S. Neves, J. A. S. Cavaleiro, R. Fernandes and J. P. C. Tome, *Chem. Commun.*, 2014, **50**, 8363-8366.

117. S. Singh, A. Aggarwal, S. Thompson, J. P. C. Tomé, X. Zhu, D. Samaroo, M. Vinodu, R. Gao and C. M. Drain, *Bioconjugate Chem.*, 2010, **21**, 2136-2146.
118. S. Achelle, P. Couleaud, P. Baldeck, M.-P. Teulade-Fichou and P. Maillard, *Eur. J. Org. Chem.*, 2011, **2011**, 1271-1279.
119. H. Mori, T. Tanaka, N. Aratani, B. S. Lee, P. Kim, D. Kim and A. Osuka, *Chem. - Asian J.*, 2012, **7**, 1811-1816.
120. K. A. D. F. Castro, S. Silva, P. M. R. Pereira, M. M. Q. Simões, M. d. G. P. M. S. Neves, J. A. S. Cavaleiro, F. Wypych, J. P. C. Tomé and S. Nakagaki, *Inorg. Chem.*, 2015, **54**, 4382-4393.
121. K. A. D. F. Castro, M. M. Q. Simoes, M. G. P. M. S. Neves, J. A. S. Cavaleiro, F. Wypych and S. Nakagaki, *Catal. Sci. Technol.*, 2014, **4**, 129-141.
122. J. F. B. Hall, X. Han, M. Poliakoff, R. A. Bourne and M. W. George, *Chem. Commun.*, 2012, **48**, 3073-3075.
123. M. E. Lipińska, S. L. H. Rebelo and C. Freire, *J. Mater. Sci.*, 2014, **49**, 1494-1505.
124. H. J. Mackintosh, P. M. Budd and N. B. McKeown, *J. Mater. Chem.*, 2008, **18**, 573-578.
125. J. Yang, B. Gabriele, S. Belvedere, Y. Huang and R. Breslow, *J. Org. Chem.*, 2002, **67**, 5057-5067.
126. A. Varotto, C.-Y. Nam, I. Radivojevic, J. P. C. Tomé, J. A. S. Cavaleiro, C. T. Black and C. M. Drain, *J. Am. Chem. Soc.*, 2010, **132**, 2552-2554.
127. J. Rutishauser and P. Kopp, *Eur J Endocrinol*, 1998, **138**, 623-624.
128. M. A. Carvalho de Medeiros, K. Gorgy, A. Deronzier and S. Cosnier, *Mater. Sci. Eng., C*, 2008, **28**, 731-738.
129. S. F. Liu, A. R. Petty, G. T. Sazama and T. M. Swager, *Angew. Chem., Int. Ed.*, 2015, **54**, 6554-6557.
130. J. L. Worlinsky, S. Halepas and C. Bruckner, *Org. Biomol. Chem.*, 2014, **12**, 3991-4001.
131. A. Yella, H.-W. Lee, H. N. Tsao, C. Yi, A. K. Chandiran, M. K. Nazeeruddin, E. W.-G. Diao, C.-Y. Yeh, S. M. Zakeeruddin and M. Grätzel, *Science*, 2011, **334**, 629-634.
132. D. Batabyal, H. Li and T. L. Poulos, *Biochemistry*, 2013, **52**, 5396-5402.
133. G. Smeureanu, A. Aggarwal, C. E. Soll, J. Arijeloye, E. Malave and C. M. Drain, *Chemistry (Weinheim an der Bergstrasse, Germany)*, 2009, **15**, 12133-12140.
134. M. Eckshtain, I. Zilbermann, A. Mahammed, I. Saltsman, Z. Okun, E. Maimon, H. Cohen, D. Meyerstein and Z. Gross, *Dalton Trans.*, 2009, 7879-7882.
135. E. V. Kudrik, P. Afanasiev, L. X. Alvarez, P. Dubourdeaux, M. Clémancey, J.-M. Latour, G. Blondin, D. Bouchu, F. Albrieux, S. E. Nefedov and A. B. Sorokin, *Nat. Chem.*, 2012, **4**, 1024-1029.
136. Q. Liu and C. C. Guo, *Sci. China: Chem.*, 2012, **55**, 2036-2053.
137. J. A. R. Salvador, V. M. Moreira, R. M. A. Pinto, A. S. Leal and J. A. Paixão, *Beilstein J. Org. Chem.*, 2012, **8**, 164-169.
138. P. E. Ellis, Jr., J. E. Lyons and S. N. Shaikh, *Catal. Lett.*, 1994, **24**, 79-83.
139. L. Costa, M. A. F. Faustino, M. G. P. M. S. Neves, Â. Cunha and A. Almeida, *Viruses*, 2012, **4**, 1034-1074.
140. T. B. Zanoni, T. M. Lizier, M. d. D. Assis, M. V. B. Zanoni and D. P. de Oliveira, *Food and Chemical Toxicology*, 2013, **57**, 217-226.
141. M. M. Q. Simões, R. De Paula, M. G. P. M. S. Neves and J. A. S. Cavaleiro, *J. Porphyrins Phthalocyanines*, 2009, **13**, 589-596.
142. D.-h. Shen, L.-t. Ji, L.-l. Fu, X.-l. Dong, Z.-g. Liu, Q. Liu and S.-m. Liu, *J. Cent. South Univ.*, 2015, **22**, 862-867.
143. G. S. Machado, P. B. Groszewicz, K. A. D. d. F. Castro, F. Wypych and S. Nakagaki, *J. Colloid Interface Sci.*, 2012, **374**, 278-286.
144. R. Kato, Y. Kobayashi, M. Akiyama and T. Komatsu, *Dalton Trans.*, 2013, **42**, 15889-15892.
145. L. Li and G.-k. Hu, *Biosci. Rep.*, 2015, **35**, e00174-00171-00178.
146. I. Batinic-Haberle, I. Spasojevic, R. D. Stevens, P. Hambright, P. Neta, A. Okado-Matsumoto and I. Fridovich, *Dalton Trans.*, 2004, 1696-1702.
147. K. Durr, B. P. Macpherson, R. Warratz, F. Hampel, F. Tuczek, M. Helmreich, N. Jux and I. Ivanovic-Burmazovic, *J. Am. Chem. Soc.*, 2007, **129**, 4217-4228.
148. P. A. Gale, *Acc. Chem. Res.*, 2011, **44**, 216-226.
149. E. K. Hoffmann, *Biochim. Biophys. Acta*, 1986, **864**, 1-31.
150. J. Rutishauser and P. Kopp, *Eur. J. Endocrinol.*, 1998, **138**, 623-624.
151. P. D. Beer and P. A. Gale, *Angew. Chem., Int. Ed. Engl.*, 2001, **40**, 486-516.
152. J. M. M. Rodrigues, A. S. F. Farinha, P. V. Muteto, S. M. Woranovicz-Barreira, F. A. Almeida Paz, M. G. P. M. S. Neves, J. A. S. Cavaleiro, A. C. Tome, M. T. S. R. Gomes, J. L. Sessler and J. P. C. Tome, *Chem. Commun.*, 2014, **50**, 1359-1361.
153. P. Prados and R. Quesada, *Supramol. Chem.*, 2008, **20**, 201-216.
154. S. Kubik, C. Reyheller and S. Stüwe, *J. Inclusion Phenom. Macrocyclic Chem.*, 2005, **52**, 137-187.
155. B. Verdejo, F. Rodriguez-Llansola, B. Escuder, J. F. Miravet and P. Ballester, *Chem. Commun.*, 2011, **47**, 2017-2019.
156. E. Palomares, M. V. Martínez-Díaz, T. Torres and E. Coronado, *Adv. Funct. Mater.*, 2006, **16**, 1166-1170.
157. C. I. M. Santos, E. Oliveira, J. F. B. Barata, M. A. F. Faustino, J. A. S. Cavaleiro, M. G. P. M. S. Neves and C. Lodeiro, *J. Mater. Chem.*, 2012, **22**, 13811-13819.
158. D. P. Cormode, M. G. B. Drew, R. Jagessar and P. D. Beer, *Dalton Trans.*, 2008, DOI: 10.1039/B807153E, 6732-6741.
159. Y. Amao, *Microchim. Acta*, 2003, **143**, 1-12.
160. X.-d. Wang, X. Chen, Z.-x. Xie and X.-r. Wang, *Angew. Chem., Int. Ed.*, 2008, **47**, 7450-7453.
161. H. Hasumoto, T. Imazu, T. Miura and K. Kogure, *J. Oceanogr.*, 2006, **62**, 99-103.
162. Y. Tian, B. R. Shumway, W. Gao, C. Youngbull, M. R. Holl, R. H. Johnson and D. R. Meldrum, *Sens. Actuators, B*, 2010, **150**, 579-587.
163. M. I. J. Stich, M. Schaeferling and O. S. Wolfbeis, *Adv. Mater.*, 2009, **21**, 2216-2220.
164. S. Nagl and O. S. Wolfbeis, *Analyst*, 2007, **132**, 507-511.
165. S. M. Borisov, A. S. Vasylevska, C. Krause and O. S. Wolfbeis, *Adv. Funct. Mater.*, 2006, **16**, 1536-1542.
166. M. E. Köse, B. F. Carroll and K. S. Schanze, *Langmuir*, 2005, **21**, 9121-9129.
167. B.-H. Han, I. Manners and M. A. Winnik, *Anal. Chem.*, 2005, **77**, 8075-8085.
168. S.-K. Lee and I. Okura, *Anal. Commun.*, 1997, **34**, 185-188.
169. M. R. Friswell, 1992, European patent no. 0509818 A1.

Review

Organic & Biomolecular Chemistry

170. J. J. Krutak, M. R. Cushaman and M. A. Waever, 1996, US patent no. 5525516.
171. S. Puangmalee, A. Petsom and P. Thamyongkit, *Dyes Pigm.*, 2009, **82**, 26-30.
172. G. de la Torre, C. G. Claessens and T. Torres, *Chem. Commun.*, 2007, 2000-2015.
173. S. Honda, T. Nogami, H. Ohkita, H. Benten and S. Ito, *ACS Appl. Mater. Interfaces*, 2009, **1**, 804-810.
174. B. Walker, A. B. Tamayo, X.-D. Dang, P. Zalar, J. H. Seo, A. Garcia, M. Tantiwivat and T.-Q. Nguyen, *Adv. Funct. Mater.*, 2009, **19**, 3063-3069.
175. G. Calogero, G. Di Marco, S. Cazzanti, S. Caramori, R. Argazzi, A. Di Carlo and C. A. Bignozzi, *Int. J. Mol. Sci.*, 2010, **11**, 254-267.
176. T. Bříza, R. Kaplánek, M. Havlík, B. Dolenský, Z. Kejík, P. Martásek and V. Král, *Supramol. Chem.*, 2008, **20**, 237-242.
177. Y.-H. Chan, A. E. Schuckman, L. M. Pérez, M. Vinodu, C. M. Drain and J. D. Batteas, *J. Phys. Chem. C*, 2008, **112**, 6110-6118.
178. O. Jacobson, D. O. Kiesewetter and X. Chen, *Bioconjugate Chem.*, 2015, **26**, 1-18.
179. R. Jacquemin, *Int. J. Radiat. Appl. Instrum., Part A* 1987, **38**, 1087-1089.
180. P. S. Rose, J. P. Smith, R. C. Aller, J. K. Cochran, R. L. Swanson and R. B. Coffin, *Environ. Sci. Technol.*, 2015, **49**, 10312-10319.
181. C. M. Drain, G. Smeureanu, S. Patel, X. Gong, J. Garno and J. Arijeloye, *New J. Chem.*, 2006, **30**, 1834-1843.
182. C. M. Drain, F. Nifiatis, A. Vasenko and J. D. Batteas, *Angew. Chem., Int. Ed.*, 1998, **37**, 2344-2347.
183. C. M. Drain and J.-M. Lehn, *J. Chem. Soc., Chem. Commun.*, 1994, DOI: 10.1039/C39940002313, 2313-2315.
184. M. J. Jurow, C. Mayr, T. D. Schmidt, T. Lampe, P. I. Djurovich, W. Brutting and M. E. Thompson, *Nat. Mater.*, 2015, advance online publication.