

Photochemical & Photobiological Sciences

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Kinetics of photochemical reactions of biphotochromic compounds based on spironaphthopyran and enamine - conjugation effect

Peter P. Levin,^{a,b} Alexander S. Tatikolov,^a Natalia L. Zaichenko,^{*b} Andrey I. Shienok,^b Liubov S. Koltsova,^b Irina M. Sherbakova,^b Irina R. Mardaleishvili,^b and Alexander A. Berlin^b

Novel biphotochromic compound (BPC) with two photochromic fragments, namely spironaphthopyran and hydroxyazomethine, was synthesized and studied by the nanosecond laser flash photolysis using the excitation wavelengths of 337, 430, 470 and 500 nm in methanol and toluene. The photoexcitation of BPC results in the formation of at least two colored transients. The first one is the merocyanine form B (the maximum in the absorption spectrum is near 600 nm and the lifetime is 0.1 and 0.05 s in methanol and toluene, respectively) due to the spiro-bond break followed by isomerization. The second one is the trans-keto form A^{Kt} (the maximum in the absorption spectra is near 480 nm and the lifetime is 0.1 and 5 ms in methanol and toluene, respectively) as a result of cis-enol or cis-keto tautomers transformations. The relative yields of B and A^{Kt} depends essentially on the wavelength of excitation. The form A^{Kt} is the key transient formed under the excitation with the visible light, whereas its yield under the excitation with UV light is comparable with that of B. The specific solvation by methanol molecules favors the spirocycle opening even under the visible light excitation. The results obtained for novel BPC were compared with those for other BPC where the same fragments are combined in such a way that the form B is the major one under the excitation with UV light whereas it virtually is not observed under visible light excitation. The difference in both BPC is discussed in terms of conjugation (π -coupling) between photochromic fragments.

^a N.M. Emanuel Institute of Biochemical Physics, Russian Academy of Sciences, Kosygina 4, Moscow 119334, Russia

^b N.N. Semenov Institute of Chemical Physics, Russian Academy of Sciences, Kosygina 4, Moscow 119991, Russia

† Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

1. Introduction

In recent years, bi- and multiphotochromic systems were widely studied owing to their properties which can be attractive for practical purposes. One of their potential applications is the development of a new generation of materials for molecular electronics, such as molecular switching devices, molecular logic gates, etc.^{1,2}. The important features required for the design of such systems is a different photochemical behavior of the constituent elements implying the sensitivity to the wavelength of the excitation and the molecular environment. There is a large diversity of structures described in literature, using several switches connected through a variety of organic and inorganic structures³⁻¹².


Nevertheless, effective multi-addressable and multi-active compounds remain a minority and a search of novel effective systems is an important task. We have developed the synthetic route to a new type of biphotochromic compounds on the basis of spironaphthoxazine linked to a substituted salicylideneaniline (SA)¹³. The essential sensitivity to the excitation wavelength was detected for the triphotochromic compound of this series having the photochromic azobenzene substituent in the hydroxyazomethine fragment : only merocyanine form was observed at the excitation by UV light, whereas three intermediates appeared after the visible light excitation¹³.


In the present work in order to proceed our research further we studied the spectral-kinetics properties of novel BPC composed of the naphthopyran and hydroxyazomethine (1,3-dihydro-8'-hydroxy-7'-(4-methylphenyliminomethyl)-1,3,3-trimethylspiro[(2H)-

indole-2,3'-[3H]-naphtho[2,1-b]pyran] (1) (Scheme 1, where 1A^E is (1) in the closed form of spiro-cycle linked to cis-enol form of hydroxyazomethine fragment) in two solvents of different nature (toluene and methanol). The ns laser flash photolysis techniques with different wavelengths of excitation (337, 430, 470 and 500 nm) was used with the aim to vary the ratio between the intermediates (photoinduced isomeric species) of different nature.

The photochemical behavior of (1) is compared with that of other BPC (2) -1,3-dihydro-5-(2-hydroxy-1-naphthylmethylideneimino)-1,3,3-trimethylspiro[(2H)-indole-2,2'-(3H)-naphtho[2,1-b]pyran] (Scheme 1, where 2A^E is (2) in the closed form of spiro-cycle linked to cis-enol form of hydroxyazomethine fragment) studied earlier by us^{14,15}.

The first structural difference between BPC (1) and (2) is that in (1) there is a conjugation between π -electronic systems of two photochromic fragments – spirocycle (naphthopyran) and hydroxyazomethine and in (2) the conjugation is expected to be between π -electronic systems of the indole and the hydroxyazomethine moieties. The second difference is that in (2) the indole fragment acts as “aniline” part of azomethine moiety whereas in (1) the naphthopyran fragment is a “phenol” type substituent (Scheme 2).






Scheme 2

It was supposed that the comparison of the photochemical behavior of (1) and (2) can be useful to clarify the role of conjugation between different photochemical fragments in their selective photoexcitation. The compounds (3) and (4) (Scheme 3) were used as references representing constituent fragments of the BPC (1) and (2).

2. Experimental

2.1. Materials

Toluene and methanol of spectroscopic grade and CDCl_3 were purchased from Aldrich and used as received. 1,3,3-trimethyl-2-methylene-3H-indole purchased from Acrus Organics was additionally purified by vacuum distillation. All other chemicals were reagent grade and used without additional purification.



Scheme 3

2.2. Devices

The steady-state absorption spectra of solutions (using 10- mm quartz cells) were recorded in the UV and visible regions on a MultiSpec-1501 spectrophotometer.

The absorption spectra and kinetics of formation and decay of the intermediates were recorded in the time range ≥ 10 ns in the $400 \leq \lambda \leq 800$ nm spectral region by the nanosecond laser flash photolysis techniques with registration of electronic absorption¹³. A nitrogen laser (PRA LN 1000 with 1 ns pulse duration and radiation wavelength of 337 nm, operating at ≤ 10 Hz), or a dye laser (PRA LN 102 with 0.5 ns pulse duration and 430, 470 or 500 nm radiation wavelength) pumped by a nitrogen laser were used as an excitation sources. The acquisition and averaging of kinetic curves were performed by the sequence of pulses followed at a repetition rate which was long enough to insure the complete decay of the intermediates. The samples were sufficiently stable in the sense of photodegradation so that at least one hundred laser pulses can be used in experiments with a sample. The effect of molecular oxygen was not found neither on the resistance of compounds nor on the decay kinetics of the intermediates. The data presented in this paper are average values obtained by processing at least ten kinetic curves under the same selected conditions. The absorbance at excitation wavelength was adjusted to 10 - 40%. All measurements were carried out at ambient temperature.

The ^1H NMR spectra were recorded using Bruker WM 400 spectrometer, operating at 400 MHz, at 25°C in

CDCl_3 . Elemental analyses were performed using EuroVector analyzer Euro EA 3000.

2.3. Preparation of biphotochromic compounds

The synthetic scheme 4 was carried out for BPC (**1**) synthesis. In short, we carried out the diformylation of the starting 2,6-dihydroxynaphthaline by a modified Daff method¹⁶, then 2,6-dihydroxy-1,5-naphthalindicarboxaldehyde was condensed with 1 equivalent of 1,3,3-trimethyl-2-methylene-3H-indol giving 7'-formyl-8'-hydroxy-spiro[naphthopyran], which was condensed with p-toluidine .1-(4-Methylphenyliminomethyl)-2-naphthol (**3**) was synthesized according¹⁷.

BPC (**2**) was synthesized as described in¹⁸. Purity of the compounds was controlled by TLC, melting points, NMR (Bruker WM 400) and elemental analysis data.

1,3-Dihydro-7'-formyl-8'-hydroxy-1,3,3-trimethylspiro[2H-indole-2,3'-[3H]naphtho[2,1-b][1,4]pyran] (5)

The solution of 1,3,3-trimethyl-2-methylene-3H-indol (0.9 ml, 5 mmol) in 10 ml of dry dioxane was added dropwise to a solution of 2,6-dihydroxy-1,5-naphthalinedicarboxaldehyde (4 g, 18.5 mmol) in 200 ml of dry dioxane and refluxed with stirring for 5.5 h. Then, the mixture was cooled to room temperature and the excessive dialdehyde was filtered off. The solvent was evaporated and the residue was separated from concomitant dispiro-compound using column chromatography on SiO_2 with benzene as eluent. The fractions with the desired product (R_f 0.6 in


chloroform) were combined, the solvent was evaporated, and 5.37 g (26 %) of green precipitate, mp 240 °C (benzene) was obtained.

$^1\text{H NMR}$ (400 MHz, CDCl_3 , δ , ppm): 1.22, 1.34 (both s, 3 H each, $\text{C}(\text{CH}_3)_2$); 2.75 (s, 3H, NCH_3); 5.88 (d, 1H, $\text{H}(2')$, $^3J_{1,2'}$ 10.5 Hz); 6.55 (d, 1 H, $\text{H}(7)$, $^3J_{6,7}$ 7.9 Hz); 6.88 (t, 1 H, $\text{H}(5)$, $^3J_{4,5} = ^3J_{5,6}$ 7.4 Hz); 7.10 (d, 1 H, $\text{H}(4)$, $^3J_{4,5}$ 7.4 Hz); 7.14 (d, 1 H, $\text{H}(5')$, $^3J_{5,6}$ 9.4Hz); 7.20 (1H, $\text{H}(6)$); 7.20 (1H, $\text{H}(9')$); 7.54 (d, 1 H, $\text{H}(1')$ $^3J_{1,2'}$ 10.5 Hz); 8.15 (d 1H, $\text{H}(10')$, $^3J_{9',10'}$ 9.2 Hz); 8.25 (d, 1H, $\text{H}(6')$, $^3J_{5',6'}$ 9.4 Hz); 10.75 (s, 1H, $\text{C}(\text{O})\text{H}$); 12.89 (s, 1H, OH).

Found: C, 77.53; H, 5.81; N, 3.88. $\text{C}_{24}\text{H}_{21}\text{NO}_3$ requires C, 77.61; H, 5.70; N, 3.77%.

1,3-Dihydro-8'-hydroxy 7'-(4-methylphenyliminomethyl)-1,3,3-trimethylspiro[2H-indole-2,3'-[3H]naphtho[2,1-b][1,4]pyran] (BPC 1)

Solution of 95 mg (0,256 mmol) 1,3-dihydro-7'-formyl-8'-hydroxy-1,3,3-trimethylspiro[2H-indole-2,3'-[3H]naphtho[2,1-b][1,4]pyran] (**5**) and 27,5 mg (0,256 mmol) of p-toluidine in 10 ml toluene was refluxed for 6 h. Then the mixture was cooled to room temperature



Scheme 4

and filtered. Solvent was evaporated and the residue was purified by preparative TLC on SiO₂ with 19:1 benzene- acetone mixture as an eluent. The photochromic strip with *R_f* 0.7 was collected, the product desorbed by 5:1 chloroform- acetone mixture, and the solvent was evaporated on a rotary evaporator to give 0.57 g (yield 48 %) of an orange precipitate, mp 282 °C (benzene- acetone mixture).

¹H NMR (400 MHz, CDCl₃, δ, ppm):

1.22, 1.34 (both s, 3 H, C(CH₃)₂); 2.40 (s, 3H, CH₃); 2.75 (s, 3H, NCH₃); 5.85 (d, 1H, H(2'), ³J_{1,2'} 10.5 Hz); 6.54 (d, 1H, H(7), ³J_{6,7} 7.7 Hz); 6.87 (t, 1H, H(5), ³J_{4,5} 7.0 Hz, J_{5,6} 7.4 Hz); 7.10 (d, 1H, H(4), ³J_{4,5} 7.0 Hz); 7.18-7.22 (3H, H(6), H(9'), CHN); 7.25 (2H, H(2'')); 7.26 (2H, H(3'')); 7.04 (d, 1H, H(5'), ³J_{5',6'} 9.3 Hz); 7.55 (d, 1H, H(1'), ³J_{1',2'} 10.5 Hz); 7.94 (d, 1H, H(10'), ³J_{9',10'} 9.2 Hz); 8.09 (d, 1H, H(6'), ³J_{5',6'} 9.4 Hz); 9.32 (s, 1H, OH).

Found: C, 80.62; H, 6.34; N, 5.82. C₃₁H₂₈N₂O₂ requires C, 80.84; H, 6.13; N, 6.08%.

2.4. Abbreviations

The following symbols are used for marking the state of the components of BPC molecules. The capital letters A and B designate the initial closed and the final opened colored spirocycle fragment states, respectively. The upper case indexes E^c, E^t, K^c and K^t denote cis- and trans- enol, cis-keto- and trans-keto states of hydroxyazomethine fragment, respectively. The same abbreviations are used for the corresponding model compounds.

3. Results and discussion

3.1. Ground State Absorption

The UV-vis absorption spectra of (1) and (3) in methanol and toluene are presented in Fig. 1. The model (4) in A form absorbs in UV region¹⁹. The absorption spectrum of (3) in toluene is characterized by the long-wavelength maximum at 380 nm and relatively weak ones at 440 and 460 nm (Fig.1, curve 4). It is well-known for anil-type compounds that their molecules may exist in E^c and K^c tautomeric forms (Scheme 5) depending on the molecule structure and the environmental conditions²⁰⁻²⁶. The E^c absorbs in near UV and it is usually more stable in aprotic solvents. The model (3) in toluene exists for the most part in E^c (maximum at 380 nm) although the presence of small amount of K^c (maxima at 440 and 460 nm) is observed (approximately 10% if to assume the equality of the corresponding extinction coefficients).






Fig.1 UV-vis absorption spectra of PBC (1) (curves 1, 2) and model (3) (curves 3, 4) in toluene (curves 2, 4) and methanol (curves 1, 3) solutions.

The intensive absorption bands with a maxima at 440 and 460 nm are observed for (**3**) in methanol (Fig.1, curve 3) indicating that (**3**) exists in methanol solution mainly in K^c form (at least 70%). This difference between the absorption of (**3**) in toluene and methanol should be attributed to the specific solvation effect rather than to the polarity. Indeed the absorption spectrum of (**3**) in acetonitrile is similar to that in toluene (the contribution of K^c is 20%).

The long-wavelength parts of the spectrum of (**1**) are considerably shifted to the longer wavelength with respect to that of the reference compound (**3**) in both solvents (Fig.1, spectra 1, 2). The absorption at 600 nm of the open B form was not observed, showing that (**1**) exists in the closed form A of spirocycle fragment in both solvents. The absorption of A^{Kc} (Scheme 6) in methanol (maxima at 470 and 490 nm) is pronounced as compared with that of A^{Ec} (maximum at 420 nm) (see Fig.1, curve 1). The estimated contribution of A^{Kc} is at least 50%. The absorption of A^{Kc} (**1**) in toluene in the region ≥ 470 nm is relatively weak (see Fig.1, curve 2, the estimated contribution of A^{Kc} is 10%). Very similar regularities were found for the solvent effect on the spectrum of BPC (**2**).¹⁸ It is worthwhile mentioning that the arrangement and locations of the relevant bands in spectra of both (**1**) and (**2**) resemble each other rather well. The equilibrium between cis and trans forms in the ground state was not found for both compounds on the basis of UV-vis absorption spectra investigation as it is the case for the most of anil-type compounds²⁰⁻²².




Scheme 5 Tautomeric forms of N-naphthylsalicylideneimine molecules



Scheme 6

3.2. Intermediates of model compounds produced by the excitation with UV and visible light

The spectra and kinetics of the intermediates obtained by the laser flash photolysis of the model compounds (**3**) and (**4**) representing constituent fragments of BPC are required for the recognition of BPC intermediates structure. It was shown earlier that the photoexcitation of (**4**) in methanol (Fig.2, spectrum 2) and toluene by UV light (337 nm) produces in the time interval ≥ 10 ns the only one intermediate – merocyanine form B (Scheme 7) with the absorption maximum at 560 and 550 nm and first order decay rate constant $k = 14$ and 42 s^{-1} respectively¹⁸. Photoexcitation of (**4**) by the visible light has no result since it does not absorb at those wavelengths¹⁹.



Scheme 7 Photochromic process in spironaphthopyran (4).





Fig.2 Visible absorption spectra of the intermediates produced by the 337 nm laser flash photolysis of (3) - (spectrum 1) and (4) - (spectrum 2) in methanol solution in 0.05 μ s after the laser pulse.

The photoexcitation of (3) in both solvents with UV or visible light (except 500 nm where (3) practically does not absorb) results in the formation of the single intermediate with the characteristic absorption spectrum with a maxima at 450 and 480 nm in toluene and methanol (Fig.2, spectrum 1), respectively¹⁸. Its decay kinetics is first order with the rate constants equal to $9.9 \cdot 10^2$ and $9.5 \cdot 10^3 \text{ s}^{-1}$ in toluene and methanol, respectively. We have identified this intermediate as *trans*-keto form K^t , which was formed due to the photoisomerization of K^c (Scheme 8) or proton transfer coupled with isomerization after the

photoexcitation of E^c . The K^c decay is accelerated significantly in the hydrogen-bonding environment. The similar results were found for unsubstituted 1-(phenyliminomethyl)-2-naphthol¹⁸ and N,N-bis-(salicylidene)-p-phenylenediamines²² demonstrating that one and the same keto-tautomer is created either from the enol form via the excited state proton transfer coupled with isomerization or via the direct photoisomerization of the keto-form. The dynamics of K^t formation process is in the subpicosecond time scale as it is for spiro-compounds^{15, 21, 22, 27-31}.

3.3. Intermediates of BPC (1) produced by the excitation with UV and visible light in methanol

The photoexcitation of BPC (1) by visible light 500 nm (where only A^{K^c} absorbs) leads immediately after the laser pulse to the formation of two intermediates which absorption spectra contain characteristic bands with maxima at 480, 520 and 600 nm (Fig. 3 spectrum 1). The intermediate with absorption at 480 - 520 nm is relatively short-lived (first order rate constant ($9.7 \cdot 10^3 \text{ s}^{-1}$) is very similar to that measured for model (3), whereas that one with absorption at 600 nm is long-lived (Fig. 3 spectrum 2, first order rate constant (11 s^{-1}) is very similar to that found for model (4).



Scheme 8 *Cis*-ketone to *trans*-ketone photoisomerization of model compound (3).






Fig.3. Visible absorption spectra of the intermediates produced by the 500 nm laser flash photolysis of BPC (**1**) in methanol solution in 0.05 μ s (spectrum 1) and 1 ms (spectrum 2) after the laser pulse. Insert - decay kinetics of the intermediates, observed at 480 nm (1) and 600 nm (2) under flash photolysis of (**1**) with 500 nm light in MeOH.

On the basis of flash-photolysis data for the model compounds (**3**) and (**4**) the relatively short lived intermediate can be identified as A^{Kt} . The long-lived one seems to be B^{Kc} since there is no visible changes in absorption at $\lambda \leq 500$ nm. The form A^{Kt} is appeared as a result of ketone cis-trans photoisomerization in hydroxyazomethine fragment according to Scheme 9 and B^{Kc} arises via spiro-bond cleavage and corresponding isomerization according to Scheme 10. The yield of B^{Kc} was estimated to be 4% relative to that of A^{Kt} assuming that the ratio of the corresponding molar absorption coefficients of B and A^{Kt} at the maxima is 3¹⁷.



Scheme 9 Photochromic process in hydroxyazomethine fragment of BPC (**1**).



Scheme 10 Spiro-bond cleavage and formation of the merocyanine form B from A^{Kc} in BPC (**1**).

The same transients with the same yields were obtained under the photolysis with light 470 and 430 nm, where A^{Ec} also absorbs. Therefore, A^{Kt} is formed via the excited state proton transfer coupled with isomerization as well. The yield of A^{Kt} in comparison with that of Kt (**3**) was estimated to be 80%. Laser photoexcitation of (**1**) by UV light (337 nm) yields the same two intermediates A^{Kt} and B^{Kc} , but the yield of merocyanine form B^{Kc} increases 2.5 times relative to that of A^{Kt} . The yields of A^{Kt} and B^{Kc} as compared with those of Kt (**3**) and B (**4**) measured for the samples with the same absorption at 337 nm were estimated to be 60 and 2%, respectively.

The interconversion of A^{Kt} and B^{Kc} was not observed in the time interval studied. Therefore, two photoprocesses in BPC (**1**) – cis- trans isomerization of

azomethine fragment and merocyanine formation from spirocycle – occur simultaneously in the course of relaxation of singlet excited states¹⁵.

3.4. Intermediates of BPC (1) produced by the excitation with UV and visible light in toluene

Pulse photoexcitation of (1) in toluene with visible light results in the formation of the only single isomer A^{Kt} with the absorption maximum at 470 nm (Fig.4) and first-order decay rate constant being $2.0 \cdot 10^2 \text{ s}^{-1}$. The yield of A^{Kt} in comparison with that of Kt (3) was estimated to be 80% as it is in methanol.

However the small amount of B^{Ec} is observed (maximum at 580 nm, first-order decay rate constant 18 s^{-1}) under the UV light photoexcitation in addition to A^{Kt} . The estimated yield of B^{Kc} relative to that of A^{Kt} (assuming that the ratio of the corresponding molar absorption coefficients of B and A^{Kt} at the maxima is 3¹⁷) was found to be 3% that is three times smaller than that in methanol. The absorption spectrum of B^{Ec} (1) has maximum which is slightly shifted to the long-wavelength region with respect to that of form B of model compound (4).




Fig. 4. Visible absorption spectra of the intermediates produced by the 500 nm laser flash photolysis of BPC (1) in toluene solution in $0.05 \mu\text{s}$ (1), and 30 ms after the laser pulse (2). Insert - decay kinetics of the intermediates, observed at 480 nm (1) and 600 nm (2) under laser flash photolysis of BPC (1) with 500 nm light in toluene

3.5. The comparison between the photochemical behaviors of BPC (1) and (2)

Pulse photoexcitation of BPC (2) with the visible light in toluene as well as in methanol results in cis- trans isomerization of azomethine fragment whereas the form B formation was not found at all¹⁸. The rise time of the precursor of trans isomers (corresponding non relaxed singlet excited states) was 0.2 ps ¹⁵. Since the excitation with the visible light corresponds to the electronic transitions localized in the hydroxynaphthaldehydeimine moiety of (2), which is insufficiently coupled with spiro fragment, then the spiro bond rupture is inefficient in time interval of the localized singlet excited state relaxation. In the case of BPC (1) the more efficient corresponding coupling results in formation of small amount of form B even under the visible light excitation. It is worthwhile

mentioning that the hydrogen bonds formation between (1) and methanol favor the spiro bond scission in (1) probably due to the decrease of the corresponding energy required and/or the retardation of hydroxyazomethine fragment isomerization. On the one hand, photoexcitation of BPC (2) with UV light results in the formation of form B as a major product in both solvents¹⁸. The relatively poor coupling between spiro and hydroxynaphthalideneimine parts of (2) ensures opportunity of spiro bond break in the course of high singlet excited states relaxation (the rise time of B precursor after UV excitation of (2) was 0.15 ps¹⁵) rather than it would be relaxed to the low-lying singlet excited states localized on the hydroxynaphthalideneimine fragment. On the other hand, the most part of high singlet excited states of (1) relax principally to the low-lying singlet excited states localized in the azomethine part of (1) due to the efficient coupling and the form B still remains the minor product even under UV excitation of (1).

Conclusion

Our previous works were directed towards the search of biphotochromic compounds which photochemical behavior would be dependent on the wavelength of excitation light^{13,14}. We had synthesized two series of BPCs, which molecules combine spirocycle and hydroxyazomethine fragments in different ways in order to modify the conjugation (π -coupling) between photochromic fragments. No excitation wavelength effect was observed in the first series of compounds with conjugation between the fragments, however one compound from the second series, namely BPC (2),

demonstrated pronounced excitation wavelength effect¹⁸. In order to insight into the effect of conjugation on the photochemistry of biphotochromic compounds we have synthesized novel BPC (1) with other type of the connection between spirocycle and hydroxyazomethine fragments. The coupling in compound (1) seems to be somewhat intermediate between those insensitive and considerably sensitive to the wavelength of excitation. The specific solvation effect was found. The hydrogen bonding with methanol molecules favors the spirocycle opening even under the visible light excitation whereas it was not observed in toluene. However the spirocycle opening in (1) remains the minor pathway even under UV light excitation while the cis-trans photoisomerization coupled to the excited state intramolecular proton transfer in hydroxyazomethine fragment is the basic process.

Acknowledgements

This work was financially supported by the Russian Academy of Sciences (under Presidium Program No. 8) and by the RFBR (Project No. 13-03-00396a)

References

- 1 V. I. Minkin, Bistable organic, organometallic, and coordination compounds for molecular electronics and spintronics, *Russ. Chem. Bull. Int. Ed.*, 2008, **57**, 673 - 703.
- 2 A. P. De Silva, N.D. Mc Clenaghan, Molecular-scale logic gates, *Chem. Eur. J.*, 2004, **10**, 574 - 586.

- 3 A. Fihey, A. Perrier, W. R. Browne and D. Jacquemin, Multiphotochromic molecular systems, *Chem. Soc. Rev.*, 2015, **44**, 3719 - 3759.
- 4 K. Higashiguchi, K. Matsuda, N. Tanifuji and M. Irie, Full-color photochromism of a fused dithienylethene trimer, *J. Am. Chem. Soc.*, 2005, **127**, 8922 - 8923.
- 5 S. Delbaere, G. Vermeersch, M. Frigoli and G. H. Mehl, Bridging the visible: the modulation of the absorption by more than 450 nm, *Org. Lett.*, 2010, **12**, 4090 - 4093.
- 6 J. Andreasson, S. D. Straight, T. A. Moore, A. L. Moort and D. Gust, Molecular all-photonic encoder-decoder, *J. Am. Chem. Soc.*, 2008, **130**, 11122 - 11128.
- 7 J. Andreasson, U. Pischel, S. D. Straight, T. A. Moore, A. L. Moore and D. Gust, All- photonics multifunctional molecular logic device, *J. Am. Chem. Soc.*, 2011, **133**, 11641 - 11648.
- 8 G. Sevez, J. Gan, S. Delbaere, G. Vermeersch, L. Sanguinet, E. Levillain and J.-L. Pozzo, Photochromic performance of a dithienylethene-indolinoxazolidine hybrid, *Photochem. Photobiol. Sci.*, 2010, No9, 131 - 135.
- 9 A. Samat, V. Lokshin, K. Chamontin, D. Levi, G. Pepe and R. Guglielmetti, Spirooxazine photoisomerization and relaxation in polymer matrixes, *Tetrahedron*, 2001, **57**, 7349 - 7359.
- 10 J. Berthet, J.-C. Micheau, A. Metelitsa, G. Vermeersch and S. Delbaere, Multistep Thermal Relaxation of Photoisomers in Polyphotochromic Molecules, *J. Phys. Chem. A*, 2004, **108**, 10934 - 10940.
- 11 M. Frigoly and G. H. Mehl, Multiplrid biphotochromic system, *Angew. Chem. Int. Ed.*, 2005, **44**, 5048 - 5052.
- 12 K. Szacilowski, Digital information processing in molecular systems, *Chem. Rev.*, 2008, **108**, 3481 -3548 .
- 13 P. P. Levin, A. S. Tatikolov, N. L. Zaichenko, A. I. Shienok, L. S. Koltsova, O. Yu. Oskina, I. R. Mardaleishvili, L. D. Popov, S. I. Levchenkov and A. A. Berlin, Kinetics of photochemical reactions of multifunctional hybrid compounds based on spironaphthoxazines upon photoexcitation with light of different wavelength *J. Photochem. Photobiol. A* , 2013, **251**, 141 - 147.
- 14 N. L. Zaichenko, A. I. Shienok, L. S. Koltsova , I. R. Mardaleishvili, A. S. Tatikolov , P. P. Levin and A. A. Berlin, Bifunctional photosensitive compound for optical procssrs, *Physica Status Solidi A* , 2010, 2746 - 2748.
- 15 V. A. Nadtochenko, P. P. Levin, N. L. Zaichenko, F. E. Gostev, I. V. Shelaev, A. I. Shienok, L. S. Koltsova, O. M. Sarkisov and A.A. Berlin, Spectral and kinetic parameters of transient species in the photolysis of naphthylmethylideneiminospironaphthopyran by excitation at different wavelengths: nano_ and femtosecond laser photolysis, *High Energy Chem.*, 2013, **47**, 121 - 128.
- 16 H. Houjou, T. Motoyama, S. Banno, I. Yoshikawa and K. Araki, *J. Org. Chem.*, 2000, **974**, 520.
- 17 N. L. Zaichenko, P. P. Levin, I. R. Mardaleishvili, A. I. Shienok, L. S. Koltsova, O. Yu. Oskina and A. S. Tatikolov, Synthesis and kinetics of photochemical


- reactions of novel bifunctional salicylideneiminospironaphthoxazines, *Russian Chemical Bulletin International Edition*, 2008, **57**, 2394 - 2401.
- 18 P. P. Levin, N. L. Zaichenko, A. I. Shienok, L. S. Koltsova, I. R. Mardaleishvili and A. S. Tatikolov, Spectral kinetic characteristic of the photoisomerization products of naphthylmethylideneiminospironaphthopyran undused by photolysis at different wavelength, *Russ. Chem. Bull. Int. Ed.*, 2012, **61**, 532 -538 .
- 19 Photochromism: Molecules and Systems, Ed. H. Dürr, H. Bouas – Laurent, Elsevier, London, 2003, 1218 pp.
- 20 M. Z. Zgiersk and A. Grabowska, *J. Chem. Phys.*, Photochromism of salicylideneaniline .2000, **112**, 6329 - 6337.
- 21 W. M. F. Fabian, L. Antonov, D. Nedeltcheva, F. S. Kamounah and P. J. Taylor, Tautomerism in Hydroxynaphthaldehyde Anils and Azo Analogues: a Combined Experimental and Computational Study, *J. Chem. Phys. A*, **108**, 2004, 7603 - 7612.
- 22 M. Ziótek, J. Kubicki, A. Maciejewski, R. Naskrecki and A. Grabowska, Enol-keto tautomerism of aromatic photochromic Schiff base N,N`-bissalicylidene-p-phenylenediamine: Ground state equilibrium and excited state deactivation studied by solvatochromic measurements on ultrafast time scale, *J. Chem.Phys.*, 2006, **124**, 124518.
- 23 P. Fita, E. Luzina, T. Dziembowska, Cz. Radzewicz and A. Grabowska, Chemistry, photophysics, and ultrafast kinetics of two structurally related Schiff bases containing the naphthalene or quimoline ring, *J. Chem. Phys.*, 2006, **125**, 184508.
- 24 M. Ziótek, J. Kubicki, A. Maciejewski, R. Naskrecki, W. Luniewska, and A. Grabowska, Unusual conformational effects in proton transfer kinetics of an excited photochromic Schiff base, *J. Photochem. Photobiol. A*, 2006, **180**, 101 - 108.
- 25 M. Ziótek, M. Gil, J. A. Organero and A. Douhal, What is the difference between the dynamics of anion- and keto- type of photochromic salicylaldehyde azine ?, *Phys. Chem. Chem. Phys.*, 2010, **12**, 2107 - 2115.
- 26 R. S. Becker and W. F. Richey, Photochromic aniles. Mechanism and products of photoreactions and thermal reactions, *J. Am. Chem. Soc.*, 1967, **89**, 1298.
- 27 S. Mitra and N. Tamai, Femtosecond spectroscopic study on photochromic salixylidene anilinem, *Chem. Phys. Lett.*, 1998, **282**, 391 - 397.
- 28 S. Mitra and N. Tamai, Dynamics of photochromism in salicylideneanilin: a femtosecond spectroscopic study, *Phys. Chem. Chem. Phys.*, 2003, **5**, 4647 - 4652.
- 29 C. Okabe, T. Nakabayashi, Y. Inokuchi, N. Nishi and H. Sekiya, Ultrafast excited-state dynamics in photochromic N-salicylideneaniline studied by femtosecond time-resolved REMPI spectroscopy, *J. Chem. Phys.*, 2004, **121**, 9436 - 9442.
- 30 M. Ziótek, J. Kubicki, A. Maciejewski, R. Naskręcki and A. Grabowska, Proton transfer reaction and photochromism of the new family of

Journal Name

ARTICLE

Schiff bases, *Phys. Chem. Chem. Phys.*, 2004, **6**, 4682 - 4689.

31 M. Sliwa, N. Mouton, C. Ruckebusch, L. Poisson, A. Idrissi, S. Aloise, L. Potier, J. Dubois, O. Poizat and G. Buntinx, Investigation of ultrafast photoinduced processes for salicylidene aniline in solutions and gas phase: towards a general photo-dynamic scheme, *Photochem. Photobiol. Sci.*, 2010, **9**, 661 - 669.



Novel biphotochromic compound with spironaphthopyran and hydroxyazomethine moieties was synthesized and studied by the nanosecond laser flash photolysis using different excitation wavelengths in methanol and toluene.