



ORGANIC CHEMISTRY

FRONTIERS

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

ARTICLE

The Mitsunobu Reaction in the 21st Century

Cite this: DOI: 10.1039/x0xx00000x

S. Fletcher

Received 00th January 2012,

Accepted 00th January 2012


DOI: 10.1039/x0xx00000x

www.rsc.org/

The Mitsunobu reaction was first described almost fifty years ago and has enjoyed immense popularity since its inception. The purpose of this review is to focus on the more recent advances and applications of Mitsunobu chemistry, particularly from the 1990s to the present day. In addition to a discussion of newer reagents that facilitate purification, we will describe more contemporary applications of this chemistry, especially as applied to the synthesis of pharmaceuticals and their precursors.

Introduction

The Mitsunobu reaction is the dehydrative coupling of a primary or secondary alcohol (occasionally, tertiary alcohols have been used) to a pronucleophile (NuH), which is mediated by the reaction between a dialkyl azodicarboxylate and a trialkyl- or triarylphosphine (Scheme 1). The reaction is named after its discoverer Oyo Mitsunobu who first reported this chemistry in 1967.^{1,2} When chiral, secondary alcohols are employed, complete inversion of stereochemistry is observed in all but a few cases.³ In “conventional” Mitsunobu chemistry, there is a wide range of pronucleophiles that can participate in the reaction. Suitable pronucleophiles include (thio)carboxylic acids, (thio)phenols, imides and sulfonamides, permitting the formation of C-O, C-S and C-N bonds (and C-C bonds as will be seen shortly). It is generally agreed that the p*K*_a of the pronucleophile must be around 12 or below for a successful reaction with the typical Mitsunobu reagents diethyl azodicarboxylate (DEAD; R³ = Et in Scheme 1) or diisopropyl azodicarboxylate (DIAD; R³ = *i*Pr in Scheme 1) and triphenylphosphine (PPh₃). This “p*K*_a rule” stems from the mechanism of the reaction wherein the betaine that results from the reaction between DEAD and PPh₃ has a p*K*_a of about 12 and removes the acidic proton from the pronucleophile, otherwise alkylation of DEAD will occur, which is a known side reaction of the Mitsunobu reaction. The Mitsunobu reaction occurs under mild, essentially neutral conditions, and typically at 0 °C to room temperature. Standard solvents for the reaction include THF, diethyl ether, dichloromethane and toluene, although sometimes more polar solvents including ethyl acetate, acetonitrile and DMF have been used. The Mitsunobu reaction has been reviewed several times before.²⁻⁵ The purpose of this review is to focus on newer applications and more recent innovations associated with Mitsunobu chemistry.



Scheme 1: The Mitsunobu reaction.


Contemporary Mitsunobu Chemistry

Oxygen Pronucleophiles

Perhaps the single, most important feature of the Mitsunobu reaction is that secondary alcohols undergo clean inversion of their stereogenic centres.¹⁻⁵ Thus, hydrolysis of the ester that is yielded from the dehydrative coupling of a carboxylic acid and a secondary alcohol will furnish the inverted alcohol generally in high enantiomeric purity. In addition to carboxylic acids, other oxygen nucleophiles that have become popular coupling partners in the Mitsunobu reaction include phenols and also alcohols themselves in intramolecular reactions. Below are some newer oxygen nucleophiles that have been successfully introduced into this reaction.


Oximes. En route to some *O*-vinyl penicillin derivatives, such as **4**, Stachulski at SmithKline Beecham demonstrated that oximes function well as oxygen nucleophiles (Scheme 2).⁶ In place of the usual DEAD or DIAD, dimethyl acetylenedicarboxylate (DMAD) was used alongside PPh₃. Akin to DIAD, DMAD can also undergo a Michael-type reaction with phosphines to yield an activated betaine. Although the Mitsunobu reaction proceeded in only moderate

yield (64%), it is unclear if attempts were made to optimize the reaction.



Scheme 2. Oximes as oxygen pronucleophiles in the Mitsunobu reaction.


2-Amidophenols: Synthesis of Benzoxazoles. Motivated by the prevalence of the benzoxazole motif in a range of anti-bacterials, anti-parasitics and H₂-antagonists,⁷ Wang and Hauske wanted to explore this pharmacophoric scaffold through combinatorial synthesis on solid phase.⁸ However, the traditional approaches of thermal cyclization of a 2-amidophenol derivative with acidic reagents such as SOCl₂ proved incompatible with a solid support. Thus, the authors looked to the Mitsunobu reaction which they reasoned would accomplish the same goal of cyclodehydration but under mild and neutral conditions. As illustrated in Scheme 3, this new application of Mitsunobu chemistry proceeded in high yield, providing a milder alternative to the existing methods for benzoxazole synthesis.



Scheme 3. 2-Amidophenols undergo cyclodehydrations with DIAD-PPh₃ to furnish benzoxazoles.


2-Ureidophenols: Synthesis of N²-Substituted Benzoxazoles. Yan and colleagues inadvertently took Wang's work further. Initially intending to alkylate the phenol OH of 2-ureidophenol **8**, the isolated product was instead the N²-substituted benzoxazole **9** (Scheme 4).⁹ The authors proposed that the hydroxyl within the -N=COH-NH- tautomeric form of the urea group plays the role of the alcohol, which is turned into a good leaving group during the course of the reaction and is subsequently displaced by the phenoxide ion to generate the benzoxazole scaffold. A second Mitsunobu reaction at elevated temperature then alkylates the exocyclic nitrogen to deliver the observed product **9**. The chemistry appears quite general, and

was also successful with the corresponding tosyl-activated anilines (as a substitute for the phenol OH) to yield the analogous N²-substituted benzimidazoles, for example **11**.




Scheme 4. 2-Ureidophenols and 2-ureidoanilines undergo intramolecular Mitsunobu reactions to deliver N²-substituted benzoxazoles and benzimidazoles, respectively.

Salicylaldoximes: Synthesis of 1,2-Benzisoxazoles. 1,2-Benzisoxazoles are useful species for monitoring the efficiency of artificial enzyme systems since the presence of a general base and an appropriately-positioned hydrogen bond donor will catalyze a ring-opening reaction, provided there is an abstractable hydrogen at the 3-position, to afford the corresponding 2-cyanophenol (salicylonitrile).¹⁰ This β-elimination reaction is referred to as the Kemp elimination,¹¹ and is one of the most widely studied reactions in biomimetic catalysis.¹⁰ A new synthetic route to 1,2-benzisoxazoles (e.g. **13**) has been accomplished by dehydrative ring closure of salicylaldoximes (e.g. **12**) with DEAD-PPh₃, as depicted in Scheme 5.¹² It was noted that the reactions must be run dilute (1 mM) to achieve the good to excellent yields reported. Also, DEAD must be added slowly. Attempts to expand the chemistry to *ortho*-hydroxyarylketoimines as a means of preparing 3-substituted, 1,2-benzisoxazoles were met with mixed results. Methyl and ethyl ketoimines proved excellent substrates but the phenyl ketoxime **14** afforded the corresponding 3-phenyl-1,2-benzisoxazole **15** in only moderate yield (53%). The low yield was partially due to a Beckmann rearrangement that resulted in the isomeric 2-phenylbenzoxazole **16**. Modification of the reaction conditions to reduce formation of the benzoxazole were unsuccessful.




Scheme 5. The intramolecular attack of activated oximes by phenols: synthesis of 1,2-benzisoxazoles.

Salicylaldoximes: Synthesis of Salicylonitriles. In connection with our research program to develop Mcl-1 inhibitors,¹³ we sought more cell penetrable bioisosteres of the carboxylic acid function in our compounds.¹⁴ Exhibiting pK_a values of around 7, we considered that salicylonitriles might serve this role.¹⁵ We reasoned that treating salicylaldoxime with an excess of the Mitsunobu reagents would effect the expected cyclodehydration as in the previous sub-section, and then, by virtue of the basicity of the betaine, a subsequent β -elimination might ensue to afford the corresponding salicylonitrile. Pleasingly, this was found to be the case, as illustrated by the reaction of 5-nitrosalicylaldoxime (**17**) in Scheme 6 to generate 5-nitrosalicylonitrile (**19**) via the 1,2-benzisoxazole **18**.¹⁶ The chemistry is compatible with a variety of functional groups to yield the salicylonitriles in good to excellent yields. Notably, reactions are swift, typically being complete within 30 min at room temperature. Of great significance is the cleanliness of the reactions coupled with the acidity of the products that allowed their isolation through simple acid-base work-ups.




Scheme 6. One-pot synthesis of salicylonitriles: a Mitsunobu-mediated cyclodehydration- β -elimination cascade.

Sulfonic Acids. A group at Bristol-Myers Squibb showed that sulfonic acids are suitable nucleophiles in the Mitsunobu reaction.¹⁷ As with carboxylic acids, the routine inversion of stereochemistry of alcoholic secondary substrates was observed (Scheme 7). The chemistry is scalable, working just as well on a multi-kilo scale.



Scheme 7. Synthesis of sulfonate esters employing sulfonic acids as the pronucleophile.

3-Hydroxyisoxazoles. As already alluded to, carboxylic acids often limit the cellular activity of drug molecules through inhibiting cell penetration. More lipophilic bioisosteres, such as tetrazoles, have served as effective surrogates of carboxylic acids that aid in cell entry.¹⁴ The 3-hydroxyisoxazole motif is also bioisosteric with the carboxylic acid function.¹⁸ What is especially noteworthy about 3-hydroxyisoxazoles is, like the acyl sulfonamides, the acidic group can be functionalized bilaterally.¹⁹ This is in contrast to carboxylic acids and tetrazoles where the acidic group occurs at the terminus of a drug molecule. Chemical transformation of the 3-hydroxyisoxazole scaffold, for example in protecting group strategies, is hampered by its ambidentate nature, and classical alkylation leads to a mixture of N and O functionalizations;²⁰ we have demonstrated that Mitsunobu chemistry predominantly yielded the *O*-alkylated products ($\geq 95\%$ *O*-selectivity in many cases); an example is given in Scheme 8.²¹ Moreover, the Mitsunobu reaction is a more convenient means to functionalize the motif than the classical alkylation procedure: reactions proceeded quickly at room temperature, rather than requiring heat for several days.¹⁹ Also, in the context of methylation, this chemistry avoids the use of explosive diazomethane.




Scheme 8. Regioselective *O*-alkylation of 3-hydroxyisoxazoles under Mitsunobu control.

Nitrogen Pronucleophiles


Sulfonamides, exhibiting pK_a values of about 10, are excellent nucleophiles in the Mitsunobu reaction.²⁻⁵ Azide is also a superb nucleophile and may be administered in a variety of forms: hydrazoic acid, metal azide or diphenylphosphoryl azide (DPPA).² Care must be taken, however, as an excess of PPh_3 might result in transformation of the azide into the corresponding iminophosphorane, which constitutes the first step of the Staudinger reaction to generate primary amines.

Phthalimide: Synthesis of Primary Amines. Protected primary amines can be synthesized from alcohols using phthalimide (**25**) as the nucleophile.²² Reminiscent of the Gabriel synthesis, the resulting *N*-alkylated phthalimide is then subjected to hydrazinolysis to reveal the free primary amine. An example of this chemistry in action is shown in Scheme 9 in which β -naltrexamine derivatives were prepared from dihydrocodeine.




Scheme 9. Phthalimide in the Mitsunobu reaction permits a two-step conversion of alcohols to primary amines.

Nitrobenzenesulfonamides: Synthesis of Secondary Amines. In 1989, Weinreb reported that protected sulfonamides such as BocNHTs are proficient pronucleophiles in the Mitsunobu reaction, in concordance with their acidic pK_a s of around 5.²³ Sequential removal of the Boc protecting group followed by another Mitsunobu reaction permits the synthesis of tertiary sulfonamides, while concomitant removal of both protecting and sulfonyl groups reveals the primary amine. In a similar vein to Weinreb's work, Fukuyama developed a strategy to synthesize secondary amines through the use of nitrobenzenesulfonamides (e.g. **27** and **28**) as both the activating group and protecting group. The first step is a Mitsunobu reaction on the sulfonamide, and the second step is a deprotection of the sulfonyl moiety. Colloquially, this reaction is referred to as the Fukuyama–Mitsunobu reaction, and an example is given in Scheme 10.²⁴ Yields are typically very good to excellent for the Mitsunobu step. When *ortho*- or *para*-nitrobenzenesulfonamides (Ns) are employed, the sulfonyl group can be removed by treatment with thiophenol and K₂CO₃ for 1 h to deliver the secondary amines in very good yields. Use of *ortho,para*-dinitrobenzenesulfonamide (DNs) in place of Ns as the activating/protecting group is also effective and, due to its greater lability, the DN group may be deprotected selectively over the Ns group by stirring with thioglycolic acid in a mildly basic medium.²⁵ Notably, the resulting by-product carrying a carboxylic acid can be removed by a basic wash of the ethereal layer during reaction work-up. Fukuyama later applied his Ns protection/activation strategy to the synthesis of some of the iboga alkaloids.²⁶




Scheme 10. Fukuyama-Mitsunobu synthesis of secondary amines employing thiolate-labile Ns protecting/activating groups.

Activated Hydroxylamines: Synthesis of Ureas. Acylated hydroxylamines are sufficiently acidic to couple to alcohols.²⁸ In the total synthesis of the 5-lipoxygenase inhibitor CMI-977 (Scheme 11), the alcohol of lactone **31** underwent a Mitsunobu reaction with 4-fluorophenol to deliver ether **32**. After further synthetic transformations, resulting molecule **33** was subjected to a second Mitsunobu reaction with *N,O*-bis(phenoxycarbonyl)hydroxylamine (**34**), which, upon aminolysis, furnished CMI-977 (**36**).²⁷




Scheme 11. Synthesis of ureas with activated hydroxylamines.

Synthesis of optically active α,α -disubstituted amino acids. Chiral tertiary α -hydroxy esters can be converted to α -azido esters with hydrazoic acid as the pronucleophile. As complete inversion of the stereochemical configuration is observed, this chemistry provides, after reduction and ester hydrolysis, an efficient and reliable strategy to synthesize α,α -disubstituted amino acids.²⁸ An example is illustrated in Scheme 12. However, it is unclear how scalable is the chemistry for several reasons. First, hydrazoic acid is toxic and potentially explosive. Second, reactions were optimal with ADDP as the azodicarbonyl species but ADDP is not cheap, and 2 equiv are required. And finally, PMe₃ is required as the phosphine partner of the redox system. Notwithstanding its stench, PMe₃ is pyrophoric and must be handled with extreme care. Nevertheless, the total inversion of the tertiary alcohol's chiral centre renders this chemistry a powerful tool in the synthesis of hindered amino acids, at least on a laboratory scale.




Scheme 12. Synthesis of an α,α -disubstituted amino acid from tertiary α -hydroxy esters.

Activated Aminoazoles. Kim and Kahn from Molecumetics Ltd have demonstrated that Boc-protected 2-aminoazoles, such as **41**, are sufficiently activated to smoothly undergo the Mitsunobu reaction to furnish some lysine and arginine analogues (Scheme 13).²⁹ Given their high yields, it appears there was negligible interference, if any, by the aliphatic Boc and Cbz carbamates, which is consistent with their higher pK_a values of around 16-18.




Scheme 13. Alkylation of a Boc-protected/activated 2-aminoazole.

2-Nitrobenzenesulfonylhydrazine: Synthesis of Allenes. The stereospecific synthesis of allenes has been accomplished by Myers and colleagues.³⁰ As shown in Scheme 14, the Mitsunobu amination of a propargyl alcohol with the sulfonyl hydrazide NsNHNH₂ (**44**) afforded the expected product **45**, which was not isolated. Upon allowing the reaction solution to warm to RT, N₂ was lost as was the arylsulfinic acid to generate allene **46**.




Scheme 14. Stereospecific synthesis of an allene.

Activated Guanidines. Kozikowski and Dodd showed that bis-Boc-protected guanidine **47** coupled in excellent yields to a variety of alcohols, delivering, after deprotection, mono-substituted guanidines, such as **49** (Scheme 15, upper panel).³¹ As expected, the reactions proceeded with inversion of configuration. The analogous bis-Cbz-protected guanidine was also a suitable pronucleophile in this reaction. Later, Kim and colleagues at Molecumetics, Ltd. demonstrated that similarly activated guanylation agents, such as **50**, are also capable of coupling to primary and secondary alcohols under Mitsunobu conditions.³² Subsequent reaction with an amine then affords disubstituted guanidines. When suitable alcohols are used, spontaneous cyclization occurs, as shown in Scheme 15 (lower panel).



Scheme 15. Synthesis of substituted guanidines exploiting the Boc protecting group as an activating group.

Purines. *N*⁹-Regioselective alkylation of 2-amino-6-chloropurine was first reported in 2005, although the substrate scope was limited and yields were poor to modest.³³ Shi and colleagues were keen to expand on this work to aid in their discovery of self-assembled nucleoside molecular architectures.³⁴ In particular, the authors focused on the guanine precursor **53**. Like guanine, *O*-carbamate-*N*-acetate protected guanine **53** proved only partially soluble in THF and so, to promote solubility, the Mitsunobu reaction was run at 70 °C. Good to excellent yields of the N9 product were reported, which was confirmed by X-ray crystallography, but the high temperature necessitated two rounds of addition of the Mitsunobu reagents due to their decomposition in the reaction.³⁴ Deprotection of the products with NH₃ in MeOH delivered the desired *N*⁹-substituted guanines (Scheme 16). It is interesting to note that no alkylation at the N2 position of the NHAc group was reported, even at 70 °C, because chemists at Utah showed that this is achievable using the DIAD-triisopropyl phosphite redox system at 50 °C.³⁵




Scheme 16. *N*⁹-Regioselective alkylation of the guanine precursor **53**.


This chemistry is also compatible with a variety of other purine bases, primary and secondary alcohols, and X-ray crystal structures confirmed complete inversion of stereochemistry. Overall, Shi and colleagues introduced a reproducible and high-yielding approach to effect the *N*⁹-regioselective alkylation of purines,³⁴ in contrast to the classical alkylation strategy of R-Hal/K₂CO₃ in DMF that often delivers an approximate 1:1 mixture of N9:N7 products.³⁶

We also required an efficient means of synthesizing N^9 -alkylated guanines towards some novel PNA molecules, and considered that Shi's work might be improved upon by Boc-protecting the N2 position of 2-amino-6-chloropurine. It was predicted this would greatly enhance the solubility of the purine in THF, allowing the Mitsunobu reaction to proceed at room temperature. Moreover, the greater steric bulk of the Boc group versus Ac was postulated to offset any undesired N^2 -alkylation that might have been experienced in Shi's chemistry. Accordingly, N^2 -Boc-2-amino-6-chloropurine (**58**) was prepared and then exposed to Mitsunobu conditions: **58** coupled very efficiently to various primary and secondary alcohols to deliver the N^9 -alkylated products in good to excellent yields at room temperature within 15 min, in stark contrast to Shi's conditions (Scheme 15), and we observed no N^2 -alkylation.³⁷ The products were then subjected to heat in formic acid to effect Boc group deprotection and hydrolysis of the 6-chloro group in one pot to furnish the N^9 -alkylated guanines (e.g. **59**). Note: since our original report of this chemistry, we have found the nature of the NaH to be critical in the Boc transfer reaction from **57** to **58** – NaH in mineral oil provides capricious results, occasionally with little to no Boc transfer, whilst dry NaH consistently drives complete transfer. We extended this chemistry further and showed that the N2 position could be alkylated under modified Mitsunobu conditions, which could then undergo treatment with formic acid to afford N^2,N^9 -disubstituted guanines (e.g. **61**), or nucleophilic aromatic substitution with various amines followed by treatment with TFA to yield N^2,N^6,N^9 -tri-substituted purines, including the CDK inhibitor bohemine (**62**) (Scheme 18).^{37,38}


Later, we similarly activated the N6 position of adenine and adenosines with a Boc group (**63**), and demonstrated it, too, was an effective pronucleophile in the Mitsunobu reaction, providing a milder alternative to the traditional nucleophilic aromatic substitution that is employed to functionalize the 6-position of the purine nucleus (Scheme 19).³⁹ In summary, the practicality, efficiency and adaptability (guanines as well as other purines may be synthesized) suggest this chemistry will be of great use in the regioselective functionalization of nucleobases.



Scheme 17. N^9 -Regioselective alkylation of an alternative and more soluble guanine precursor than **53**.




Scheme 18. The versatility of N^9 -substituted- N^2 -Boc-2-amino-6-chloropurine.




Scheme 19. Alkylation of the exocyclic N^6 -amino group of adenine.

Benzodiazepine-2,5-diones. Given their roles in proteomimicry,^{40,41} coupled with our interest in this area of research to develop new inhibitors of protein–protein interactions, we asked if Mitsunobu chemistry could be used to regio- and chemoselectively alkylate benzodiazepine-2,5-diones (and benzodiazepine-2-ones). Classical alkylation of benzodiazepine-2,5-diones can lead to a mixture of N - and O -alkylations of the anilide function, and an excess of reagents can also result in alkylation of the benzamide group.⁴² Given that K_2CO_3 is sufficiently basic for such alkylations, we rationalized that the anilide NH group must exhibit a pK_a of around 10. Fittingly, subjecting benzodiazepine-2,5-dione **65** to an excess of DIAD, PPh_3 and $BnOH$ resulted in alkylation of the anilide functional group with no detectable alkylation of the benzamide moiety (Scheme 20).⁴³ We conducted a solvent, phosphine and temperature study to identify the optimum conditions for N -alkylation of the anilide functional group. It was discovered that DIAD and PBu_3 in $CHCl_3$ at room temperature afforded the highest ratios of N -alkylation with excellent yields ($\geq 95\%$) observed with primary alcohols. Isopropanol required heating to 40 °C to consume all of the benzodiazepine-2,5-dione, delivering the N -isopropyl product in a yield of 85% and a selectivity of 6.1:1, N:O. This chemistry was also successfully extended to the selective N -alkylation of a 1,4-benzodiazepin-2-one.



Scheme 20. Regio- and chemoselective N^1 -alkylation of benzodiazepine-2,5-diones.

Activated Hydrazones. Chemists at Johnson and Johnson investigated the reactivity of tosyl- and Boc-hydrazones in the Mitsunobu reaction.⁴⁴ Exhibiting a sulfonamide-type acidic NH group, tosyl hydrazones proved excellent substrates, coupling to primary, secondary, aliphatic and benzylic alcohols in mostly very good to excellent yields (Scheme 21). On the other hand, Boc hydrazones were unreactive in the absence of electron-withdrawing groups to lower the pK_a of the BocNH group.




Scheme 21. Alkylation of Ts and Boc hydrazides.

Carbon Pronucleophiles


Carbon-carbon bond forming reactions are highly sought after in the synthesis of natural products and pharmaceuticals, and new strategies to fashion such connections could present a paradigm shift in traditional drug retrosynthesis.⁴⁵ There are limited reports of utilizing Mitsunobu chemistry to effect such transformations; the relatively high pK_a of carbon acids is largely responsible for this.⁴⁶ Falck and Manna were the first to show that this chemistry was possible by using lithium cyanide to displace an activated alcohol to deliver nitrile products.⁴⁷ In more recent years, the Mitsunobu reaction has played a bigger role in fashioning carbon-carbon bonds.

Triethyl methanetricarboxylate. Palmisano's laboratory reported that triethyl methanetricarboxylate (**72**) is adequately acidic to couple to alcohols under Mitsunobu conditions with no erosion of optical purity.⁴⁸ For example, (*S*)-2-methylsuccinic acid (**74**) was prepared by alkylation of (*S*)-ethyl lactate (**71**) followed by acidic hydrolysis and decarboxylation (Scheme 22).




Scheme 22. Carbon-carbon bond formation in the Mitsunobu reaction with the triply-activated methane of triethyl methanetricarboxylate (**72**).

Meldrum's Acid. Meldrum's acid is a versatile synthon in organic synthesis.⁴⁹ Bearing an "anomalously low" pK_a of 4.97,⁵⁰ it was anticipated that Meldrum's acid would be an effective pronucleophile in the Mitsunobu reaction. In fact, with two acidic hydrogen atoms, it can be difficult to control the alkylation step.⁵¹ Although mono-*C*-alkylation failed, Shing *et al.* showed that the *C*-alkylation of mono-substituted Meldrum's acids proceeds in very good yields to furnish the 5,5-disubstituted derivatives (Scheme 23). In addition, double *C*-alkylation of Meldrum's acid is also possible with moderate to very good yields reported (Scheme 24).⁵¹ Notably, 2-phenylethanol and isopropanol did not react with Meldrum's acid. For successful reactions to occur, the more reactive primary allylic and arylmethyl alcohols had to be employed. In reactions with secondary allylic alcohols, a catalytic amount of Pd(0) was required in order to promote the regioselectivity of *C*- over *O*-alkylation.




Scheme 23. Alkylation of C^5 -mono-substituted Meldrum's acids.



Scheme 24. Double C^5 -alkylation of Meldrum's acid.


Bis(2,2,2-trifluoroethyl) Malonates. Although structurally similar to Meldrum's acid, diesters of malonic acids are poor nucleophiles in the Mitsunobu reaction⁴⁶ since the activated methylene protons exhibit a much higher pK_a of about 13.⁵² Takacs rationalized that making the alkyl components of dialkyl malonate esters more electron-withdrawing would lower the pK_a of the activated methylene flanked by the two carbonyls.⁵³ Accordingly, his group prepared the bis(2,2,2-trifluoroethyl) malonate **79**, which underwent *C*-alkylations with primary alcohols in good to excellent yields (Scheme 25),⁵³ with no reported *O*-alkylation. Secondary alcohols

coupled in moderate yield at best. Importantly, however, in addition to the observed regiocontrol, Takacs found that monoalkylation could be controlled, in contrast to analogous chemistry described with Meldrum's acid.⁵¹




Scheme 25. Activation of the malonate methylene function with greater electron-withdrawing 2,2,2-trifluoroethyl groups allows *C*-alkylation to proceed smoothly.

Acetone Cyanohydrin. The direct conversion of primary alcohols to nitriles has been achieved with acetone cyanohydrin (**83**) as the HCN equivalent (Scheme 26).⁵⁴ Szántay and colleagues showed that this application of the Mitsunobu reaction was compatible with secondary alcohols, too, provided they were not sterically hindered. With its avoidance of toxic HCN and metal cyanide salts, this represents a particularly attractive protocol to synthesize organic nitriles.



Scheme 26. Conversion of alcohols to nitriles with acetone cyanohydrin as a latent source of the pronucleophile HCN.

Carbamic Acids. The Mitsunobu reaction has recently been used to trap carbamic acids generated *in situ* as their carbamates, wherein the alcohol may be intramolecular or intermolecular.⁵⁵ Interestingly, the stereochemical course of the Mitsunobu step is dependent on the substitution of the nitrogen atom: when R is carbon, the expected inversion is observed; when R is hydrogen, retention is detected (Scheme 27). The authors have proposed a mechanism that accounts for this deviance. When R = H, the activated carbamic acid undergoes a β -elimination to afford an isocyanate that subsequently reacts with the alcohol to deliver the carbamate. Since attack does not happen at the alcohol's carbon atom, retention is observed.




Scheme 27. Synthesis of carbamates. Note the nature of the R group dictates the stereochemical course of the reaction.

New Azodicarbonyl Species

DEAD or its hydrazine by-product DEAD-H₂ can often contaminate the Mitsunobu product owing to similar *R_f* values.³ In the last couple of decades in particular, new DEAD analogues have emerged that facilitate purification of the reaction mixture, and these are presented in Scheme 28. The majority are commercially available but may, alternatively, be synthesized by reaction of hydrazine with the appropriate chloroformate, followed by oxidation to the target azo species.³

Di-4-chlorobenzyl azodicarboxylate (DCAD). Replacement of the ethyl groups of DEAD with 4-chlorobenzyl groups furnishes the orange solid DCAD, which is more convenient to handle than the liquids DEAD and DIAD, can be stored at room temperature and is almost as effective.⁵⁶ Most of the hydrazine by-product of DCAD generated in the Mitsunobu reaction can be removed by precipitation from dichloromethane, and, owing to a different polarity to the hydrazine by-product of DIAD, purification by column chromatography may be facilitated.



Scheme 28. DEAD/DIAD alternatives to facilitate reaction mixture purification.

Di-4-nitrobenzyl azodicarboxylate (DNAD). Dai and colleagues recently introduced DNAD as another alternative to DEAD and DIAD.⁵⁷ Like DCAD, DNAD is a crystalline solid that is stable at room temperature. DNAD is at least as efficient as DIAD but, significantly, the hydrazine by-product exhibits poor solubilities in CH₂Cl₂ and THF, and so can be precipitated from these solvents.


Di-cyclopentyl azodicarboxylate (DCpAD). DCpAD represents another variant of DEAD and DIAD.⁵⁸ Once again, DCpAD is an orange solid and can be stored at room temperature. DCpAD appears to be as effective as DEAD in the Mitsunobu reaction.

Di-2-methoxyethyl azodicarboxylate (DMEAD). Chemists in Japan devised yet another analogue of DEAD in the guise of DMEAD, wherein the isopropyl groups have been replaced with 2-methoxyethyl moieties.⁵⁹ DMEAD is about as effective as DIAD in Mitsunobu reactions. Importantly, though, the reduced hydrazine by-product can be removed in an aqueous work-up. Moreover, in our hands, excess DMEAD itself can also be extracted into the aqueous phase. Since $\text{Ph}_3\text{P}=\text{O}$ can be removed by filtration, column chromatography may be completely avoided. DMEAD is now commercially available and, although it is somewhat more expensive than DIAD, time and money saved through evasion of chromatography render this new azodicarbonyl affordable.

Di-tert-butyl azodicarboxylate (DTBAD). In contrast to the other DIAD variants, DTBAD's "muscle" comes from its inherent instability to acid, which results in its conversion to gaseous by-products.⁶⁰ The combination of DTBAD with diphenyl(2-pyridyl)phosphine (Ph_2PPy) provides an especially convenient protocol since, in addition to decomposing DTBAD (and its hydrazine by-product), work-up with acid will extract the phosphine (and its oxide) into the aqueous, indicating column chromatography might be avoidable altogether. DTBAD is also commercially available.

Adamantyl-tagged derivatives of DEAD. Curran and co-workers prepared two analogues of DEAD in which the ethyl groups have been replaced with adamantyl-based groups.⁶¹ The azo parents and hydrazine by-products have longer retention times on cyclodextrin-bonded silica gel than do DEAD and DEAD- H_2 , respectively, owing to host-guest chemistry between the adamantyl tags (guests) and the cyclodextrin (hosts). The different retention times facilitates purification of the Mitsunobu products.

***N,N,N',N'*-tetramethyldicarboxamide (TMAD), 1,1'-(azodicarbonyl)dipiperidine (ADDP) and 4,7-dimethyl-3,5,7-hexahydro-1,2,4,7-tetrazocin-3,8-dione (DHTD).** Conversion of the esters of the azodicarboxylates to amides affords more active coupling agents;⁶² upon reaction with phosphines, the betaine intermediates that are generated are more basic and can deprotonate pronucleophiles exhibiting $\text{p}K_{\text{a}}$ values greater than 11. An impressive example of *C*-alkylation of diethyl malonate with 1,1'-(azodicarbonyl)dipiperidine (ADDP), *N,N,N',N'*-tetramethyldicarboxamide (TMAD) and 4,7-dimethyl-3,5,7-hexahydro-1,2,4,7-tetrazocin-3,8-dione (DHTD) demonstrates their superiority over DEAD (Scheme 29).⁶³ An interesting note about ADDP is that it precipitates from a THF reaction medium, and can also be precipitated out by the addition of hexanes or ether.




DEAD-PPh ₃	ADDP-PBu ₃	TMAD-PBu ₃	DHTD-PBu ₃
2%	56%	66%	75%

Scheme 29. Comparison of the efficiencies of Mitsunobu co-reagents in the *C*-alkylation of diethyl malonate.

Azodicarbonyl dimorpholide (ADDM). We recently showed that the morpholine analogue of ADDP, azodicarbonyl dimorpholide (ADDM; Scheme 29), is also an effective azo species for the Mitsunobu reaction.⁶⁴ An important distinction from ADDP, however, is that ADDM and its hydrazine by-product can both be completely removed from the reaction mixture by a neutral, aqueous work-up. Moreover, we demonstrated that the combination of PPh_3 on resin and ADDM resulted in a chromatography-free Mitsunobu reaction.


4,4'-Azopyridine. 4,4'-Azopyridine (**93**) has been introduced as an alternative to DEAD (Scheme 30).⁶⁵ Although reaction times are typically longer, yields are comparable. One of the main advantages of 4,4'-azopyridine over DEAD is that its by-product 4,4'-bispyridine hydrazine largely precipitates from the reaction mixture and so can mostly be removed by filtration. It is noteworthy that the hydrazine by-product can be re-oxidized to the azo form by treatment with iodosobenzene diacetate in DMSO at RT for 6 h.



Scheme 30. 4,4'-Azopyridine (**93**) is a suitable alternative to DIAD.

New Phosphine Species

Contamination of reaction products with triphenylphosphine or triphenylphosphine oxide is arguably the biggest thorn in the Mitsunobu reaction's side.² One of the earliest approaches to simplify the purification of the reaction mixture was to replace triphenylphosphine with tributylphosphine, since excess phosphine may be removed via evaporation and the tributylphosphine oxide is extractable into water during aqueous work-up. Although switching triphenylphosphine with tributylphosphine appears to have little effect on product yield, the latter is pyrophoric and so care should be taken during its use, especially on a large scale. 1,2-bis(diphenylphosphino)ethane (DPPE) has also been used as a surrogate of PPh₃; due to its greater polarity, the resulting bis-oxide precipitates from the reaction mixture and so can be removed by simple filtration prior to chromatography.⁶⁶ Finally, in addition to multiple reports of triphenylphosphine on resin² as a means to facilitate the purification of Mitsunobu reactions, new solution phase phosphine species have also been developed (Scheme 31).




Scheme 31. PPh₃ alternatives that simplify purification of the Mitsunobu reaction.

2-Pyridyl-diphenylphosphine (PPh₂Py). Replacement of one of the phenyl rings with a pyridine ring to give 2-pyridyl-diphenylphosphine (PPh₂Py) is tolerated with no detrimental impact on the reaction yield.⁶⁷ Significantly, however, the introduction of a basic nitrogen allows the removal of excess PPh₂Py and its oxide by-product into the acidic aqueous layer, considerably improving reaction mixture purification.


Tris(dimethylamino)phosphine (Tris-DAP). The introduction of dimethylamino groups into the *para* positions of each of the phenyl rings of PPh₂Py affords tris(dimethylamino)phosphine, or Tris-DAP.⁶⁸ Akin to PPh₂Py, Tris-DAP and its phosphine oxide by-product can also be removed by a simple acidic aqueous work-up.

Phosphoranes/Ylides. Tsunoda and colleagues have prepared cyanomethylenephosphorane (CMMP) and cyanomethylenetriethylphosphorane (CMBP) as “all-in-one” Mitsunobu reagents (Scheme 32). CMMP and CMBP serve as both the azo-type reagent and as the phosphine.^{69,70} The neutral phosphorane deprotonates the pronucleophile to afford a positively charged phosphonium ion, which then turns the alcohol into a good leaving group. A particular strength of these phosphoranes is that they are capable of deprotonating pronucleophiles that are normally not acidic enough for the Mitsunobu reaction. On the other hand, a couple of caveats include their limited commercial availabilities and the need for reaction temperatures in excess of the solvent boiling points, which requires re-sealable reaction vessels.



Scheme 32. Phosphoranes/ylides that function as combined equivalents of phosphines and azodicarbonyls.


4-(Diphenylphosphino)benzoic acid. Sugimura's group recently creatively combined a reducing agent and a pronucleophile in a single compound, 4-(diphenylphosphino)benzoic acid (**96**), which is commercially available.⁷¹ As illustrated in Scheme 33, secondary alcohols coupled to this phosphine in the presence of DMEAD to afford, after saponification, the corresponding inverted secondary alcohols. In addition to exhibiting high stereochemical purities, the compounds were sufficiently chemically pure owing to the extractability of DMEAD/DMEAD-H₂ and 4-(diphenylphosphoryl)benzoic acid into the aqueous layer during work-up.



Scheme 33. Built-in phosphine: inversion of a secondary alcohol with a carboxylic acid that also carries the requisite triarylphosphine functionality.

Organocatalytic Mitsunobu Reaction


One of the drawbacks of the Mitsunobu reaction is poor atom economy. The “Holy Grail” of Mitsunobu chemistry would be to require only catalytic quantities of the azo and phosphine species. Towards this end, Toy and co-workers have introduced a modification to the procedure that requires only 10 mol% of DEAD whose hydrazine by-product is constantly re-oxidized *in situ* by the additive PhI(OAc)₂ (Scheme 34).⁷² However, stoichiometric amounts of PhI(OAc)₂ are required, negating the reduction of DIAD in terms of atom economy (moreover, prices of PhI(OAc)₂ and DIAD are comparable), and acetic acid is generated during oxidation of DEAD-H₂, which could inadvertently participate in the Mitsunobu reaction as the pronucleophile. This represents an exciting piece of work in Mitsunobu history, and it is hoped that further improvements will be made in this area soon.



Scheme 34. Towards an organocatalytic Mitsunobu reaction. Note the sub-stoichiometric use of DIAD.

Other Strategies to Facilitate Purification

Fluorous Mitsunobu Reaction. Curran introduced fluorous reverse phase silica gel at the turn of this Century, and demonstrated that materials partition over the silica at different rates according to their fluorine content.⁷³ His group later applied this as a means to facilitate purification of Mitsunobu reactions.⁷⁴ Although F-DEAD-1 and F-TPP, the fluorous analogues of DEAD and PPh₃, respectively (Scheme 35), successfully promoted the coupling of alcohols to acidic pronucleophiles, yields were typically very low. Whilst this shortcoming may be considered offset by the improved purification of the reaction mixture through fluorous solid-phase extraction (FSPE) relative to a non-fluorous Mitsunobu reaction, Curran and colleagues sought to improve the efficiency of their fluorous reagents. Accordingly, a library of second generation fluorous azodicarboxylates was prepared.⁷⁵ Pleasingly, F-DEAD-2 and F-DEAD-3 demonstrated greatly improved reactivities, which has been attributed to the propyl spacer in place of an ethyl spacer that renders the resulting betaine sufficiently basic to deprotonate the pronucleophile.




Scheme 35. Fluorous Mitsunobu reagents to facilitate purification of the reaction mixture.

Hydrogen peroxide. The use of a 15 wt% solution of hydrogen peroxide (H₂O₂) has been described to oxidize all remaining PPh₃ to Ph₃P=O.⁷⁶ Interestingly, the hydrazine by-product of DIAD did not get oxidized back to DIAD. Nevertheless, this exposure to H₂O₂ significantly facilitated purification with the requirement of column chromatography simply over a plug of silica gel. It is anticipated that reactions performed on a large scale will particularly benefit from this purification strategy.


Miscellaneous

Sonication promotes the Mitsunobu reaction of phenols with sterically encumbered alcohols. He and Lepore have demonstrated that sonication affords dramatic rate enhancements in the coupling of alcohols to phenols in which either or both reaction partners is/are sterically hindered (Scheme 36).⁷⁷ The authors note that the increased rate of reaction might be related to the generation of free radicals, which suggests there may be an erosion of stereospecificity in the Mitsunobu reaction; this was not explored further. Also, the reader is cautioned against performing this reaction on a multi-gram scale, as the heat generated by sonication might cause the azodicarboxylate species to explode. Nevertheless, the chemistry appears useful for the milligram scale.




Scheme 36. Sonication assists in difficult Mitsunobu etherifications.

Reductive Deoxygenation. The traditional approach to accomplish the reductive deoxygenation of alcohols involves first the acylation of the alcohol with thiocarbonyldiimidazole followed by reductive cleavage mediated by tri-*n*-butyltin hydride.⁷⁸ Myers and colleagues have introduced a single-step procedure to achieve the same goal.⁷⁹ Particularly, conducting the Mitsunobu reaction with *o*-nitrobenzenesulfonylhydrazine (**44**) at -30 °C yields the corresponding 1-alkyl-1-sulfonylhydrazine. Warming to 0 °C resulted in gentle extrusion of *o*-nitrobenzenesulfinic acid and the concomitant formation of a diazene intermediate that slowly lost N₂ gas to furnish the deoxygenated product, which appears to occur through a free-radical mechanism (Scheme 37).⁷⁹ The chemistry works well for sterically unencumbered substrates.⁷⁹



Scheme 37. Deoxygenation of alcohols to alkanes using *o*-nitrobenzenesulfonylhydrazine **44**.

Synthesis of Thiocyanates (and Isothiocyanates). Introduction of NH₄SCN into the Mitsunobu protocol permits the conversion of alcohols, thiols, carboxylic acids, as well as TMS ethers and esters, to their corresponding thiocyanates and isothiocyanates, as depicted in Scheme 38. Yields are good, although 2 eq of DEAD and PPh₃ are required to consume all of the starting materials.⁸⁰



Scheme 38. Synthesis of thiocyanates and isothiocyanates with the thiocyanate anion as the nucleophile.

Conclusions

The Mitsunobu reaction between an alcohol and a pronucleophile has become a standard chemical reaction in many laboratories, which is largely due to its mildness, occurring essentially under neutral conditions, speed, often many reactions are complete within 15 min, and clean inversion of the alcohol's stereocentre. The scope has been expanded

beyond the traditional pronucleophiles to encompass oximes, purines and β -ketoesters to name but a few, permitting the construction of C-O, C-N and C-C bonds, respectively. At the same time as the expansion of the pronucleophile cache, research into new reagents to facilitate purification of the reaction mixture has been rife. In particular, new azodicarbonyls, such as DMEAD and ADDM, may be extracted into water during a standard work-up, whilst PPh_2Py may be extracted into the acidic aqueous. Some progress has also been made towards a catalytic Mitsunobu reaction, widely regarded as the "Holy Grail" of Mitsunobu chemistry. We hope that further advancements are made in this regard as it would address two caveats of the Mitsunobu reaction: problematic purification and poor atom economy. Finally, an area of Mitsunobu chemistry that would benefit greatly from further research is C-C bond formation, as this has been largely neglected.

Acknowledgements

The University of Maryland School of Pharmacy is acknowledged for supporting our work in this area.

Notes and references

^a Department of Pharmaceutical Sciences, University of Maryland School of Pharmacy, 20 N. Pine St., Baltimore, MD 21201, USA

^b University of Maryland Greenebaum Cancer Center, 22 S. Greene St., Baltimore, MD 21201, USA.

- O. Mitsunobu, M. Yamada, *Bull. Chem. Soc. Jpn.*, 1967, **40**, 2380.
- O. Mitsunobu, *Synthesis*, 1981, 1.
- K. C. Kumara Swamy, N. N. Bhuvan Kumar, E. Balaraman, K. V. P. Pavan Kumar, *Chem. Rev.*, 2009, **109**, 2551.
- D. L. Hughes, *Org. Prep.*, 1996, **28**, 127.
- T. Y. S. But, P. H. Toy, *Chem. Asian J.*, 2007, **2**, 1340.
- A. V. Stachulski, *J. Chem. Soc., Perkin Trans. 1*, 1991, 3065.
- J. W. Westly, J. W. Liu, J. F. Blount, L. H. Sello, N. Troupe, P. A. Miller, *J. Antibiotic.*, 1983, **36**, 1275; R. D. Hsugwitz, R. G. Angel, G. A. Jacobs, B. V. Maurer, V. L. Narayanan, L. R. Cruthers, J. Szanto, *J. Med. Chem.*, 1982, **25**, 969; Y. Katsura, S. Nishino, Y. Inoue, M. Tomoi, H. Tskasugi, *Chem. Pharm. Bull.* 1992, **40**, 371.
- Wang, F.; Hauske, J. R. *Tetrahedron Lett.*, 1997, **38**, 6529.
- Y. Yan, Q. Zhong, N. Zhao, G. Liu, *Mol. Divers.*, 2012, **16**, 157.
- D. R hlisberger, O. Khersonsky, A. M. Wollacott, L. Jiang, J. DeChancie, J. Betker, J. L. Gallaher, E. A. Althoff, A. Zanghellini, O. Dym, S. Albeck, K. N. Houk, D. S. Tawfik, D. Baker, *Nature*, 2008, **453**, 190.
- M. L. Casey, D. S. Kemp, K. G. Paul, D. D. Cox, *J. Org. Chem.*, 1973, **38**, 2294.
- G. Poissonnet, *Synth. Commun.* 1997, **27**, 3839.
- L. Chen, M. E. Lanning, S. Fletcher, *Aust. J. Anal. Pharm. Chem.*, 2014, **1**, 1015.
- C. Ballatore, D. M. Huryn, A. B. Smith 3rd, *ChemMedChem*, 2013, **8**, 385.
- W. M. Haynes, Ed. *CRC Handbook of Chemistry and Physics*, 95th ed.; CRC Press: Boca Raton, FL, 2014–2015.
- E. Whiting, M. E. Lanning, J. A. Scheenstra, S. Fletcher, *J. Org. Chem.*, 2015, **80**, 1229.
- N. G. Anderson, D. A. Lust, K. A. Colapret, J. H. Simpson, M. F. Malley, J. Z. Gougoutas, *J. Org. Chem.*, 1996, **60**, 7955.
- L. Matzen, A. Engesgaard, B. Ebert, M. Didriksen, B. Fr lund, P. Krogsgaard-Larsen, J. W. Jaroszewski, *J. Med. Chem.*, 1997, **40**, 520.
- B. Fr lund, A. T. Jorgensen, L. Tagmose, T. B. Stensbol, H. T. Vestergaard, C. Engblom, U. Kristiansen, C. Sanchez, P. Krogsgaard-Larsen, T. Liljefors, *J. Med. Chem.*, 2002, **45**, 2454.
- G. Schlewier, P. Krogsgaard-Larsen, *Acta Chem. Scand. B*, 1984, **38**, 815.
- L. Chen, S. Fletcher, *Tetrahedron Lett.*, 2014, **55**, 1693.
- C. Simon, S. Hosztafi, S. Makleit, *Tetrahedron*, 1994, **50**, 9757.
- J. R. Henry, L. R. Marcin, M. C. McIntosh, R. M. Scola, G. D. Harris, Jr., S. M. Weinreb, *Tetrahedron Lett.*, 1989, **30**, 5709.
- T. Fukuyama, C.-K. Jow, M. Cheung, *Tetrahedron Lett.*, 1995, **36**, 6373.
- T. Kan, T. Fukuyama, *Chem. Commun.*, 2004, 353.
- S. Kobayashi, G. Peng, T. Fukuyama, *Tetrahedron Lett.*, 1999, **40**, 1519.
- X. Cai, M. S. Chorghade, A. Fura, G. S. Grewal, K. A. Jauregui, H. A. Lounsbury, R. T. Scannell, C. G. Yeh, M. A. Young, S. Yu, *Org. Proc. R&D*, 1999, **3**, 73.
- J. E. Green, D. M. Bender, S. Jackson, M. J. O'Donnell, J. R. McCarthy, *Org. Lett.*, 2009, **11**, 807.
- H.-O. Kim, M. Kahn, *Synlett*, 1999, 1239.
- A. G. Myers, B. Zheng, *J. Am. Chem. Soc.*, 1996, **118**, 4492.
- K. S. Dodd, A. P. Kozikowski, *Tetrahedron Lett.* 1994, **35**, 977.
- H.-O. Kim, F. Matthew, C. Ogbu, *Synlett*, 1999, 193.
- Y. Kitade, T. Ando, T. Yamaguchi, A. Hori, M. Nakanishi, Y. Ueno, *Bioorg. Med. Chem.*, 2006, **14**, 5578; M. Yang, W. Ye, S. W. Schneller, *J. Org. Chem.*, 2004, **69**, 3993.
- W. Lu, S. Sengupta, J. L. Petersen, N. G. Akhmedov, X. Shi, *J. Org. Chem.* 2007, **72**, 5012.
- E. A. V  iz, P. A. Beal, *Tetrahedron Lett.*, 2006, **47**, 3153.
- J. Kjellberg, N. G. Johansson, *Tetrahedron*, 1986, **42**, 6541.
- S. Fletcher, V. M. Shahani, P. T. Gunning, *Tetrahedron Lett.*, 2009, **50**, 4258.
- S. Fletcher, V. M. Shahani, A. J. Lough, P. T. Gunning, *Tetrahedron*, 2010, **66**, 4621.
- S. Fletcher, *Tetrahedron Lett.*, 2010, **51**, 2948.
- W. C. Ripka, G. V. De Lucca, A. C. Bach II, R. S. Pottorf, J. M. Blaney, *Tetrahedron*, 1993, **49**, 3593.
- M. D. Cummings, C. Schubert, D. J. Parks, R. R. Calvo, L. V. LaFrance, J. Lattanze, K. L. Milkiewicz, T. Lu, *Chem. Biol. Drug Des.*, 2006, **67**, 201.
- B. E. Blass, T. M. Burt, S. Liu, D. E. Portlock and E. M. Swing, *Tetrahedron Lett.*, 2000, **41**, 2063.
- K.-Y. Jung, S. Fletcher, *Med. Chem. Commun.*, 2012, **3**, 1160.
- J. M. Keith, L. Gomez, *J. Org. Chem.*, 2006, **71**, 7113.
- E. J. Corey, X.-M. Cheng, X.-M. The Logic of Chemical Synthesis (Wiley, 1995).
- M. Wada, O. Mitsunobu, *Tetrahedron Lett.* 1972, **13**, 1279.
- S. Manna, J. R. Falck, *Synth. Commun.*, 1985, **15**, 663.

Journal Name

- 1 48 G. Cravotto, G. B. Giovenzana, M. Sisti, G. Palmisano, *Tetrahedron*,
2 1996, **52**, 13007.
- 3 49 Y. Oikawa, H. Hirasawa, O. Yonemitsu, *Tetrahedron Lett.*, 1978, **19**,
4 1759.
- 5 50 S. Nakamura, H. Hirao, T. Ohwada, *J. Org. Chem.*, 2004, **69**, 4309.
- 6 51 T. K. M. Shing, L.-H. Li, K. Narkunan, *J. Org. Chem.*, 1997, **62**,
7 1617.
- 8 52 R. G. Pearson, R. L. Dillon, *J. Am. Chem. Soc.* 1953, **75**, 2439.
- 9 53 J. M. Takacs, Z. Xu, X.-T. Jiang, A. P. Leonov, G. C. Theriot, *Org.*
10 *Lett.*, 2002, **4**, 3843.
- 11 54 M. C. Aesa, G. Ba n, L. Nov k, C. Sz ntay, *Synth. Commun.*, 1996,
12 **26**, 909.
- 13 55 C. J. Dinsmore, S. P. Mercer, *Org. Lett.*, 2004, **6**, 2885.
- 14 56 B. H. Lipshutz, D. W. Chung, B. Rich, R. Corral, *Org. Lett.*, 2006, **8**,
15 5069.
- 16 57 J. Yang, L. Dai, X. Wang, Y. Chen, *Tetrahedron*, 2011, **67**, 1456.
- 17 58 J. Yang, L. Dai, X. Z. Wang, Y. Chen, *Chin. Chem. Lett.*, 2011, **22**,
18 1047.
- 19 59 K. Hagiya, N. Muramoto, T. Misaki, T. Sugimura, *Tetrahedron*,
20 2009, **65**, 6109.
- 21 60 M. Kiankarimi, R. Lowe, J. R. McCarthy, J. P. Whitten, *Tetrahedron*
22 *Lett.*, 1999, **40**, 4497.
- 23 61 S. Dandapani, J. J. Newsome, D. P. Curran, *Tetrahedron Lett.*, 2004,
24 **45**, 6653.
- 25 62 T. Tsunoda, Y. Yamamiya, S. It , *Tetrahedron Lett.*, 1993, **34**, 1639.
- 26 63 T. Tsunoda, M. Nagaku, C. Nagino, Y. Kawamura, F. Ozaki, H.
27 Hioki, S. It , *Tetrahedron Lett.*, 1995, **36**, 2531.
- 28 64 M. E. Lanning, S. Fletcher, *Tetrahedron Lett.*, 2013, **54**, 4624.
- 29 65 N. Iranpoor, H. Firouzabadi, D. Khalili, S. Motevalli, *J. Org. Chem.*,
30 2008, **73**, 4882.
- 31 66 I. A. O'Neil, S. Thompson, C. L. Murray, S. B. Kalindjian,
32 *Tetrahedron Lett.*, 1998, **39**, 7787.
- 33 67 D. Camp, I. D. Jenkins, *Aust. J. Chem.* 1988, **41**, 1835.
- 34 68 G. Grynkiewicz, J. Jurczak, A. Zamojski, *Tetrahedron*, 1975, **31**,
35 1411.
- 36 69 T. Tsunoda, F. Ozaki, S. It , *Tetrahedron Lett.*, 1994, **35**, 5081.
- 37 70 T. Tsunoda, C. Nagino, M. Oguri, S. It , *Tetrahedron Lett.*, 1996, **37**,
38 2459.
- 39 71 N. Muramoto, K. Yoshino, T. Misaki, T. Sugimura, *Synthesis*, 2013,
40 **45**, 931.
- 41 72 T. Y. But, P. H. Toy, *J. Am. Chem. Soc.*, 2006, **128**, 9636.
- 42 73 D. P. Curran, S. Hadida, A. Studer, M. He, S.-Y. Kim, Z. Luo, M.
43 Larhed, M. Hallberg, B. Linciau, "Fluorous Synthesis: A User's
44 Guide", In *Combinatorial Chemistry: A Practical Approach*; H.
45 Fenniri, Ed.; Oxford Univ Press: Oxford, 2000; **2**; 327.
- 46 74 S. Dandapani, D. P. Curran, *Tetrahedron*, 2002, **58**, 3855.
- 47 75 S. Dandapani, D. P. Curran *J. Org. Chem.*, 2004, **69**, 8751.
- 48 76 A. J. Proctor, K. Beaument, J. M. Clough, D. W. Knight, Y. Li,
49 *Tetrahedron Lett.*, 2006, **47**, 5151.
- 50 77 S. D. Lepore, Y. He, *J. Org. Chem.*, 2003, **68**, 8261.
- 51 78 D. H. R. Barton, S. W. J. McCombie, *Chem. Soc., Perkin Trans.*
52 *1*, 1975, 1574.
- 53 79 A. G. Myers, M. Movassaghi, B. Zheng, *J. Am. Chem. Soc.*, 1997,
54 **119**, 8572.
- 55 80 N. Iranpoor, H. Firouzabadi, B. Akhlaghinia, R. Azadi, *Synthesis*,
56 2004, 92.