

RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Dissociative ionization and Coulomb explosion of ethyl bromide under a near-infrared intense femtosecond laser field

Chong Teng, Hua Wu, Jian Zhang, Yan Yang, Tianqing Jia, Shian Zhang[#] and Zhenrong Sun[#]

State Key Laboratory of Precision Spectroscopy, and Department of Physics, East China Normal University, Shanghai 200062, People's Republic of China

[#]Correspondence to sazhang@phy.ecnu.edu.cn and zrsun@phy.ecnu.edu.cn

The multi-photon dissociation and Coulomb explosion of ethyl bromide C_2H_5Br under near-infrared (800 nm) femtosecond laser field are experimentally investigated by a DC-sliced ion imaging technique. The sliced images of fragment ions $C_2H_5^+$, Br^+ , CH_3^+ , CH_2Br^+ , H_2^+ and $C_2H_3Br^+$ are obtained, and their dissociative pathways are assigned by observing their corresponding kinetic energy release (KER) and angular distribution. It is shown that low-KER components of these fragment ions result from multi-photon dissociation of singly charged parent ion $C_2H_5Br^+$, while high-KER components come from Coulomb explosion of doubly charged parent ion $C_2H_5Br^{2+}$. It is also shown that the precursor specie $[C_2H_5^+...Br^+]$ has the longer lifetime than $[C_2H_3Br^+...H_2^+]$ and $[CH_3...CH_2Br^+]$. In addition, the probable H_2 and H_2^+ elimination channels are theoretically simulated by Gaussian 09 software packages, and the results show that the former is an asynchronous concerned process while the latter is a synchronous concerned process.

Keywords: dissociative ionization, Coulomb explosion, femtosecond laser pulses, ab initio calculation, laser-induced elimination

I. Introduction

When the atoms or molecules are exposed to a femtosecond laser field with the laser intensities in the range of $10^{13}\sim 10^{15}$ W/cm², this intense femtosecond laser field will modify the potential energy surface (PES) of the electronic state in the atomic or molecular systems, and thus a variety of novel physical phenomena will occur, such as multi-photon ionization (MPI) [1-4], above threshold ionization (ATI) [5, 6], high-order harmonic generation (HHG) [7, 8], Coulomb explosion (CE) [9-12], and so forth. On the basis of these related phenomena, one can obtain the atomic or molecular dynamical processes. Alkyl bromide C₂H₅Br was usually taken as study example in the field of molecular dynamics under the intense laser field, and its photodissociation behaviors have been regarded as the reference for understanding photodissociation dynamics of polyatomic molecules [13-15]. In the photodissociation process of alkyl bromide, the dominant dissociation channel is C-Br bond fission to form an alkyl cation and a ground or spin-orbit excited state bromine atom [16-20], and such observation has an important significance on the photochemical study of atmospheric bromine. Zhang et al. have experimentally studied the photodissociation dynamics of C-Br bond fission in a series of alkyl bromide by ultraviolet (UV) laser excitation [13, 21-23], and shown that fragments Br* (²P_{1/2}) and Br (²P_{3/2}) can be attributed to the perpendicular transition to the ¹Q₁ and ³Q₁ states and the parallel transition to the ³Q₀ state. In addition to C-Br bond cleavage, Xu et al. have demonstrated other photodissociation pathways of ethyl bromide cation C₂H₅Br⁺ based on 355-nm laser excitation [24], and concluded that both C₂H₅Br⁺ → C₂H₃⁺ + H₂ + Br and C₂H₅Br⁺ → C₂H₅⁺ + HBr channels are produced through multi-center intermediates according to the anisotropy parameter. Gardiner et al. also confirmed above two dissociation channels by 266 and 355-nm

laser excitations [25], and theoretically simulated the lowest thermochemical threshold of accessible photofragmentation pathways from ethyl bromide cation $C_2H_5Br^+$, involving C-C, C-Br and C-H bond fissions.

In these previous studies, fragmentation dynamics of ethyl bromide by C-Br, C-C or C-H bond cleavage were usually performed by UV laser excitation, while studying them by near-infrared (IR) laser field was seldom reported. In this work, we present an investigation on the photodissociation process of ethyl bromide C_2H_5Br by 800-nm intense femtosecond laser field based on a DC sliced ion imaging technique. These fragment ions H_2^+ , CH_3^+ , $C_2H_5^+$, Br^+ , CH_2Br^+ and $C_2H_3Br^+$ by C-C, C-Br and C-H bond fissions are observed, and low-KER components of these fragment ions are attributed to multi-photon dissociation (MPD) of singly charged parent ion $C_2H_5Br^+$ while high-KER components are assigned to Coulomb explosion (CE) of doubly charged parent ion $C_2H_5Br^{2+}$. Furthermore, elimination channels of H_2 and H_2^+ in MPD and CE processes are also experimentally observed, and the theoretical simulations show that the former is an asynchronous concerned process while the latter is a synchronous concerned process.

II. Experimental arrangement

Our experimental setup of DC sliced ion imaging system has been described in our earlier publications [26, 27], and here the simple statement is given. The multi-stage ion lens installed in the main chamber has the similar geometry designed by Suits et al. [28]. According to ion trajectory simulation by Simon 8.0 software, the voltages applied on these electrodes are set to be $U_{\text{Repeller}}=2000$ V, $U_1=1768$ V, $U_2=1661$ V and $U_3=0$ V, respectively. The ethyl bromide vapor seeded in helium gas at a backing pressure of 0.3 atm is expanded into the source chamber through a pulsed

valve with a repetition rate of 100 Hz, and the supersonic molecular beam is skimmed before entering the reaction chamber. The femtosecond laser outputted from a regenerative amplification laser system has the center wavelength of 800 nm, pulse width of about 72 fs and repetition rate of 1 kHz, and the linearly polarized femtosecond laser pulse is focused into the reaction area by a lens with the focal length of 400 mm. The ethyl bromide is ionized and dissociated by the femtosecond laser field in the reaction chamber. These fragment ions are accelerated by multi-stage ion lens velocity apparatus and detected by a pair of 77 mm micro-channel plates (MCP) coupled to a P47 phosphor screen, and finally the sliced images of the fragment ions are recorded by an intensified charge-coupled device (ICCD) camera. To identify the fragment ion species, the time-of-flight (TOF) mass spectra are measured by a photomultiplier tube connected to a digital oscilloscope. In this experiment, all the timing sequence is controlled by Stanford Instrument Digital Delay/Pulse Generation (DG535).

III. Results and discussion

Figure 1 shows three typical TOF mass spectra of ethyl bromide C_2H_5Br with the laser intensities of 1.8×10^{13} (a), 4.9×10^{13} (b) and 1.5×10^{14} W/cm^2 (c), here the laser polarization direction is kept to be perpendicular to TOF axis. Since there are two different isotopes of bromine (^{79}Br and ^{81}Br) in nature, those fragment ions that contain bromine atom will exhibit multiple peak structure. The fragment ions $C_2H_m^+$ ($m=2-5$) and singly charged parent ion $C_2H_5Br^+$ are observed with the laser intensity of 1.8×10^{13} W/cm^2 (see Fig. 1(a)), which means that C-Br bond fission is the dominant dissociative channel in the lower laser intensities. When the laser intensity is increased to 4.9×10^{13} W/cm^2 , these fragment ions H^+ , H_2^+ , CH_n^+ ($n=0-3$) and Br^+ are observed (see Fig.1(b)), and this illustrates the participation of C-C and C-H bond

fissions. Here, the fragment ion H_3^+ is not observed, which may be due to the more complicated formation process comparing with the fragment ion H_2^+ , which includes the breakage of three C-H bonds and formation of three H-H bonds, and therefore its formation need more demanding experimental condition. When the laser intensity is further increased to $1.5 \times 10^{14} \text{ W/cm}^2$, highly charged fragment ions C^{2+} and Br^{2+} are observed (see Fig.1(c)), which shows that the CE process will occur in the higher laser intensities. In this CE process, the ionization probability can be greatly enhanced when the bond length of the molecular ions elongates to a critical distance, which triggers the Coulomb repulsion, and then the Coulomb repulsive energy accelerates the fragmentation process [29, 30]. Therefore, the fragment ions produced by the Coulomb explosion process are easy to be observed in the higher laser intensities. In this work, our goal is to explore the photodissociation process via C-Br, C-C and C-H bond fissions.



DC sliced images can provide the related information of fragment ions, involving the KER and angular distributions. To further investigate the fragmentation dynamics of ethyl bromide under the intense femtosecond laser field, DC sliced images of fragment ions C_2H_5^+ , Br^+ , CH_3^+ , CH_2Br^+ , H_2^+ and $\text{C}_2\text{H}_3\text{Br}^+$ via C-Br, C-C and C-H bond fissions are obtained with the laser intensity of $1.1 \times 10^{14} \text{ W/cm}^2$, and their corresponding velocity distributions are calculated, as shown in Fig. 2. Here, all velocity distributions are fitted with Gaussian function, and the peak velocities and kinetic energy releases are also labeled. It is noted that the fragment ion $\text{C}_2\text{H}_3\text{Br}^+$ cannot be observed in the mass spectrum because of its weak signal intensity, but it can be identified by our dc-sliced ion imaging system with high detection efficiency and sensitivity. In general, the fragment ions with low KER come from the MPD process of parent ions, while those with high KER come from the CE process [26, 31].

Thus, these fragment ions with low KER in Fig. 2 should result from the MPD process of singly parent ion $C_2H_5Br^+$, and these corresponding MPD channels can be assigned as follows

C-Br bond cleavage:



C-C bond cleavage:



C-H bond cleavage:




Under the intense femtosecond laser field, the ethyl bromide can also be ionized to the multi-charged parent ions, and then dissociates into fragment ions with high KER by Coulomb repulsive force. As well known, in the two-body CE model, the two fragment ions produced by the two-body CE process should have a relationship as follows [26, 27, 31, 32]

$$\frac{KER(X^{p+})}{KER(Y^{q+})} = \frac{M(Y^{q+})}{M(X^{p+})} \quad (7)$$


where X and Y denote the two fragment ions, M is mass of fragment ions, and p and q are charge numbers of the ions. According to Eq. (7), the two fragment ions Br^+ (1.29 eV) and $C_2H_5^+$ (3.55 eV) can be attributed to the CE channel as follows



Utilizing the same method, the two fragment ions CH_3^+ (3.85 eV) and CH_2Br^+ (0.59 eV) by C-C bond cleavage are also attributed to the following CE channel



Similarly, the two fragment ions H_2^+ (8.79 eV) and $\text{C}_2\text{H}_3\text{Br}^+$ (0.16 eV) by C-H bond cleavage can be assigned to such a CE channel



To further investigate the CE process of doubly charged parent ion $\text{C}_2\text{H}_5\text{Br}^{2+}$, the angular distributions of these fragment ions from DC sliced images in Fig. 2 are extracted, and the calculated results are shown in Fig. 3. These experimental data are theoretically fitted by Legendre polynomial function [33], and such a function is given as follows

$$I(\theta) = 1 + \sum_L a_L P_L(\cos^2\theta) \quad (L=2, 4, 6) \quad (11)$$

where θ is the ejection angle of fragment ions measured from the laser polarization direction, which represents the instantaneous spatial distribution at the moment of CE process, a_L is the Legendre expansion coefficient, and the expectation value $\langle \cos^2\theta \rangle$ is the characterization of fragmentation anisotropy. The values of a_L and $\langle \cos^2\theta \rangle$ are summarized in Table 1. It is noted that the two fragment ions that come from the same CE channel show the similar expectation value $\langle \cos^2\theta \rangle$ (i.e., fragmentation anisotropy), which can further confirm above photodissociation channel assignments in the CE process.,

As known to all, when the short-lived precursor specie rotates as a whole in the space, its corresponding fragment ions will have stronger angular anisotropy. In Table 1, the leading expansion coefficient a_2 is an important anisotropy parameter, which characterizes the lifetime of the precursor specie state, and the precursor specie with

the longer lifetime will induce the smaller a_2 value of the fragment ions [33-35]. In our experiment, the a_2 value by C-Br bond fission is determined to be $a_2=0.46$, which is smaller than that by C-C bond fission ($a_2=0.61$) as well as C-H bond fission ($a_2=0.61$). That is to say, the precursor specie $[C_2H_5^+ \dots Br^+]$ has the longer lifetime than $[C_2H_3Br^+ \dots H_2^+]$ and $[CH_3 \dots CH_2Br^+]$. This means that the precursor specie $[C_2H_5^+ \dots Br^+]$ is prepared in a deeper well of potential energy surface, and dissociates into the fragment ions $C_2H_5^+$ and Br^+ by Coulomb repulsive force following by the C-Br bond stretch.

As shown in Eqs. (5), (6) and (10), the elimination of H_2 or H_2^+ is the most complicated dissociative process that includes two C-H bond fissions and H-H bond formation. To understand how the fragment H_2 or H_2^+ is produced from the parent ion $C_2H_3Br^+$ or $C_2H_5Br^{2+}$, the probable photodissociation pathways are theoretically simulated by Gaussian 09 software packages [36]. The geometry of parent ions, transition states (TS) and dissociative products are optimized at the MP2 level with a 6-311++G(d,p) basis set. Meanwhile, the energies of related fragment ions are further refined by CCSD/6-311++G (d,p) with MP2 level zero-point energy corrections (ZPE). Figure 4 shows the structure evolution during the formation of H_2 from the singly charged parent ion $C_2H_3Br^+$. The dissociation pathways of H_2 elimination from the same C atom and two C atoms are both calculated at the same theoretical level, and it is found that the energy barrier of H_2 elimination from two C atoms is much higher than from the same C atom, and so here only the H_2 elimination from the same C atom is presented. As shown in Fig. 4, the optimized geometry of parent ion $C_2H_3Br^+$ is with C-H bond lengths of 1.10 Å, $\angle H-C-H$ of 109° and H-H distance of 1.78 Å. The elimination of H_2 starts with the C-H bond stretch to form TS. In the TS structure, two C-H bond lengths are elongated asymmetrically to 1.78Å and 1.55 Å,

$\angle\text{H-C-H}$ is quickly decreased to 26° , and the distance between two H atoms is reduced to 0.79 \AA , which is much shorter than that of parent ion $\text{C}_2\text{H}_5\text{Br}^+$. After surpassing TS with energy of 2.30 eV , the singly charged parent ion $\text{C}_2\text{H}_5\text{Br}^+$ dissociates into two parts $\text{C}_2\text{H}_3\text{Br}^+$ and H_2 . The values of the three main parameters $\angle\text{H-C-H}$, two C-H bond lengths and H-H distance along reaction coordinates are listed in Table 2. As can be seen, $\angle\text{H-C-H}$ is decreased, H-H distance is shortened, and the two C-H bond lengths are asymmetrically elongated due to the uneven distribution of surface charges in the two H atoms, which indicate that the $\text{C}_2\text{H}_5\text{Br} \rightarrow \text{H}_2 + \text{C}_2\text{H}_3\text{Br}^+$ elimination channel is an asynchronous concerted dissociation.

The energy path of H_2^+ elimination channel from doubly charged parent ion $\text{C}_2\text{H}_5\text{Br}^{2+}$ is shown in Fig. 5. It is noted that the optimized geometry of $\text{C}_2\text{H}_5\text{Br}^{2+}$ is quite different from the structure of $\text{C}_2\text{H}_5\text{Br}^+$. The two C-H bonds are stretched to 1.37 \AA , which are increased by 0.27 \AA comparing with $\text{C}_2\text{H}_5\text{Br}^+$. Meanwhile, the distance between two H atoms is much closer, only 0.87 \AA , which is almost half of the stable structure of $\text{C}_2\text{H}_5\text{Br}^+$. Such an observation indicates that Coulomb repulsive force will greatly change the molecular structure when the two electrons are stripped off under the intense femtosecond laser field. The elimination of H_2^+ also starts with the C-H bond stretch. In the TS structure, $\angle\text{H-C-H}$ is decreased from 37° to 26° , the two C-H bond lengths are symmetrically elongated to 1.61 \AA , and H-H distance is reduced to 0.81 \AA . After surpassing TS, the doubly charged parent ion $\text{C}_2\text{H}_5\text{Br}^{2+}$ dissociates into the fragment ions H_2^+ and $\text{C}_2\text{H}_3\text{Br}^+$. It is noticeable that the energy of TS is lower (0.03 eV) than that of the doubly charged parent ion $\text{C}_2\text{H}_5\text{Br}^{2+}$, and this phenomenon has been demonstrated in previous studies [37, 38]. That is to say, the doubly charged parent ion $\text{C}_2\text{H}_5\text{Br}^{2+}$ is trapped in a very shallow quasi-bound well of the potential

energy surface, and can produce a barrierless reaction process. This conclusion can also be confirmed by the experimental observation that no doubly charged parent ion $C_2H_5Br^{2+}$ is observed in the full range of laser intensity. The values of the three main parameters $\angle H-C-H$, two C-H bond lengths and H-H distance are listed in Table 3. It is noted that the two C-H bonds are simultaneously broken and H-H bond is simultaneously formed, which shows that H_2^+ elimination from the doubly charged parent ion $C_2H_5Br^{2+}$ is a synchronous concerted process.

IV. Conclusions

In summary, we have experimentally and theoretically studied the photodissociation process of ethyl bromide C_2H_5Br under the near-IR (800nm) intense femtosecond laser field by the DC-slice imaging technique. These fragment ions $C_2H_5^+$, Br^+ , CH_3^+ , CH_2Br^+ , H_2^+ and $C_2H_3Br^+$ by C-C, C-Br and C-H bond fissions were measured and their corresponding KER and angular distributions were calculated. We confirmed that these fragment ions with low KER are attributed to the MPD process of singly charged parent ion $C_2H_5Br^+$, while those components with high KER result from doubly charged parent ion $C_2H_5Br^{2+}$ via the CE process. The angular anisotropy of the fragment ions by the CE process suggested that the precursor specie $[C_2H_5^+...Br^+]$ has a longer lifetime than $[C_2H_3Br^+...H_2^+]$ and $[CH_3...CH_2Br^+]$. Moreover, the elimination channels of H_2 and H_2^+ were observed, and ab initio calculations showed that H_2 elimination is an asynchronous concerted reaction while H_2^+ elimination is a synchronous concerted reaction.

Acknowledgements

This work was partly supported by National Natural Science Fund (Nos. 51132004 and 11474096), and Shanghai Municipal Science and Technology Commission (No.

14JC1401500).

References

- [1] P. B. Corkum, Plasma Perspective on Strong-Field Multiphoton Ionization, *Phys. Rev. Lett.* 71, 1994 (1993).
- [2] J. E. Bayfield, P. M. Koch, Multiphoton Ionization of Highly Excited Hydrogen Atoms, *Phys. Rev. Lett.* 33, 258 (1974).
- [3] A. J. Gotch, T. S. Zwier, Multiphoton Ionization Studies of Clusters of Immiscible Liquids. I. $C_6H_6-(H_2O)_n$, $n=1, 2$. *J. Chem. Phys.* 96, 3388 (1992).
- [4] M. Castillejo, S. Couris, E. Koudoumas, M. Martín, Subpicosecond Ionization and Dissociation of Benzene and Cyclic Alkanes at 800 and 400 nm. *Chem. Phys. Lett.* 289, 303 (1998).
- [5] A. G. Suzor, X. He, O. Atabek, Above-Threshold Dissociation of H_2^+ in Intensity Laser Fields. *Phys. Rev. Lett.* 64, 515 (1990).
- [6] R. R. Freeman, P. H. Bucksbaum, H. Milchberg, S. Darack, D. Schumacher, M. E. Geusic, Above-Threshold Ionization with Subpicosecond Laser Pulses. *Phys. Rev. Lett.* 59, 1092 (1987).
- [7] J. J. Macklin, J. D. Kmetec, C. L. Gordon, High-Order Harmonic Generation Using Intense Femtosecond Pulses. *Phys. Rev. Lett.* 70, 766 (1993).
- [8] I. P. Christov, M. M. Murnane, H. C. Kapteyn, High-Harmonic Generation of Attosecond Pulses in the “Single-Cycle” Regime. *Phys. Rev. Lett.* 78, 1251 (1997).
- [9] X. G. Xue, C. Wu, Y. R. Liu, W. Huang, Y. K. Deng, Y. Q. Liu, Q. H. Gong, C. Y. Wu, Identifying Isomers of Carbon-dioxide Clusters by Laser-Driven Coulomb Explosion. *Phys.*

- Rev. A* 90, 033411 (2014).
- [10] K. Harumiya, H. Kono, Y. Fujimura, Intense Laser-field Ionization of H₂ Enhanced by Two-electron Dynamics. *Phys. Rev. A* 66, 043403 (2002).
- [11] J. P. Nibarger, S. V. Menon, G. N. Gibson, Comprehensive Analysis of Strong-field Ionization and Dissociation of Diatomic Nitrogen. *Phys. Rev. A* 63, 053406 (2001).
- [12] S. Das, P. M. Badani, P. Sharma, R. K. Vatsa, D. Das, A. Majumder, A. K. Das, Multiphoton Ionization and Coulomb Explosion of C₂H₅Br Cluster: A Mass Spectrometric and Charge Density Study. *Rapid Commun. Mass Spectrom.* 25, 1028 (2010).
- [13] B. F. Tang, R. S. Zhu, Y. Tang, L. Ji, B. Zhang, Photodissociation Dynamics of C₂H₅Br and n-C₃H₇Br in UV Region. *Chem. Phys.* 303, 37 (2004).
- [14] R. Montero, A. P. Conde, A. Longarte, F. Castaño, M. E. Corrales, R. de Nalda, L. Bañares, Femtosecond Time-resolved Photophysics and Photodissociation Dynamics of 1-Iodonaphthalene. *Phys.Chem.Chem.Phys.* 12, 7988 (2010).
- [15] A. Giuliani, F. M. Tollet, J. Delwiche, N. J. Mason, N. C. Jones, J. M. Gingell, I. C. Walker, Franksin, M. J. H., Electronic Excitation and Oscillator Strength of Ethyl Bromide by Vacuum Ultraviolet Photoabsorption and Electron Energy Loss Spectroscopy. *J. Chem. Phys.* 112, 6285 (2000).
- [16] B. Urban, V. E. Bongybey, One-color Multiphoton Threshold Photoelectron Spectra of Methyl Bromide, and Their Comparison with Methyl Iodide. *J. Chem. Phys.* 116, 4938 (2002).
- [17] A. G. Sage, T. A. A. Oliver, G. A. King, D. M. Murdock, J. N. Harvey, M. N. R. Ashfold, UV Photolysis of 4-iodo-, 4-bromo-, and 4-chlorophenol: Competition between C–Y (Y = halogen) and O–H Bond Fission. *J. Chem. Phys.* 138, 164318 (2013).
- [18] T. Gougousi, P. C. Samartzis, T. N. Kitsopoulos, Photodissociation Study of CH₃Br in the

- First Continuum. *J. Chem. Phys.* 108, 5742 (1998).
- [19] S. Eden, P. L. Vieira, S. V. Hoffmann, N. J. Mason, VUV Photoabsorption in CF₃X (X = Cl, Br, I) Fluoro-Alkanes. *Chem. Phys.* 323, 313 (2005).
- [20] P. Zou, J. N. Shu, T. J. Sears, G. E. Hall, S. W. Northe, Photodissociation of Bromoform at 248 nm: Single and Multiphoton Processes. *J. Phys. Chem. A* 108, 1482 (2004).
- [21] C. H. Zhang, Z. Z. Cao, L. Q. Hua, Y. Chen, B. Zhang, Study on the Photodissociation Mechanisms of m-Bromotoluene at 234 and 267 nm Using Velocity Ion Imaging. *Chem. Phys. Lett.* 454, 171 (2008).
- [22] P. J. Liu, B. F. Tang, B. Zhang, Velocity Map Imaging of the Photolysis of n-C₄H₉Br in UV Region. *Chem. Phys.* 340, 141 (2007).
- [23] Y. M. Wang, S. Zhang, Q. S. Zheng, B. Zhang, C-Br bond fission dynamics in Ultraviolet Photodissociation of 1,2-dibromoethane. *Chem. Phys. Lett.* 423, 106 (2006).
- [24] D. D. Xu, R. J. Price, J. H. Huang, W. M. Jackson, Photodissociation of the Ethyl Bromide Cation at 355 nm by Means of TOF-MS and Ion Velocity Imaging Techniques. *Phys. Chem.* 215, 253 (2001).
- [25] S. H. Gardiner, T. N. V. Karsili, M. L. Lipciuc, E. Wilman, M. N. R. Ashfold, C. Vallance, Fragmentation Dynamics of the Ethyl Bromide and Ethyl Iodide Cations: A Velocity-Map Imaging Study. *Phys. Chem. Chem. Phys.* 16, 2167 (2014).
- [26] Y. Yang, L. L. Fan, S. Z. Sun, J. Zhang, Y. T. Chen, S. A. Zhang, T. Q. Jia, Z. R. Sun, Dissociation Double Ionization of 1-Bromo-2-Chloroethane Irradiated by an Intense Femtosecond Laser Field. *J. Chem. Phys.* 135, 064303 (2011).
- [27] H. Wu, Y. Yang, S. Z. Sun, J. Zhang, L. Deng, S. A. Zhang, T. Q. Jia, Z. G. Wang, Z. R. Sun, Concerted Elimination of Br₂⁺ Resulting from the Coulomb Explosion of 1,2-Dibromoethane

- in an Intense Femtosecond Laser Field. *Chem. Phys. Lett.* 607, 70 (2014).
- [28]D. Townsend, M. P. Miniti, A. G. Suits, Direct Current Slice Imaging. *Re. Sci. Instrum.* 74, 2530 (2003).
- [29]D. Normand, M. Schmidt, Multiple Ionization of Atomic and Molecular Iodine in Strong Laser Fields. *Phys. Rev. A* 53, R1958 (1995).
- [30]I. Bocharova, R. Karimi, E. F. Penka, J. P. Brichta, P. Lassonde, X. Q. Fu, J. C. Kieffer, A. D. Bandrauk, I. Litvinyuk, J. Sanderson, F. Légaré, Charge Resonance Enhanced Ionization of CO₂ Probed by Laser Coulomb Explosion Imaging. *Phys. Rev. Lett.* 107, 063201 (2011).
- [31]H. Wu, S. A. Zhang, Y. Yang, S. Z. Sun, J. Zhang, L. Deng, T. Q. Jia, Z. G. Wang, Z. R. Sun, Coulomb Explosion and Dissociative Ionization of 1,2-Dibromoethane under an Intense Femtosecond Laser Field. *RSC Adv.* 4, 45300 (2014).
- [32]J. Zhang, S. A. Zhang, Y. Yang, S. Z. Sun, H. Wu, J. Li, Y. T. Chen, T. Q. Jia, Z. G. Wang, F. A. Kong, Z. R. Sun, Photodissociation of Br₂ Molecules in an Intense Femtosecond Laser Field. *Phys. Rev. A* 90, 053428 (2014).
- [33]T. Okino, Y. Furukawa, P. Liu, T. Ichikawa, R. Itakura, K. Hoshina, K. Yamanouchi, H. Nakano, Coincidence Momentum Imaging of Ejection of Hydrogen Molecular Ions From Methanol in Intense Laser Fields. *Chem. Phys. Lett.* 419, 223 (2006).
- [34]S. Z. Sun, Y. Yang, J. Zhang, H. Wu, Y. T. Chen, S. A. Zhang, T. Q. Jia, Z. G. Wang, Z. R. Sun, Ejection of Triatomic Molecular Ion H₃⁺ from Methyl Chloride in an Intense Femtosecond Laser Field. *Chem. Phys. Lett.* 581, 16 (2013).
- [35]Y. Furukawa, K. Hoshina, K. Yamanouchi, H. Nakano, Ejection of Triatomic Hydrogen Molecular Ion from Methanol in Intense Laser Fields. *Chem. Phys. Lett.* 414, 117 (2005).
- [36]M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G.

Scalmani, V. Barone, B. Mennucci, G. A. Petersson, et al. Gaussian 09, revision B.01; Gaussian, Inc.; Wallingford, CT, 2010.

[37] Y. X. Zhao, X. N. Wu, J. B. Ma, S. G. He, X. L. Ding, Experimental and Theoretical Study of the Reactions between Vanadium-Silicon Heteronuclear Oxide Cluster Anions with n-Butane. *J. Phys. Chem. C* 114, 12271 (2010).

[38] X. N. Wu, Y. X. Zhao, W. Xue, Z. C. Wang, S. G. He, X. L. Ding, Active Sites of Stoichiometric Cerium Oxide Cations ($\text{Ce}_m\text{O}_{2m}^+$) Probed by Reactions with Carbon Monoxide and Small Hydrocarbon Molecules. *Phys. Chem. Chem. Phys.* 12, 3984 (2010).

Table 1 Legendre expansion coefficients a_L ($L=2, 4, 6$) and expectation value $\langle \cos^2\theta \rangle$ of fragment ions $C_2H_5^+$, Br^+ , CH_3^+ , CH_2Br^+ , H_2^+ and $C_2H_3Br^+$ by Coulomb explosion.

Channels	Ions	a_2	a_4	a_6	$\langle \cos^2\theta \rangle$
(7)	$C_2H_5^+$	0.46	0.15	0.07	0.35
	Br^+	0.46	0.15	0.06	0.35
(8)	CH_3^+	0.59	0.10	0.03	0.41
	CH_2Br^+	0.61	0.04	-0.05	0.40
(9)	H_2^+	0.59	-0.01	0.05	0.39
	$C_2H_3Br^+$	0.61	0.04	-0.02	0.39

Table 2 Molecular structure evolution from the singly charged parent ion $C_2H_3Br^+$ characterized by the three main parameters $\angle(H-C-H)$, two C-H bond lengths and H-H distance along the reaction coordinates calculated at MP2/6-311++G(d,p) level.

$\angle H-C-H$ (°)	C-H bond lengths (Å)	H-H distance (Å)
109	1.10, 1.10	1.78
77	1.12, 1.16	1.42
46	1.20, 1.32	0.99
36	1.30, 1.48	0.88
26	1.55, 1.78	0.79
25	1.60, 1.83	0.78
21	1.91, 2.12	0.75
16	2.46, 2.61	0.74

Table 3 Molecular structure evolution from the doubly charged parent ion $C_2H_5Br^{2+}$ characterized by the three main parameters ($\angle(H-C-H)$, two C-H bond lengths and H-H distance) along the reaction coordinates calculated at MP2/6-311++G(d,p) level.

$\angle(H-C-H)$ (°)	C-H bond lengths (Å)	H-H distance (Å)
37	1.37, 1.37	0.87
36	1.40, 1.40	0.86
33	1.48, 1.48	0.84
31	1.52, 1.52	0.83
26	1.61, 1.61	0.81
25	1.80, 1.80	0.78
21	2.07, 2.07	0.76
17	2.50, 2.50	0.75

Figure captions

Figure 1. TOF mass spectra of ethyl bromide C_2H_5Br under the femtosecond laser field with the laser intensities of 1.8×10^{13} (a), 4.9×10^{13} (b) and 1.5×10^{14} W/cm^2 (c).

Figure 2. DC sliced images of fragment ions $C_2H_5^+$ (a), Br^+ (b), CH_3^+ (c), CH_2Br^+ (d), H_2^+ (e) and $C_2H_3Br^+$ (f) with the laser intensity of 1.1×10^{14} W/cm^2 , together with their corresponding KER distributions ((g)~(l)). The open circles (○) denote the experiment data, the short dot lines (.....) are the fitting Gaussian peaks, and the solid lines (—) are the simulated results.

Figure 3. Angular distributions of fragment ions $C_2H_5^+$ (a), Br^+ (b), CH_3^+ (c) and CH_2Br^+ (d). The open circles (○) are experimental data and the solid lines (—) are the simulated results by Legendre polynomial function.

Figure 4. The dissociation pathway for H_2 elimination from the singly charged parent ion $C_2H_5Br^+$, and here the theoretical simulation is performed at MP2/6-311++G (d,p) level.

Figure 5. The dissociation pathway for H_2^+ elimination from the doubly charged parent ion $C_2H_5Br^{2+}$, and here the calculation level is MP2/6-311++G (d,p).

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5

