



Labelling of brassinosteroids by isotopes of hydrogen and carbon

Journal:	<i>RSC Advances</i>
Manuscript ID:	RA-REV-03-2015-004081.R1
Article Type:	Review Article
Date Submitted by the Author:	24-Apr-2015
Complete List of Authors:	Patil, Mahadeo; Jain University, centre for nano and material science, Jain University, centre for nano and material science Elbert, Tomáš; Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, v.v.i., Flemingovo nám. 2, 16610 Prague 6, CZECH REPUBLIC., Laboratory of radioisotopes Keri, Rangappa; Jain University, centre for nano and material science; Jain University, , Centre for Nano and Material Sciences

Labelling of brassinosteroids by isotopes of hydrogen and carbon

Mahadeo R. Patil^a, Tomáš Elbert^{*b} and Rangappa S. Keri^{*a}

^aCentre for Nano and Material Sciences, Jain University, Jain Global Campus, Bangalore 562112, Karnataka, INDIA.

^bInstitute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, v.v.i., Flemingovo nám. 2, 16610 Prague 6, CZECH REPUBLIC.

Abstract

The brassinosteroids (BRs) are a class of native plant growth regulating substances with high biological activity even at very low concentration. These compounds have been rigorously explored and found that they are not only growth regulators in plants but also are promising antiviral agents. Recently, it has been reported that natural BRs exhibit relatively interesting anticancer activities. Up to now basic anticancer potential of BRs against several normal and human cancer cell lines has been determined. Natural BRs, at micromolar concentrations, impart cell growth-inhibitory responses in several human cancer cell lines without affecting the normal cells. To study the mechanism of action of BRs at molecular level, the corresponding isotopically labelled compounds are essential. The latter BRs are essential for the investigation of biosynthesis, metabolism, transport and distribution in plants. This venture ultimately led us to explore the labeling of BRs by isotopes of hydrogen and carbon and all about the related technique. The present review will shed light to provide a glimpse on the synthetic avenues in this field from the time of their discovery of labelled BRs till their recent advancements.

Keywords: Labelled brassinosteroids, epibrassinolide, castasterone, deuterium, tritium, carbon isotopes.

*Corresponding author: elbert@uochb.cas.cz (Dr. T. Elbert)
keriphd@gmail.com, sk.rangappa@jainuniversity.ac.in (Dr. R. S. Keri)

Content

1. Introduction
2. Synthesis of deuterium-labelled BRs
 - 2.1. BRs with deuterium label in the side chain
 - 2.2. BRs with deuterium label in ring A
 - 2.3. BRs with deuterium in ring B
3. Synthesis of tritium-labelled BRs
4. Synthesis of ^{14}C -labelled BRs
5. Summary
6. Acknowledgement
7. References
8. List of Abbreviations

1. Introduction

Brassinosteroids (BRs) represent a class of naturally occurring phytohormones with various physiological activities and ubiquitous distribution in the plant kingdom.¹ The vicinal diol grouping on ring A is typical for BR-plant hormones discovered thirty years ago.² In Figure 1, the formulae of two typical BRs- 24-epibrassinolide **1** and castasterone **2** - are given. These compounds have been intensively studied and found that they exhibit not only growth regulation functions in plants but also some promising antiviral activities.³




Figure 1 Structures of 24-epibrassinolide (24-epiBL) and 24-epicastasterone (24-epiCS)

Recently, molecular studies directed towards the essential role of BRs in plant growth and development⁴ and their chemical synthesis, biological mode of action, and practical application in agriculture and horticulture^{5,6} have been greatly intensified. The essentialities of isotopically labelled BRs in the investigation of the biosynthesis, metabolism, transport and distribution of endogenous BRs in plants have been documented. In such studies, BRs labelled with isotopes of hydrogen are most frequently used to explain the biosynthesis, metabolism and mode of action at a molecular level.

Therefore, the numerous strategies for the synthesis of BRs labelled with deuterium (²H or D) and tritium (³H or T) in the side chain or in the ring system have been developed.⁷ BRs are formed by a biosynthetic network of alternative pathways and sub-pathways. A number of

feeding experiments using labelled BRs precursors as a substrate are necessary to elucidate these pathways. In these reactions, labelled BRs are being used as internal standards for qualitative analysis of endogenous BRs and they have been employed in biosynthetic experiments.⁸⁻¹¹ Some of the deuterium-labelled 24-methylenecholesterol and related C₂₈ steroids were used as both, internal standards in quantitative analyses and substrate, for metabolic studies in BRs deficient mutants of *Arabidopsis thaliana* and *Pisumsativum*. The biosynthetic sequence from campesterol¹² to campestanol in *Arabidopsis thaliana* was determined by identification of each intermediate followed by feeding experiments with deuterium-labelled intermediates.^{13,14} The dwarf pea (*Pisumsativum*) mutants Ika and Ikb are BRs insensitive and deficient, respectively. The latter mutant was rescued to wild type by exogenous application of labelled brassinolide and its precursors. Feeding experiments using deuterium labelled 24-methylenecholesterol indicated that an Ikb mutant is unable to isomerize and/or reduce the double bond.

Recently, brassinolide biosynthetic pathways have been elucidated by feeding deuterium-labelled intermediate to suspension cultures of *Catharanthus roseus*.^{15,16} Identification of cathasterone with one of the hydroxyl groups lacking and application of its labelled analogue in feeding experiments⁹ demonstrated that the brassinolide biosynthesis proceeds *via* initial hydroxylation at C-22 followed by an introduction of the hydroxyl group at C-23. Deuterium labelled of secasterol, teasterone and typhasterol, upon administration to rye seedlings, were incorporated into secasterone and 2,3-diepisecasterone, indicating a biosynthetic route *via* teasterone/typhasterol to secasterol to 2,3-epoxybrassinosteroids, secasterone in seedlings of *Secale cereal*. Similarly, deuterated secasterone upon administration resulted in deuterated castasterone and 2-epicastasterone used in biosynthetic sub-pathways from typhasterol/teasterone *via* 2,3-epoxybrassinosteroids intermediate to castasterone (CS).¹⁷ Some of the ¹⁴C labelled BRs,

namely (22*R*, 23*R*)-and (22*S*, 23*S*)-[4-¹⁴C]-24-epiBL (A and B in Scheme 11, respectively) were used to facilitate metabolic and distribution studies of (22*R*, 23*R*)-24-epiBL and (22*S*, 23*S*)-24-epiBL as well as to study their role in the growth of grain and vegetables as they are promising candidates for agriculture application.¹⁸ Predominantly the ¹⁴C-labelled epiBL (Scheme 10) was used in the uptake and the transport study of exogenously applied epibrassinolide on seedlings of cucumber and wheat. When applied to roots, ¹⁴C-epiBL was readily taken up and swiftly transported throughout both the plant species. When ¹⁴C-epiBL was applied to the adaxial surface of a young cucumber leaf, it was readily taken up, however, transported very slowly compared with the previous case. In wheat leaves, ¹⁴C-epiBL was transported only in the apical direction from the treated spot after 3 days of treatment; however, it was not transported from the treated leaf to the other leaves or organs even after seven days. Recently, isotopically labelled [7,7-²H₂] epibrassinolide was used for biosynthetic transformation studies.¹⁹ This compound found useful for biochemical and physiological investigation in the plant.²⁰ Simultaneously, it was also discovered that natural BRs exhibit relatively interesting anticancer activities. So far, potential anticancer activities of 24-epiBL **1** and 24-epicastasterone **2** on several human cancer cell lines have been determined.¹⁹ 24-Episcasterol was prepared and found cytotoxic against human breast carcinoma MCF-7 (Michigan Cancer Foundation) cells.²¹ It was also demonstrated that non-plant cells, yeast WAT21, generates a steroidal plant hormone castasterone. To understand how castasterone is generated in WAT21 cells, deuterium labelled 6-deoxo-[26,28-²H₆]teasterone(TE), 6-deoxo-[26,28-²H₆]typhasterol (TY) and 6-deoxo-[26,28-²H₆](CS) were fed to WAT21 cells and their metabolites were isolated. [26,27-²H₆] labelling of brassinolide, castasterone, typhasterol, and teasterone were reported by Takatsuto and Ikekawa in 1986.²² Deuterium labelled 6-deoxo-BRs were identified as biosynthetic precursors of CS in WAT21

cells.²³ In this review, a variety of labelling procedures of BRs with stable and radioactive isotopes of hydrogen and carbon will be presented.


2. Synthesis of deuterium-labelled BRs

2.1. BRs with deuterium labels in the side chain


2.1.1. [26-²H₃], [26,27-²H₆] and [26,28-²H₆] labelled BRs

Brassinolide (BL) and related C-28 BRs as castasterone (CS), typhasterol (TY), and teasterone (TE) occur in a wide variety of higher plants.^{24,25} The synthesis of [26,28-²H₆] brassinolide **1**, [26,28-²H₆]CS **2**, [26,28-²H₆]TY **3** and [26,28-²H₆]TE **4** as an internal standards for GC-MS assays of BRs was reported in the literature.^{26,27} [26,28-²H₆]crinosterol²⁸ **5** served as a starting material for the synthesis of four deuterated BRs (**1-4**) (Scheme 1).

Alternatively, labelled BRs containing three or six deuterium atoms appended in the terminal methyl groups of the side chain (in a position ensuring lack of isotopic exchange) were prepared from stigmasterol or bisnorcholenic acid. There are main strategies for the construction of the side chain containing an asymmetric centre at C-24 of the compound **18** and these are based on the coupling of **16**-aldehyde with an appropriate chiral sulfone synthon **17**. These methodologies were also used for the stereoselective construction of the side chain labelled BRs. Preparation of trideuterated fragment **22** in the side chain of BRs^{29,30} through Claisen rearrangement for stereoselective construction of the steroidal side chain. Construction of the side chain of hexa-deuterated BR was performed *via* coupling of sulfone **17** with aldehyde **16** (Scheme 2)



Scheme 1 Synthesis of [26,28-²H₆]BL (**1**), [26,28-²H₆]CS (**2**), [26,28-²H₆]TY (**3**) and [26,28-²H₆]TE (**4**)



Scheme 2 Construction of the side chain of hexa-deuterated BRs


Introduction of functional groups at the cyclic part of the steroids could be done prior to the construction stages of the side chain based on various 22-aldehydes.^{31,32} As a result, the aldehyde **25** was the best choice for the synthesis of labelled BR derivatives with functional groups tethered to the cyclic part and it was suitable for the preparation of many highly functionalized BRs. The aldehyde **25** was prepared either from stigmasterol **24** by traditional method^{33,34} in 3 steps or from 23,24-bisnorcholeonic acid³⁵ **26** (Scheme 3). An important part of the convergent synthesis was the preparation of the chiral intermediate **17**, which was

accompanied through (2*R*)-3-hydroxy-2-methylpropanoate.^{35,36} The synthesis of precursor of [26-²H₃]-brassinosteroids **34** for biochemical studies from the starting material **25** is depicted in Scheme 3. In the first convergent strategies based on the Claisen rearrangement of a single isomer of allylic alcohol is considered useful in this rearrangement to access certain product.^{27,28,37,38} A Claisen rearrangement has been widely used in the preparation of Δ^{22} -steroids containing an alkyl substituent with predictable stereochemistry at C-24. Then the ester compounds **32** were prepared *via* isomeric acetylenic alcohols **28** and **30** and the allylic alcohol intermediates **29** and **31** (Scheme 3). Successive reduction of the ester **32** with LiAlD₄ followed by tosylation and deuteride reduction gave compound **33** containing three deuterium atoms in the terminal part of the side chain.³⁰ Regeneration of the cyclic part of **33** by acid treatment to yield the deuterated crinosterol **34** (Scheme 3).


After formation of a 3-membered ring and oxidation on C-6 by standard sequence of reactions starting from the **34** and then the Sharpless oxidation of side chain double bond using (DHQD)₂-PHAL as a ligand and osmium tetroxide as catalyst led to expected alcohol and followed by acetylation to obtained the compound **35** and then to bromide **36**. Nucleophilic substitution of the bromide in **36** offered 3 α -acetoxyketone **37** and after deacetylation [26-²H₃]TY **38** was obtained. Baeyer-Villiger oxidation of **37** yielded lactone **39** along with its regioisomer **40**. Deprotection of the hydroxyl groups furnished 2-dehydroxy-[26-²H₃] brassinolide **41** (Scheme 4).

Bromide **36** found useful as an intermediate for the preparation of deuterated BRs with an epoxy group in ring A (Scheme 5). Dehydrobromination of **36** offered 6-ketone **42** as the main product of the reaction with formic ester **43** as a byproduct. Epoxidation of **42** ensued stereoselectively to yield 2 α ,3 α -epoxyacetate **44** that was then deprotected to 2,3-[26-

$^2\text{H}_3$]episcasterone **47**. Compound 2,3-[$^{26}\text{-}^2\text{H}_3$] secasterone **48** was obtained *via* bromohydrin **45** from **42**. Formate **43** can be used for the preparation of 3,6-diketo BRs **49** via 3-dehydroteasterone **46** (Scheme 5).




Scheme 3 Construction of the side chain of [$^{26}\text{-}^2\text{H}_3$]BRs




Scheme 4 Synthesis of [26-²H₃]TY **38** and related compounds

The epoxide **44** was used as key intermediate for the synthesis of [26-²H₃] BRs **53-55** as depicted in Scheme 6.³⁰ *Trans*-diaxial ring opening of epoxide **44** with the action of HBr led to bromohydrin **50**. Jones oxidation ensued with the formation of a ketonic group at C-3 and inversion of configuration at C-2 (bromoketone **51**). After nucleophilic substitution of bromine by a hydroxyl group, the hydroxyl ketone **52** was isolated. The regiospecific and stereoselective reduction of **52** followed by saponification to remove acetate groups in the side chain furnished a mixture of 3-epi[26-²H₃]CS **53** and [26-²H₃]CS **55**. Bayer-Villiger oxidation of **53** with trifluoroperoxoacetic acid offered 3-[26-²H₃]epiBL **54**.


An important convergent approach using the side chain fragment was used for the preparation of the labelled BRs containing six deuterium atoms at C-26 and C-27.^{35,36} Coupling of the appropriate aldehyde with lithium salt of chiral sulfone led to compounds **56** (Scheme 7).



Scheme 5 Synthesis of deuterated BRs with an epoxy group in ring A




Scheme 6 Synthesis of deuterated tetra-hydroxyl BRs



Scheme 7 Construction of the side chain of [26,27-²H₆]BRs


Then alcohol **56** is functionalized by modified Julia olefination protocol *via* acetylation followed by reduction elimination with magnesium amalgam to furnish the olefin **57**. Introduction of a (22*R*, 23*R*)-diol function was attained by using asymmetric dihydroxylation (AD) as discussed in the earlier case for [26-²H₃] BRs. Synthesis of a derivative **61** demonstrated the methodology applied in the preparation of BRs containing one hydroxyl group in the side chain. Swern oxidation of hydroxyl sulfones **56** directed to the ketosulfones **59** and then desulfurization with aluminium amalgam was smoothly resulted in the formation of the ketones **60**. The further reduction of **60** with lithium aluminium hydride to offered desired (22*S*)-alcohols **61** as the main product (Scheme 7). Mono- and dihydroxy derivatives **62** and **63**, respectively allowed easy access to a variety of

corresponding [26,27-²H₆]BRs as shown in Scheme 8. Δ^5 -3 β Alcohols **64** and **65** were synthesized from **62** and **63** in a dioxane-water solution in presence of toluenesulfonic acid at elevated temperature. Hydrogenation of double bond over palladium bequeathed 6-deoxoteasterone **67** and 6-deoxocathasterone **68**. The preparation of 3 α -alcohol **66** involved inversion configuration at C-3 *via* nucleophilic substitution of intermediate mesylate with potassium superoxide.



Scheme 8 Modification of rings A and B of [26,27-²H₆]BRs

An attempt to prepare labelled brassinolide biosynthetic precursors having 3,6-diketone moiety was first reported with by-products similar to **42** obtained from dehydrobromination of 3 β -bromides (see also Scheme 5). It is proved incompatible with the removal of the acetate protecting groups in the side chain and the problem was solved by using aldehyde **69** as a key intermediate (Scheme 9). Essentially, the same protocol as that described in Scheme 7 was used for the preparation of 3-dehydro-[26,27-²H₆]cathasterone **72** and 3-dehydro-[26,27-²H₆] TE **73**.³⁵



Scheme 9 Synthesis of 3,6-diketo-[26,27- $^2\text{H}_6$] BRs

2.1.2 Synthesis of deuterio-labelled 24-methylenecholesterol and related steroids

Anastasia *et al.* reported the synthesis of BRs with one deuterium atom,³⁹ however, usually four and more deuterium atoms in the molecule are required for the MS standards. Contemporarily, Takatsuto *et al.* reported labelling of the side chain of targeted steroid by more than one deuterium

atom. It is described in the work that the synthesis of deuterium labelled C₂₈ steroids **80** to **86** from the known 3 β -tetrahydropyranyloxychol-5-en-24-al **74** (Scheme 10).⁴⁰

2.1.3. Synthesis of [25, 26, 27-²H₇]BL from parent brassinolide.

Several routes for the synthesis of labelled BRs have been reported so far,⁴¹ however, all are quite lengthy multi step reactions which are neither eco-friendly nor cost-effective. The easy way to synthesize the side chain labelled BRs from parent BRs includes three basic steps; C-25 hydroxylation, dehydration to create double bond between C25-C26 and deuteriogenation or tritiation of the double bond. Based on these basic steps, the preparation of [25, 26, 27-²H₇]BL **97** from brassinolide **88** as a starting material was achieved⁴² (Scheme 11). 25-Oxy-functionalization of tetra-*O*-acetyl BL **89** with TFD offered predominantly 25-hydroxy compound **90** while 14-hydroxy,25-hydroxy-15-oxo- and 14,25-dihydroxy derivatives **91**, **92**, **93**, respectively were identified as byproducts. Dehydration of **90** headed to the formation of a mixture of the compounds **94** and **95** (which were unseparable) and then the mixture of isomers were subjected to deprotection of tetra-*O*-acetyl groups. Subsequently the mixture of isomers were treated with KOH in aqueous MeOH at room temperature until the lactone ring completely opened to the carboxylate and then at refluxing temperature. After recyclization of lactone ring and desalting the crystalline compound **96** was obtained. Reduction of the double bond of $\Delta^{25(26)}$ BL **96** using deuterium gas and 5% Pd/C as catalyst was preceded with extensive isotopic exchange, which resulted in high deuterium incorporation leading to the formation of [25,26,27-²H₇]BL **97** with 60% isotopic purity. The partial migration of double bond provided the minor byproduct $\Delta^{24(25)}$ [26,27-²H₆]BL **98** with 68% isotopic purity.


2.1.4. Alternative way to [26,28-²H₆]brassinolide and [26,28-²H₆]castasterone

Kolbe *et al.*⁴³ reported the synthesis of deuterium labelled BRs [26,28-²H₆]CS **107** and [26,28-²H₆]BL **108** starting from 6,6-ethylenedioxy-20-formyl-2 α ,3 α -isopropylidenedioxy-5 α -pregnane **99** and 3-[methyl-²H₃]methyl-[4,4,4-²H₃]but-1-yne **100**, respectively (Scheme 12). The acetylenic alcohol **101** was oxidized by pyridinium chlorochromate and stereoselective reduction of ketone **102** yielded stereospecifically the acetylenic alcohol **103**. Partial hydrogenation on Raney Ni gave an allylic alcohol **104**. The double bond was epoxidized by 3-chloroperbenzoic acid and then epoxide **105** was treated by trimethylaluminium/*n*-butyllithium reagent. Opening of the epoxide ring in **105** leads to the unusual methyl group migration from C-26 to C-25 took place during methylation and then subjected to hydrolysis to furnish [26, 28-²H₆]CS **107** was obtained instead of [26,27-²H₆]-labelled isomer. Baeyer-Villiger oxidation of [26,28-²H₆]castasterone **107** gave mixture of regioisomers **109** and **108** in which [26,28-²H₆]BL **109** prevailed (**109:108** = 6:1).


2.1.5. Synthesis of [26-²H₃]-epibrassinolide and its precursors

A further study towards the synthesis of labelled BRs containing three deuterium atoms in the terminal part of the side chain starting from commercially available stigmaterol **24** is reported by Khripach *et al.*³² 3,5-cyclo ester **110** was prepared from stigmaterol **24** in 5 steps. Labelling with deuterium was done in two steps. Reduction of 3,5-cyclo ester **110** with lithium aluminium deuteride gave alcohol **111** (Scheme 13) and then mesylation of the alcohol **111** followed by reduction with LiAlD₄ was introduced the third deuterium to the side chain end methyl group and thus Δ^{22} -derivative **112** was obtained. Asymmetric dihydroxylation of the double bond in **112** followed by acetylation yielded (22*R*, 23*R*)-diacetate **113**. A methoxy group at C-6 was replaced by an oxo group in four steps to give cycloketone **115** that was used for further construction of cyclic part of brassinosteroids (Scheme 13).


Treatment of compound **115** in boiling dimethyl acetamide in the presence of Py.HBr directed to the preparation of compound **116** containing 2,3 double bond in the ring A (Scheme 14).



Scheme 11 Synthesis of [25, 26, 27-²H₇]BL.




Scheme 12 Synthesis of [26, 28-²H₆]CS and [26, 28-²H₆]BL




Scheme 13 Synthesis of cycloketone **115** from stigmasterol **24**

Removal of acetyl masking groups revealed the presence of 24-[26- $^2\text{H}_3$]episcasterol **117**. Recently, 24-episcasterol was found cytotoxic against MCF-7 cells.⁴⁴ Epoxidation of the olefin **117** gave deuterated epoxide **118** and dihydroxylation of olefin **117** gave 24-[26- $^2\text{H}_3$]epiCS **119** (Scheme 14). 24-[26- $^2\text{H}_3$]epiBL **123** was eventually prepared from protected 24-[26- $^2\text{H}_3$]episcasterol **116** (Scheme 15). The diacetate of [26- $^2\text{H}_3$] epics **120** obtained by dihydroxylation of **116** was acetylated and converted by Baeyer-Villiger oxidation to tetracetate of 24-[26- $^2\text{H}_3$]epiBL **121**. After alkaline hydrolysis of acetate groups and acidic

workup to restore the lactone ring in 24-[26- $^2\text{H}_3$] epibrassinolide **123**. It is worth mentioning here that this paper is reporting the protocols enabling the direct Baeyer-Villiger oxidation of 24-epicastasterone to 24-epibrassinolide with more than satisfactory yields and without need of the protection of hydroxy groups.^{45,46}




Scheme 14 Synthesis of 24-[26- $^2\text{H}_3$] epiCS **119**



Scheme 15 Synthesis of deuterated 24-[26- $^2\text{H}_3$] epiBL **123**

2.1.6. Synthesis of 6-deoxo-24-[26- $^2\text{H}_3$]epiCS

The position and reaction sequence for introducing the label was selected based on the previous experience with the synthesis of deuterated BRs (Scheme 16).



Scheme 16 Synthesis of 6-deoxo-24-[26-²H₃] epiCS 132

Stigmasterol **24** was a starting compound that was converted in five steps to 22-aldehyde **124**. Claisen rearrangement was the key step in conversion of 22-aldehyde **124** to the ester **125**. Using the reduction with LiAlD₄-mesylation-reduction with LiAlD₄ sequence as described above the ester group of **125** was converted to [²H₃] methyl group in **128**. Rearrangement of 3,5-cyclo

derivative **128** to Δ^2 derivative **129** was achieved by boiling with pyridinium hydrobromide. Deoxygenation of at C-6 was done *via* methyl xanthate **130** and for final *cis* hydroxylation potassium osmate was used. Thus, the 6-deoxo-24-[26- $^2\text{H}_3$]epiCS **132** was prepared⁴⁷ in overall yield 18% calculated on ester **125**.


2.2. BRs with deuterium label in ring A

2.2.1. Stereospecific labelling of brassinosteroids with hydrogen isotope.

As the deuterium labelled BRs must have at least 4 deuterium atoms in metabolically stable positions to be useful as MS internal standards the deuterium labelling in ring A is not very attractive. On the other hand ring A is good target for labelling with tritium especially if the aim is to prepare suitable precursor for tritiation from the non-labelled target BR. It is the rule to model the tritiation reactions first with deuterium and therefore we developed recently general stereospecific method of introduction of deuterium to ring A of BRs having $2\alpha,3\alpha$ -dihydroxy group.

In quest for the suitable precursor for tritiation we discovered a stereospecific reaction of α -hydroxy ketone **135**, prepared in three steps from 24-epicastasterone **2** in 47% yield, with triphosgene⁴⁸ giving in 99% yield 3β -chloro- $2\alpha,3\alpha$ -(carbonyldioxy) derivative **136** (Scheme 17).⁴⁵ While catalytic reductive dechlorination is frequently used for the introduction of hydrogen isotopes on aromatic rings⁴⁹ to our best knowledge there are no literature data available for either aliphatic or alicyclic substrates. Catalytic dehalogenation of chlorocarbonate **136** with deuterium gas on 5% PdO/CaCO₃ in the presence of triethylamine in ethyl acetate afforded the $2\alpha,3\alpha$ -(carbonyldioxy)[3β - ^2H]derivative **137** in 65% yield and 80% isotopic enrichment. Removal of protecting groups in two-step one pot sequence gave 24-[3β - ^2H] epicastasterone **138** in 91%

yield on **137**. Baeyer-Villiger oxidation of [3 β -²H]epiCs **138** with trifluoroperoxyacetic acid gave 24-[3 β -²H]epiBL **139** in a 65% yield with no loss of deuterium label.




Scheme 17 Synthesis of 24-[3 β -²H]epiCS and 24-[3 β -²H]epiBL

2.3. BRs with deuterium label in ring B

2.3.1. Synthesis of 24-[5, 7, 7-²H₃]epiBL


For BRs containing keto group the base catalyzed exchange of α -hydrogens with ²H₂O can be exploited for deuterium introduction. Kolbe *et al.*²⁰ reported the preparation of 24-[5,7,7-²H₃]epiBL starting from tetracetate of 24-epiCS **140** (Scheme 18) or alternatively from 2,3,22,23-bis-isopropylidenedioxy-24-epiCS **133** (see Scheme 17). The exchange with ²H₂O in DMF was catalyzed by triethyl amine. Bayer-Villiger oxidation with CF₃CO₃H of the obtained labelled 24-epiCS derivative **141** and its final deprotection gave 24-[5,7,7-²H₃]epiBL **143**. This procedure has the advantage of specific introduction of labelling at a late stage, whereas three deuterium atoms were introduced in a stable position at C-5 and C-7. A procedure reported by Allevi *et al.*⁵⁰ is less convenient because it uses high excess of labelled water and methanol and there is the need of separation of side chain epimers.



Scheme 18 Synthesis of 24-[5, 7, 7-²H₃] epiBL

2.3.2. Synthesis of 24-[7,7-²H₂]epiBL


The intended transformation of lactone group in epibrassinolide **146** was carried out after protecting both diol functions by isopropylidene. The key transformation is the preparation of 6, 7-seco diacid **153** from diacetone **147** in three step and then formation of the cyclic anhydride **154** from 6, 7-seco diacid **153** (Scheme 19). Regioselective reduction of anhydride **154** with NaBD₄ gave deuterated diacetone **155**. Acidic hydrolysis of acetone groups on **155** provided [7,7-²H₂]epiBL **156** with 82% isotopic enrichment.⁵¹

Scheme 19 Synthesis of 24-[7,7- $^2\text{H}_2$]epiBL

3. Synthesis of tritium-labelled BRs

3.1. Synthesis of 24-[5,7,7-³H]epiBL


By the same method as described above for the preparation of corresponding deuterium labelled 24-epiBL **143** tritium was introduced to tetracetate of 24-epiCS **140** by base catalyzed exchange reaction with tritiated water having specific activity 1.1 Ci/mL. After Bayer-Villiger oxidation of **157** and removal of acetyl protecting groups 24-[5,7,7-³H] epiBL with specific activity (S.A.) 6 mCi/mmol.²⁰ The same sequence was performed also with 2,3,22,23-bis-isopropylidenedioxy-24-epiCS **133** as starting compound. The advantage of the use of isopropylidene protecting groups is the more simple deprotection step - the lactone ring is not opened under acidic hydrolysis conditions as the opposite is true for basic hydrolysis needed for the acetate group cleavage.



Scheme 20 Synthesis of 24-[5,7,7-³H]epiBL

3.2. Synthesis of 24-[5,7,7-³H]epiTE, 6-oxo-24 β -methyl-22-dehydro[5,7,7,³H]cholestanol and 6-oxo-24-[5,7,7-³H]epicampestanol

All title compounds were prepared by base catalysed exchange of α -keto hydrogens with tritiated water⁵² with S.A. = 0.8 Ci/mmol (Scheme 21). 24-[5,7,7-³H]epiteasterone **162** was prepared with S.A. = 1.5 mCi/mmol. 6-oxo-24 β -methyl-22-dehydro[5,7,7,³H]cholestanol **164** was labelled without protection of hydroxyl groups and its S.A. was not indicated. Cholestanol derivative **164** was further converted *via* hydrogenation of the double bond in the side chain catalysed by 10% Pd/C and then 6-oxo-24-[5,7,7-³H]epicampestanol purified by column chromatography had S.A. = 3.5 mCi/mmol.




Scheme 21 Labelling of biogenetic BRs precursors

4. Synthesis of ^{14}C -labelled BRs


4.1. Synthesis of (22*R*, 23*R*)-and (22*S*, 23*S*)-24-[4- ^{14}C] epiBL

There is only one report available in the literature on ^{14}C -labelled brassinosteroids so far. Seo *et al.*⁵³ have reported the synthesis of [4- ^{14}C]-labelled epiBL **183** and **184**. The C-4 position in epiBL **183** was selected for ^{14}C labelling because of its stability to metabolic loss and easy way to do the preparation. According to the established method reported in the literature for

incorporation of ^{14}C into the C-4 position of steroids, the enol lactone **172** was synthesized from the starting material brassicasterol **167** in five steps (Scheme 22). This lactone **172** was then treated with [^{14}C] methyl iodide (prepared by known method from barium [^{14}C]carbonate *via* [^{14}C]methanol) to give bridged ketone **173**. Alkaline treatment of **173** in MeOH gave [4- ^{14}C]brassicasterone **174**. Acetylation of **174** with isopropenyl acetate under acid catalysis gave the enol acetate **175** that was reduced with sodium borohydride in methanol to give [4- ^{14}C]brassicasterol **176** as the major product. 3,5-cyclo-6-ol **178** was obtained in 91.7% yield by mesylation to give **177** followed by treatment with sodium carbonate in acetone (Scheme 23). Jones oxidation of **178** gave 3,5-cyclo-6-one **179**, which was treated with lithium bromide and camphor sulfonic acid in dimethyl acetamide to rearrange to 2,22-diene-6-one **180** in quantitative yield. Oxidation of **180** with osmium tetroxide gave a stereoisomeric mixture of 2,3,22,23-tetraols **181** and **182** which is separated through the repeated chromatography and recrystallization. Bayer-Villiger oxidation with TFA in dichloromethane gave the (22*R*, 23*R*)-7-oxa-lactone **183** contaminated with a small amount of its 6-oxa isomer. (22*S*, 23*S*)-Tetraol also gave the (22*S*, 23*S*)-7-oxa-lactone **184** contaminated with its 6-oxa isomer. The final products (22*R*, 23*R*)-24-[4- ^{14}C]epiBL **183** and (22*S*, 23*S*)-24-[4- ^{14}C]epiBL **183** were obtained in 3.20 % and 4.46 % radiochemical yield (based on $\text{Ba}^{14}\text{CO}_3$), respectively. Specific radioactivity of **183** and **183** was 56.8 mCi/mmol.



Scheme 22 Synthetic pathway to (22*R*, 23*R*)- and (22*S*, 23*S*)-24-[4-¹⁴C] epiBL I



Scheme 23 Synthetic pathway to (22*R*, 23*R*)- and (22*S*, 23*S*)-24-[4-¹⁴C] epiBL II

5. Summary

The number of published synthesis of BRs labelled either by stable isotopes or by radioisotopes reflects their importance for the biochemical studies of this interesting group of plant growth regulators. The main effort was devoted to the synthesis of multideuterated BRs in side chain as internal standards for MS. For labelling with tritium the methods developed for multideuterated BRs are not useful. The methods using exchange with tritiated water were described. However, of in this way prepared BRs have specific activities only in order of several mCi/mmol. What remains to be done is to prepare the BRs ligands with the specific activities of order tens of Ci/mmol to enable the search for BRs receptors. Only one example of the synthesis of ^{14}C -labelled BR reflects the shortage and fast increase of price of $\text{Ba}^{14}\text{CO}_3$ in recent years. Notwithstanding, some BRs are considered as potential drugs and if they will pass the preclinical sieve, there will be certainly need for their labelling with ^{14}C .

6. Acknowledgements

This work was realized at IOCB, Prague, Czech Republic and supported by Academy of Sciences of the Czech Republic (RVO 61388963), by Grant IAA400550801 of the Grant Agency of the Academy of Sciences of the Czech Republic and by CNMS, Jain University, Bangalore. The authors wish to express thanks to all for financial support.

7. References

- 1 S. Fujioka and A. Sakurai, *Nat. Prod. Rep.*, 1997, **14**, 1-10.
- 2 A. Bajguz and A. Tretyn, *Phytochemistry*, 2003, **62**, 1027-1046.
- 3 V. Castilla, J. Ramirez and C. E. Coto, *Curr. Med. Chem.*, 2010, **17**, 1858-1873.
- 4 S. D. Clouse, *Plant. J.*, 1996, **10**, 1-8.
- 5 V. A. Khripach and V. N. Zhabinski, A. de Groot, *Brassinosteroids: a new class of plant hormones*, San Diego, Academic Press, 1999.
- 6 A. Sakurai, T. Yokota and S.D. Clouse, ed, *Brassinosteroids: steroidal plant hormones*. Tokyo: Springer, 1999.
- 7 V. Marquardt and Adam.-in: W. Ebing , (ed.- in- chief) *Chemistry of plant protection* , Springer, Berlin 1991, **7**, 103-139 and the references sited therein.
- 8 H. Suzuki, T. Inoue, S. Fujioka, T. Saito, S. Takatsuto, T. Yokota, N. Murofushi, T. Yanagisawa and A. Sakurai, *Photochemistry*, 1995, **40**, 1391-1397.
- 9 S. Fujioka, T. Inoue, S. Takatsuto, T. Yanagisawa, T. Yokota and A. Sakurai, *Biosci. Biotech. Biochem.*, 1995, **59**, 1543-1547.
- 10 S. Takatsuto, C. Gotoh, T. Noguchi, T. Nomura, S. Fujioka and T. Yokota, *J. Chem. Res. Synop.*, 1998, 206-207.
- 11 H. Suzuki, S. Fujioka, S. Takatsuto, T. Yokota, N. Murofushi and A. Sakurai, *J. Plant Growth Regul.*, 1994, **13**, 21-26.
- 12 A. Sakurai and S. Fujioka, *Biosci. Biotech. Biochem.*, 1997, **61**, 757-762.
- 13 S. Fujioka, J. Li, Y-H. Choi, H. Seto, S. Takatsuto, T. Noguchi, T. Watanabe, H. Kuriyama, T. Yokota, J. Chory and A. Sakurai, *Plant Cell*, 1997, **9**, 1951-1962.

- 14 T. Noguchi, S. Fujioka, S. Takatsuto, S. Yoshida, A. Sakurai, J. Li and J. Chory, Abstract Papers of 32nd Annual Meeting of The Japanese Society for Chemical Regulation of Plants, Tokyo, 1997, p. 93.
- 15 S. Fujioka and A. Sakurai, *Physiol. Plant*, 1997, **100**, 710-715.
- 16 T. Yokota, *Trends Plant Sci.*, 1997, **2**, 137-143.
- 17 A. Antonchick, A. Svatoš, B. Schneider, O. V. Konstantinova, V. N. Zhabinskii and V.A. Khripach, *Phytochemistry*, 2005, **66**, 65-72.
- 18 S. Takatsuto and N. Ikekawa, *Chem. Pharm. Bull.*, 1984, **32**, 2001-2004.
- 19 a) J. Malíková, J. Swaczynová, Z. Kolář and M. Strnad, *Phytochemistry*, 2008, **69**, 418-426.
b) J. Steigerová, J. Oklešťková, M. Levková, L. Rárová, Z. Kovář and M. Strnad, *Chem.-Biol. Interact.* 2010, **188**, 487-496.
- 20 A. Kolbe, V. Marquardt and G. Adam, *J. Labelled Compd. Rad.*, 1992, **31**, 801-805.
- 21 V. A. Khripach, V. N. Zhabinskii, O. V. Gulyakevich, O. V. Konstantinova, A. Y. Misharin, A. R. Mekhtiev, V. P. Timofeev and Y. V. Tkachev, *Russ. J. Bioorg. Chem.*, 2010, **36**, 746-754.
- 22 S. Takatsuto and N. Ikekawa, *Chem. Pharm. Bull.*, 1986, **34**, 4045-4049.
- 23 S. C. Lee, S. Joo, C. Park, S. Son, J. Youn, M. Kim, S. Jeong, and S. Kim, *Bull. Korean Chem. Soc.* 2011, **32**, 332-334.
- 24 G. Adam, A. Porzel, J. Schmidt, B. Schneider and B. Voigt, in *Studies in Natural Products Chemistry*, Atta-ur-Rahman, ed., Elsevier: Amsterdam, 1996, **18**, pp. 495-549.
- 25 M. D. Grove, G. F. Spencer, W. K. Rohwedder, N. Mandava, J. F. Worley, J. D. Warthen, Steffens, J. L. Flippen-Anderson and J. C. Cook, *Nature*, 1979, **225**, 1065-1066.
- 26 S. Takatsuto and N. Ikekawa, *Chem. Pharm. Bull.*, 1986, **34**, 1415-1418.
- 27 S. Takatsuto and N. Ikekawa, *Chem. Pharm. Bull.*, 1986, **34**, 4045-4049.

- 28 S. Takatsuto and N. Ikekawa, *J. Chem. Soc. Perkin Trans.1*, 1986, 591-593
- 29 V.A. Khripach, V. N. Zhabinskii, O. V. Konstantinova and N. B. Khripach, *Tetrahedron Lett.*, 2000, **41**, 5765-5767.
- 30 V. A. Khripach, V. N Zhabinskii, O. V Konstantinova, N. B. Khripach, A. P. Antonchick and B. Schneider, *Steroids*, 2002, **67**, 587-595.
- 31 K. Mori, M. Sakakibara, Y. Ichikawa, H. Ueda, K. Okada, T. Umemura, G. Yabuta, S. Kuwahara, M. Kondo, M. Minobe and A. Sogabe, *Tetrahedron*, 1982, **38**, 2099-2109.
- 32 V. A. Khripach, V. N. Zhabinskii, V. K. Olkhovick and F. A. Lakhvich, *Zhurn. Org Khim.*, 1990, **26**, 1966-1976
- 33 J. A. Steele and E. Mosettig, *J. Org. Chem.*, 1963, **28**, 571-572.
- 34 J. J. Pappas, W. P. Keaveney, E. Gancher and M. Berger, *Tetrahedron Lett.*, 1966, **7**, 4273-4278.
- 35 A. P. Antonchick, B. Schneider, V. N. Zhabinskii and V. A. Khripach, *Steroids*, 2004, **69**, 617-628.
- 36 V. A. Khripach, V.N. Zhabinskii, A. P. Antonchick, O.V. Konstantinova and B. Schneider, *Steroids*, 2002, **67**, 1101-1108
- 37 M. Anastasia, P. Allevi, P. Ciuffreda and A. Fiecchi, *J. Chem. Soc., Perkin Trans. 1*, 1983, 2365-2367.
- 38 M. Anastasia, P. Ciuffreda, M. Delpuppo and A. Fiecchi, *J. Chem. Soc., Perkin Trans. 1*, 1983, 379-382.
- 39 M. Anastasia, P. Allevi, P. Ciuffreda, A. Fiecchi, P. Gariboldi and A. Scala, *J. Chem. Soc., Perkin Trans. 1*, 1985, 595-599.

- 40 M. Koizumi, M. Ishiguro, M. Yasuda and N. Ikekawa, *J. Chem. Soc., Perkin Trans. 1*, 1983, 1401-1410.
- 41 V. Marquardt and G. Adam, (Ebing, W., ed.-in-chief), *Chemistry of plant protection*, Springer, Berlin 1991, 7, 103.
- 42 H. Seto, S. Fujioka, H. Koshino, S. Yoshida, T. Watanabe and S. Takatsuto, *Tetrahedron Lett.*, 1998, **39**, 7525-7528.
- 43 A. Kolbe, A. Porzel, J. Schmidt and G. Adam, *J. Labelled. Compd. Rad.*, 2003, **46**, 231-242
- 44 V. A. Khripach, V. N. Zhabinskii, O. V. Gulyakevich, O. V. Konstantinova, A. Y. Misharin, A. R. Mekhtiev, V. P. Timofeev and Y. V. Tkachev, *Russ. J. Bioorg. Chem.*, 2010, **36**, 746-754.
- 45 A. Marek, M. R. Patil, B. Klepetářová, L. Kohout and T. Elbert, *Tetrahedron Lett.*, 2012, **53**, 2048-2050.
- 46 L. Kohout, *Collect. Czech. Chem. Commun.* 1994, **59**, 457-460.
- 47 V. A. Khripach, V. N. Zhabinskii and O. V. Gulyakevich, *Chem. Nat. Compd.*, 2012, **48**, 601-605.
- 48 H. Eckert and B. Forster, *Angew. Chem. Int. Ed. Engl.*, 1987, **26**, 894-895.
- 49 R. Voges, J. R. Heys and T. Moenius, Preparation of compounds labelled with tritium and Carbon-14, John Wiley & Sons: Chichester, 2009, pp 133-144.
- 50 P. Allevi, M. Anastasia, R. Cerana, P. Ciuffreda and P. Lado, *Phytochemistry.*, 1988, **27**, 1309-1313.
- 51 V. A. Khripach, N. B. Khripach, V. N. Zhabinskii, Y. Y. Zhiburtovich, B. Schneider and A. De Groot, *J. Labelled Compd. Rad.*, 2007, **50**, 1153-1158.
- 52 A. Kolbe, B. Schneider, B. Voigt and G. Adam, *J. Labelled Compd. Rad.*, 1998, **41**, 131-137.

53 S. Seo, T. Nagasaki, Y. Katsuyama, F. Matsubara, T. Sakata, M. Yoshioka and Y. Makisumi,
J. Labelled Compd. Rad., 1989, **27**, 1383-1393.

8. List of Abbreviations

Ac ₂ O	acetic anhydride
AIBN	azobisisobutyronitrile
aq	aqueous
BL	brassinolide
BR	Brassinosteroid
BuLi	butyllithium
CS	castasterone
D	deuterium
DBU	1,8-diazabicyclo[5.4.0]undec-7-ene
DCC	dicyclohexylcarbodiimide
DCM	dichloromethane
(DHQ) ₂ -PHAL	dihydroquinine 1,4-phthalazinediyl diether
(DHQD) ₂ -PHAL	dihydroquinidine 1,4-phthalazinediyl diether
DMD	dimethyldioxirane
DMF	dimethylformamide
DMP	4-dimethylamino pyridine
DMSO	dimethyl sulfoxide
epiBL	epibrassinolide
Et ₃ N	triethyl amine
EtOAc	ethyl acetate
Hg(OAc) ₂	mercury(II) acetate
LiAlD ₄	lithium AluminumDeuteride
<i>m</i> -CPBA	<i>meta</i> -chloroperoxybenzoic acid
MCF	Michigan Cancer Foundation
MeOH	methanol
MsCl	methanesulfonyl chloride
Na/Hg	sodium amalgam
NBS	<i>N</i> -bromosuccinic imide
PCC	pyridinium chlorochromate
<i>p</i> -TsOH	<i>p</i> -toluenesulfonic acid
Py	pyridine
T	tritium
TE	teasterone
TEMPO	2,2,6,6-tetramethylpiperidiny-1-oxyl
TFD	trifluoromethyldioxirane
THF	tetrahydrofuran
TsCl	4-toluenesulfonyl chloride
TY	typhasterol