

RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Synthesis of 6-membered cyclic carbonates from 1,3-diols and low CO₂ pressure : a novel mild strategy to replace phosgene reagents

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,
Accepted 00th January 2012Georgina L. Gregory,^a Marion Ulmann^a and Antoine Buchard^{a,*}

DOI: 10.1039/x0xx00000x


www.rsc.org/

Low pressure carbon dioxide is used as the carbonation agent in a simple, safe and efficient procedure for the synthesis of 6-membered cyclic carbonates from 1,3-diols. Using readily available reagents and proceeding at room temperature, this route offers a novel mild alternative to phosgene derivatives.

Unlike aromatic polycarbonates, aliphatic polycarbonates (APCs) have been little explored commercially. However, their low toxicity and biodegradability make them excellent candidates for biomedical applications.¹ Recently, they have also been investigated as thermoplastics,² binders for photovoltaics,³ polymer electrolytes⁴ and adhesives.⁵

Part of this renewed attention stems from the possibility to obtain APCs via the copolymerisation of CO₂ and epoxides,⁶ an attractive reaction which uses an abundant and virtually free source of carbon, but is limited by the scope of usable epoxides. APCs can also be synthesised by polycondensation of diols with phosgene derivatives or dialkyl carbonates,⁷ but the control of the polymer molecular weight is highly sensitive to the reaction conditions. Hence, Ring-Opening Polymerisation (ROP) of cyclic carbonates has become the method of choice for APC synthesis,^{1b, 8} as the development of ROP catalysts⁹ has enabled controlled polymerisation of highly functionalised monomers¹⁰ under mild conditions (usually 6-, 7-,¹¹ or strained 5-membered rings¹²).

A common preparation of cyclic carbonate monomers involves the transesterification of diols with phosgene,^{8, 13} a toxic reagent synthesised from CO and Cl₂ in an energy intensive process. Nevertheless, phosgene is still widely employed due to its efficiency and the lack of a more versatile and sustainable alternative. Safer phosgene derivatives¹⁴ include di-*tert*-butyl dicarbonate,¹⁵ 1,1'-carbonyldiimidazole,¹⁶ and aromatic carbonates,¹⁷ but these reagents all derive from phosgene, and can lead to unwanted side reactions, low reactivity and difficult workups.




Scheme 1 Strategy for the synthesis of cyclic carbonate from 1,3-diols and CO₂ (R¹-R⁴ are the various substituents of 1,3-diols used, see Table 2)

Alternative methods, such as the Pd-catalysed oxidative carbonylation of diols using CO pressure,¹⁸ or the catalytic coupling of CO₂ with oxetanes,¹⁹ have also been reported. The latter however, is limited by the availability of oxetanes.²⁰ Transesterification of diols with urea, industrially produced from CO₂, is also described but with moderate success.²¹

Using CO₂ as a C1-carbonation agent is an attractive goal for phosgene related emissions mitigation and direct CO₂ utilisation. The direct coupling of CO₂ with diols, where water is the sole by-product, would be an attractive process. However, the reaction is generally not kinetically and thermodynamically favoured. Therefore, a catalyst and an efficient stoichiometric dehydrating strategy (*e.g.* nitriles, zeolites, orthoesters, or Mitsunobu reagents) are necessary for the reaction to proceed.²² Despite recent advances, such as the use of CeO₂ catalyst in tandem with a large excess of 2-cyanopyridine as a dehydrating agent,²³ or the *in situ* introduction of a leaving group based on dibromomethane,²⁴ high CO₂ pressures (10-50 bar) and high temperatures (70-140 °C) are still required, particularly for the synthesis of 6-membered rings.

Herein, we report a safe and efficient one-pot procedure for the synthesis of 6-membered cyclic carbonates directly from 1,3-diols, which uses CO₂ as the carbonation agent instead of phosgene derivatives. This methodology proceeds at room temperature, needs only 1 atm. of CO₂ and common lab reagents (Scheme 1), and does not require the preliminary preparation of oxetanes or chloroalcohols.²⁵

Table 1 Reaction of 1,3-butanediol **1a** with DBU under 1 atm of CO₂^a

Entry	Solvent	T (°C)	[1a] (M)	Conv. into carbonated products (%) ^b				
				Total	1'a_{I+II}	1'a_I	1'a_{II}	1'abis
1	Neat	25	n/a	47	44	30	14	3
2	C ₇ D ₈	25	1.7	60	56	37	19	4
3	CDCl ₃	25	1.7	78	69	46	23	9
4	CD ₃ CN	25	1.7	73	58	40	18	15
5 ^c	CDCl ₃	25	1.7	69	52	35	17	17
6	CDCl ₃	25	0.1	78	73	49	24	5
7 ^d	CDCl ₃	-78	1.7	77	68	45	23	9
8 ^e	CDCl ₃	25	1.7	62	56	37	19	6
9 ^f	CDCl ₃	25	1.7	>99	57	40	17	43

^a Reactions conditions: diol (5.6 mmol, 1.7M), DBU (5.6 mmol), room temperature, CO₂ (1 atm), 2 h (saturation). ^b Based on diol conversion and determined by relative integration of methine signals in ¹H NMR (e.g. 4.64 ppm for **1'a_{II}**/see ESI Fig S1-S3). ^c Addition of 1-butyl-3-methylimidazolium hexafluorophosphate (bmimPF₆, 5.6 mmol); [diol]=0.4 M. ^d Addition of DBU at T= -78 °C then rt. ^e 1 equiv. of TBD was used instead of DBU. ^f 3 equiv. of DBU were used.

In 2005, Jessop and coworkers reported the reversible carbonation of alcohols with CO₂ promoted by 1,8-diazabicyclo-[5.4.0]-undec-7-ene (DBU).²⁶ The conversion was later found to depend on the choice of solvent with almost full conversion of 1-hexanol being achieved in CDCl₃.²⁷ Inspired by these reports, we investigated the selective mono-insertion of CO₂ into one alcohol moiety of 1,3-diols in various solvents. The slow addition of DBU under 1 atm of CO₂ to solutions of (+/-)-1,3-butanediol (**1a**) was monitored and the conversion to alkyl carbonate species assessed by ¹H NMR (Table 1). Regardless of the solvent used, the carbonation of **1a** by CO₂ proceeded quickly at low pressure and room temperature, with good selectivity towards the mono-insertion products (ca. 90% of products). A slight preference for the primary alcohol moiety was also observed, in agreement with DFT calculations (see ESI Figure S4). In neat diol, the conversion was limited by the increased viscosity upon CO₂ insertion (Table 1, Entry 1), but even in solution and with the addition of ionic liquid bmimPF₆, as used by Lim *et al.* (Table 1, Entry 5),²⁴ full conversion of **1a** was not observed with only one DBU equivalent. Apolar toluene-*d*₈ gave less conversion than more polar solvents, whereas more dicarbonated product was obtained in acetonitrile-*d*₃ (Table 1, Entry 2 and 4). Overall, optimal results were obtained in CDCl₃, and despite it not being a desirable solvent which should be replaced in the future, chloroform was used in the rest of our study. Addition of DBU at low temperature resulted in no further improvement in mono-insertion selectivity (Table 1, Entry 7), while increasing the temperature promoted decarboxylation. A lower concentration of diol slightly decreased the amount of disubstituted product (Table 1, Entry 6) but did not affect the global conversion. The

use of stronger base 1,5,7-diazabicyclo[4.3.0]-undec-7-ene (TBD) in place of DBU resulted in less conversion, which is attributed to an increased sensitivity to moisture (Table 1, Entry 8). The use of triethylamine resulted in no carbonation of **1a**.


Further DFT calculations showed that the direct coupling of **1a** and CO₂ to form cyclic carbonate **2a** is slightly thermodynamically disfavoured ($\Delta\Delta G = +3.0$ kcal.mol⁻¹). Following the DBU-aided insertion of CO₂, the activation barriers are then too high for the cyclisation of **1'a_I** (or **1'a_{II}**) to proceed under mild conditions (61.1 and 42.2 kcal.mol⁻¹ via an S_N2 or addition/elimination mechanism, respectively) (see Fig S4). An *in situ* leaving group strategy was thus applied experimentally to overcome the kinetic as well as the thermodynamic limitation of the reaction. After the selective mono-insertion of CO₂ into **1a** in CDCl₃ at low concentration, 1 equivalent of tosyl chloride and triethylamine were added to the reaction mixture and stirred at room temperature overnight. Rapidly, the cyclic carbonate **2a** was detected by ¹H NMR (addition of triethylamine alone did not result in any product). The pure product was later isolated by column chromatography in a 44% yield, *i.e.* 60% conversion based on the mono-CO₂ inserted products (Table 2, Entry 1). A higher concentration of diol (1.7 M), despite being slightly detrimental to the first step of the procedure, proved to lead to a higher isolated yield of cyclic carbonate (68%, *i.e.* 99% conversion based on CO₂ mono-insertion, Table 2, Entry 2). This compares well with the traditional phosgene-based method (50%)²⁸ as well as oxidative carbonylation methods (45%).^{18a} The procedure was found to be robust: CO₂ from sublimed dry ice could be used and yielded cyclic carbonate **2a**, albeit in lower yield (48%, Table 2, Entry 3). The reaction also proceeded without solvent (30% yield). Investigation into the scope of the procedure was carried out with various 1,3-diols. The cyclisation step proceeded efficiently from the CO₂-mono insertion products and all cyclic carbonates were isolated in moderate yields (Table 2), comparative with phosgene-based and alternative methods. For example, the isolation of **2d** from 2,2-dimethyl-1,3-propanediol (53%, Table 2, Entry 8) was previously reported using phosgene derivatives (60%),²⁹ oxidative carbonylation (50%),^{18b} and metal free cyclisation (50%).²⁴ *O*-isopropylidene-xylose was also successfully transformed, though isolation proved challenging (11%, Table 2, Entry 12).

As cyclisation happens readily, reaction intermediates could not be isolated, but DFT calculations on model compound (*R*)-**1a** and additional experiments with optically active diols were carried out to investigate the reaction mechanism. After insertion of CO₂, tosylation can occur at the carbonate or at the remaining alcohol group, so that cyclisation proceeds via either an addition/elimination or a S_N2 pathway, leading to retention or inversion of stereochemistry (see Scheme 2). However, the exclusive formation and isolation of (*R,R*)-cyclic carbonate **2b** from (*R,R*)-2,4-pentanediol, as well as the optical activities of the cyclic carbonates obtained from enantiopure (*R*) and (*S*)-1,3-butanediol (Table 2, Entries 4-6),³⁰ both indicated an addition/elimination pathway, with no racemisation or inversion of stereochemistry being observed.

Table 2 Synthesis of various cyclic carbonates from 1,3-diols and CO₂[†]

Entry	Substrate	% Yield ^a [conv. _{1'→2}] ^b	Product
1 ^c		44 [60]	
2		68 [99]	
3 ^d		48	
4		68	
5		70	
6		53 [71]	
7		55 [96]	
8		53 [82]	
9		49 [78]	
10		46 [71]	
11		41 [70]	
12		11	

^a Isolated yield based on starting diol. ^b Spectroscopic conversion of CO₂ mono-insertion products **1'** into cyclic carbonate **2**. ^c [diol]=0.1M. ^d CO₂ from the sublimation of dry ice was used.



DFT calculations supported these conclusions and gave supplementary insight into the selectivity. Scheme 2 illustrates the computed pathways for the cyclisation of **1'a₁** to **2a**. As expected with a tosyl leaving group, the formation of the cyclic carbonate is strongly favoured ($\Delta\Delta G = -36.7$ kcal.mol⁻¹). Additionally, the activation energy barriers for both the nucleophilic addition/elimination and S_N2 ring-closing mechanisms are similar and low enough to happen at room temperature ($\Delta\Delta G^{\text{TS}} = +18.9$ and $+22.3$ kcal.mol⁻¹ respectively). The tosylation of the second alcohol moiety is however more favoured thermodynamically than that of the carbonate ($\Delta\Delta G = -25.4$ vs -2.3 kcal.mol⁻¹). Yet, the activation barrier for the tosylation of the carbonate is lower than that of the alcohol moiety ($\Delta\Delta G^{\text{TS}} = +23.8$ and $+29.7$ kcal.mol⁻¹ respectively). Hence, we believe that the addition/elimination pathway, resulting in stereochemical retention, is kinetically favoured over the S_N2 ($\Delta\Delta G^{\text{TS}} = +5.9$ kcal.mol⁻¹).

In conclusion, in the wider context of an ever-growing need for innovative materials and CO₂ emissions mitigation and utilisation, a novel one-pot procedure that uses safe and readily available reagents yielded 6-membered cyclic carbonates directly from abundant 1,3-diols and carbon dioxide under mild conditions (room temperature and low pressure), and compared well with toxic or expensive alternative methods. Computational and experimental evidence shows that the reaction proceeds through an addition/elimination pathway. Ongoing work is now aiming at replacing chloroform for a more environmental benign solvent and making the procedure catalytic in order to reduce the amount of salts produced.

We thank Sue and Roger Whorrod, the University of Bath, and the UK Engineering and Physical Sciences Research Council (EPSRC) (NSCCS/CHEM752 and CDT in Sustainable Chemical Technologies) for financial support.

Notes and references

^a Department of Chemistry, Centre for Sustainable Chemical Technologies, University of Bath, Claverton Down, Bath BA2 7AY, UK. E-mail: a.buchard@bath.ac.uk; Tel: +44 (0) 1225 386122.

† Typical procedure for the synthesis of cyclic carbonate monomers: Under a CO₂ atmosphere, DBU (1 equiv.) was added dropwise to a stirring solution/suspension of diol (0.5 g, 1 equiv.) in dry chloroform (1.7 M). After stirring at room temperature for 2 hours, triethylamine (1 equiv.) was added dropwise to the resulting viscous solution. A solution of TsCl in chloroform (1 equiv., 0.5 M) was then added slowly and the mixture stirred overnight. Removal of volatiles *in vacuo* afforded an oily residue, which was purified by column chromatography.

Electronic Supplementary Information (ESI) available: full experimental and computational details. See DOI: 10.1039/c000000x/

- (a) J. Xu, E. Feng and J. Song, *J. Appl. Polym. Sci.*, 2014, **131**, 39822-39838; (b) J. Feng, R.-X. Zhuo and X.-Z. Zhang, *Prog. Polym. Sci.*, 2012, **37**, 211-236.
- (a) K. Kobayashi, S. Kanmuri, Y. Kimura and K. Masutani, *Polym. Int.*, 2014, DOI: 10.1002/pi.4839; (b) G. A. Luinstra and E. Borchardt, in *Synthetic Biodegradable Polymers*, eds. B. Rieger, A. Künkel, G. W. Coates, R. Reichardt, E. Dinjus and T. A. Zevaco, Springer Berlin Heidelberg, 2012, vol. 245, pp. 29-48; (c) C. Koning, J. Wildeson, R. Parton, B. Plum, P. Steeman and D. J. Darensbourg, *Polymer*, 2001, **42**, 3995-4004.
- S.-Y. Ku, C. D. Liman, J. E. Cochran, M. F. Toney, M. L. Chabinyk and C. J. Hawker, *Adv. Mater.*, 2011, **23**, 2289-2293.
- Y. Tominaga, T. Shimomura and M. Nakamura, *Polymer*, 2010, **51**, 4295-4298.
- N. Fritz, H. Dao, S. A. B. Allen and P. A. Kohl, *Int. J. Adhes. Adhes.*, 2012, **38**, 45-49.
- M. R. Kember, A. Buchard and C. K. Williams, *Chem. Commun.*, 2011, **47**, 141-163.
- (a) J. H. Park, J. Y. Jeon, J. J. Lee, Y. Jang, J. K. Varghese and B. Y. Lee, *Macromolecules*, 2013, **46**, 3301-3308; (b) W. Zhu, X. Huang, C. Li, Y. Xiao, D. Zhang and G. Guan, *Polym. Int.*, 2011, **60**, 1060-1067; (c) G. Rokicki, J. Pawlicki and W. Kuran, *Polym. J.*, 1982, **14**, 839-845; (d) Y. Ono, *Catal. Today*, 1997, **35**, 15-25; (e) D. J. Brunelle, in *Advances in Polycarbonates*, eds. D. J. Brunelle and M. R. Korn, American Chemical Society, 2005, vol. 898, pp. 1-5.
- G. Rokicki, *Prog. Polym. Sci.*, 2000, **25**, 259-342.
- S. M. Guillaume and J.-F. Carpentier, *Catal. Sci. Technol.*, 2012, **2**, 898-906.
- (a) S. Tempelaar, L. Mespouille, O. Coulembier, P. Dubois and A. P. Dove, *Chem. Soc. Rev.*, 2013, **42**, 1312-1336; (b) D. P. Sanders, D. J. Coady, M. Yasumoto, M. Fujiwara, H. Sardon and J. L. Hedrick, *Polym. Chem.*, 2014, 327-329.
- G. Rokicki and P. G. Parzuchowski, in *Polymer Science: A Comprehensive Reference*, eds. K. Matyjaszewski and M. Möller, Elsevier, Amsterdam, 2012, pp. 247-308.
- (a) M. Azechi, K. Matsumoto and T. Endo, *J. Polym. Sci. Part A: Polym. Chem.*, 2013, **51**, 1651-1655; (b) W. Guerin, A. K. Diallo, E. Kirilov, M. Helou, M. Slawinski, J.-M. Brusson, J.-F. Carpentier and S. M. Guillaume, *Macromolecules*, 2014, **47**, 4230-4235; (c) W. Guerin, M. Helou, M. Slawinski, J.-M. Brusson, J.-F. Carpentier and S. M. Guillaume, *Polym. Chem.*, 2015, DOI: 10.1039/c1034py01660b; (d) A. K. Diallo, E. Kirilov, M. Slawinski, J.-M. Brusson, S. M. Guillaume and J.-F. Carpentier, *Polym. Chem.*, 2015, DOI: 10.1039/c1034py01713g.
- A.-A. G. Shaikh and S. Sivaram, *Chem. Rev.*, 1996, **96**, 951-976.
- H. A. Staab, *Angew. Chem., Int. Ed. Engl.*, 1962, **1**, 351-367.
- Y. Basel and A. Hassner, *J. Org. Chem.*, 2000, **65**, 6368-6380.
- J. Mindemark and T. Bowden, *Polymer*, 2011, **52**, 5716-5722.
- (a) H. Matsukizono and T. Endo, *J. Appl. Polym. Sci.*, 2015, 41956-41966; (b) D. P. Sanders, K. Fukushima, D. J. Coady, A. Nelson, M. Fujiwara, M. Yasumoto and J. L. Hedrick, *J. Am. Chem. Soc.*, 2010, **132**, 14724-14726; (c) D. P. Sanders, D. J. Coady, M. Yasumoto, M. Fujiwara, H. Sardon and J. L. Hedrick, *Polym. Chem.*, 2014, **5**, 327-329; (d) K. Mikami, A. T. Lonacker, T. P. Gustafson, N. F. Zinnel, P. J. Pai, D. H. Russell and K. L. Wooley, *J. Am. Chem. Soc.*, 2013, **135**, 6826-6829; (e) *US Pat.*, US20100280242, 2010.
- (a) B. Gabriele, R. Mancuso, G. Salerno, L. Veltri, M. Costa and A. Dibenedetto, *ChemSusChem*, 2011, **4**, 1778-1786; (b) D. M. Pearson, N. R. Conley and R. M. Waymouth, *Adv. Synth. Catal.*, 2011, **353**, 3007-3013.
- C. Martín, G. Fiorani and A. W. Kleij, *ACS Catal.*, 2015, **5**, 1353-1370.
- J. A. Burkhard, G. Wuitschik, M. Rogers-Evans, K. Müller and E. M. Carreira, *Angew. Chem. Int. Ed.*, 2010, **49**, 9052-9067.
- (a) Q. Li, W. Zhang, N. Zhao, W. Wei and Y. Sun, *Catal. Today*, 2006, **115**, 111-116; (b) Q. Li, N. Zhao, W. Wei and Y. Sun, *J. Mol. Catal. A: Chem.*, 2007, **270**, 44-49.
- (a) D. Chaturvedi, N. Mishra and V. Mishra, *Tetrahedron Lett.*, 2007, **48**, 5043-5045; (b) T. Sakakura and K. Kohno, *Chem. Commun.*, 2009, 1312-1330; (c) M. Tamura, M. Honda, Y. Nakagawa and K. Tomishige, *J. Chem. Technol. Biotechnol.*, 2014, **89**, 19-33.
- M. Honda, M. Tamura, K. Nakao, K. Suzuki, Y. Nakagawa and K. Tomishige, *ACS Catal.*, 2014, **4**, 1893-1896.
- Y. N. Lim, C. Lee and H.-Y. Jang, *Eur. J. Org. Chem.*, 2014, **9**, 1823-1826.
- M. R. Reithofer, Y. N. Sum and Y. Zhang, *Green Chem.*, 2013, **15**, 2086-2090.
- P. G. Jessop, D. J. Heldebrant, X. Li, C. A. Eckert and C. L. Liotta, *Nature*, 2005, **436**, 1102-1102.
- (a) P. M. Mathias, K. Afshar, F. Zheng, M. D. Bearden, C. J. Freeman, T. Andrea, P. K. Koech, I. Kutnyakov, A. Zwoster, A. R. Smith, P. G. Jessop, O. G. Nik and D. J. Heldebrant, *Energy Environ. Sci.*, 2013, **6**, 2233-2242; (b) L. Phan, J. R. Andreatta, L. K. Horvey, C. F. Edie, A.-L. Luco, A. Mirchandani, D. J. Darensbourg and P. G. Jessop, *J. Org. Chem.*, 2008, **73**, 127-132.
- H. Yasuda, M.-S. Aludin, N. Kitamura, M. Tanabe and H. Sirahama, *Macromolecules*, 1999, **32**, 6047-6057.
- B. J. Ludwig and E. C. Piech, *J. Am. Chem. Soc.*, 1951, **73**, 5779-5781.
- Low volatility and UV sensitivity prevented chiral GC/HPLC study.