

# RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

# Effect of different substituents on the one-pot formation of 3,4,5-substituted furan-2(5H)-ones: A kinetics and mechanism study

Mehdi Shahraki, Sayyed Mostafa Habibi-Khorassani\*, Maryam Dehdab

*Department of Chemistry, Faculty of Science, University of Sistan and Baluchestan, P O Box 98135-674, Zahedan, Iran*

*\*Corresponding author: Tel: +98 541 2446565; fax: +98 541 2446565; E-mail: smhabibi@chem.usb.ac.ir*

## Abstract

Kinetics of reactions between benzaldehyde and diethyl acetylenedicarboxylate with para- substituted anilines for the one-pot formation of 3,4,5-substituted furan-2(5H)-ones have been studied spectrally at different temperatures in formic acid. For all substituents on para position of aniline ring, the reaction followed the second-order kinetics. The partial orders with respect to substituted aniline and the diethyl acetylenedicarboxylate was one and one, and also the reactions revealed zero-order kinetics in relation to benzaldehyde. Para-Electron donating substituents on aniline ring increased the rate of reaction. On the basis of experimental data, a three steps mechanism has been proposed. Kinetic values ( $k$  and  $E_a$ ) and associated activation parameters ( $\Delta H^\ddagger$ ,  $\Delta G^\ddagger$  and  $\Delta S^\ddagger$ ) of the reactions were determined.

**Keywords:** Kinetics; Mechanism; furan-2(5H)-ones; para- substituted aniline


## 1. Introduction

A multicomponent reaction (MCR) is a process in which three or more accessible components combining together to produce a final product that display features of all inputs and thus offers greater possibilities for molecular diversity per step with a minimum of synthetic time and effort in a single reaction.

Synthesis of furan-2(5H)-ones as a subunit in a large number of natural products [1–5] is a significant target for organic and medicinal chemists. Highly substituted furans are a structural component of a vast number of biologically active natural and synthetic compounds such as anticancer [6, 7], anti-inflammatory [8], antifungal [9, 10], antimicrobial [11, 12], and antiviral HIV-1 [13–15]. The reactions leading to the formation of furanones are frequently reported by several research groups, however, they have proposed the reactions mechanism without offering evidence to support them [16–20]. There are a number of reasons for tendency to know reaction mechanisms. Knowledge of mechanisms aids the synthetic chemist in predicting by products and improving reaction conditions, the most powerful tool for the experimental study of reaction mechanisms is chemical kinetics. Kinetics is the study of the speed with which a chemical reaction occurs, and with all of the factors which influence these rates [21].

The spectroscopic technique is the largest and important techniques used in chemistry. The most widely method used for kinetic and mechanistic studies of chemical reactions is ultraviolet-visible (UV-visible) spectroscopy. Several research studies have documented over a large area of different reactions which have previously been reported using the UV-visible spectrophotometry technique [22–30]. In our previous work, experimental and theoretical kinetics studies and mechanism investigation have fully been discussed for various reactions [31–37]. In continuing our investigations on the kinetic of the organic reactions, we focused on 3, 4, 5-substituted furan-2(5H)-ones by one-pot multicomponent reaction that was reported by MT Maghsoodlou and coworkers

[38]. Depending upon the stoichiometry and conditions, other products have also been reported for this reaction [39-42]. However, under optimized conditions reported by MT Maghsoodlou and et al (**Fig. 1**) [38], 3, 4, 5-substituted furan-2(5H)-ones are obtained in high yields (%90-98). In the current work for the first time, we describe kinetic results together with the detailed mechanistic studies for the mentioned reactions based on a global kinetic analysis methodology using the UV-vis spectrophotometry apparatus. Also, this work focused especially on the effects of temperature, different para-substituted anilines on the activation parameters of preparation reaction of 3,4,5-substituted furan-2(5H)-ones.



**Fig. 1:** The three-component synthesis of furan-2(5H)-ones [38]

## 2. Experimental

### 2.1. Chemicals and Apparatus Used

All chemicals were used without further purification. Substituted-aniline **1**, diethyl acetylenedicarboxylate **2** and benzaldehyde **3** were supplied by Merck (Darmstadt, Germany), Acros (Geel, Belgium) and Fluka (Buchs, Switzerland). Extra pure formic acid and acetic acid were also obtained from Merck (Darmstadt, Germany).


A Cary UV-vis spectrophotometer model Bio-300 with a 10 mm light-path quartz spectrophotometer cell equipped with thermostated housing cell was used to record absorption spectra in order to follow reaction kinetics.

### 2.2. General procedure


**The first experiment** is performed with  $10^{-2}$  M solutions of aniline **1**, diethyl acetylenedicarboxylate **2** and benzaldehyde **3** in formic acid solvent. The progress of reaction is monitored by the absorbance changes as a function of wavelengths (**Fig. 2**). Since at wavelength of 420 nm, reactants **1**, **2** and **3** have no relatively absorbance values, it provides the opportunity to fully investigate the kinetics and mechanism of the reaction. Hence, the product absorbance changes with time are undertaken under the same condition at wavelength of 420 nm (**Fig. 3**) at 20°C. As can be seen in **Fig. 3**, the original experimental absorbance curve versus time (dotted line) is exactly fitted to second order fit curve (solid line). So the reaction is second order ( $\alpha + \beta + \gamma = 2$ ), the overall reaction rate is described by the kinetic equation:

$$\text{rate} = k_{\text{ovr}}[1]^{\gamma}[2]^{\beta}[3]^{\alpha} \quad (1)$$

Then the second-order rate constant ( $k_{\text{ovr}} = 3.50 \text{ min}^{-1} \cdot \text{M}^{-2}$ ) is automatically calculated using the standard equations within the program at  $20^\circ\text{C}$  [34].



**Fig. 2.** Absorption changes versus wavelength for the reaction Between **1** ( $10^{-2} \text{ M}$ ), **2** ( $10^{-2} \text{ M}$ ) and **3** ( $10^{-2} \text{ M}$ ) in formic acid for generation of product **4**



**Fig. 3.** The original experimental absorbance curve versus time at selected wavelength of 420 nm for the reaction between **1** ( $10^{-2} \text{ M}$ ) and **2** ( $10^{-2} \text{ M}$ ) with **3** ( $10^{-2} \text{ M}$ ) in formic acid. Dotted line is experimental values and the solid line is a fit curve

### 3. Results and Discussion:

#### 3.1. Effects of concentration

In order to obtain partial order with respect to aniline **1**, pseudo-order condition is performed for the reaction. So **in the second experiment**, we followed the reaction kinetics by plotting the UV-vis absorbance versus time at wavelength of 420 nm with **1** ( $10^{-3} \text{ M}$ ), **2** ( $10^{-2} \text{ M}$ ) and **3** ( $10^{-2} \text{ M}$ ) at  $20.0^\circ\text{C}$  in formic acid. For this case the rate law can be expressed:

$$\begin{aligned} \text{rate} &= k_{\text{ovr}}[1]^{\alpha}[2]^{\beta}[3]^{\gamma} \\ \text{rate} &= k_{\text{obs}}[3]^{\gamma} \\ k_{\text{obs}} &= k_{\text{ovr}}[1]^{\alpha}[2]^{\beta} \end{aligned} \quad (2)$$

All kinetic mathematical fits (the zero, first or second fit curve) are done using the software [43] associated with the UV-Vis spectrophotometer instrument. The original experimental absorbance against time data make a pseudo-first-order available fit curve at 420 nm, which exactly fits the experimental curve, **Fig. 4-A**. Herein, observation rate constant ( $k_{\text{obs}}$ ) is automatically calculated for the equation (2) by the software associated within the UV-vis instrument. It is obviously that the reaction is of the first order type with respect to aniline **1** ( $\alpha=1$ ).

Also, to gain the partial order of reaction with respect to reactant **3**, **third experiment** is done under pseudo-order condition [ $(10^{-2} \text{ M}, \mathbf{1})$ , [ $(10^{-2} \text{ M}, \mathbf{2})$ ] and [ $(10^{-3} \text{ M}, \mathbf{3})$ ]. In this experiment the original experimental absorbance versus time provides a second-order curve ( $k_{\text{ovr}} = 3.49 \text{ min}^{-1} \cdot \text{M}^{-2}$ , **Fig. 4-B**) like the first experiment. Due to the concentration changes of reactant **3** had no effect on the reaction rate, then reaction is zero-order with respect to reactant **3** ( $\gamma=0$ ). Also, in **fourth experiment**, we followed kinetics under pseudo order condition of reactant **2**. As can be seen in **Fig. 4-C**, the reaction shows first order kinetics that it means the partial order with respect to reactant **2** is one ( $\beta=1$ ). So, the reaction between aniline **1** and diethyl acetylenedicarboxylate **2** with benzaldehyde **3** follows second-order kinetics.

According to the obtained results from the four experiments, the rate law of reaction can be written:


from the first experiment:  $\alpha + \beta + \gamma = 2$

from the second experiment:  $\alpha = 1$


from the third experiment:  $\alpha + \beta = 2$  and  $\gamma = 0$

from the fourth experiment:  $\beta = 1$


rate =  $k_{\text{ovr}} [1] [2]$



**Fig. 4-A.** Plot of absorbance versus time at 420 nm for the reaction between **1** ( $10^{-2}$  M), **2** ( $10^{-2}$  M) and **3** ( $10^{-3}$  M) in formic acid. Dotted line is experimental values, and the solid line is fit curve



**Fig. 4-B.** Plot of absorbance versus time at 420 nm for the reaction between **1** ( $10^{-3}$  M), **2** ( $10^{-2}$  M) and **3** ( $10^{-2}$  M) in formic acid. Dotted line is experimental values, and the solid line is fit curve



**Fig. 4-C.** Plot of absorbance versus time at 420 nm for the reaction between **1** ( $10^{-2}$  mol.L $^{-1}$ ), **2** ( $10^{-3}$  mol.L $^{-1}$ ) and **3** ( $10^{-2}$  mol.L $^{-1}$ ) in formic acid, dotted line is experimental values, and the solid line is fit curve

### 3.2. Effect of solvents and temperature

The solvent dielectric constant and polarity are important parameters that influence rate constants. Solvent effects on the rate of reaction depend on the relative stabilization of the starting materials and the corresponding transition state through solvation [44, 45]. For investigation of the effect of medium dielectric constant on the rate of reaction, the same kinetic procedure is followed in the presence of acetic acid in 20 °C.

It is found that as medium dielectric constant decreased, the reaction rate reduced from formic acid ( $k_{\text{ovr}} = 3.50 \text{ min}^{-1} \cdot \text{M}^{-2}$ ) to acetic acid ( $k_{\text{ovr}} = 0.713 \text{ min}^{-1} \cdot \text{M}^{-2}$ ).

Temperature is considered a major factor that affects the rate of a chemical reaction. The influences of temperature on the reaction rate were studied in the range of 20° C to 60 °C with 10 °C intervals for each

reaction and the values of second-order rate constants are determined. Kinetic data are shown in **Table 1**. Reaction rate constant has a temperature dependency, which is usually given by the Arrhenius equation

Most reactions are found to obey the Arrhenius equation:

$$k = A \exp\left(\frac{-E_a}{RT}\right) \quad (3)$$

Taking the natural logarithm of both sides of the equation provides:

$$\ln k = \ln A - \frac{E_a}{RT} \quad (4)$$

Where  $E_a$  is the activation energy of reaction,  $R$  is universal gas constant ( $8.314 \text{ J K}^{-1} \text{ mol}^{-1}$ ), and  $A$  is frequency factor.

The activation energy of reaction is usually determined experimentally by plotting the graph of  $\ln k$  versus the reciprocal of the temperature  $1/T$ , which will yield a straight line with a slope of  $-\frac{E_a}{RT}$  and an intercept  $\ln A$  (**Fig. 5**). The highest activation energy ( $E_a = 80.12 \text{ kJ.mol}^{-1}$ ) is obtained for  $\text{NO}_2$ -aniline, that means reaction will occur slower than alone aniline and para-donating substituent-aniline. Nevertheless, the differences in activation energies are not much, because the substituents on aniline ring are far from the center of reaction. The Eyring equation is used in chemical kinetics to describe the variance of the rate of a chemical reaction with temperature and was used to evaluate activation parameters involving  $\Delta H^\ddagger$  (activation enthalpy),  $\Delta S^\ddagger$  (activation entropy) [46].

The linear form of the Eyring equation is given below:

$$\ln \frac{k}{T} = -\frac{\Delta H^\ddagger}{RT} + \frac{\Delta S^\ddagger}{R} + \ln \frac{k_B}{h} \quad (5)$$

That  $k_B$  = Boltzmann's constant,  $T$  = temperature,  $h$  = Planck's constant and  $R$  = universal gas constant

The values for  $\Delta H^\ddagger$  (activation enthalpy),  $\Delta S^\ddagger$  (activation entropy) can be determined from slopes and intercepts of  $\ln \frac{k}{T}$  versus  $1/T$  plot (**Fig. 6**). Obtained activation parameters for para- substituted aniline are given in **Tables 2** and **3**.

Also, the transition state equation was used to calculate activation parameters  $\Delta H^\ddagger$ ,  $\Delta S^\ddagger$  in order to check the results obtained from the two methods (Eyring and the transition state theory). According to the transition state theory, activation energy can be related to activation enthalpy (equation 6) and also pre-exponential factor is relevant to activation entropy according to equations (8) and (9), respectively [46]:

$$E_a = \Delta H^\ddagger + 2RT \quad (6)$$

$$A = \frac{e k_B T}{h} e^{\frac{\Delta S^\ddagger}{R} + 2} \quad (7)$$

$$\Delta S^\ddagger = R(\ln A - \ln\left(\frac{e^2 k_B T}{h}\right)) \quad (8)$$

The values obtained on the basis of transition state theory inferred that the activation enthalpy is the lowest for donating substituent- aniline that have the higher reaction rates and the highest for withdrawing substituent-aniline with the lowest reaction rates (**Table 3**). Note that the negative activation entropy indicated a greater degree of ordering in the transition state than in the initial state. These results suggest that the transition state in first step of reaction, **Fig. 8**, involving donating substituent-aniline is ordered more effectively stabilized than the transition state containing withdrawing substituent-aniline (**Tables 2 and 3**). The activation Gibbs free energy has also been evaluated from the following form of the Gibbs–Helmholtz equation:

$$\Delta G^\ddagger = \Delta H^\ddagger - T\Delta S^\ddagger \quad (9)$$

The activation free energy is essentially the energy requirement to get a molecule (or a mole of them) to undergo the reaction. It is of interest to note that activation free energies  $\Delta G^\ddagger$  for the different substituents are positive. The value of the activation free energy for donating substituent-aniline is lower than withdrawing substituent-aniline that means the reaction will occur harder for withdrawing substituent-aniline.

To investigate the impact of the different substituents on the rate constant of reaction and activation parameters, we changed the substituents R on aniline from the electron donating (Methoxy, Ethyl) to the electron withdrawing (Fluorine, Nitro) group. The reaction provides respective Schiff base and is reversible in absence of reactant **3**. The first step of reaction consists of Michael addition. According to the experimental observations, the first step is slow (please see mechanism section 3.3). Then, electron donating groups cause the aniline more nucleophilic and accelerate the reaction.

**Table 1.** Reaction rate constant ( $k_{\text{ovr}}$  ( $\text{min}^{-1} \cdot \text{M}^{-2}$ )) in different temperatures and different substituted-aniline under the same conditions for the reaction between **1** ( $10^{-2}$  M) and **2** ( $10^{-2}$  M) with **3** ( $10^{-2}$  M) in formic acid

( $\lambda$ ) (nm)	substituted-aniline	20.0 °C	30.0 °C	40.0 °C	50.0 °C	60.0 °C
420	OMe- aniline	8.10 (0.0028) <sup>a</sup>	24.31 (0.0057)	52.53 (0.0063)	115.42(0.0023)	210.31 (0.0039)
420	Ethyl- aniline	6.16 (0.0026)	15.9 (0.0068)	34.88 (0.0074)	91.97 (0.0064)	174.10 (0.0075)
420	aniline	3.50 (0.0046)	9.06 (0.0038)	20.40 (0.0038)	48.45 (0.0025)	115.35 (0.0037)
420	F- aniline	0.75 (0.0029)	2.10 (0.0057)	5.12 (0.0037)	12.87 (0.0010)	32.938 (0.0037)
420	NO <sub>2</sub> - aniline	0.44 (0.0022)	1.40 (0.0029)	3.42 (0.0026)	9.33 (0.0021)	23.84 (0.0072)


<sup>a</sup>: Standard deviation (SD)

**Table 2.** Activation parameters ( $\Delta S^\ddagger$ ,  $\Delta H^\ddagger$ ,  $\Delta G^\ddagger$  and  $E_a$ ) using the Arrhenius equation (3) and Eyring equation (5 and 6) in 20 °C


substituted-aniline	$\Delta H^\ddagger$ (kJ.mol <sup>-1</sup> )	$\Delta S^\ddagger$ (J.mol <sup>-1</sup> K <sup>-1</sup> )	$\Delta G^\ddagger$ (kJ.mol <sup>-1</sup> )	$E_a$ (kJ.mol <sup>-1</sup> )
<b>OMe- aniline</b>	63.08	-11.20	59.24	65.67
<b>Ethyl- aniline</b>	65.91	-4.67	64.92	68.51
<b>aniline</b>	68.17	-2.01	68.76	70.76
<b>F- aniline</b>	73.46	3.40	74.83	76.05
<b>NO<sub>2</sub>- aniline</b>	77.53	13.10	80.81	80.12

**Table 3.** Activation parameters ( $\Delta S^\ddagger$ ,  $\Delta H^\ddagger$ ,  $\Delta G^\ddagger$  and  $\ln A$ ) using the Arrhenius equation (3) and The transition state equation (6, 7 and 8) in 20 °C

substituted-aniline	$\Delta H^\ddagger$ (kJ.mol <sup>-1</sup> )	$\Delta S^\ddagger$ (J.mol <sup>-1</sup> K <sup>-1</sup> )	$\Delta G^\ddagger$ (kJ.mol <sup>-1</sup> )	$\ln A$
<b>OMe- aniline</b>	60.71	-18.87	66.34	29.15
<b>Ethyl- aniline</b>	63.55	-12.36	67.24	29.94
<b>aniline</b>	65.80	-9.68	68.69	30.26
<b>F- aniline</b>	71.09	-4.27	72.37	30.91
<b>NO<sub>2</sub>- aniline</b>	75.16	5.42	73.55	32.08



**Fig. 5.** Dependence of second order ( $\ln k_{\text{ovr}} = \ln k_1$ ) on reciprocal temperature for the reaction between reactants **1**, **2** and **3** in formic acid measured at wavelength 420 nm in accordance with Arrhenius equation




**Fig. 6.** Eyring plot ( $\ln \frac{k_{\text{ovr}}}{T}$  versus  $T$ ) according to equations (5) for the reaction between **1**, **2** and **3** in formic acid

### 3.3. Mechanism

In separate experiments, reactions between aniline **1** and diethylacetyldicarboxylate **2** (Re.1), aniline **1** and benzaldehyde **3** (Re. 2), and then diethylacetyldicarboxylate **2** and benzaldehyde **3** (Re. 3) are performed under the same concentration of each reactant ( $10^{-2}$  M) at  $20^\circ\text{C}$ . The Re. 1 is monitored by recording scans of the entire spectra with 5 min intervals in 45 min reaction time at  $20^\circ\text{C}$  (**Fig. 7-A**). The Re. 2 was monitored by recording scans of the entire spectra with 1 min intervals during the whole reaction time (5 min) at  $20^\circ\text{C}$  (**Fig. 7-B**). There are no reactions between reactants **2** and **3** due to the lack of any progress. Moreover, some  $^1\text{H}$ NMR information about the reaction products is presented in supplementary file, and also the about first intermediate **I<sub>1</sub>** was reported previously [47, 48]. According to these references, the first step of the reaction consists of Michael addition and is slow which respective diethyl 2-aryloamino-fumarate **I<sub>1</sub>** can be readily synthesize and characterized according to known protocols and can be purified by convenient distillation, then, **I<sub>1</sub>** can be used for subsequent steps [49, 50]. As a result of these preliminary investigations and experimental observations, the speculative mechanism containing three steps with starting reaction between reactants **1** and **2** (the more rapidly occurring reaction amongst competing reactions) is proposed in **Fig. 8**.





**Fig. 7-A:** Changes in absorption spectra vs. wavelength in 20°C for the reaction between aniline **1** and diethylacetylenedicarboxylate **2** in formic acid as reaction proceeds into a 10 mm light-path cell. The total acquisition time was 45 min; the time increment was set to 5 min. Herein, the upward of direction of the arrow indicate that the progress of reaction

**Fig. 7-B:** Changes in absorption spectra vs. wavelength in 20°C for the reaction between aniline **1** and benzaldehyde **3** in formic acid as reaction proceeds into a 10 mm light-path cell. The total acquisition time was 5 min; the time increment was set to 1min. Herein, the upward of direction of the arrow indicate that the progress of reaction

For the reaction, a steady state treatment is coupled with the kinetic observations. So, the rate law is written using the final step of the proposed mechanism in **Fig. 8** for the generation of product **4**:

$$\text{rate} = k_5[I_4] \quad (10)$$

Also, the overall reaction can be written as:

$$\text{rate} = k_1[1][2] \quad (11)$$

This makes it possible to accept that the first step is the rate-determining step (*RDS*) of the reaction and the overall order of reaction is two (since  $k_1$  is the only rate constant which appeared in the equation (11)). This equation, obtained from the proposed mechanism, is same with the rate law that resulted from UV-vis experiment ( $\text{rate} = k_{\text{ovr}}[1][2]$ ). It is possible to analyze the effect of the solvent on the reaction, because the first step ( $k_1$ ) is the rate determining step, in this case in both solvents the transition state partially charged, in solvent with higher dielectric constant the transition state is highly solvated relative to the reactant molecules that have no any charge.

If the second step to be a rate determining step, therefore, the first step is fast with forward rate constant ( $k_1$ ) and return rate constant ( $k_{-1}$ ). We can write the rate law:

$$\text{rate} = k_2[I_1][3], \frac{d[I_1]}{dt} = k_1[1][2] - k_{-1}[I_1] - k_2[I_1][3] = 0 \quad (12)$$

$$[I_1] = \frac{k_1[1][2]}{k_2[3] + k_{-1}} \quad (13)$$


$$\text{rate} = \frac{k_2 k_1 [1][2][3]}{k_2[3] + k_{-1}} \quad (14)$$

According to the above assumption,  $k_2$  is a rate determining step, therefore it is logical to express  $k_{-1} \gg k_2[3]$ .

As a result, rate law becomes:

$$\text{rate} = \frac{k_2 k_1 [1][2][3]}{k_{-1}} \quad (15)$$

Herein, the overall order of recent rate law is three and is not compatible with the rate law arising from UV-vis experiment ( $\text{rate} = k_{\text{ovr}} [\mathbf{1}] [\mathbf{2}]$ ), so the second step could not be a rate determining step.



**Fig. 8.** Speculative mechanism for the reaction between aniline **1** and diethylacetylenedicarboxylate **2** with para-substituted benzaldehyde **3** for generation of product **4** in formic acid

#### 4. Conclusions

Kinetics and mechanism of the reactions between para-substituted anilines **1** and diethyl acetylenedicarboxylate **2** with benzaldehyde **3** were investigated in formic acid and the below results were obtained:

A mechanism involving three steps was proposed for the reactions. The overall order of reaction followed second-order kinetics and the partial orders with respect to para-substituted aniline, diethyl acetylenedicarboxylate and benzaldehyde were one, one and zero, respectively. Also, the first step of the reaction is recognized as a rate-limiting step according to the experimental data. The electron donating group on

substituted aniline more stabilized the transition state than the electron withdrawing group and with decreasing the activation energy the value of rate constant ( $k_1$ ) was speed up. The rate of reaction was also accelerated by increasing the temperature and was in agreement with Arrhenius, Eyring equations and the transition state equations.

### Acknowledgment

We gratefully acknowledge financial support from the Research Council of the University of Sistan and Baluchestan.

### References

- [1] Carter NB, Nadany AE, Sweeney JB, (2002) *J Chem Soc Perkin Trans 1*: 2324.
- [2] Chen QH, Praveen PN, Knaus EE, (2006) *Bioorg Med Chem* 14:7898.
- [3] Boukouvalas J, Cote S, Ndzi B, (2007) *Tetrahedron Lett* 48: 105.
- [4] Zarghi A, Praveen PN, Knaus EE, (2007) *Bioorg Med Chem* 15: 1056.
- [5] Braun M, Hohmann A, Rahematpura J, Buhne C , Grimme S, (2004) *Chem Eur J* 10:4584.
- [6] Bellina F, Falchi E, Rossi R, (2003) *Tetrahedron* 59:9091.
- [7] Takahashi S, Kubota A, Nakata T, (2002) *Tetrahedron Lett* 43: 8661.
- [8] Padakanti S, Pal M, Yeleswarapu KR, (2003) *Tetrahedron* 59: 7915.
- [9] Hein SM, Gloer JB, Koster B, Malloch D J, (2001) *Nat Prod* 64: 809.
- [10] Pour M, Spulak M, Balsanek V, Kunes J, Kubanova P, Butcha V, (2003) *Bioorg Med Chem* 11:2843.
- [11] Levy LM, Cabrera GM, Wright JE, Seldes AM, (2003) *Phytochemistry* 62:239.
- [12] Grossmann G, Poncioni M , Bornand M, Jolivet B, Neuburger M, Sequin U, (2003) *Tetrahedron* 59:3237.
- [13] Hanessian S, Park RY, Yang RY, (1997) *Synlett* 4:351.
- [14] Hanessian S, Park RY, Yang RY, (1997) *Synlett* 4:353.
- [15] Choudhury A , Jin F , Wang D , Wang Z , Xu G, Nguyen D , Castoro J , Pierce ME , Confalone PN, (2003) *Tetrahedron Lett* 44: 247.
- [16] Hazeri N, Maghsoodlou M T, Mahmoudabadi N, Doostmohammadi R, Salahi S, (2014) *Current Organocatalysis*, 1: 45-50.
- [17] Doostmohammadi R, Maghsoodlou M T, Hazeri N, Habibi-Khorassani S M, (2013) *Research on Chemical Intermediates*, 39:4061.
- [18] Doostmohammadi R, Hazeri N, (2013) *Letters in Organic Chemistry*, 10:199.
- [19] Narayana M, S. et al, (2009) *Tetrahedron*, 65:5251.
- [20] Mohammad RM et al, (2014) *Comptes Rendus Chimie*, 17:131.
- [21] Carey FA, Sundberg RJ *Advanced Organic Chemistry, Part A: Structure Mechanisms* Kluwer Academic Press, New York, 2000.
- [22] Nielsen OJ, Sehested J, Langer S, Ljungström E, Wangberg I (1995) *Chem Phys Lett* 238:359.
- [23] Ivanov VV, Decker C (2001) *Polym Int* 50:113.
- [24] Almandoz MC, Da'vila YA, Sancho MI, Gasull EI, Blanco SE (2010) *Spectrochim Acta Part A* 77:51.
- [25] Lu LL, Li YH, Lu XY (2009) *Spectrochim Acta Part A* 74:829.

- [26] Erasmus JJC, Conradie MM, Conradie J (2012) *React Kinet Mech Cat* 105:233.
- [27] Kalyan BP, Gupta KS (2012) *Transit Met Chem* 37:671.
- [28] Chandra AK (2012) *J Mol Model* 18:4239.
- [29] Sandhiya L, Kolandaivel P, Senthilkumar K (2012) *Struct Chem* 23:1475.
- [30] Pandeewaran M, Elango KP (2009) *J Solut Chem* 38:1558.
- [31] Habibi Khorassani SM, Ebrahimi A, Maghsoodlou MT, Shahraki M, Price D, (2011) *Analyst*, 136:1713.
- [32] Dehdab M, Habibi-Khorassani SM, Shahraki M (2014) *Catalysis Letters*, 144:1790.
- [33] Habibi-Khorassani SM, Maghsoodlou MT, Aghdaei E, Shahraki M (2012) *Prog React Kinet Mech* 37:301.
- [34] Shahraki M, Habibi-Khorassani SM, Ebrahimi A, Maghsoodlou MT, Paknahad A (2012) *Prog React Kinet Mech* 37:321.
- [35] Shahraki M, Habibi-Khorassani SM, (2015) *J. Phys. Org. Chem.* doi: 10.1002/poc.3424
- [36] Habibi-Khorassani S M, Ebrahimi A, Maghsoodlou M T, Asheri O, Shahraki M, Akbarzadeh N, Ghalandarzehi, Y (2013) *Int J Chem Kinet*, 45:596.
- [37] Pourpanah SSh, Habibi-Khorassani SM, Shahraki M (2015) *Chinese Journal of Catalysis* 36:757.
- [38] Doostmohammadi R, Maghsoodlou MT, Habibi-Khorassani SM (2012) *Iranian Journal of Organic Chemistry* 4:939.
- [39] Lv, Longyun et al, (2013) *ACS Combinatorial Science*, 15:183.
- [40] Sarkar R, Mukhopadhyay Ch, (2013) *Tetrahedron Letters*, 54:3706.
- [41] Zhang M, Huan-Feng J, (2008) *European Journal of Organic Chemistry*, 20: 3519.
- [42] Zhang M et al, (2007) *Synlett*, 20:3214.
- [43] Nielsen OJ, Sehested J, Langer S, Ljungström E, Waˆngberg I (1995) *Chem Phys Lett* 238:359.
- [44] Lerveny C, Ruzicka V, (1981) *Adv Catal* 30:335.
- [45] Mortimer M and Taylor P, *Chemical Kinetics and Mechanism*, The Royal Society of Chemistry, Thomas Graham House, Science Park, Milton Road, Cambridge, CB4 0WF, UK, 2002.
- [46] Laidler KJ, *Theories of Chemical Reaction Rates*; McGraw-Hill: New York, 1941.
- [47] Mashaly M M, El-Gogary S R, Kosbar TR, (2014) *J. Heterocyclic Chem* 51:1078.
- [48] Huang C, Yin Y, Guo J, Wang J, Fan B, Yang L, (2014) *RSC Adv* 4:10188.
- [49] Safaei-Ghomi J, Salimi F, Ramazani A, Nasrabadi FZ, Ahmadi Y, (2012) *Turk J Chem* 36:485.
- [50] Domschke G, (1980) *Z Chem* 20:16.