

RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Journal Name

COMMUNICATION

p-Nitrobenzenesulfonamides and their Fluorescent Dansylsulfonamides derived from *N*-Alkylated *o*-(Purine-Methyl)Anilines as Novel Antitumour Agents

 Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

 Fátima Morales,^a Ana Conejo-García,^{a,b} Alberto Ramírez,^c Cynthia Morata,^{d,e} Juan Antonio Marchal,^{d,e,*} and Joaquín M. Campos^{a,b,*}

We have designed and synthesized a novel series of fourteen alkylated purines with notable *in vitro* anti-proliferative activities against three cancerous cell lines. *In vivo* studies show absorption by the oral route of administration with systemic distribution and low toxicity for the two more active derivatives.

Cancer is one of the most important health issues and the leading cause of death worldwide. Collaboration between researchers, epidemiologists and clinicians is essential to improve cancer control. Recent advances in the knowledge of cancer development have allowed researchers to determine that tumours are constituted by differentiated tumour cells and their precursor cells, also named as tumour-initiating cells or cancer stem cells (CSCs). These CSCs are a small tumour subpopulation responsible for initiating and maintaining tumour growth and are characterized by their self-renewal capacity and differentiation ability, the high metastatic potential and their quiescent state makes them resistant to chemo- and radio-therapy.¹ Conventional anticancer therapies kill the rapidly proliferating bulk differentiated cancer cells but obviate the quiescent CSCs. Therefore, development of specific therapies targeted at CSCs holds hope to gain the fight against cancer.

We have previously described a series of cyclic and acyclic *O,N*-acetals (**1** and **2**, Figure 1) as potent anti-tumour compounds.^{2–5} The IC₅₀ values against the MCF-7 cell line were below 1 μM, **1** also called bozepinib, and **2** being the most potent structures so far reported (IC₅₀ = 0.355 μM in both cases). Moreover, they




Figure 1. The arrows and the legends on top of them show the corresponding structural modifications carried out for the design of the target molecules.

^a Department of Pharmaceutical and Organic Chemistry, c/ Campus de Cartuja s/n, Faculty of Pharmacy, University of Granada, 18081 Granada, Spain. Fax: +34 958 24 38 45, Tel: +34 958 24 38 50. E-mail: jmcampos@ugr.es..

^b Biosanitary Institute of Granada (ibs.GRANADA), SAS-Universidad de Granada, 18071 Granada, Spain.

^c Department of Health Sciences, Paraje de las Lagunillas s/n, University of Jaén, Jaén, Spain.

^{d,e} Biopathology and Medicine Regenerative Institute (IBIMER), University of Granada, Granada, Spain; Biosanitary Institute of Granada (ibs.GRANADA), 18071 Granada, Spain. Fax: +34 958 24 62 96, Tel: +34958 24 93 21. E-mail: jarmarchal@ugr.es.

Electronic Supplementary Information (ESI) available: See DOI: 10.1039/x0xx00000x

are the most selective compounds against the MCF-7 cell line [TIs (Therapeutic Index) = 5.14 and 5.25, respectively] in relation to the normal one.² cDNA microarray technology reveals potential drug targets, mainly centred on positive apoptosis regulatory pathway genes, and the repression of genes involved in carcinogenesis, proliferation, tumour invasion and CSCs related pathways.³ Bozepinib is a potent anti-tumour agent that induces apoptosis in breast and colon cancer cells mediated by the double-stranded RNA-dependent protein kinase (PKR).⁴ The combination with IFN α potentiates the apoptosis induced by bozepinib and also enhances autophagy and senescence,⁴ both processes of great importance in tumour cells that show resistance to conventional chemotherapy. Bozepinib is able to exert an inhibitory activity against the formation of both mammo- and colono-spheres and eliminates aldehyde dehydrogenase enriched (ALDH⁺) CSC subpopulations at a low micromolar range. Moreover, the inhibitory effect of bozepinib on CSC pathways was emphasized by the down-regulation of c-Myc, β -CATENIN and SOX2 proteins and the up-regulation of the hedgehog-signalling repressor form of GLI-3. Bozepinib also shows *in vivo* anti-tumour and anti-metastatic efficacy in xenotransplanted nude mice without presenting sub-acute toxicity.⁵

In the present paper we present a new series of acyclic compounds with the aim to develop new anti-tumour agents based on the following modifications of the acyclic analogue of bozepinib (**2**), which in spite of being obtained as a by-product during the formation of bozepinib (**1**), its remarkable anti-proliferative properties² justify to be considered as a new prototype: in short, the change of position of the purine base to the benzylic carbon and concomitant replacement of the *O,N*-acetalic function for an *O,O*-acetalic one, in the form of the 1,3-dioxolane group (Figure 1). The anti-tumour activity has recently been described of a series of derivatives having a 1,3-dioxolane moiety in their structures (**3**). Such compounds are inhibitors of the proliferation and eradication of cancer cells, besides being inhibitors of CSCs.⁶ We maintain the *p*-nitrobenzenesulfonyl group and we introduce a linker between the sulfonamide and the dioxolane ($n = 1, 2$). We also include the 5-FU derivative (**13**) in order to act as a reference (Figure 2).

The target compounds **4–13** (Figure 2) are obtained following the synthetic procedure described in Scheme S1 ($R_1 = A$; $R_2 = C, D, E$ and F). Compounds **18–20** were synthesized as previously described.⁷ When **21** and **22** are treated with tetrabutylammonium fluoride (TBAF) in anhydrous THF to give the de-protected compounds **25** and **26**, by-products **29** and **30** are also obtained in smaller yields respectively (Scheme S2). Compounds **29** and **30** are the result of a Smiles rearrangement,⁸ in which the *p*-nitrophenyl group migrates to the hydroxyl group through the formation of the mono-nitro spiro-complex (Scheme S2).⁹ Target compounds **4–13** were obtained by microwave-assisted Mitsunobu reaction. Microwave-assisted organic synthesis is becoming instrumental for the rapid synthesis of drugs.¹⁰ The two following facts must be underlined: a) Reaction between

26 and adenine yields, apart from the *N*-9 derivative **11**, the *N*-3 derivative **12**;¹¹ and b) Reaction between **27** (and **28**) and 2,6-dichloropurine yields both the *N*-9 derivative **14** (and **15** when starting from **28**), and the *N*-7 derivative **16** (and **17** when starting from **28**).




Figure 2. The fourteen molecules studied in this communication.

The effects on cell proliferation were investigated on the human breast cancer cell line MCF-7, human colon carcinoma cell line HCT-116 and the human malignant melanoma cell line A-375. The anti-proliferative activities for **4–17** against the three tumour cell lines are shown in Table S1 in μM units. The three most promising compounds are **8**, **9** and **10** (with IC_{50} values ranging from 0.134 μM against A-375, and 4.110 μM against HCT-116 for **10**).

Fluorescence is a very useful technique to elucidate the mechanism of action of drugs inside the cell and to know their distribution in tissues and organs. Knowing drug distribution is critical for the design of more selective and efficient agents in cancer therapy, although this knowledge is neither always easy nor cheap. However, fluorescence technique provides a comprehensive tool for investigating these aspects in single cells and whole tissue.¹² Some drugs are inherently auto-fluorescent and therefore can be tracked using microscopy techniques, such as topotecan and camptothecin¹³ but others

need to be labelled with chromophores such as the ones shown in Figure S1. As an example, the mechanism of action of the anti-tumour drug paclitaxel by the stabilization of the microtubules was disclosed by the fluorescence techniques using fluorescein as a label in the drug.¹⁴ We have chosen the *p*-nitrobenzene sulfonyl derivative with $n=2$ and the 2,6-dichloropurine moiety (**10**), because it is the most active compound against the A-375 cell line, in order to study the *in vitro* and *in vivo* drug distribution by fluorescence. Epidemiological studies showed that the incidence of melanoma is increasing at a rate faster than that of any other cancers worldwide.¹⁵ Therefore we need to design a fluorescent analogous of the target compound **10**. The *p*-nitrobenzenesulfonyl group was interchanged with the dansyl group as a chromophore because of its similarity (**15**, Figure 2). Moreover, a fluorescence study of some anti-tumour drugs with a dansyl group in their structure that present no toxicity *in vivo*¹⁶ confirms the benefit of the use of this chromophore in the design. We also include the dansyl derivative with $n=1$ (**14**) in order to complete the SAR studies. For the synthesis, see Scheme S1, $R_1 = B$, and $R_2 = C$ and D . We can establish the following SARs for all the target compounds **4-17**: The IC_{50} values of compounds with $n=2$ (**8-10**, **15**, **17**) are lower than those of compounds with $n=1$ (**4-6**, **14-16**) and this fact might be attributed to the increase of lipophilicity.¹⁷ The most active compounds against all cell lines assayed are the 6-chloro (**8**), 6-bromo (**9**) and 2,6-dichloropurine (**10**, **15-17**) derivatives, as it has been previously reported.¹⁸ The *N*-3 adenine isomer (**12**) is slightly more active than the *N*-9 one (**11**). The 5-FU derivative (**13**) is the least active as anti-proliferative agent against all the cell lines. It should be pointed out that all the target compounds show better anti-proliferative activity against the human melanoma cell line with sub-micromolar IC_{50} values. Compound **10** is the best anti-proliferative agent of these *p*-nitrobenzenesulfonyl derivatives with an IC_{50} of 134 nM against the A-375 cell line. Dansyl derivatives (**14-17**) are also potent anti-proliferative agents against all cell lines with IC_{50} values between 0.3–3 μ M, with the sole exception of **14** against the MCF-7 cell line ($IC_{50} = 10.40 \mu$ M). Interestingly, when we tested the anti-proliferative activities for **8** and **10** on the A-375 isolated cell line, based on its ALDH activity, it was found that **10** shows a potent anti-proliferative activity against this isolated subpopulation (Table S2).

The effects on cell cycle distribution of the most active compounds (**4-6**, **8-10**, **14-17**) were analyzed by flow cytometry for 24 and 48 h at $3 \times IC_{50}$ in the A-375 cell line (Table S3). Compounds **5** and **8-10** provoked a G_2/M cell cycle arrest at the expense of the other two phases (26.60%, 34.40%, 37.27% and 21.36%, respectively, in contrast to the 8.71% of the control). The induction of the G_2/M cell cycle arrest by three compounds previously synthesized by us was associated with increased phosphorylation of eIF2 α in human breast cancer cells.¹⁹

We then analyzed the apoptotic effects of the most active compounds (**4-6**, **8-10**, **14-17**) in the A-375 cell line by flow cytometry after 24 and 48 h of treatment, in order to study the mechanisms of their anti-tumour activity. The results are

shown in Table S4. The most apoptotic compounds are **8-10**, that are also the most active as anti-proliferative agents.

Microscopy is a useful tool to study drug distribution in biological tissues.²⁰ We subsequently investigated the *in vitro* and *in vivo* distribution of **15**, which is the fluorescent analogous of **10**, in three tumour cell lines, in mice organs and tissues. MCF-7 breast cancer cells, HCT-116 colon cancer cells and A-375 melanoma cells were treated with compound **15** and absorbance was measured from 30 min to 2 h. Our results showed an increase in uptake quantities in treated cells that was time-dependent, being higher for MCF-7 at 30 min and 1 h. However, not significant differences were found after 2 h of treatment with **15** for the three tumour cell lines (Figure 3A).




Figure 3A. *In vitro* distribution of **15** in MCF-7 breast cancer cells, HCT-116 colon cancer cells and A-375 melanoma. **B.** Confocal imaging in A-375 melanoma cells treated with a concentration of 10 μ M of **15**.

Clearly, it is important to directly monitor the means by which **15** penetrates the cell membrane and whether it penetrates cell nuclei. To study the entry and distribution of **15**, confocal imaging was used in the A-375 melanoma treated-cells. (Figure 3B, Movies S1 and S2). Images show that after 1 h, compound **15** was internalized in a high percentage of cells and it was distributed in the perinuclear zone. After 2 h most of the cells showed enhanced fluorescence intensity and in some of them, **15** was located inside the nucleus. After 4 h a significant fraction of **15** was identified inside the cell nuclei with additional portions surrounding the perinuclear region (Figure 4). Molecules observed in the cytoplasm appear as bright blue

spots, suggesting that they were embedded by cells via endocytosis.




Figure 4. Dansyl distribution inside the cell nuclei (arrows) of **15** after 4 h.

Toxicity was determined after selecting the most active compound (**10**) that presents an IC_{50} value of $0.134 \mu M$ against A-375 cells. Its fluorescent analogue **15** was included in this study. We examined the acute-toxicity profile of **10** and **15** in BALB/c mice when they were administered in a single oral bolus each one ($n = 40$) at doses levels of 50-250 mg/kg for a week. Compounds **10** and **15** were non-toxic, even at the highest bolus dose of 250 mg/kg after 2 weeks. All the 40 **10**-treated and **15**-treated mice remained healthy throughout the 14-day observation period with no evidence of morbidity. For *in vivo* organ distribution analysis, mice were treated with **10** and **15** (100 mg/kg) by oral administration and fluorescence was detected in the organs extracted by necropsy after 6 h (Figure 5A).




Figure 5. *In vivo* distribution study of **10** and **15** using the IVIS spectrum imaging system. A. Distribution of **10** in several organs. A drop of **15** was used as positive control. B. Distribution of **15** in several organs.

Fluorescence intensity was high in all organs and tissues of mice treated with **15**, with the exception of pancreas and lungs (Figure 5B). Fluorescence intensity was not detected in mice treated with **10**, used as a control (Figure 5A). Results confirm that the drug is absorbed by the oral route of administration with systemic distribution. Furthermore the drug accumulates mainly in liver and kidney, the organs associated to drug metabolism and excretion, respectively.

Conclusions

A series of *p*-nitrobenzenesulfonyl and dansyl derivatives have been designed, synthesized and biologically evaluated as anti-tumour agents. Derivatives with a 6-bromo-, 6-chloro- and 2,6-dichloro-purine show better anti-proliferative activities against all the assayed cell lines than the adenine and 5-FU derivatives. Compounds with a dimethylene linker ($n = 2$) are better anti-proliferative agents than those with a methylene linker ($n = 1$). 2,6-Dichloropurine derivative with $n = 2$ and the *p*-nitrobenzenesulfonyl moiety (**10**) is the most active compound as an anti-proliferative agent with an IC_{50} of 134 nM against the A-375 cell line. In the A-375 ALDH⁺ CSC subpopulation **10** shows a high anti-proliferative activity. The most apoptotic compounds are the purine derivatives with $n = 2$ and they provoked a G₂/M cell cycle arrest. Moreover, the 6-bromopurine derivative (**9**) is the most potent apoptosis inductor so far reported by us.

The fluorescence of *N*-9 2,6-dichloropurine derivative with $n = 2$ (**15**) was detected by fluorimetric assays in cell cultures. Confocal imaging showed fluorescent *N*-9 dichloropurine derivative with $n = 2$ distribution in cell. The fluorescent compound seems to be incorporated to cell by an endocytic process. We observed vesicle transportation through cell cytoplasm and these vesicles accumulate in the perinuclear zone, and at late time (4 h) inside nuclei. It has been demonstrated that **10** and its fluorescent analogue **15** are nontoxic to BALB/c mice, even at the highest bolus dose of 250 mg/kg after 2 weeks. Compound **15** is absorbed by the oral route of administration with systemic distribution. Furthermore the drug accumulates mainly in liver and kidney, the organs associated to drug metabolism and excretion respectively. Nevertheless, deeper studies such as ADME assays are necessary to obtain a more accurate distribution of **15**. Our findings support the potent anti-tumour ability of the fluorescent derivative (**15**) which has permitted to conduct single-cell imaging of this drug upon binding to its targets in differentiated bulk tumour cells and CSCs. These results encourage further studies in molecular CSC pathways and suggest that this molecule could be a new CSC-targeted drug.

We thank the Instituto de Salud Carlos III through projects no. PI10/00592 and PI10/02295 and the European Regional Development Fund for financial supports. The award of a grant from the Ministerio de Educación to F. M. is gratefully acknowledged.

Notes and references

- 1 V. Tirino, V. Desiderio, F. Paino, A. De Rosa, F. Papaccio, M. La Noce, L. Laino, F. De Francesco, G. Papaccio, *FASEB J*, 2013, **27**, 13.
- 2 L. C. López-Cara, A. Conejo-García, J. A. Marchal, G. Macchione, O. Cruz-López, H. Boulaiz, M. A. García, F. Rodríguez-Serrano, A. Ramírez, C. Cativiela, A. I. Jiménez, J. M.

- García-Ruiz, D. Choquesillo-Lazarte, A. Aránega, J. M. Campos, *Eur J MedChem*, 2011, **46**, 249.
- 3 O. Caba, M. Díaz-Gavilán, F. Rodríguez-Serrano, H. Boulaiz, A. Aránega, M. A. Gallo, J. A. Marchal, J. M. Campos, *Eur J MedChem*, 2011, **46**, 3802.
- 4 J. A. Marchal, E. Carrasco, A. Ramírez, G. Jiménez, C. Olmedo, M. Perán, A. Agil, A. Conejo-García, O. Cruz-López, J. M. Campos, M. A. García, *Drug Des Devel Ther*, 2013, **7**, 1301.
- 5 A. Ramírez, H. Boulaiz, C. Morata-Tarifa, M. Perán, G. Jiménez, M. Picón-Ruiz, A. Agil, O. Cruz-López, A. Conejo-García, J. M. Campos, A. Sánchez, M. A. García, J. A. Marchal, *Oncotarget*, 2014, **5**, 3590.
- 6 M. Annette, S. Patil, N. S. Malik, K. D. Panghavan, *PCT Int Appl.*, 2012, WO 2012081039 A1 20120621.
- 7 M. Díaz-Gavilán, F. Rodríguez-Serrano, J. A. Gómez-Vidal, J. A. Marchal, A. Aránega, M. A. Gallo, A. Espinosa, J. M. Campos, *Tetrahedron*, 2004, **60**, 11547.
- 8 K. Bowden, P. R. Williams, *J Chem Soc Perkin Trans*, 1991, **2**, 215.
- 9 L. Fitzgerald, R. L. Blakeley, B. Zerner, *ChemLett*, 1984, 29.
- 10 A. Conejo-García, M. C. Núñez, J. A. Marchal, F. Rodríguez-Serrano, A. Aránega, M. A. Gallo, A. Espinosa, J. M. Campos. *Eur J MedChem*, 2008, **43**, 1742.
- 11 M. E. García-Rubiño, M. C. Núñez-Carretero, D. Choquesillo-Lazarte, J. M. García-Ruiz, Y. Madrid, J. M. Campos. *RSC Adv*, 2014, **4**, 22425.
- 12 N.S. Whitea, R. J. Erringtonb, *Adv Drug Deliv Rev*, 2005, **57**, 17.
- 13 H. M. Coley, W. B. Amos, P. R. Tewntyman, *Br J Cancer*, 1993, **67**, 1316.
- 14 M. P. Lillo, O. Cañadas, R. E. Dale. *Biochemistry*, 2002, **41**, 12436.
- 15 M. S. Goldberg, J. T. Doucette, H. W. Lim, J. Spencer, J. A. Carucci, D. S. Rigel. *J Am Acad Dermatol*, 2007, **57**, 60.
- 16 C. H. Chui, Q. Wang, W. C. Chow, M. C. Yuen, K. L. Wong, W. M. Kwok, G. Y. Cheng, R. S. Wong, S. W. Tong, K. W. Chan, F. Y. Lau, P. B. Lai, K. H. Lam, E. Fabbri, X. M. Tao, R. Gambari, W. Y. Wong, *ChemCommun*, 2010, **46**, 3538.
- 17 R. Sánchez-Martín, J. M. Campos, A. Conejo-García, O. Cruz-López, M. Báñez-Coronel, A. Rodríguez-González, M. A. Gallo, J. C. Lacal, A. Espinosa. *J Med Chem*, 2005, **48**, 3354.
- 18 M. C. Núñez, F. Rodríguez-Serrano, J. A. Marchal, O. Caba, A. Aránega, M. A. Gallo, A. Espinosa, J. M. Campos, *Tetrahedron*, 2007, **63**, 183.
- 19 A. Conejo-García, M. E. García-Rubiño, J. A. Marchal, M. C. Núñez, A. Ramírez, S. Cimino, M. A. García, A. Aránega, M. A. Gallo, J. M Campos, *Eur J Med Chem*, 2011, **46**, 3795.
- 20 C. A. Thorling, Y. Dancik, C. W. Hupple, G. Medley, X. Liu, A. V. Zvyagin, T. A. Robertson, F. J. Burczynski, M. S. Roberts, *J Biomed Opt*, 2011, **16**, 086013.