

RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



ARTICLE

One-Pot C-C/C-O Bonds Formation: Synthesis of Spirocyclic Lactones

Pedireddi Niharika and Gedu Satyanarayana*

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

An efficient and a practical method for the synthesis of novel spiro(tri or tetra)cyclic lactones via the formation of C–C and C–O bonds in one-pot is presented. The method was successful under Lewis acidic (FeCl₃) conditions and enabled the formation of spiro(tri or tetra)cyclic lactones from simple aliphatic exocyclic enoate esters as reacting partners with phenols. Remarkably, the usual self-aromatization of cyclohexanone based enoate esters, under such Lewis acidic conditions is overridden by intermolecular coupling. Significantly, the method was amenable on indanone derived esters as well and furnished novel spiro(tetra or penta)cyclic lactones bearing simple to dense functionalities on the aromatic rings. Notably, these novel spirocyclic systems constitute core structures of natural/unnatural compounds that show good biological properties.

Introduction

Organic chemistry has always dealt with exploring innovative, alternate and new methods for the construction of C–C and/or C–heteroatom bonds. The focus has always been to perform multiple transformations in a single vessel employing one or more catalysts.¹ In this context, domino one-pot methods² have acclaimed immense interest as they involve the construction of two or more bonds in a single vessel reducing the chemicals for a sequence of product purifications. It also lowers the loss in yield thus saves time and resources.

One-pot methods involving the Lewis acid mediated domino reactions are highly promoted in the synthetic field. As a part of our ongoing research interest in domino or sequential one-pot transformations, recently, we reported an efficient method for the synthesis of dihydrocoumarins.³ The process was based on the Lewis acid (FeCl₃) promoted dual bond (C–C and C–O) formation in one-pot. Also, our group reported some interesting acid mediated transformations for the synthesis of indanones⁴, indanes^{5a} and fused indenoindole tetracyclics.⁶ Significantly, in the dihydrocoumarin's report³, for the first time, we synthesized the novel spiro(tetra or penta)cyclic compounds starting from tetralone derived enoate ester (Scheme 1). Quite interestingly, this is not usual under either strong acidic or basic conditions, because, tetralone derived systems generally undergo rapid self-aromatization under such harsh acidic conditions. Inspired by the versatility of the method to furnish dihydrocoumarins and the spiro(tetra or

penta)cyclic compounds dexterously, it instigated us to apply this method on the aliphatic exocyclic α,β -unsaturated esters, wherein, again, aromatization was a serious concern, particularly, for those derived from cyclohexanone. It was envisioned that one-pot intermolecular Michael addition and intramolecular condensation of the esters with the phenols, would result in the direct construction of the spirocyclic lactones. Although there are a reasonable number of reports on the synthesis of aryl-dihydrocoumarins, from cinnamates/cinnamic acids in conjunction with the external phenols under acidic conditions,^{7,8,9} significantly, to the best of our knowledge there are no reports for direct access to the synthesis of spirocyclic lactones from simple aliphatic exocyclic enoate esters.

Often, it is reported that the spirocyclic compounds have an advantage and the twist around the centrally placed carbon atom discloses some unique properties mostly applicable in the pharmaceuticals. The conformational rigidity of these spirocyclic systems has an added advantage in binding to the protein molecules and as they form the integral part of various naturally occurring compounds having biological significance, their synthesis has been a big source of interest. Some of the spirocyclic compounds are reported to display narcotic, skeletal muscle relaxant, hypotensive and anti-viral properties.¹⁰ Very commonly used antihypertensive, diuretic drug (aldactone) and aldosterone antagonist (eplerenone) comprise the spirocyclic-core responsible for depicting these properties. Significantly, the spiro chiral ligands are also extensively implemented in the asymmetric catalysis for the synthesis of spiro-heterocyclic compounds.¹¹ Amongst the various spirocyclic systems, the heterocyclic spiro compounds are known to be highly significant for their drug like properties¹² and also for their structural novelty.¹³


^a Address here.

^b Address here.

^c Address here.

† Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x



Scheme 1: Present study vs our previous results for the synthesis of spirocyclic lactones.

The nitrogen containing spirocyclic compounds like the spirolactams¹⁴ and spirooxindols^{14g} have been synthesized diversely owing to their restricted conformational flexibility and interesting biological properties. The oxygen containing spirocyclics like the spiro-ketals and spirolactones form an important class of compounds possessing biological significance. The spiroketal ring system is a structural subunit that is often found in many naturally occurring substances of bacterial, insect, marine or plant origin¹⁵ and these assemblies have triggered immense interest in the synthesis owing to their pharmacological benefits.^{12a-d} Also, spirolactone based compounds are reported to show antibacterial,¹⁶ birth control,¹⁷ dehydrogenase type 2 inhibition¹⁸ formulations etc. Due to these benefits, numerous methods are reported for synthesis of spirolactones. Bach et al. reported spirocyclic formation via photochemical cycloaddition reactions in the presence of chiral Lewis acid,^{19a} and also by acid mediated cyclization of the corresponding synthons.^{19b} Very recently, free radical mechanism was used by implementing ACCN as the radical initiator to construct γ -lactones using Beckwith-Dowd ring expansion.^{19d} We became very fascinated by the abundance of the spirolactone structure in many of the naturally occurring compounds of which a few are mentioned in Figure 1. Acorenone B **1**, β -vetivone **2**, shizuca-acordienol **3**, are some of the naturally occurring spirocyclic compounds depicting some unique structural properties.²⁰ Upon taking a closer look at the compounds **4**, **5** and **6**, they possess the bicyclic spirolactone as a core structure (which is in close resemblance to that present in our designed molecules) and

are known to mimic neurotrophin activity useful in treating Alzheimer's disease & Parkinson's disease and show antifungal & antibacterial properties (Figure 1).^{19b} Quite interestingly, our further investigations for compounds with similar structural resemblances, disclosed the natural product^{13,19c} yuccaone A **7**, a spirobenzopyran-4-cyclopentan-3-one derivative, and we were highly motivated to design the entire tetracyclic core of the compound as it is one of the very rare naturally occurring phenolic spiro derivative having C-17 core (Figure 1). Herein, we report an efficient and practical method for the synthesis of novel spiro(tri or tetra)cyclic lactones (Table 2). To our delight, the method was applied to the synthesis of novel spiro(tetra or penta)cyclic systems as well (Table 3). Significantly, this method enabled us to accomplish the entire tetracyclic carbon core of yuccaone A.





Figure 1: Natural products containing the spiro-bicyclic lactone core and the major spirotricyclic core.

Results and Discussion

To initiate the study, the requisite cyclic esters **8a-8c** were prepared from the commercially available ketones by implementing the standard Wittig-Horner-Wadsworth-Emmons reaction. Initially, the acid promoted lactonization was explored on cyclohexanone ester **8b** with *para*-cresol **9d** as an external arene, under the conditions that were best suited for the synthesis of dihydrocoumarins reported by our group previously.³ However, the reaction did not proceed and resulted only in the starting material isolation (Table 1, entries 1 & 2). This directed us to explore the other conditions to optimize the reaction. We began our investigations using catalytic loading of Lewis acid FeCl₃ with increased phenol **9d** concentration (3 equiv), but it did not assist the reaction and led to the recovery of starting materials (Table 1, entries 3 & 4). Other Lewis acids like AlCl₃, Sc(OTf)₃, BF₃·OEt₂ and AuCl₃ also did not result in any product formation when used in the catalytic quantities (Table 1, entries 5 to 8). Even the reaction with 1 equiv of the Lewis acid FeCl₃ proved inefficient to initiate the reaction (Table 1, entry 9). Further, we explored the reaction with the increased quantity of the acid. When the reaction was performed with 3 equiv of acid at 50 °C, BF₃·OEt₂ showed only starting material (Table 1, entry 10), but quite interestingly, all starting material disappeared in case with FeCl₃, however, the product **10bd** was not observed (Table 1, entry 11). Hence, we performed the reaction at room temperature using FeCl₃ (3 equiv) in DCE, the required product

10bd was obtained in 40% yield (Table 1, entry 12). Delightfully, the yield of tricyclic lactone **10bd** was improved to 67% when benzene was used as the solvent (Table 1, entry 13). However, further increase in FeCl₃ to 5 equiv, did not increase the yield of product **10bd** (Table 1, entry 14). We also planned to experiment toluene in place of benzene, and it is worth mentioning that the reaction was successful in toluene as well, albeit, in poor yield (Table 1, entry 15).


Table 1: Optimizations for the formation of spirotricyclic lactone **10bd**.



entry ^a	acid (equiv)	phenol 9d (equiv)	Solvent (1.5 mL)	temp (°C)	yield of 10bd (%) ^b
1.	FeCl ₃ (3)	1.5	DCE	80	0 ^c
2.	FeCl ₃ (3)	1.5	DCE	rt	0 ^c
3.	FeCl ₃ (0.1)	3	DCE	80	0 ^c
4.	FeCl ₃ (0.5)	3	DCE	50	0 ^c
5.	AlCl ₃ (0.5)	3	DCE	50	0 ^c
6.	Sc(OTf) ₃ (0.1)	3	DCE	50	0 ^c
7.	BF ₃ ·OEt ₂ (0.5)	3	DCE	50	0 ^c
8.	AuCl ₃ (0.1)	3	DCE	50	0 ^c
9.	FeCl ₃ (1)	3	DCE	50	0 ^c
10.	BF ₃ ·OEt (3)	3	DCE	50	0 ^c
11.	FeCl ₃ (3)	3	DCE	50	0 ^d
12.	FeCl ₃ (3)	3	DCE	rt	40
13.	FeCl ₃ (3)	3	Benzene	rt	67
14.	FeCl ₃ (5)	3	Benzene	rt	60
15.	FeCl ₃ (3)	3	Toluene	rt	52

^a All reactions were performed on a 0.5 mmol scale of ester **8b** and 3 equiv of phenol **9d**. ^b Yields mentioned are of chromatographically purified compounds. ^c Only starting material was isolated. ^d Neither starting material nor the product were seen on TLC.


Among all the above screened conditions, those mentioned in entry 13 of Table 1 were best with regard to yield of **10bd**. Therefore, to check the scope and limitations of the method, these established conditions were applied to other homologous esters **8** with different phenols **9**. The method was highly amenable on different cyclic enoate esters **8** and furnished the corresponding spiro(tri or tetra)cyclic lactones **10** that are tabulated in Table 2. It is important to mention that the similar methodology when attempted with the ester derived from cyclooctanone failed to furnish the desired product. This might be probably due to the peculiar tub shaped conformational structure of the cyclooctanone moiety which might have hindered the product formation or may facilitate some unwanted reactions/rearrangements under these Lewis acidic conditions.

Table 2^{a,b}: Synthesis of spiro(tri/tetra)cyclic lactones **10** from the exocyclic α,β -unsaturated esters **8**.

^a All reactions were carried out on 0.5 mmol scale of **8** and 3 equiv of **9**, in solvent benzene (1.5 mL). ^b Isolated yields of chromatographically pure products **10**.

After successful accomplishment of novel spiro(tri or tetra)cyclic lactones **10aa-10ce**, to further improve the scope of the methodology, we applied it on various indanone based esters **11**. Various indanones were prepared from the corresponding aldehydes using Wittig Horner protocol to give the cinnamates, followed by reduction of the double bond and subsequent acid mediated cyclization to form the indanones, which were subjected to Horner-Wadsworth-Emmons reaction to yield esters **11**. Delightfully, it was found that the method suited to these systems as well and furnished the spiro(tetra or penta)cyclic compounds **12aa-12ec** containing simple to dense

functionalities on the aromatic rings. Interestingly, the products **12aa-12ec** were obtained in good to excellent yields, as summarized in Table 3. It is worth mentioning that the reaction of the phenyl substituted indanone **11e**, furnished the spiro-tetracyclic product **12ec** wherein it is expected that the phenol moiety attacks the double bond from the less hindered exo-face of the ester. Hence, the stereochemistry of **12ec** would be as shown in Table 3. Significantly, all these spirocyclic lactones **12aa-12ec** constitute the major spiro-tetracyclic carbon core of the natural product yuccaone A.

Table 3: Synthesis of novel spiro(tetra/penta)cyclic lactones **12** from the indanone based esters **11**.

^a All reactions were carried out on 0.5 mmol scale of **11** and 3 equiv of **9**, in solvent benzene (1.5 mL). ^b Isolated yields of chromatographically pure products **12**.





Figure 2: X-ray crystal structure of **12ec**. Ellipsoids are drawn at 50% probability factor.

In addition to the spectroscopic evidence for structural elucidation of the spirocyclic compounds, the complete structure and expected stereochemistry of **12ec** was further confirmed from the single crystal x-ray diffraction analysis (Figure 2).

A plausible mechanism for the synthesis of spirocyclic compounds is delineated in Scheme 2. The formation

of spiro-coumarins might be possible in two ways (path a & b). It may be either initial Friedel Crafts acylation followed by intramolecular Michel addition (path a) or Michael addition followed by intramolecular acylation (path b) (Scheme 2). The carbonyl of the enoate may be activated by Lewis acid FeCl_3 , triggering the attack of phenol at the carbonyl carbon to give **B** (Friedel-Crafts acylation). The adduct **B** would further rearrange to form **C** in which the carbonyl is again activated by the Lewis acid and internal Michael addition at the β -carbon of α,β -unsaturated ester might lead to the **D**, which stabilizes to form the required product **10aa**. On the other hand, in path b, initially FeCl_3 mediated Michael addition might lead to **E** which undergoes an intramolecular Friedel-Crafts acylation via **F** to form **10aa**.



Scheme 2: Plausible mechanism for the formation of spirocyclic compounds 10.

Conclusions

In summary, we have developed an efficient and practical method for the synthesis of novel spiro(tri or tetra)cyclic lactones via the construction of C-C and C-O bonds in one-pot using the Lewis acidic (FeCl_3). It enabled the formation of spirocyclic lactones starting from simple aliphatic exocyclic enoate esters as reacting partners to phenols. Significantly, the method was successfully applied to the synthesis of novel spiro(tetra or penta)cyclic lactones bearing simple to dense functionalities on the aromatic rings.

Acknowledgements

G.S. and N.P. are grateful to the Department of Science and Technology - Science and Engineering Research Board (DST-SERB) [NO.:SB/S1/OC-39/2014], New Delhi, for the financial support.

RSC Advances

ARTICLE

References

- ¹ K. C. Nicolaou, D. J. Edmonds and P. G. Bulger, *Angew. Chem., Int. Ed.*, 2006, **45**, 7134.
- ² L. F. Tietze, *Chem. Rev.*, 1996, **96**, 115.
- ³ P. Niharika, B. V. Ramulu and G. Satyanarayana, *Org. Biomol. Chem.*, 2014, **12**, 4347.
- ⁴ (a) B. V. Ramulu, A. G. K. Reddy and G. Satyanarayana, *Synlett* 2013, **24**, 868; (b) B. V. Ramulu, P. Niharika and G. Satyanarayana, *Synthesis* 2015, **26**, 1255; (c) A. Das, A. G. K. Reddy and G. Satyanarayana, *RSC Adv.*, 2014, **4**, 26662.
- ⁵ (a) B. V. Ramulu and G. Satyanarayana, *RSC Adv.*, 2015, **5**, 70972; (b) D. H. Dethe, R. Boda and G. M. Murhade, *Org. Chem. Front.*, 2015, **2**, 645; (c) D. H. Dethe and G. M. Murhade, *Chem. Commun.*, 2015, **51**, 10891; (d) D. H. Dethe and G. M. Murhade, *Chem. Commun.*, 2013, **49**, 8051.
- ⁶ A. G. K. Reddy and G. Satyanarayana, *Synthesis*, 2015, **47**, 1269.
- ⁷ (a) C. Jia, T. Kitamura and Y. Fujiwara, *Acc. Chem. Res.*, 2001, **34**, 633; (b) V. Ritzleng, C. Sirlin and M. Pfeffer, *Chem. Rev.*, 2002, **102**, 1731; (c) K. Li, L. N. Foresee and J. A. Tunge, *J. Org. Chem.*, 2005, **70**, 2881; (d) A. R. Jagdale and A. Sudalai, *Tetrahedron Lett.*, 2007, **48**, 4895; (e) S. Aoki, C. Amamoto, J. Oyamada and T. Kitamura, *Tetrahedron*, 2005, **61**, 9291.
- ⁸ (a) E. Fillion, A. M. Dumas, B. A. Kuropatwa, N. R. Malhotra and T. C. Sitler, *J. Org. Chem.*, 2006, **71**, 409; (b) C. E. Rodrigues-Santos and A. Echevarria, *Tetrahedron Lett.*, 2007, **48**, 4505; (c) S. Duan, R. Jana and J. A. Tunge, *J. Org. Chem.*, 2009, **74**, 4612.
- ⁹ (a) Y. Gu and K. Xue, *Tetrahedron Lett.*, 2010, **51**, 192; (b) A. Kumar, P. Kumar, V. D. Tripathi and S. Srivastava, *RSC Adv.*, 2012, **2**, 11641; (c) M. C. Laufer, H. Hausmann and W. F. Holderich, *J. Catal.*, 2003, **218**, 315; (d) E. Tang, W. Li, Z. Y. Gao and X. Gu, *Chin. Chem. Lett.*, 2012, **23**, 631; (e) D. P. Kamat, S. G. Tilve and V. P. Kamat, *Tetrahedron Lett.*, 2012, **53**, 4469; (f) C. R. Reddy, B. Srikanth, N. N. Rao and D.-S. Shin, *Tetrahedron*, 2008, **64**, 11666.
- ¹⁰ F. W. Bell, M. Hoberg and X.-X. Zhou, *J. Med. Chem.* 1995, **38**, 4959.
- ¹¹ (a) A. K. Franz, N. V. Hanhan and N. R. B.-Jones, *ACS Catal.* 2013, **3**, 540. (b) K. Ding, Z. Han and Z. Wang, *Chem. Asian J.*, 2009, **4**, 32.
- ¹² (a) Y. Zheng, C. M. Tice and S. B. Singh, *Bioorg. Med. Chem. Lett.*, 2014, **24**, 3673; (b) E. Prusov and M. E. Maier, *Tetrahedron*, 2007, **63**, 10486.
- ¹³ S. Kumar, P. D. Thornton and C. Santini, *ACS Comb. Sci.* 2013, **15**, 564.
- ¹⁴ (a) N. A. Braun, M. A. Ciufolini, K. Peters, E.-M. Peters, *Tetrahedron Lett.*, 1998, **39**, 4667; (b) D. Cheng, Y. Ishihara, B. Tan and C. F. Barbas, *ACS Catal.* 2014, **4**, 743; (c) G. Satyanarayana and M. E. Maier, *J. Org. Chem.*, 2008, **73**, 5410; (d) M. M. Fernandez, A. Diez, M. Rubiralta, E. Montenegro and N. Casamitjana, *J. Org. Chem.* 2002, **67**, 7587; (e) T. Dohi, A. Maruyama, Y. Minamitsuji, N. Takenaga and Y. Kita, *Chem. Commun.*, 2007, 1224; (f) G. Satyanarayana, C. M.-Mo'ssmerz and M. E. Maier, *Chem. Commun.*, 2009, 1571; (g) A. Patra, A. Bhunia, S. R. Yetra, R. G. Gonnade and A. T. Biju, *Org. Chem. Front.*, 2015, DOI: 10.1039/C5QO00242G.
- ¹⁵ (a) F. Perron and K. F. Albizati, *Chem. Rev.*, 1989, **89**, 1617; (b) J. E. Aho, P. M. Pihko and T. K. Rissa, *Chem. Rev.*, 2005, **105**, 4406; (c) B. Liu and J. K. De Brabander, *Org. Lett.*, 2006, **8**, 4907; (d) J. Liu and R. P. Hsung, *Org. Lett.*, 2005, **7**, 2273; (e) J. S. Potuzak, S. B. Moilanen and D. S. Tan, *J. Am. Chem. Soc.*, 2005, **127**, 13796.
- ¹⁶ (a) B. Bister, D. Bischoff, M. Strobele, J. Riedlinger, A. Reicke, F. Wolter, A. T. Bull, H. Zahner, H. P. Fiedler and R. D. Süßmuth,

Angew. Chem., Int. Ed., 2004, **43**, 2574; (b) J. Riedlinger, A. Reicke, H. Zahner, B. Krismer, A. T. Bull, L. A. Maldonado, A. C. Ward, M. Goodfellow, B. Bister, D. Bischoff, R. D. Süßmuth and H. P. Fiedler, *J. Antibiot.*, 2004, **57**, 271. (c) S. B. Singh, *Bioorg. Med. Chem. Lett.*, 2014, **24**, 3683.

¹⁷ W. Elger, S. Beier, K. Pollow, R. Garfield, S. Q. Shi and A. Hillisch, *Steroids*, 2003, **68**, 891.


¹⁸ K. Cimanga, N. Hermans, S. Apers, S. V. Miert, H. Van den Heuvel, M. Claeys, L. Pieters and A. Vlietinck, *J. Nat. Prod.*, 2003, **66**, 97.

¹⁹ (a) R. Brimiouille, H. Guo and T. Bach, *Chem. Eur. J.*, 2012, **18**, 7552; (b) H. Guo, E. Herdtweck and T. Bach, *Angew. Chem. Int. Ed.*, 2010, **49**, 7782; (c) T. J. Brocksom, F. Coelho, J.-P. Depre's, A. E. Greene, M. E. F. de Lima, O. Hamelin, B. Hartmann, A. M. Kanazawa and Y. Wang, *J. Am. Chem. Soc.*, 2002, **124**, 15313; (d) J. Hierold and D. W. Lupton, *Org. Lett.*, 2012, **14**, 3412.

²⁰ (a) R. Rios, *Chem. Soc. Rev.*, 2012, **41**, 1060; (b) N. Ingavat, C. Mahidol, S. Ruchirawat and P. Kittakoop, *J. Nat. Prod.*, 2011, **74**, 1650; (c) S. Piacente, G. Bifulco, C. Pizza, A. Stochmal and W. Oleszek, *Tetrahedron Lett.*, 2002, **43**, 9133.

One-Pot C-C/C-O Bonds Formation: Synthesis of Spirocyclic Lactones

Graphical Abstract



An efficient and a practical method for the synthesis of novel spiro(tri or tetra)cyclic lactones via the formation of C-C and C-O bonds in one-pot is presented. The method was successful under the Lewis acidic (FeCl_3) conditions and enabled the formation of spiro(tri or tetra)cyclic lactones from simple aliphatic exocyclic enoate esters as reacting partners with phenols. Remarkably, the usual self-aromatization of cyclohexanone based enoate esters, under such Lewis acidic conditions is overridden by intermolecular coupling. Significantly, the method was amenable on indanone derived esters as well and furnished novel spiro(tetra or penta)cyclic lactones bearing simple to dense functionalities on the aromatic rings. Notably, these novel spirocyclic systems constitute core structures of natural/unnatural compounds that show good biological properties.