

# RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

## Efficient lipid extraction and quantification of fatty acids from algal biomass using accelerated solvent extraction (ASE)

Received 00th January 20xx,  
Accepted 00th January 20xx

Yuting Tang,<sup>a,b,†</sup> Yue Zhang<sup>b,†</sup>, Julian N. Rosenberg,<sup>b</sup> Nadia Sharif,<sup>b</sup> Michael J. Betenbaugh,<sup>b,\*</sup> Fei

DOI: 10.1039/x0xx00000x

Wang<sup>a</sup>

www.rsc.org/

Accelerated solvent extraction (ASE), a commercially available pressurized fluid extraction technique and conventional manual extraction were compared to identify the most effective chloroform-methanol extraction method for algal lipids. Using optimal ASE operating conditions (methanol/chloroform = 2:1 by vol., 100°C, static time of 5 min, and four static cycles), the lipid contents of *Chlorella vulgaris*, *C. sorokiniana*, *C. zofingiensis* and *Nannochloropsis gaditana* were 27.5%, 25.8%, 15.2%, 29.8% of dry biomass, respectively. The total fatty acid methyl ester (FAME) content of dry biomass from ASE extraction was found to be 1.3–2.7 fold higher than that from conventional manual extract from these species, demonstrating that ASE exhibited significant improvement for lipid and FAME recovery. Furthermore, ASE showed the capacity to extract methyl all-cis-5,8,11,14,17-eicosapentaenoate (EPA) as 3.0% of dry biomass from *Nannochloropsis gaditana* suggesting that ASE has the potential to obtain polyunsaturated fatty acids (PUFAs) as well.

### 1 Introduction

The demand for biofuels is increasing due to limitations in the oil supply, rising greenhouse gas emissions, and the associated impact on global climate. Third generation biofuels, such as microalgae, represent one potential alternative energy source going forward, since they do not compete with food and crops.<sup>1</sup> Microalgal photosynthetic efficiency, growth performance and biomass productivity are also higher compared to other oleaginous cultures.<sup>2</sup> Moreover, microalgae species can adapt to a variety of environmental conditions and can be genetically manipulated.<sup>3,4</sup>

*Chlorella* and *Nannochloropsis* genera of microalgae are known for producing significant amounts of lipid. *C. vulgaris* has shown to produce biodiesel precursors, and nitrogen limitation contributes to increased lipid content.<sup>5</sup> *C. sorokiniana* is another promising freshwater alga for accumulating high amounts of lipids and has been found to be resistant to heat and high light intensity.<sup>6,7</sup> *C. zofingiensis* is a particular green alga that can grow well photoautotrophically

as well as heterotrophically and a candidate to generate astaxanthin and lutein.<sup>8</sup> *Nannochloropsis* spp. are promising algae for biodiesel production due to their successful cultivation at larger scales using natural sunlight. These species can achieve high levels of biomass production and contain high lipid contents, including polyunsaturated fatty acids.<sup>9–11</sup>

Lipids existing in algae can be classified as:<sup>12,13</sup> (i) neutral lipids, which serve as energy storage products, including triacylglycerols (TAG), diacylglycerols (DAG), monoacylglycerols (MAG), free fatty acids, hydrocarbons, wax, sterols etc.; (ii) phospholipids, which are mainly in cell membranes, including phosphatidylglycerol (PG), phosphatidylinositol (PI), Phosphatidylethanolamine (PE), phosphatidylcholine (PC); and (iii) glycolipids, which are mainly in cell membranes, including sulfoquinovosyldiacylglycerol (SQDG), monogalactosyldiacylglycerol (MGDG), and digalactosyldiacylglycerol (DGDG). The major fatty acids in microalgae are comprised of 12–22 carbon atoms with between 0–6 double bonds.<sup>12</sup> Esterifiable lipids can be converted into fatty acid methyl esters (FAME), which are the main components of biodiesel. The FAME profiles of microalgae have a significant effect on the quality of biodiesel. Microalgae with FAME profiles containing predominantly C:16 and C:18 are best suited for biodiesel production, which has been investigated previously.<sup>7,14–16</sup>

Lipid extraction is an important, yet time-consuming and cost intensive, step of algal species selection and biodiesel production. Organic solvent extraction is widely used to extract lipid from microalgae, but the use of organic solvents is associated with

<sup>a</sup> College of Chemical Engineering, Nanjing Forestry University, Jiangsu Key Lab of Biomass-based Green Fuels and Chemicals, Nanjing 210037, China

<sup>b</sup> Department of Chemical & Biomolecular Engineering, Johns Hopkins University 3400 N. Charles St, Baltimore, MD 21218, USA.

\*Corresponding author: Michael J. Betenbaugh, Department of Chemical & Biomolecular Engineering, Johns Hopkins University, 3400 N. Charles St, Baltimore, MD 21218, United States. E-mail address: mjbeten@gmail.com

These authors contributed equally to this work.<sup>‡</sup>

† Electronic Supplementary Information (ESI) available. See DOI: 10.1039/x0xx00000x

ith health, security and regulatory issues. Cell wall disruption technologies have been reported along with organic solvent extraction in literature reports in order to achieve higher lipid extraction efficiencies,<sup>17</sup> but the method can add costs and take time. Different disruption methods including sonication, osmotic shock, autoclaving, microwave, bead-beating treatments have been tested in *Chlorella* sp., *Nostoc* sp. and *Tolypothrix* sp. Sonication performed best among those treatments in *Chlorella* sp. and *Nostoc* sp., with efficiencies of 21% and 19% of dry biomass, respectively. Microwave treatment showed the highest efficiency (17%) in *Tolypothrix* sp.<sup>18</sup> These findings indicate that the most efficient treatment of breaking microalgal cell wall depends on the particular microalgal species.

Accelerated solvent extraction (ASE) is a promising automated extraction method that offers various extraction parameters.<sup>19</sup> Due to the use of elevated temperatures and pressures, the ASE method reduces consumption of solvents compared to other methods. Previous research suggested that as much as 90% of organic solvent usage could be reduced by using ASE.<sup>20</sup> Higher amounts of lipids can be recovered using an optimal ASE method from cereal, egg yolk and chicken breast muscle samples compared to a modified Folch procedure.<sup>21</sup> In addition to lipid extraction, ASE can be a promising technology to effectively extract valuable compounds under optimized conditions,<sup>12</sup> such as polycyclic aromatic hydrocarbons, polychlorinated biphenyls and total petroleum hydrocarbons from solid samples<sup>22</sup> and pigments from *Haematococcus pluvialis* and *Dunaliella salina*.<sup>23</sup> Furthermore, ASE can be used to extract antioxidants from microalga *Spirulina platensis* and the solvent system used in ASE was found to have a significant effect on recovery.<sup>24</sup>

In this work, we report the utilization of ASE to extract lipids from *C. vulgaris*, *C. sorokiniana*, *C. zofingiensis*, and *Nannochloropsis gaditana* and compare the lipid recoveries from ASE to conventional manual extraction methods across these four commercially-relevant microalgal species. The optimal ASE conditions, including solvents, reaction temperature, static time and static cycles were investigated. The fatty acid methyl ester (FAME) and polyunsaturated fatty acid (PUFA) recovery was also compared between ASE and conventional manual extraction in order to evaluate the full potential of these methods.

## 2. Materials and Methods

### 2.1 Materials

Stock samples of *C. vulgaris* UTEX 395, *C. sorokiniana* UTEX 1230, *C. zofingiensis* UTEX 32 were obtained from the Culture Collection of Algae at University of Texas at Austin (<http://web.biosci.utexas.edu/utex/>). *Nannochloropsis* strain number Chl-2 was purchased from IOCAS (Institute of Oceanology, Chinese Academy of Sciences), Qingdao, China and identified as *Nannochloropsis gaditana* by the colony PCR and phylogenetic analysis methods of Rosenberg *et al.*<sup>7</sup> *Chlorella* powders were purchased from Hoosier Hill Farm, USA. HPLC-grade organic solvents including chloroform, methanol, ethanol, acetone, butanol, heptane

and hexane were obtained from Sigma, USA. FAME standards including methyl heptadecanoate, methyl palmitate, methyl palmitoleate, methyl oleate, methyl linoleate, methyl linolenate, cis-5,8,11,14,17-eicosapentaenoic acid methyl ester, heptadecanoic acid were purchased from Sigma, USA. Analytical-grade hydrochloric acid was supplied by Sigma, USA.

### 2.2 Microalgal cultivation conditions

*C. vulgaris*, *C. sorokiniana*, and *C. zofingiensis* were cultured in Bold's basal medium (BBM)<sup>25</sup> containing the following components: 0.25 g L<sup>-1</sup> NaNO<sub>3</sub>, 25 mg L<sup>-1</sup> CaCl<sub>2</sub>·2H<sub>2</sub>O, 75 mg L<sup>-1</sup> MgSO<sub>4</sub>·7H<sub>2</sub>O, 75 mg L<sup>-1</sup> K<sub>2</sub>HPO<sub>4</sub>, 175 mg L<sup>-1</sup> KH<sub>2</sub>PO<sub>4</sub>, 25 mg L<sup>-1</sup> NaCl, and 1 mL of each of the trace element stock solutions in 1 L dH<sub>2</sub>O: 50 g EGTA and 31 g KOH in 1 L dH<sub>2</sub>O; 4.98 g FeSO<sub>4</sub>·7H<sub>2</sub>O, 1.0 mL H<sub>2</sub>SO<sub>4</sub> in 1 L dH<sub>2</sub>O; 11.42 g L<sup>-1</sup> H<sub>3</sub>BO<sub>3</sub>, 1.44 g MnCl<sub>2</sub>·4H<sub>2</sub>O, 8.82 g ZnSO<sub>4</sub>·7H<sub>2</sub>O, 1.57 g CuSO<sub>4</sub>·5H<sub>2</sub>O, 0.71 g MoO<sub>3</sub>, and 0.49 g Co(NO<sub>3</sub>)<sub>2</sub>·6H<sub>2</sub>O in 1 L dH<sub>2</sub>O. *Nannochloropsis gaditana* was cultured in BG-11 medium containing the following components: 1.5 g L<sup>-1</sup> NaNO<sub>3</sub>, 40 mg L<sup>-1</sup> KH<sub>2</sub>PO<sub>4</sub>·3H<sub>2</sub>O, 75 mg L<sup>-1</sup> MgSO<sub>4</sub>·7H<sub>2</sub>O, 36 mg L<sup>-1</sup> CaCl<sub>2</sub>·2H<sub>2</sub>O, 6.0 mg L<sup>-1</sup> citric acid, 6.0 mg L<sup>-1</sup> ferric ammonium citrate, 1.0 mg L<sup>-1</sup> EDTA, 20 mg L<sup>-1</sup> Na<sub>2</sub>CO<sub>3</sub>, and 1.0 mL L<sup>-1</sup> A<sup>5+</sup> Co\* solution. The A<sup>5+</sup> Co\* solution contained 2.86 g L<sup>-1</sup> H<sub>3</sub>BO<sub>3</sub>, 1.81 g L<sup>-1</sup> MnCl<sub>2</sub>·H<sub>2</sub>O, 222 mg L<sup>-1</sup> ZnSO<sub>4</sub>·7H<sub>2</sub>O, 79 mg L<sup>-1</sup> CuSO<sub>4</sub>·5H<sub>2</sub>O, 390 mg L<sup>-1</sup> Na<sub>2</sub>MoO<sub>4</sub>·2H<sub>2</sub>O, and 49 mg L<sup>-1</sup> Co(NO<sub>3</sub>)<sub>2</sub>·6H<sub>2</sub>O.<sup>26</sup> Growth experiments were carried out in 3-L batch cultures with ambient air bubbled at 1 L min<sup>-1</sup>. Cultures were illuminated with white fluorescent light at 100 μE m<sup>-2</sup> s<sup>-1</sup> with 12/12 hr light/dark photoperiod at 25°C throughout the experiment. Liquid cultures were harvested using a high-speed centrifuge (Beckman J2-21, Baltimore, USA) at 4,000 × g for 10 min and lyophilized for 24 hr at -40°C under vacuum.

### 2.3 Lipid extraction using ASE method

Lipid extraction was carried out using the ASE instrument (Dionex ASE 150). In brief, dry biomass was filled in 22 mL stainless steel cells and cellulose fiber filters (Dionex, USA) were used to prevent blockage of the frit in the bottom cap. The ASE program was set at different experimental conditions: extraction solvent was methanol and chloroform (2:1 and 1:2, v/v), methanol, ethanol, acetone, butanol, butanol and heptane (1:4, v/v) and hexane separately, extraction temperature was from 60 to 120°C, static time was from 5 to 15 min, and a static cycle number were from 2 to 5. The extraction procedure included the following steps: (1) the cell was rinsed using the extraction solvent (2) extraction cell with biomass samples were loaded into the machine; (3) cells were filled with solvents up to a pressure of 1500 psi; (4) the cell was preheated for 5 min (5) the extraction was performed for 35 min (flush volume: 100% of cell volume, purge time: 60 s); (6) the cell was rinsed again using the extraction solvent. The extracts were then collected in 60 mL collection vials for analysis.

The obtained extracts were air-dried and lipid content was determined gravimetrically. Purchased *Chlorella* powder was used to identify the optimal ASE conditions. Fresh *C. vulgaris*, *C. sorokiniana*, *C. zofingiensis* and *Nannochloropsis gaditana* biomass samples were cultivated for lipid extraction experiments under the

optimized ASE conditions. All experiments were performed in duplicate.

#### 2.4 Lipid extraction using conventional manual extraction

For the conventional extraction method, 102 mg of lyophilized biomass (*C. vulgaris*, *C. sorokiniana*, *C. zofingiensis* and *Nannochloropsis gaditana*) was blended with 10 mL solvent mixture of methanol and chloroform (2:1, v/v) in a flask.<sup>27</sup> The mixture was ground using mortar and pestle for 10 min, then the mixture was transferred into a conical test tube and stirred for 30 min. After that, the mixture of the algal cells and the solvent was separated by centrifugation at 8,000 × g for 10 min, and the extraction procedure was repeated. The supernatant containing lipid was collected in an amber vial and dried with airflow.

#### 2.5 Transesterification of lipids and quantification by gas chromatography

Transesterification of pre-weighted and air-dried lipids was executed in the presence of 500 µg heptadecanoic acid (C17-fatty acid) as an internal standard. The procedure was conducted 3 mL H<sub>2</sub>SO<sub>4</sub>/MeOH (5%, v/v) at 70°C for 1.5 h. The resulting FAME fraction was extracted by hexane at room temperature for 10 min and the process was performed duplicated.

Fatty acid methyl esters were ultimately analyzed by GC (Shimadzu 2010) with discharge ionization detection equipped with a capillary column (Stabilwax-DA, 30 m × 0.25 mm ID, film thickness 0.25 µm). The injector temperature of GC was set at 250°C and the detector was set at 260°C. The temperature program was started at 50°C and then increased to 170°C at a rate of 20°C min<sup>-1</sup>, plateau for 1 min. The rate was decreased to 4°C min<sup>-1</sup> from 170 to 220°C and kept constant remained for 14 min. Helium was used as the constant carrier gas.

The major FAME components (methyl palmitoleate (C16:1), methyl palmitate (C16:0), methyl cis-9,12-hexadecadienoate(16:2), methyl cis-6,9,12-hexadecatrienoate (C16:3), methyl oleate (18:1), methyl linoleate(C18:2), methyl linolenate (18:3) and methyl all-cis-5,8,11,14,17-eicosapentaenoate (C20:5n3, EPA)) were quantified by comparing the GC peak areas of the FAME corresponding standards. The 5-point calibration curves (0.5-2.5 mg mL<sup>-1</sup>) of each FAME standard were established.

### 3. Results and discussion

#### 3.1 Optimization of ASE parameters

##### 3.1.1 Effect of solvents on lipid recovery using ASE method.

Total lipids from commercially available *Chlorella* powder were extracted using the ASE method under various operating parameters including solvent systems, temperature, static time and static cycles in order to identify the optimized ASE operating conditions. Initially, different solvents in the ASE instrument were compared. Methanol and chloroform (2:1 and 1:2, v/v), methanol, ethanol, acetone, butanol, butanol and heptane (1:4, v/v) and hexane were used as shown in Figure 1A. It was found that methanol and chloroform (2:1, v/v) extracted up to 25.1 ± 0.2% lipid from the dry biomass, which was greater than the 19.7 ± 1.4% obtained using methanol and chloroform (1:2 v/v). These results

show that higher chloroform content, and accordingly lower polarity solvent system, did not improve extraction efficiency. The lipid content was found to be 24.8 ± 0.3% using methanol alone, which was very close to that using methanol and chloroform (1:2,v/v). This finding indicates that an increase in the polarity of the solvent system may not be effective in extracting significantly more lipids. This result is consistent with a previous study that concluded that extraction efficiency may be inversely proportional to the polarity of the solvent.<sup>28</sup> In addition, the carrying capacity of the solvent might also decline with a more polar solvent due to lower miscibility.<sup>12</sup> Based on these trends, further increases in the methanol levels in the methanol/chloroform system were not evaluated in the present study. Compared to the chloroform/methanol solvent system, none of the other solvents were as effective in extracting lipids. While ethanol extracted 18.2 ± 2.0%, all other solvents (acetone, butanol, butanol and heptane (1:4, v/v), and hexane) were relatively ineffective at lipid extraction using ASE. The non-polar solvent hexane presented the lowest lipid content of 1.1 ± 0.1%, likely due to its low polarity. Indeed, lipid recovery depends strongly on the polarity of the solvent and our results were consistent with this.<sup>29</sup> Polar solvents likely penetrate the microalgal cell wall more easily than non-polar solvents and have better contact between lipid and solvent for increased overall lipid extraction. Given these findings, polar and selective solvents are preferred and, in particular, methanol and chloroform (2:1, v/v) was chosen as the extraction solvent for subsequent evaluation of the ASE system.

Mulbry *et al.* also compared the lipid extraction efficiency using three solvents (chloroform/methanol, isopropanol/hexane, and hexane) in the ASE system with the green algae *Rhizoclonium hieroglyphicum* cultivated in an outdoor and indoor system with dairy manure effluent and fresh water aquarium. This study demonstrated that the lipid content was 6.1-11.0% of dry biomass using chloroform/methanol, 5.3-6.5% of dry biomass using isopropanol/hexane, while only 0.9-2.8% of dry biomass using hexane,<sup>30</sup> which is consistent with our findings.

To design a rational extraction system, the polarity index and the solubility of the solvent and the lipid should be matched. Several factors such as volatility, solubility, selectivity, toxic, cost, safety etc. should be considered. There are some other solvent systems to extract lipid. For example, the isopropanol:hexane (2:3, v/v) system has been studied.<sup>21, 31, 32</sup> Ryckebosch *et al.* found that total lipid yield obtained from the chloroform/methanol system was higher than that from the isopropanol and hexane system.<sup>31</sup> Though the isopropanol and hexane system has lower toxicity compared to the standard chloroform/methanol system, it has a lower selectivity for membrane lipids such as polar lipids and glycolipids, and the chloroform/methanol system provides the most quantitative and reproducible recovery of all lipid classes.<sup>32</sup> Shafer also reported that the extracted muscle lipids by the chloroform/methanol system has a higher fatty acid content compared to the isopropanol and hexane system.<sup>21</sup> Therefore, further investigation of the isopropanol and hexane system was not conducted in this work. Jones *et al.* indicated that 2-ethoxyethanol (2-EE) provides for excellent lipid recovery compared to the chloroform/methanol system,<sup>33</sup> but 2-ethoxyethanol (2-EE) belongs to ionic liquids, which have high viscosity and are sensitive to impurities. These require strict



reaction conditions and, thus, were not considered as a commonly used extraction solvent. However, ionic liquids (ILs) have low toxicity, flexible solubility and polarity and can serve as green solvents to extract lipids.<sup>34, 35</sup> Further study on a more environmentally friendly solvent system that could be applied in the ASE system will be worthwhile.

In terms of solvent ratios used with other biomass matrices, Dufreche *et al.* established that the oil yield of dried municipal sewage sludge by conventional manual extraction using a hexane, methanol and acetone (3:1:1, v/v/v) solvent system was 27.4% of the dry biomass and the biodiesel generated from the extraction oil was 4.4% of the dry biomass. Dried sewage sludge could serve as another promising feedstock to generate biodiesel.<sup>36</sup> Furthermore, Revellame *et al.* investigated the ASE method using hexane, methanol and acetone (3:1:1, v/v/v) solvent system to extract lipid from activated municipal sludge. The lipid content was around 12% of the dry biomass and the FAME yield generated from ASE was around 3.5% of the dry biomass.<sup>37</sup> Compared to our study, it shows that the lipid contents of microalgae are generally higher than that of sludge using the ASE method, and more FAME was recovered from microalgae. The optimal ASE parameters and the superior characteristics of microalgal feedstocks could both contribute to the higher recovery of lipid and FAME. Revellame *et al.* also compared the ASE method to the Bligh & Dyer method and found there was no significant difference between the two methods.<sup>37</sup> Lipid recovery could have been higher if the optimal ASE parameters were applied, and the biodiesel yield from sewage sludge might be improved as well. Collectively, these studies suggest that the ASE method has the potential to be widely used to extract lipids from different biomass samples.

**3.1.2 Effect of temperature on lipid recovery using ASE method.** The reaction temperature chosen for ASE also exhibited some effects on the efficacy of lipid extraction. Lipid contents obtained at 60°C, 80°C, 100°C and 120°C are summarized in Figure 1B. The lipid content was  $23.2 \pm 0.4\%$  at 60°C. When the reaction temperature was increased to 80°C, a higher lipid content of  $23.9 \pm 0.7\%$  could be achieved and content increased progressively with temperature up to  $25.1 \pm 0.2\%$  at 100°C. The higher temperature may contribute to the disruption of the cell wall to enable the solvent contact the lipids more thoroughly. When the temperature was further increased to 120°C, the lipid content did not increase significantly. Temperatures higher than 120°C were not studied because the highest lipid content was very close at 100°C and 120°C to suggest that higher temperature may not contribute to higher lipid recovery. Hence, a reaction temperature of 100°C was chosen as the optimum value in this study.

**3.1.3 Effect of static time on lipid recovery using ASE method.** The lipid contents achieved by adopting different static time periods are shown in Figure 1C. Static time represents the number of minutes that the sample and solvent are maintained at the set temperature. The lipid content was  $25.1 \pm 0.2\%$  at a 5 min static time and the content decreased slightly to  $23.6 \pm 0.8\%$  at 10 min and increased slightly to  $24.6 \pm 1.0\%$  at 15 min. Thus, it was found that lipid recoveries are nearly the same upon changing the static time, which means that static time has little effect on extraction

efficiency. This result is consistent with a previous study.<sup>24</sup> Thus, 5 min is sufficient for extracting lipids under the high temperature and pressure of ASE. Moreover, increased extraction times under high temperature and pressure could lead to partial degradation of some components. Since the longer contacting time was not found to not benefit lipid recovery, a shorter static time (5 min) was chosen as the optimal condition for this system.

### 3.1.4 Effect of static cycle on lipid recovery using ASE method.

The static cycle is the number of times the static heating and rinsing steps are implemented. In the present study, the positive correlation found between static cycle on the lipid recovery is shown in Figure 1D. As the number of static cycles increased, the lipid recovery was also enhanced. When static time was 2, the resulting lipid content was low at only  $22.9 \pm 0.02\%$ . As static cycle increased to 4, the lipid content increased to  $25.1 \pm 0.1\%$ . However, upon further increase of static cycles to 5, no obvious improvement in lipid recovery was observed. Considering the total time and the efficiency of the extraction process, a static cycle number of 4 was chosen as the optimum value for this system.

### 3.2 Comparison of optimized ASE method and conventional manual method for lipid extraction across different four algae species

After the optimal ASE operating conditions were determined using commercially purchased *Chlorella* powder, a comparison of the optimized ASE method and conventional manual method for lipid extraction were examined across four algae species: *C. vulgaris*, *C. sorokiniana*, *C. zofingiensis* and *Nannochloropsis gaditana*. These strains were cultivated photoautotrophically in 3-L batch cultures, then harvested and lyophilized for lipid extraction.

The lipid recovery using optimal ASE method and conventional manual extraction method are shown in Figure 2. The ASE extraction method showed higher efficiencies than manual extraction across all the four algae strains. It was found to extract 6.9% more lipid from dry biomass of *C. vulgaris* than conventional manual extraction. Previous literature indicated that the lipid content from *C. vulgaris* can be roughly 10% by dry weight using autoclaving and microwave oven method, whereas the bead-beating, sonication and osmotic shock showed lower efficiencies.<sup>17</sup> These prior results are all lower than the value obtained in the present study. The ASE method also showed a higher efficiency for *C. sorokiniana* than conventional manual extraction at  $25.8 \pm 1.4\%$  of dry biomass. In the literature, total lipid content from *C. sorokiniana* analyzed by the gravimetric analysis with chloroform and methanol solvents (2:1, v/v) was 23.5% of dry biomass,<sup>38</sup> and it was slightly lower than using the ASE method in this study. Conventional manual extraction method for *C. zofingiensis* was  $8.8 \pm 2.1\%$ , 1.7-fold lower than the ASE method. For *Nannochloropsis gaditana*, the lipid content was found to be  $29.8 \pm 2.1\%$  of dry biomass using ASE compared to  $21.3 \pm 3.1\%$  using conventional manual extraction method, also higher than that reported in the literature using the Bligh and Dyer method at 25.3%.<sup>39</sup>

### 3.3 Comparison of individual FAME recovery using ASE and conventional manual extraction across different four algae species

In order to evaluate the potential of the four strains as candidates for biodiesel production, detailed FAME compositions were elucidated in Figure 3.

**3.3.1 Individual FAME content of *C. vulgaris*.** ASE extraction was performed followed by GC analysis of FAME and the FAME contents of *C. vulgaris* using the ASE method and conventional manual extraction are shown in Figure 3A. Sixteen- to eighteen-carbon fatty acids with no more than three degrees of unsaturation were observed in *C. vulgaris* grown photoautotrophically. The total FAME content obtained using the ASE extraction method followed by transesterification was  $11.4 \pm 0.2\%$  of dry biomass, which is 2.4-fold higher than that using conventional manual extraction followed by transesterification. The ASE extraction method also exhibited a higher recovery for each individual FAME compared to conventional manual extraction method. The predominant FAME compositions in *C. vulgaris* were found to be C16:0 and C18:1 at  $2.4 \pm 0.02\%$  and  $3.2 \pm 0.04\%$  of dry biomass, respectively, from ASE extracted materials and the contents were at least 2.4- to 2.7-fold higher in extracted materials by conventional manual extraction. Previous research has shown that alternative treatment methods including high pressure steaming (HPS), similar to ASE, can alter the levels of FAME obtained.<sup>40</sup>

**3.3.2 Individual FAME content of *C. sorokiniana*.** The FAME compositions of *C. sorokiniana* are shown in Figure 3B. The total FAME content was  $11.8 \pm 0.1\%$  of dry biomass from the ASE extraction followed by transesterification, which is 1.3-fold higher than that from conventional manual extraction followed by transesterification. Small amounts of C16:2 and C17:1 were found using ASE extraction, but were not recovered by conventional manual extraction. Higher percentages of individual FAMES were also obtained from the ASE extraction compared to conventional manual extraction, while the difference was not as pronounced in *C. vulgaris*. The principal FAME components in *C. sorokiniana* were found to be C16:0 and C18:2, accounting for  $2.5 \pm 0.01\%$  and  $3.9 \pm 0.03\%$  of dry biomass, respectively, from ASE extraction. The FAME profiles are quite similar to the results reported by Rosenberg et al.<sup>7</sup>

Compared to *C. vulgaris*, the total FAME content of *C. sorokiniana* was similar, with  $11.8 \pm 0.1\%$  and  $11.4 \pm 0.2\%$  of dry biomass, respectively. Interestingly, *C. sorokiniana* exhibited greater enrichment in C18:2 relative to C18:1 and more unsaturated FAME such as C16:1, C16:3, C18:2 than *C. vulgaris*.

**3.3.3 Individual FAME content of *C. zofingiensis*.** The FAME composition of *C. zofingiensis* is shown in Figure 3C. The ASE extraction clearly generated more total and individual FAME than conventional manual extraction for *C. zofingiensis*. The total FAME content was 2.7-fold higher with ASE extraction than with conventional manual extraction. *C. zofingiensis* produced C18:3, C18:2 and C16:0 as the principal FAME, which is a different distribution than the other *Chlorella* strains. In addition, a higher relative proportion of unsaturated FAME was produced in *C. zofingiensis*. The total FAME content of *C. zofingiensis* obtained using the ASE method was 2.9- and 3.0-fold lower compared to *C. vulgaris* and *sorokiniana*, respectively. Kobayashi et al.<sup>41</sup> also observed that the FAME content of *C. zofingiensis* under ambient

CO<sub>2</sub> was only around 3% by dry weight— much lower than *Chlorella vulgaris*—and the study deduced that *C. zofingiensis* requires higher concentrations of nutrients in the medium to support its growth and accumulate more FAME.

**3.3.4 Individual FAME content of *Nannochloropsis gaditana*.** *Nannochloropsis gaditana* is another strain widely used to generate lipids and a candidate for the production of valuable polyunsaturated fatty acids (PUFAs). A comparison of the FAME profiles from ASE and conventional manual extraction is shown in Figure 3D. ASE extraction resulted in more total and individual FAME than observed with the three other *Chlorella* species. The total FAME recovered from ASE extraction was  $11.5 \pm 0.1\%$  of dry biomass, while only  $6.7 \pm 0.3\%$  of dry biomass was obtained by manual extraction. The predominant FAME in *Nannochloropsis gaditana* were found to be C16:0, C16:1 and C20:5 (EPA).

It should be noted that  $3.0 \pm 0.01\%$  of the dry *Nannochloropsis gaditana* biomass was found to be EPA using ASE extraction— higher than that recovered by manual extraction. Previously, Pieber et al. achieved an EPA content of 3.7% of dry biomass from *Nannochloropsis oculata* using ASE method with ethanol as the extraction solvent.<sup>29</sup> The study maintained a moderately lower temperature of 60°C for ASE cycles recovering EPA and claimed that further optimization of ASE conditions could increase lipid recovery. Conversely, research by Islam et al. employed more optimal elevated temperatures between 90-120°C with ASE for extraction of lipids that are amenable to biodiesel production, but not polyunsaturated fatty acids.<sup>42</sup> In the current study, our optimized process was able to utilize high temperatures for efficient extraction of lipids while simultaneously yielding high levels of PUFAs such as EPA. The efficiency of supercritical CO<sub>2</sub> extraction at optimized conditions from *Nannochloropsis granulate* was only 1.3% of dry biomass reported by Bjornsson et al.<sup>43</sup> This demonstrates that ASE method has a potential for commercial extraction of microalgal PUFAs and is potentially superior to supercritical CO<sub>2</sub> extraction method.

Of the four microalgae stains, *C. vulgaris* and *C. sorokiniana* appear to be good candidates for biodiesel production and *Nannochloropsis gaditana* has the potential to be used for EPA production.

## Conclusions

The ASE method is an effective technique to extract lipids from algal biomass. Optimal ASE operating parameters for *Chlorella* powder were found to be: an extraction solvent system of 2:1 methanol:chloroform by volume, an extraction temperature of 100°C, static time of 5 min, a static cycle number of 4. Compared with conventional manual extraction method, the ASE method can recover more lipid and FAME from *C. vulgaris*, *C. sorokiniana*, *C. zofingiensis* and *Nannochloropsis gaditana*. Furthermore, ASE showed the capacity to extract a significant component of the *Nannochloropsis gaditana* lipid fraction as EPA.

While ASE has shown to be efficient for the extraction of lipids and PUFAs such as EPA on the bench-top, commercial implementation of this approach at large scale will have to consider the energy input and other cost considerations of using the ASE. A

major advantage of the ASE is that only minutes are needed for ASE method due to automated steps, whereas other extraction method such as sonication and soxhlet can take hours to extract the target lipid. Also, the ASE does not require completely dry samples<sup>44</sup> and this could reduce the number of dewatering steps applied prior to this method. Nonetheless, the use of ASE for algal lipid extraction in the present study support its utility at the laboratory scale for tasks ranging from strain selection based on total lipid content to detailed assessments of algal species based on fatty acid composition. Future work with ASE should examine additional parameters and applications in order to identify the optimal conditions for using the ASE for research, development, and commercial applications.

### Acknowledgements

The authors gratefully acknowledge the financial support from grant number NSF-EFRI-1332344 from the National Science Foundation (MJB), DOE DE SC0012658 grant (MJB), NSF CBET-1236691 and a fellowship to JNR from the Johns Hopkins Environment, Energy, Sustainability & Health Institute (E<sup>2</sup>SHI, <http://e2shijhu.edu/>). Partial support was also provided by the Natural Science Foundation of Jiangsu universities (11KJA480001) and the national Natural Science Foundation of China (31170537). And this work was also supported by the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD).

### References

1. L. Gouveia and A. C. Oliveira, *Journal of industrial microbiology & biotechnology*, 2009, **36**, 269-274.
2. Y. Chisti, *Biotechnol Adv*, 2007, **25**, 294-306.
3. T. M. Mata, A. A. Martins and N. S. Caetano, *Renewable and Sustainable Energy Reviews*, 2010, **14**, 217-232.
4. J. N. Rosenberg, V. H. Oh, G. Yu, B. J. Guzman, G. A. Oyler and M. J. Betenbaugh, in *Handbook of Marine Microalgae*, ed. S.-K. Kim, Academic Press, Boston, 2015, pp. 331-352.
5. Y. X. Li, F. J. Zhao and D. D. Yu, *Brazilian Archives of Biology and Technology*, 2015, **58**, 462-467.
6. M. Morita, Y. Watanabe and H. Saiki, *Applied biochemistry and biotechnology*, 2000, **87**, 203-218.
7. J. N. Rosenberg, N. Kobayashi, A. Barnes, E. A. Noel, M. J. Betenbaugh and G. A. Oyler, *PLoS ONE*, 2014, **9**, e92460.
8. P.-F. Ip and F. Chen, *Process Biochemistry*, 2005, **40**, 733-738.
9. Y. Ma, Z. Wang, C. Yu, Y. Yin and G. Zhou, *Bioresource Technology*, 2014, **167**, 503-509.
10. P. C. Hallenbeck, M. Grogger, M. Mraz and D. Veverka, *Bioresource Technology*, 2015, **184**, 161-168.
11. A. San Pedro, C. González-López, F. Acien and E. Molina-Grima, *Bioresource Technology*, 2014, **169**, 667-676.
12. M. Cooney, G. Young and N. Nagle, *Separation & Purification Reviews*, 2009, **38**, 291-325.
13. V. Astrid, W. Christian, G. Reimund, S. Rosmarie and S. Jürgen, *Chemistry & Physics of Lipids*, 2008, **150**, 143-155.
14. H. Wu and X. Miao, *Bioresource Technology*, 2014, **170**, 421-427.
15. M. J. Ramos, C. M. Fernández, A. Casas, L. Rodríguez and Á. Pérez, *Bioresource Technology*, 2009, **100**, 261-268.
16. I. A. Nascimento, I. T. D. Cabanelas, J. N. d. Santos, M. A. Nascimento, L. Sousa and G. Sansone, *Algal Research*, 2015, **8**, 53-60.
17. J.-Y. Lee, C. Yoo, S.-Y. Jun, C.-Y. Ahn and H.-M. Oh, *Bioresource Technology*, 2010, **101**, S75-S77.
18. P. Prabakaran and A. D. Ravindran, *Letters in Applied Microbiology*, 2011, **53**, 150-154.
19. D. Luthria, D. Vinjamoori, K. Noel and J. Ezzell, in *Oil Extraction and Analysis: Critical Issues and Comparative Studies.*, ed. D. Luthria, AOCS, Champaign, IL, 2004, pp. 25-38.
20. K. Pilaniya, H. K. Chandrawanshi, U. Pilaniya, P. Manchandani, P. Jain and N. Singh, *Journal of Advanced Pharmaceutical Technology & Research*, 2010, **1**, 302-310.
21. K. Schäfer, *Analytica Chimica Acta*, 1998, **358**, 69-77.
22. B. E. Richter, B. A. Jones, J. L. Ezzell, N. L. Porter, N. Avdalovic and C. Pohl, *Analytical Chemistry*, 1996, **68**, 1033-1039.
23. J. R. Denery, K. Dragull, C. S. Tang and Q. X. Li, *Analytica Chimica Acta*, 2004, **501**, 175-181.
24. M. Herrero, P. J. Martín-Álvarez, F. J. Señoráns, A. Cifuentes and E. Ibáñez, *Food Chemistry*, 2005, **93**, 417-423.
25. H. W. Nichols and H. C. Bold, *Journal of Phycology*, 1965, **1**, 34-38.
26. D.-Y. Kwon, J.-H. Kwon and G.-J. Jo, *Desalination and Water Treatment*, 2013, **52**, 1007-1013.
27. E. G. Bligh and W. J. Dyer, *Can J Biochem Phys*, 1959, **37**, 911-917.
28. P. Hidalgo, G. Ciudad and R. Navia, *Bioresource Technology*, 2016, **201**, 360-364.
29. S. Pieber, S. Schober and M. Mittelbach, *Biomass and Bioenergy*, 2012, **47**, 474-482.
30. W. Mulbry, S. Kondrad, J. Buyer and D. Luthria, *J Am Oil Chem Soc*, 2009, **86**, 909-915.
31. E. Ryckebosch, K. Muylaert and I. Foubert, *J Am Oil Chem Soc*, 2012, **89**, 189-198.
32. J. B. Guckert, K. E. Cooksey and L. L. Jackson, *Journal of Microbiological Methods*, 1988, **8**, 139-149.
33. J. Jones, S. Manning, M. Montoya, K. Keller and M. Poenie, *J Am Oil Chem Soc*, 2012, **89**, 1371-1381.
34. G. Young, F. Nippgen, S. Titterbrandt and M. J. Cooney, *Separation and Purification Technology*, 2010, **72**, 118-121.
35. Y.-H. Kim, Y.-K. Choi, J. Park, S. Lee, Y.-H. Yang, H. J. Kim, T.-J. Park, Y. Hwan Kim and S. H. Lee, *Bioresource Technology*, 2012, **109**, 312-315.
36. S. Dufreche, R. Hernandez, T. French, D. Sparks, M. Zappi and E. Alley, *Journal of the American Oil Chemists Society*, 2007, **84**, 181-187.
37. E. D. Revellame, R. Hernandez, W. French, W. E. Holmes, T. J. Benson, P. J. Pham, A. Forks and R. Callahan li, *RSC Advances*, 2012, **2**, 2015-2031.
38. B. D. Wahlen, R. M. Willis and L. C. Seefeldt, *Bioresource Technology*, 2011, **102**, 2724-2730.
39. B. P. Nobre, F. Villalobos, B. E. Barragán, A. C. Oliveira, A. P. Batista, P. A. S. S. Marques, R. L. Mendes, H. Sovová, A. F. Palavra and L. Gouveia, *Bioresource Technology*, 2013, **135**, 128-136.
40. A.-M. Aguirre and A. Bassi, *Bioresource Technology*, 2014, **164**, 136-142.


41. N. Kobayashi, A. Barnes, T. Jensen, E. Noel, G. Andlay, J. N. Rosenberg, M. J. Betenbaugh, M. T. Guarnieri and G. A. Oyler, *Bioresource Technology*, 2015, **198**, 246-255.
42. M. A. Islam, R. J. Brown, I. O'Hara, M. Kent and K. Heimann, *Energy Conversion and Management*, 2014, **88**, 307-316.
43. W. J. Bjornsson, K. M. MacDougall, J. E. Melanson, S. J. B. O'Leary and P. J. McGinn, *J Appl Phycol*, 2012, **24**, 547-555.
44. B. E. Richter, *Journal of Chromatography A*, 2000, **874**, 217-224.

**Fig. 1.** Effects of reaction parameters on lipid recovery using the ASE method (A) Effect of extraction solvents. Reaction conditions: extraction temperature of 100 °C, static time of 5 min, static cycle number of 4. (B) Effect of temperature. Reaction conditions: methanol/chloroform (2:1, v/v), static time of 5 min, static cycle number of 4. (C) Effect of static time. Reaction conditions: methanol/chloroform (2:1, v/v), extraction temperature of 100 °C, static cycle number of 4. (D) Effect of static cycles. Reaction conditions: methanol/chloroform (2:1, v/v), extraction temperature of 100 °C, static time of 5 min. Error bars designate standard deviation from the average of duplicates.


**Fig. 2.** Comparison of ASE and conventional manual extraction on lipid recovery across *Chlorella vulgaris*, *Chlorella sorokiniana*, *Chlorella zofingiensis* and *Nannochloropsis gaditana*. Reaction conditions: methanol/chloroform (2:1, v/v), extraction temperature of 100 °C, static time of 5 min, static cycle number of 4. Error bars designate standard deviation from the average of duplicates.

**Fig. 3.** Comparison of ASE and conventional manual extraction on individual FAME recovery from (A) *Chlorella vulgaris* (B) *Chlorella sorokiniana* (C) *Chlorella zofingiensis* (D) *Nannochloropsis gaditana*. Error bars designate standard deviation from the average of duplicates.






220x585mm (300 x 300 DPI)




49x29mm (300 x 300 DPI)



206x511mm (300 x 300 DPI)

Accelerated Solvent Extraction optimized for extraction of algal lipids and recovery of polyunsaturated fatty acids.



37x28mm (300 x 300 DPI)