

# RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

# Synthesis, coordination behavior and structural features of chiral iron(II) PNP diferrocene complexes

Afroz Zirakzadeh,<sup>a,\*</sup> Karl Kirchner,<sup>a</sup> Alexander Roller,<sup>b</sup> Berthold Stöger,<sup>c</sup> Maria Deus Carvalho,<sup>d</sup> Liliana P. Ferreira.<sup>e,f</sup>

<sup>a</sup> Institute of Applied Synthetic Chemistry, Vienna University of Technology, Getreidemarkt 9, A-1060 Vienna, Austria

<sup>b</sup> X-ray Structure Analysis Centre, University of Vienna, Währinger Straße 42, A-1090 Wien, Austria

<sup>c</sup> Institute of Chemical Technologies and Analytics, Vienna University of Technology, Getreidemarkt 9, A-1060 Vienna, Austria

<sup>d</sup> Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

<sup>e</sup> Biosystems and Integrative Sciences Institute, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

<sup>f</sup> Department of Physics, University of Coimbra, 3004-516 Coimbra, Portugal

## Abstract

Five new chiral PNP ferrocene ligands with either an imine or amine nitrogen coordination site were synthesized. Only the imine type ligands formed Fe(II) complexes with the general formula  $[\text{Fe}(\text{PNP})\text{X}_2]$  ( $\text{X} = \text{Cl}, \text{Br}$ ). In the solid state these complexes adopt a tetrahedral geometry with the PNP ligand coordinated in a  $\kappa^2P,N$ -fashion with the one pendant-arm and the other not coordinated, as determined by X-ray crystallography and Mössbauer spectroscopy. The complexes are paramagnetic with a quintet ground state. In solution there is an equilibrium between  $[\text{Fe}(\kappa^3P,N,P\text{-PNP})\text{X}_2]$  and  $[\text{Fe}(\kappa^2P,N\text{-PNP})\text{X}_2]$  complexes. Boronation of the non-coordinated arm shifts the equilibrium towards the four-coordinate complex  $[\text{Fe}(\kappa^2P,N\text{-PNP}^{\text{BH}_3})\text{Br}_2]$ . DFT calculations are consistent with the experimental results and indicate that the experimentally observed  $\kappa^2$  isomer is thermodynamically the most. In a CO atmosphere,  $[\text{Fe}(\text{PNP})(\text{CO})_2\text{Br}]\text{Br}$  was formed rather than  $[\text{Fe}(\text{PNP})(\text{CO})\text{Br}_2]$ .

## Introduction

At present, platinum metal-based asymmetric catalysis is considered to be a standard methodology in organic synthesis and it is frequently used in both academia and industry. In particular, asymmetric hydrogenations on various types of substrates can nowadays be carried out on a very large scale.<sup>1-3</sup> However, these processes require rather expensive ruthenium-, rhodium- or iridium-based catalysts.<sup>4,5</sup> Not only for the obvious economic reasons, but also driven by the ideas of ‘Green and environmentally friendly chemistry’, it became clear that iron-based catalysts could provide a number of benefits over platinum metal catalysts if they could be made stable, active, selective and easy to handle. In this respect, besides the lower cost and greater abundance of iron, its very low toxicity makes it an attractive candidate for the preparation of well-defined iron-based catalysts.<sup>6-14</sup> To date, only one class of efficient Chiral PNP iron catalysts for the preparation of enantioenriched alcohols and imines has been reported and this was developed by the Morris group.<sup>15</sup> Since, a large number of chiral non-racemic substituted ferrocene derivatives have been synthesized and such planar chiral derivatives have been applied in a variety of different fields, including homogeneous enantioselective catalysis.<sup>16</sup> On the basis of the information outlined above, we embarked on a program to synthesize chiral iron complexes based on ferrocene backbones.

The study reported here concerns the synthesis of chiral PNP pincer ligands with a ferrocene scaffold (Chart 1) and the structural behavior and reactivity of their corresponding Fe(II) complexes.





Chart 1.


## Results and Discussion

**Synthesis of ligand (*S*<sub>Fc</sub>,*S*<sub>Fc</sub>)-1 and (*S*<sub>Fc</sub>,*S*<sub>Fc</sub>)-2.** As outlined in Scheme 1, enantiopure aldehyde (*S*<sub>Fc</sub>)-6, which was prepared by applying Kagan's methodology,<sup>17</sup> was reacted with sodium hydroxide and hydroxylamine hydrochloride to give (*S*<sub>Fc</sub>)-7. Subsequent treatment with

$\text{LiAlH}_4$  in THF gave  $(S_{\text{Fc}})$ -**8**.<sup>18</sup> In the final step, condensation of  $(S_{\text{Fc}})$ -**6** and  $(S_{\text{Fc}})$ -**8** in ethanol yielded the desired ligands  $(S_{\text{Fc}}, S_{\text{Fc}})$ -**1**. The molecular structure of  $(S_{\text{Fc}}, S_{\text{Fc}})$ -**1a** was determined by X-ray diffraction (Figure 1). Treatment of  $(S_{\text{Fc}}, S_{\text{Fc}})$ -**1** with  $\text{NaBH}_4$  in ethanol gave  $(S_{\text{Fc}}, S_{\text{Fc}})$ -**2** in quantitative yield (Scheme 1).




Scheme 1.



**Figure 1.** Structural view of  $(S_{Fc},S_{Fc})$ -**1a** showing 50 % thermal ellipsoids (H atoms and solvent molecules are omitted for clarity). Selected bond lengths (Å) and bond angles (°): C1A–N1 1.29(1), C1B–N1 1.40(1), C1A–N1–C1B 117.5(8).


**Synthesis of ligand  $(R,R,S_{Fc},S_{Fc})$ -3.**  $(R,S_{Fc})$ -**10**, which is easily accessible from  $(R)$ -Ugi's amine  $(S_{Fc})$ -**9**, was reacted with  $Ac_2O$  at 100 °C for 2 hours to afford  $(R,S_{Fc})$ -**11** followed by treatment with a large excess of ammonia in ethanol in an autoclave at 80 °C for 6 hours to yield the primary amine  $(R,S_{Fc})$ -**12**.<sup>19</sup> A mixture of  $(R,S_{Fc})$ -**9** and  $(R,S_{Fc})$ -**12** in acetonitrile and water was heated under reflux for 30 hours to give  $(R,R,S_{Fc},S_{Fc})$ -**3** in 61% isolated yield (Scheme 2).




**Scheme 2.**

**Synthesis of iron complexes.** Treatment of ( $S_{Fc},S_{Fc}$ )-**1** with 0.97 equivalents of  $FeBr_2(THF)_2$ <sup>20</sup> or anhydrous  $FeCl_2$  in THF gave the coordinatively unsaturated complexes ( $S_{Fc},S_{Fc}$ )-**13a**, ( $S_{Fc},S_{Fc}$ )-**13b**, and ( $S_{Fc},S_{Fc}$ )-**13c** in 93%, 80% and 91% yield, respectively (Scheme 3). The molecular structures of ( $S_{Fc},S_{Fc}$ )-**13a** and ( $S_{Fc},S_{Fc}$ )-**13c** were determined by X-ray crystallography and selected bond distances and angles are given in the captions (Figures 2 and 3). Both structures have a distorted tetrahedral geometry around the Fe(II) center, with the PNP ligand coordinated in a  $\kappa^2P,N$ -fashion rather than in the typical  $\kappa^3P,N,P$ -coordination mode. This bonding mode is relatively rare for pincer type ligands. We recently described a Taddol based Fe(II) PNP pincer complex with a 2,6-diaminopyridine backbone where the ligand was also coordinated in a  $\kappa^2P,N$ -fashion.<sup>21</sup> The Fe-P bond length of about 2.4 Å is comparable with those in other high-spin tetrahedral iron(II) complexes.<sup>22</sup>


Although better coordination was expected for the ligand ( $S_{Fc},S_{Fc}$ )-**1b** which contains more strongly electron donating  $P^iPr_2$  moieties, similar results to ( $S_{Fc},S_{Fc}$ )-**1a** were obtained and also this ligand is coordinated in  $\kappa^2P,N$ -fashion. On the other hand, the ligands ( $S_{Fc},S_{Fc}$ )-**2** and ( $R,R,S_{Fc},S_{Fc}$ )-**3**, under analogous reaction conditions, did not react with the iron salts and coordination did not take place. The reason for this behavior is not clear but it could be due to the larger bite angles in these ligands. As a consequence, the following discussion is focused on complexes obtained from ( $S_{Fc},S_{Fc}$ )-**1a**.



**Figures 2.** Structural view of ( $S_{Fc},S_{Fc}$ )-**13a** showing 50% thermal ellipsoids (H atoms and solvent molecules are omitted for clarity). Selected bond lengths (Å) and bond angles (°): Fe1–Br1 2.3818(5), Fe1–Br2 2.3916(5), Fe1–P1 2.3999(9), Fe1–N1 2.090(3), P1–Fe1–Br1 110.42(3), P1–Fe1–N1 95.16(8), Br1–Fe1–N1 115.47(8), Br2–Fe1–P1 112.84(3), Br2–Fe1–N1 102.08(8).



**Figures 3.** Structural view of  $(S_{Fc},S_{Fc})$ -**13c** showing 50% thermal ellipsoids (H atoms and solvent molecules are omitted for clarity). Selected bond lengths (Å) and bond angles (°): Fe1–Cl1 2.2438(8), Fe1–Cl2 2.2472(8), Fe1–P1 2.4031(8), Fe1–N1 2.095(2), Cl1–Fe1–Cl2 119.83(3), P1–Fe1–Cl1 110.96(3), P1–Fe1–Cl2 112.63(3), Cl1–Fe1–N1 113.32(7), Cl2–Fe1–N1 106.64(7).




**Scheme 3.**

**Mössbauer spectroscopy.** The magnetic properties of  $(S_{Fc},S_{Fc})$ -**13a** were investigated by means of Mössbauer spectroscopy. The Mössbauer spectrum of complex  $(S_{Fc},S_{Fc})$ -**13a** at 78 K is well defined and was fitted with three different iron sites (Figure 4). The blue and orange sites were present in equal proportions (32.9%) and their quadrupole splitting (QS) values obtained from the fitting are very similar. The isomer shift (IS) values are slightly different, i.e., 0.47(1) and 0.55(1) mm/s, and the Fe site with the lowest IS corresponds to the one with the highest s-electron density at the nucleus. Globally, the hyperfine parameters obtained for these two sites are reminiscent of the ones for Fe(II) in ferrocene (Table 1),<sup>23</sup> a finding that is consistent with the structure determined by X-ray crystallography. For the third site (green) the expected proportion (34.2%) obtained from the fit shows that it corresponds to the central iron

in a distorted tetrahedral coordination environment. The IS value is consistent with HS Fe(II) (Table 1).

**Table 1.** Estimated hyperfine parameters from the  $^{57}\text{Fe}$  Mössbauer spectra of  $(S_{\text{Fe}}, S_{\text{Fe}})$ -**13a** collected at 78 K and related tetrahedral Fe(II) complexes. IS: isomer shift; QS: quadrupole splitting.


complex	IS (mm/s)	QS (mm/s)	Fe(II)	Ref.
[Fe( $\kappa^2P,N$ -PNP)Br <sub>2</sub> ] ( $S_{\text{Fe}}, S_{\text{Fe}}$ )- <b>13a</b>	0.782(1)	3.208(2)	HS	this work
	0.467(3)	2.203(3)	LS (FeCp <sub>2</sub> arm)	
	0.552(3)	2.190(3)	LS (FeCp <sub>2</sub> arm)	
Ferrocene <sup>a</sup>	0.531(1)	2.491(1)	LS	23
[Fe( $\kappa^2P,N$ -PNP- <i>i</i> Pr,TAD)Cl <sub>2</sub> ]	0.78(1)	2.93(1)	HS	21
[Fe( $\kappa^2P,N$ -PNP- <i>i</i> Pr,TAD)Br <sub>2</sub> ]	0.76(1)	3.14(1)	HS	21
[Fe( $\kappa^2P,N$ -PNP- <i>t</i> Bu,TAD)Cl <sub>2</sub> ]	0.79(1)	3.06(1)	HS	21
[Fe( $\kappa^2P,N$ -PNP- <i>t</i> Bu,TAD)Br <sub>2</sub> ]	0.77(1)	3.22(1)	HS	21
[Fe(SN <sup>H</sup> -Ph)Br <sub>2</sub> ]	0.855(1)	3.282(2)	HS	32
[Fe(SN <sup>Et</sup> -Ph)Cl <sub>2</sub> ]	0.899(1)	3.008(1)	HS	32
[Fe(SN <sup>Et</sup> -Ph)Br <sub>2</sub> ]	0.877(1)	3.259(2)	HS	32
[Fe(PN <sup>H</sup> - <i>i</i> Pr)Cl <sub>2</sub> ]	0.680(2)	2.871(4)	HS	33
[Fe(SN <sup>H</sup> - <i>i</i> Pr)Cl <sub>2</sub> ]	0.796(2)	2.998(4)	HS	33



**Figure 4.**  $^{57}\text{Fe}$  Mössbauer spectrum of [Fe( $\kappa^2P,N$ -PNP)Br<sub>2</sub>] ( $S_{\text{Fe}}, S_{\text{Fe}}$ )-**13a** collected at 78 K.




DFT calculations for complex  $[\text{Fe}(\kappa^2P,N\text{-PNP})\text{Br}_2]$  ( $S_{\text{Fc}},S_{\text{Fc}}$ )-**13a** are in agreement with the experimental data and this indicates that the experimentally observed isomer (denoted as **A**) is thermodynamically more stable by 9.3 and 18.1 kcal/mol, respectively, than the isomers **B** and **C** (Figure 5), which were not observed. In **A** and **B** one phosphine moiety is not coordinated and **B** is a rotamer of **A** with a long  $\text{Fe}\cdots\text{P}$  distance of 3.85 Å. Complex **C** is a five-coordinate species with a distorted square pyramidal geometry.




**Figure 5.** Free energies (B3LYP) of optimized iron PNP complexes featuring  $\kappa^2P,N$  and  $\kappa^3P,N,P$ -bound PNP ligands. The free energy values (kcal/mol) are referred to the experimentally obtained tetrahedral  $[\text{Fe}(\kappa^2P,N\text{-PNP})\text{Br}_2]$  ( $S_{\text{Fc}},S_{\text{Fc}}$ )-**13a** (**A**).

**Solution studies.** The structures of complexes ( $S_{\text{Fc}},S_{\text{Fc}}$ )-**13a** and ( $S_{\text{Fc}},S_{\text{Fc}}$ )-**13c** in the solid state were unequivocally confirmed by X-ray crystallography, but in solution all of the complexes gave  $^1\text{H}$  and  $^{13}\text{C}\{^1\text{H}\}$  NMR spectra with large paramagnetic shifts. The signals were broad and featureless and, due to the complexity of the PNP ligands it was not possible to assign signals, thus making the spectra very uninformative. The solution magnetic moment of the dibromide ( $S_{\text{Fc}},S_{\text{Fc}}$ )-**13a** was measured to be  $4.98 \mu_{\text{B}}$  (Evans' method). This value is in good agreement with the effective magnetic moment of HS Fe(II) in the spin only approximation ( $4.9 \mu_{\text{B}}$ ). Accordingly, it is not clear whether these complexes are four- or five-coordinate and contain  $\kappa^2P,N$ - or  $\kappa^3P,N,P$ -bound PNP ligands, respectively, or whether both species are present and, if this is the case, whether these are in equilibrium with one another. As a consequence, further experiments were carried out in an effort to elucidate the structure of ( $S_{\text{Fc}},S_{\text{Fc}}$ )-**13a** in solution. At room temperature  $^{31}\text{P}\{^1\text{H}\}$  NMR signals could not be detected at all, but at  $-80^\circ\text{C}$  a comparatively sharp signal was observed at around 118 ppm and, this was assigned to the non-coordinated arm of the PNP ligand. This observation allows the existence of equilibria between  $\kappa^2P,N$ - and  $\kappa^3P,N,P$ -bound species to be postulated. Moreover,


the addition of  $\text{BH}_3 \cdot \text{THF}$  to a solution of  $(S_{\text{Fc}}, S_{\text{Fc}})$ -**13a** in  $\text{CD}_2\text{Cl}_2$  led to boronation of the non-coordinated arm and a shift in the equilibrium towards the four-coordinate complex  $[\text{Fe}(\kappa^2\text{P}, \text{N}-\text{PNP}^{\text{BH}_3})\text{Br}_2]$   $(S_{\text{Fc}}, S_{\text{Fc}})$ -**14** (Scheme 4). The  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra obtained in these studies are shown in Figure 6. The molecular structure of  $(S_{\text{Fc}}, S_{\text{Fc}})$ -**14** was determined by X-ray crystallography (Figure 7) and this confirmed the pendant phosphorus moiety had been boronated.



**Scheme 4.**




**Figure 6.**  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra of (a)  $(S_{\text{Fc}},S_{\text{Fc}})\text{-1a}$ , (b)  $\text{Fe}(\kappa^2P,N\text{-PNP})\text{Br}_2$   $(S_{\text{Fc}},S_{\text{Fc}})\text{-13a}$  at  $-80\text{ }^\circ\text{C}$ , and (c)  $\text{Fe}(\kappa^2P,N\text{-PNP})\text{Br}_2$   $(S_{\text{Fc}},S_{\text{Fc}})\text{-14}$  in  $\text{CD}_2\text{Cl}_2$  (spectra of  $(S_{\text{Fc}},S_{\text{Fc}})\text{-13a}$  and  $(S_{\text{Fc}},S_{\text{Fc}})\text{-14}$  are referenced internally to  $\text{PPh}_3$  set to  $-5.6\text{ ppm}$ ).



**Figure 7.** Structural view of  $(S_{Fc},S_{Fc})$ -**14** showing 50% thermal ellipsoids (selected H atoms and solvent molecules are omitted for clarity). Selected bond lengths ( $\text{\AA}$ ) and bond angles ( $^\circ$ ): Fe1–Br1 2.3911(7), Fe1–Br2 2.3887(6), Fe1–P1 2.400(1), Fe1–N1 2.088(3), P2–B1 1.927(5), Br1–Fe1–Br2 117.74(3), P1–Fe1–Br1 110.70(3), P1–Fe1–Br2 112.80(3), Br1–Fe1–N1 115.53(9), Br2–Fe1–N1 102.68(9).


**Reactivity towards CO.** In order to obtain iron-based active hydrogenation catalysts it appears to be important to have diamagnetic complexes. Accordingly, virtually all iron complexes that are active catalysts feature the strong field ligand CO, which seems to maintain a low spin configuration throughout the catalytic cycle.<sup>24–31</sup> Therefore,  $(S_{Fc},S_{Fc})$ -**13a** was stirred overnight in  $\text{CH}_2\text{Cl}_2$  under a CO atmosphere (2 bar) in an autoclave to obtain mono-CO or *bis*-CO complexes of the types  $(S_{Fc},S_{Fc})$ -**15a** and  $(S_{Fc},S_{Fc})$ -**16a** (Scheme 5). Under these conditions mono-CO complexes could not be isolated and only  $(S_{Fc},S_{Fc})$ -**16a** was obtained. Due to the fact that CO compounds are not stable and readily release CO, NMR measurements on  $(S_{Fc},S_{Fc})$ -**16a** were carried out under a CO headspace. The  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum contained two major and three minor pairs of doublets with  $^2J_{\text{pp}}$  values in the range 165–178 Hz. These signals are indicative of a *trans*-phosphorus arrangement. The  $^1\text{H}$  NMR spectrum was not very informative due to its complexity and the presence of isomers. The HRMS revealed a fragment of  $[\text{Fe}(\text{PNP})(\text{CO})_2\text{Br}]^+$  ( $[\text{M}]^+$ ) at  $m/z$  969.9684. The IR spectra displayed two strong  $\nu_{\text{CO}}$  absorptions at 1990 and 2038  $\text{cm}^{-1}$ , which implies a *cis*-CO geometry around the iron center. However, attempts to grow crystals under a CO atmosphere were unsuccessful. The experimental results are in agreement with DFT calculations, where *bis* carbonyl complexes  $(S_{Fc},S_{Fc})$ -**16a**, in which the phosphorus moieties are in a mutual *trans* position, are more stable by 50 kcal/mol than the respective complexes  $(S_{Fc},S_{Fc})$ -**17a**, which have a *cis* phosphorus

arrangement (Scheme 5). In accordance with these results, DFT calculations revealed that addition of CO to ( $S_{Fc},S_{Fc}$ )-**13a** is a thermodynamically highly unfavorable process (Figure 8). The putative CO complexes *trans*-P, *trans*-Br-[Fe(PNP)(CO)Br<sub>2</sub>] (**D**), *cis*-Br, *trans*-P-[Fe(PNP)(CO)Br<sub>2</sub>] (**E**) and *cis*-Br, *cis*-P-[Fe(PNP)(CO)Br<sub>2</sub>] (**F**) are 35.1, 37.4, and 88.9 kcal/mol, respectively, are less stable than the initial reagent **A** ( $S_{Fc},S_{Fc}$ )-**13a**.



**Figure 8.** Free energy profile (B3LYP) for the addition of CO to form the illusive complexes *trans*-P, *trans*-Br-[Fe(PNP)(CO)Br<sub>2</sub>] ( $S_{Fc},S_{Fc}$ )-**15a** (**D**), *cis*-Br, *trans*-P-[Fe(PNP)(CO)Br<sub>2</sub>] ( $S_{Fc},S_{Fc}$ )-**15a** (**E**) and *cis*-Br, *cis*-P-[Fe(PNP)(CO)Br<sub>2</sub>] ( $S_{Fc},S_{Fc}$ )-**15a** (**F**). The energy values (kcal/mol) are referred to [Fe( $\kappa^2$ P,N-PNP)Br<sub>2</sub>] ( $S_{Fc},S_{Fc}$ )-**13a** (**A**).

In addition, ( $S_{Fc},S_{Fc}$ )-**16a** was also prepared from Fe(CO)<sub>4</sub>(Br)<sub>2</sub>, as a different iron source, and the ligand ( $S_{Fc},S_{Fc}$ )-**1a** and the structure was established by NMR, HRMS and IR. In a similar way ( $S_{Fc},S_{Fc}$ )-**16b** was prepared from ( $S_{Fc},S_{Fc}$ )-**13a** and this compound had comparable spectroscopic data to ( $S_{Fc},S_{Fc}$ )-**16a**.



Scheme 5.

In the next step, in an attempt to remove a bromide ligand,  $(S_{Fc}, S_{Fc})$ -**16a** and  $(S_{Fc}, S_{Fc})$ -**16b** were reacted with 1.07 equivalents of a halide abstractor ( $AgBF_4$  and  $AgPF_6$ ) in THF under a CO atmosphere (2 bar). The inclusion of a halide abstractor led to an immediate color change from red to red-purple. The  $^{31}P\{^1H\}$  NMR spectra of isolated compounds showed the formation of a new compound with an ABX pattern and  $^2J_{PP}$  couplings in the range 44–47 Hz. However, the mixtures were intractable and all attempts to characterize these compounds failed.

**Synthesis of hydride complexes.** Iron hydride complexes as active catalysts for the asymmetric hydrogenation of ketones and imines were synthesized by treating iron dicarbonyl complexes  $(S_{Fc}, S_{Fc})$ -**16a** and  $(S_{Fc}, S_{Fc})$ -**16b** with various hydride reagents such as  $Na[HB(Et)_3]$ ,  $NaBH_4$  and  $LiAlH_4$ . However, in all cases decomposition was observed and the main product was the reduced free ligand. As mentioned above, the cases of  $(S_{Fc}, S_{Fc})$ -**2** and  $(R,R,S_{Fc}, S_{Fc})$ -**3**, neither of which contain an imine bond, coordination with iron was not observed. This finding could explain the failure of all attempts to form iron hydride complexes and the fact that the free ligand was obtained.

**Catalysis.**  $(S_{Fc}, S_{Fc})$ -**16a** and  $(S_{Fc}, S_{Fc})$ -**16b** were used as catalyst precursors in the asymmetric hydrogenation of acetophenone (reaction conditions: 25 °C, 5 bar  $H_2$ , 0.1 mol % of precatalyst and 4% potassium *tert*-butoxide). However, it was found that these compounds are not active in catalysis.

## Conclusions

New chiral imine and amine type PNP ligands, all based on diferrocene scaffold, were developed. While the amine type ligands did not coordinate to iron with use of the imine ligands Fe(II) complexes of general formula  $[\text{Fe}(\text{PNP})\text{X}_2]$  ( $\text{X} = \text{Cl}, \text{Br}$ ) were obtained. In the solid state, as established by X-ray crystallography and supported by Mössbauer result, these complexes adopt a tetrahedral geometry with the PNP ligand coordinated in a  $\kappa^2P,N$ -fashion with one pendant-arm but the other arm is not coordinated. All complexes are paramagnetic with a quintet ground state. Equilibria exist in solution between  $[\text{Fe}(\kappa^3P,N,P\text{-PNP})\text{X}_2]$  and  $[\text{Fe}(\kappa^2P,N\text{-PNP})\text{X}_2]$  complexes. Boronation of the non-coordinated arm shifts the equilibrium towards the four-coordinate complex  $[\text{Fe}(\kappa^2P,N\text{-PNP}^{\text{BH}_3})\text{Br}_2]$ . The results of DFT calculations are in agreement with the experimental results and indicates that the experimentally observed  $\kappa^2P,N$  isomer is thermodynamically the most stable. Under a CO atmosphere  $[\text{Fe}(\text{PNP})(\text{CO})_2\text{Br}]\text{Br}$  was formed rather than  $[\text{Fe}(\text{PNP})(\text{CO})\text{Br}_2]$ . However, this complex decomposed upon removing the CO atmosphere and  $[\text{Fe}(\text{PNP})\text{Br}_2]$  was formed. All attempts to transform  $[\text{Fe}(\text{PNP})(\text{CO})_2\text{Br}]\text{Br}$  complexes into hydride derivatives resulted in decomposition of complexes and gave the reduced ligand as a main product. The reluctance of the Fe(PNP) complexes to form low-spin mono CO complexes, which seems to be a prerequisite for the subsequent formation of hydride complexes, may explain, why no hydride complexes are formed and thus why these complexes are inactive in hydrogenation catalysis. All complexes remain in the high spin state and renders the PNP ligands too labile, which upon treatment with hydride sources liberate the PNP ligands. This again shows that small electronic changes in the ligand environment of first row transition metal complexes has a drastic impact on reactivity.

**Computational Details.** Calculations were performed using the GAUSSIAN 09 software package, and the B3LYP functional without symmetry constraints. This functional was shown to perform well in mechanistic studies of spin forbidden reactions in closely related Fe systems. The optimized geometries were obtained with the Stuttgart/Dresden ECP (SDD) basis set to describe the electrons of the iron atom. For all other atoms a standard 6-31G\*\* basis set was employed. Frequency calculations were performed to confirm the nature of the stationary points the yielded no imaginary frequency for the minima.

**Acknowledgement.** Financial support from the Austrian Science Fund (FWF) is gratefully acknowledged (Projects No. T 631-N28). The authors wish to express sincere appreciation and

gratitude to Professor Walter Weissensteiner for his kind support and helpful discussions and comments during the course of this research. The NMR Center the University of Vienna - in particular Susanne Felsing - is acknowledged. The X-ray center of the University of Vienna and Vienna University of Technology is also acknowledged for financial support and for providing access to the single-crystal diffractometer. MDC and LPF acknowledge Fundação para a Ciência e Tecnologia (Projects UID/MULTI/00612/2013 and UID/MULTI/04046/2013).

†Electronic supplementary information (ESI) available.  $^1\text{H}$  and  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra of all ligands and complexes. Complete crystallographic data and technical details in CIF format for  $(S_{\text{Fc}}, S_{\text{Fc}})$ -**1a**,  $(S_{\text{Fc}}, S_{\text{Fc}})$ -**13a**,  $(S_{\text{Fc}}, S_{\text{Fc}})$ -**13c**, and  $(S_{\text{Fc}}, S_{\text{Fc}})$ -**14** (CCDC entries 1435577, 1435448, 1433174 and 1433175). For ESI and crystallographic data in CIF or other electronic format see DOI: xxxxxxxx.



**References:**

1. H.-U. Blaser, C. Malan, B. Pugin, F. Spindler, H. Steiner and M. Studer, *Adv. Synth. Catal.*, 2003, **345**, 103–151.
2. F. Naud, F. Spindler, C. J. Rueggeberg, A. T. Schmidt and H.-U. Blaser, *Org. Process Res. Dev.*, 2007, **11**, 519–523.
3. J. G. De Vries and C. J. Elsevier, Eds., *Handbook of Homogeneous Hydrogenation*, Wiley-VCH, Weinheim, 2007, **40**, Vols. 1–3.
4. I. Ojima, Ed., *Catalytic Asymmetric Synthesis*, Wiley-VCH, Weinheim, 2000.
5. H. Brunner and W. Zettlmeier, *Handbook of Enantioselective Catalysis*, VCH, Weinheim, 1993.
6. P. Bhattacharya and H. Guan, *Comments on Inorganic Chemistry*, 2011, **32**, 88–112.
7. C. Bolm, J. Legros, J. Le Paih and L. Zani, *Chem. Rev.*, 2004, **104**, 6217–6254.
8. S. Enthaler, K. Junge and M. Beller, *Angew. Chem., Int. Ed.*, 2008, **47**, 3317–3321.
9. E. B. Bauer, *Curr. Org. Chem.*, 2008, **12**, 1341–1369.
10. R. H. Morris, *Chem. Soc. Rev.*, 2009, **38**, 2282–2291.
11. L.-X. Liu, *Curr. Org. Chem.*, 2010, **14**, 1099–1126.
12. M. Zhang and A. Zhang, *Appl. Organomet. Chem.*, 2010, **24**, 751–757.
13. K. Junge, K. Schröder and M. Beller, *Chem. Commun.*, 2011, **47**, 4849–4859.
14. H. Nakazawa and M. Itazaki, *Organomet. Chem.*, 2011, **33**, 27–81.
15. P. O. Lagaditis, P. E. Sues, J. F. Sonnenberg, K. Y. Wan, A. J. Lough and R. H. Morris, *J. Am. Chem. Soc.*, 2014, **136**, 1367–1380.
16. P. Štěpnička, *Ferrocenes: Ligands, Material and Biomolecules*, Ed.; Wiley: Chichester, 2008.
17. O. Riant, O. Samuel, T. Flessner, S. Taudien and Henri B. Kagan, *J. Org. Chem.*, 1997, **62**, 6733–6745.
18. X. Hu, H. Dai, X. Hu, H. Chen, J. Wang, C. Bai and Z. Zheng, *Tetrahedron: Asymmetry*, 2002, **13**, 1687–1693.

19. W. Li, T. Wei, Y. Gao, K. Xi and X. Jia, *Polymer*, 2012, **53**, 1236–1244.
20. E. Alberico, P. Sponholz, C. Cordes, M. Nielsen, H. J. Drexler, W. Baumann, H. Junge and M. Beller, *Angew. Chem. Int. Ed.*, 2013, **52**, 14162–14166.
21. C. Holzhaecker, B. Stöger, D. M. Carvalho, L. P. Ferreira, E. Pittenauer, G. Allmaier, L. F. Veiros, S. Realista, A. Gil and M. J. Calhorda, *Dalton Trans.*, 2015, **44(29)**, 13071–13086.
22. E. J. Hawrelak, H. Wesley, E. Lobkovsky, G. T. Yee, E. Bill and P. Chirik, *Inorg. Chem.*, 2005, **44(9)**, 3103–3111.
23. H. Schottenberger, M. R. Buchmeiser, and R. H. Herber, *J. Organomet. Chem.*, 2000, **612**, 1–8.
24. P. O. Lagaditis, P. E. Sues, A. J. Lough and R. H. Morris, *Dalton Trans.*, 2015, **44**, 2119–12127.
25. S. Werkmeister, K. Junge, B. Wendt, E. Alberico, H. Jiao, W. Baumann, H. Junge, F. Gallou and M. Beller, *Angew. Chem., Int. Ed.*, 2014, **53**, 8722–8726.
26. C. Bornschein, S. Werkmeister, B. Wendt, H. Jiao, E. Alberico, W. Baumann, H. Junge, K. Junge and M. Beller, *Nat. Commun.*, 2014, **5**, DOI: 10.1038/ncomms5111.
27. S. Chakraborty, H. Dai, P. Bhattacharya, N. T. Fairweather, M. S. Gibson, J. A. Krause and H. Guan, *J. Am. Chem. Soc.*, 2014, **136**, 7869–7872.
28. S. Chakraborty, W. W. Brennessel and W. D. Jones, *J. Am. Chem. Soc.*, 2014, **136**, 8564–8567.
29. E. A. Bielinski, P. O. Lagaditis, Y. Zhang, B. Q. Mercado, C. Würtele, W. H. Bernskoetter, N. Hazari and S. Schneider, *J. Am. Chem. Soc.*, 2014, **136**, 10234–10237.
30. R. Langer, G. Leitius, Y. Ben-David and D. Milstein, *Angew. Chem., Int. Ed.*, 2011, **50**, 2120–2124.
31. R. Langer, M. A. Iron, L. Konstantinovski, Y. Diskin-Posner, G. Leitius, Y. Ben-David and D. Milstein, *Chem. – Eur. J.*, 2012, **18**, 7196–7209.
32. C. Holzhaecker, M. J. Calhorda, A. Gil, M. D. Carvalho, L. P. Ferreira, K. Mereiter, B. Stöger, E. Pittenauer, G. Allmaier and K. Kirchner, *Polyhedron*, 2014, **81**, 45–55.
33. C. Holzhaecker, C. M. Standfest-Hauser, M. Puchberger, K. Mereiter, L. F. Veiros, M. J. Calhorda, M. D. Carvalho, L. P. Ferreira, M. Godinho, F. Hartl and K. A. Kirchner, *Organometallics*, 2011, **30**, 6587–6601.

**Graphic for Table of Contents**

New chiral PNP and PNHP ferrocene ligands and the corresponding Fe(II) complexes of PNP ligands with the general formula  $[\text{Fe}(\text{PNP})\text{X}_2]$  ( $\text{X} = \text{Cl}, \text{Br}$ ) were synthesized.

