

# Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

## ARTICLE

# Electrophilic *Bis*-Fluorophosphonium Dications: Lewis Acid Catalysts from Diphosphines

Cite this: DOI: 10.1039/x0xx00000x

Michael H. Holthausen,<sup>a</sup> Rashi R. Hiranandani<sup>a</sup> and Douglas W. Stephan<sup>a,b</sup>Received 00th January 2015,  
Accepted 00th January 2015

DOI: 10.1039/x0xx00000x

www.rsc.org/

Stepwise oxidation of 1,8-*bis*(diphenylphosphino)naphthalene and a series of (oligo)methylene-linked diphosphines with XeF<sub>2</sub> followed by fluoride abstraction yields a family of compounds featuring phosphine, phosphonium and phosphorane moieties in close proximity. The *bis*phosphonium ions [(C<sub>10</sub>H<sub>6</sub>)(Ph<sub>2</sub>PF<sub>2</sub>)<sub>2</sub>]<sup>2+</sup> (**5**) and [CH<sub>2</sub>(Ph<sub>2</sub>PF<sub>2</sub>)<sub>2</sub>]<sup>2+</sup> (**9a**) exhibit remarkable Lewis acidity arising from the proximity of the phosphonium centers. The effectiveness of bisphosphonium dications (**5**, **9a-e**) is examined in a series of Lewis acid catalysed transformations.

## Introduction

Main group based Lewis acids are used as co-catalysts in polymerization catalysis, or as Lewis acid catalysts in a wide range of organic transformations.<sup>[1]</sup> While such reactivity is predicated only on being electron deficient, Group XIII compounds such as BH<sub>3</sub>, BF<sub>3</sub>, or AlMe<sub>3</sub> are the quintessential examples of main-group Lewis acids. Group XIV-centered Lewis acids (C,<sup>[2]</sup> Si<sup>[3]</sup>) have also been exploited although as these compounds are generally very strong Lewis acids and coordinatively unsaturated, they tend to be employed as stoichiometric reagents. Moving further to the right to group XV, derivatives of these elements are most commonly Lewis bases and the electrophilic character of group XV species has drawn lesser attention. Nonetheless, P(III) phosphonium ions [R<sub>2</sub>P]<sup>+</sup> have been shown to be ambiphilic effecting C–C, C–H<sup>[4]</sup> and P–P<sup>[5]</sup> bond activation. P(V) phosphonium cations of the form [R<sub>4</sub>P]<sup>+</sup> (R = aryl, alkyl) are weakly Lewis acidic and have been exploited as fluoride ion sensors<sup>[6]</sup> or as catalysts for (cyclo)addition reactions to polar or activated unsaturates.<sup>[7]</sup> Exploring more electron deficient phosphonium cations, we found that [(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>PF]<sup>+</sup> is a highly electrophilic phosphonium cation (EPC). This Lewis acidity is attributed to the presence of the strongly electron-withdrawing substituents and the energetically accessible σ\*(P–F) orbital.<sup>[8]</sup> In an alternative approach towards EPCs, the dicationic phosphonium salt [(SImes)PPh<sub>2</sub>F][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]<sub>2</sub><sup>[9]</sup> was shown to be even more Lewis acidic than [(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>PF]<sup>+</sup>, demonstrating the impact of the additional positive charge. Nonetheless both [(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>PF]<sup>+</sup> and [(SImes)PPh<sub>2</sub>F]<sup>2+</sup> were shown to be effective catalysts for the hydrodefluorination of fluoroalkanes and the hydrosilylation or transfer-hydrogenation of olefins and alkynes.<sup>[8–10]</sup>

In Group XIII<sup>[11]</sup> and group XIV chemistry,<sup>[12]</sup> the proximity of two Lewis acidic centers has been shown to enhance the Lewis acidity, providing even more potent Lewis

acids and thus impact on reactivity. Only recently, a group XV-based bidentate stiborane was introduced as strong, chelating Lewis acid for fluoride complexation.<sup>[13]</sup> Exploring these notions in phosphonium chemistry prompted us to target bidentate *bis*-phosphonium salts. In this manuscript we prepare a series of such compounds exploiting a naphthalene backbone which locks two P moieties in the close vicinity of the *peri*-positions. In addition a series of more flexible (oligo)methylene-linked derivatives ((CH<sub>2</sub>)<sub>n</sub>, n = 2, 3, 4, 5) are prepared. The impact of the proximity of two EPC centers is gauged by an examination of these species in a series of Lewis acid catalyses.


## Results and Discussion

The naphthalene-based diphosphine (C<sub>10</sub>H<sub>6</sub>)(PPh<sub>2</sub>)<sub>2</sub> reacts with XeF<sub>2</sub> at low temperatures (–30 °C) affording the selective formation of the phosphine-phosphorane species (C<sub>10</sub>H<sub>6</sub>)(Ph<sub>2</sub>PF<sub>2</sub>)(PPh<sub>2</sub>) (**1**) in high yields (87%, Scheme 1). The <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of **1** shows two triplet of doublet resonances at –55.5 ppm (<sup>1</sup>J<sub>PF</sub> = 742 Hz, <sup>4</sup>J<sub>PP</sub> = 10 Hz) and –18.5 ppm (<sup>5</sup>J<sub>PF</sub> = 17 Hz, <sup>4</sup>J<sub>PP</sub> = 10 Hz) which are assigned to the phosphorane and phosphine moieties respectively.<sup>[14]</sup> The associated doublet of doublet resonance of the fluorine atoms shows a chemical shift of –42.0 ppm in the <sup>19</sup>F{<sup>1</sup>H} NMR spectrum.

Reacting (C<sub>10</sub>H<sub>6</sub>)(PPh<sub>2</sub>)<sub>2</sub> with XeF<sub>2</sub> in a 1 : 2 stoichiometry at ambient temperature yields quantitatively the *bis*-difluorophosphorane (C<sub>10</sub>H<sub>6</sub>)(Ph<sub>2</sub>PF<sub>2</sub>)<sub>2</sub> (**2**, 95%). An AA'X<sub>2</sub>X<sub>2</sub>' spin system is observed in the <sup>31</sup>P{<sup>1</sup>H} (δ<sub>A</sub> = –49.4 ppm) and <sup>19</sup>F{<sup>1</sup>H} NMR (δ<sub>X</sub> = –34.3 ppm) spectra of **2** and its parameters were evaluated by full line shape iteration.<sup>[15]</sup> Both resonances of **2** are shifted to lower field compared to the phosphorane moiety in **1**. A relatively large <sup>5</sup>J<sub>PF</sub> (<sup>5</sup>J<sub>A'X</sub> = <sup>5</sup>J<sub>AXA'</sub> = –18 Hz) and a comparatively small <sup>4</sup>J<sub>PP</sub> coupling constant (<sup>5</sup>J<sub>AA'</sub> = 3 Hz)

might indicated a through space interaction between both phosphonium moieties.<sup>[16]</sup>

The molecular structure of **2** shows two trigonal bipyramidal phosphorus moieties with fluoride atoms occupying the axial positions (*av.* F–P–F: 177.0(1)°, Figure 1). The P–F bond length (*av.* 1.67(3) Å) is typical for difluorotriarylphosphoranes.<sup>[14, 17]</sup> The close proximity of the phosphorane moieties in **2** leads to significant steric strain. This results in a buckling of its naphthalene-backbone as evidenced by the mean C-atom deviation from the least-squares C<sub>10</sub> plane of 0.120(2) Å. The corresponding deviation in the parent diphosphine (C<sub>10</sub>H<sub>6</sub>)(Ph<sub>2</sub>P)<sub>2</sub> is 0.006 Å.<sup>[18]</sup> The steric strain also causes the phosphorus moieties of **2** to exhibit a strong bending out of the naphthalene plane (*av.* 0.73(1) Å *vs.* (C<sub>10</sub>H<sub>6</sub>)(Ph<sub>2</sub>P)<sub>2</sub> *av.* 0.42 Å). The shortest P...F interactions between the phosphorane moieties is 2.825(4) Å which is well within the sum of the respective van der Waal's radii (PF: 3.27 Å).<sup>[19]</sup> The corresponding P...P distance of 3.823(3) Å is significantly longer than that in the parent diphosphine (3.052 Å).<sup>[18]</sup> Earlier reports have described difficulties in effecting the quaternisation of both P atoms in (C<sub>10</sub>H<sub>6</sub>)(Ph<sub>2</sub>P)<sub>2</sub> due to steric congestion,<sup>[20]</sup> and, to the best of our knowledge, **2** represents the first compound featuring two penta-coordinated P atoms in *peri*-positions on a naphthalene framework.




**Figure 1.** POV-ray depiction of **2**. P: orange, F: yellow-green, C: black. Hydrogen atoms are omitted for clarity.

The reactions of phosphoranes **1** and **2** with [Et<sub>3</sub>Si][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]**2**(C<sub>7</sub>H<sub>8</sub>) in 1 : 1 stoichiometries yielded the respective salts [(C<sub>10</sub>H<sub>6</sub>)(Ph<sub>2</sub>PF)(PPh<sub>2</sub>)]<sup>+</sup>[B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]<sup>-</sup> (**3**) and [(C<sub>10</sub>H<sub>6</sub>)(Ph<sub>2</sub>PF<sub>2</sub>)(Ph<sub>2</sub>PF)]<sup>+</sup>[B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]<sup>-</sup> (**4**) in good to excellent yields (Scheme 1). The <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of **3** shows doublet of doublet resonances at -17.4 and -5.5 ppm. The former is assigned to the Ph<sub>2</sub>PF-moiety on basis of its <sup>1</sup>J<sub>PF</sub> value of 783 Hz. A very large coupling constant between both P atoms (138 Hz) indicates coordination of the lone pair of the phosphine-moiety to the fluorophosphonium centre.<sup>[21]</sup> This also results in a large coupling constant between the fluorine atom and the distal P atom (164 Hz; *vs.* <sup>5</sup>J<sub>PF</sub> = 17 Hz in **1**). The indicated PP interaction was confirmed by single crystal X-ray crystallography (Figure 2) as the P–P bond length was found to be 2.530(1) Å. This is longer than the P–P distance in P<sub>4</sub> (2.2 Å)<sup>[22]</sup> indicating that the interaction in **3** is weak. The

fluoro-substituted P atom shows a trigonal bipyramidal bonding environment with the fluorine and the coordinating phosphine-moiety adopting axial positions. The P–F bond distance (1.637(2) Å) is significantly longer than in other fluorophosphonium species (*av.* 1.54(1) Å)<sup>[8, 9, 14]</sup> which is consistent with the donation of electron density from the phosphine unit into the σ\*-orbital of the P–F bond. In contrast to **2**, the naphthalene backbone of **3** show no evidence of steric strain (mean C-atom deviation from the least-squares C<sub>10</sub> plane: 0.006(1) Å).


It is noteworthy that interactions between the P atoms of adjacent phosphine and alkyl or aryl phosphonium centers in the related cations [(C<sub>10</sub>H<sub>6</sub>)(Ph<sub>2</sub>PR)(Ph<sub>2</sub>P)]<sup>+</sup> (R = alkyl, aryl) is not observed,<sup>[17, 20a, 23]</sup> reflecting the Lewis acidity of the fluorophosphonium cation. While we have previously reported the intramolecular coordination of an amide fragment to a fluorophosphonium moiety,<sup>[24]</sup> compound **3** appears to be the first example of phosphine coordinating to a fluorophosphonium center. It is important to note that such an interaction is not observed in a 1:1 mixture of Ph<sub>3</sub>P and [Ph<sub>3</sub>PF][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] indicating that the close proximity between Lewis acid and base imposed by the naphthalene backbone is crucial for adduct formation.




**Scheme 1.** Synthetic routes to **1-5**.

The <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of [(C<sub>10</sub>H<sub>6</sub>)(Ph<sub>2</sub>PF<sub>2</sub>)(Ph<sub>2</sub>PF)]<sup>+</sup> (**4**) shows a broad triplet resonance at -51.7 ppm (<sup>1</sup>J<sub>PF</sub> = 717 Hz) assigned to the phosphorane moiety and a broad doublet at 96.6 ppm (<sup>1</sup>J<sub>PF</sub> = 1012 Hz) corresponding to the

phosphonium centre. A crystallographic study revealed that the P–F bond lengths at the penta-coordinated P atom of **4** (*av.* 1.664(2) Å) are longer than the P–F bond distance at the adjacent fluorophosphonium centre (1.543(1) Å; Figure 3). Similar to **2**, a weak P⋯F interaction between the F on the phosphonium and P of the phosphorane fragment is observed (2.995(2) Å). Compared to **2** there is a decreased buckling of the naphthalene moiety (mean C-atom deviation from the least-squares C<sub>10</sub> plane: 0.097(1) Å), a shorter P⋯P distance (3.668(2) Å), and the phosphonium P atom in **4** shows a reduced bending out of the naphthalene plane (0.057(1) Å) compared to its phosphorane neighbour (0.076(1) Å).



**Figure 2.** POV-ray depiction of the cation in **3**. P: orange, F: yellow-green, C: black. Hydrogen atoms are omitted for clarity.




**Figure 3.** POV-ray depiction of the cation in **4**\*(CH<sub>2</sub>Cl<sub>2</sub>). P: orange, F: yellow-green, C: black. Hydrogen atoms are omitted for clarity. The solvate molecule was removed by the squeeze routine of the program PLATON.

The *bis*-difluorophosphorane **2** also reacts with two equivalents of [Et<sub>3</sub>Si][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]\*2(C<sub>7</sub>H<sub>8</sub>) to give the salt [(C<sub>10</sub>H<sub>6</sub>)(Ph<sub>2</sub>PF)<sub>2</sub>][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]<sub>2</sub> (**5**) in high yields (84%, Scheme 1). The <sup>31</sup>P{<sup>1</sup>H} and <sup>19</sup>F{<sup>1</sup>H} NMR spectra of **5** reveal an AA'XX' spin system. The <sup>19</sup>F{<sup>1</sup>H} NMR spectrum reveals a very broad doublet resonance ( $\delta_X = -117.1$  ppm,  $\Delta\nu_{1/2} = 450$  Hz). The <sup>31</sup>P{<sup>1</sup>H} NMR resonance ( $\delta_A = 96.8$  ppm) shows a relatively large <sup>1</sup>J<sub>PF</sub> coupling constant (<sup>1</sup>J<sub>AX} = <sup>1</sup>J<sub>A'X'} = 1004 Hz) which has also been observed for the related cations [(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>PF]<sup>+</sup> (1062 Hz),<sup>[8]</sup> and [(SIMes)PPh<sub>2</sub>F]<sup>2+</sup> (1040 Hz)<sup>[9]</sup>. Similar to **2**, dication **5** also features a relative large <sup>5</sup>J<sub>PF</sub> (<sup>5</sup>J<sub>A'X} = <sup>5</sup>J<sub>A'X'} = -13 Hz) and a comparatively small <sup>4</sup>J<sub>PP</sub> coupling constant (<sup>5</sup>J<sub>AA'} = 3 Hz). The</sub></sub></sub></sub></sub>

related doubly protonated naphthalene-based *bis*-phosphine was previously studied by NMR spectroscopy,<sup>[25]</sup> while the dications incorporating a P–P bond<sup>[21b, 26]</sup> or a bridging substituent<sup>[23, 27]</sup> have been previously isolated.

In an effort to probe the Lewis acidity of **5**, it was reacted with an equivalent of Et<sub>3</sub>PO employing the Gutmann-Beckett method (Scheme 2).<sup>[28]</sup> The <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of the reaction mixture shows a broad resonance at 60.5 ppm corresponding to Et<sub>3</sub>PO interacting with **5**. This results in an acceptor number of 21.4, which suggests that **5** is approximately half as Lewis acidic as B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>. In comparison, combination of the salt [Ph<sub>3</sub>PF][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] and Et<sub>3</sub>PO in a 2 : 1 stoichiometry showed only a very small shift in the resonance of Et<sub>3</sub>PO. These observations infer that the Lewis acidity of a phosphonium center in **5** is enhanced by the proximity of a second fluorophosphonium moiety. Monitoring the 1 : 1 mixture of **5** and Et<sub>3</sub>PO over time revealed slow transformation to [Et<sub>3</sub>PF]<sup>+</sup> (**6**)<sup>[9]</sup> and [(C<sub>6</sub>H<sub>10</sub>)(Ph<sub>2</sub>PO)(Ph<sub>2</sub>PF)]<sup>+</sup> (**7**). The <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of (**7**) shows the expected doublet of doublet resonances at -34.5 ppm (<sup>1</sup>J<sub>PF} = 697 Hz, J<sub>PP} = 23 Hz) and 47.1 ppm (J<sub>PF} = 2 Hz). The high field shift of the fluorophosphonium moiety and the low field shift of the phosphine oxide indicate some degree of donation of electron density from the PO fragment to the electrophilic fluorophosphonium cation. A similar but faster fluoride oxide exchange reaction was previously reported for the more Lewis acidic dication [(SIMes)PPh<sub>2</sub>F]<sup>2+</sup>.<sup>[9]</sup></sub></sub></sub>




**Scheme 2.** Reaction of **5** with Et<sub>3</sub>PO.

A family of related *bis*-phosphonium dication salts was envisioned and thus reactions of (oligo)methylene-bridged diphosphines (CH<sub>2</sub>)<sub>n</sub>(Ph<sub>2</sub>P)<sub>2</sub> (*n* = 1 (**a**), 2 (**b**), 3 (**c**), 4 (**d**), 5 (**e**)) and XeF<sub>2</sub> were performed in CH<sub>2</sub>Cl<sub>2</sub> to yield the respective *bis*-difluorophosphoranes (CH<sub>2</sub>)<sub>n</sub>(Ph<sub>2</sub>PF<sub>2</sub>)<sub>2</sub> (**8a-e**) in high to almost quantitative isolated yields (Scheme 3). The species **8a-d** have been previously prepared by using a range of oxidation reagents including F<sub>2</sub>,<sup>[29]</sup> SF<sub>4</sub>,<sup>[30]</sup> Me<sub>2</sub>NSF<sub>3</sub>,<sup>[31]</sup> (CF<sub>3</sub>)<sub>2</sub>CO<sup>[32]</sup> COF<sub>2</sub>,<sup>[33]</sup> and NF<sub>3</sub>O<sup>[34]</sup> or by halide exchange from the respective bromophosphoranes<sup>[30]</sup> or by hydrofluorination of the respective silyl phosphinimines with HF.<sup>[35]</sup> However, most of these methods suffer from low yields, laborious workup procedures and the use of hazardous reagents. The reaction of phosphoranes **8a-e** with two equivalents of [Et<sub>3</sub>Si][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]\*2(C<sub>7</sub>H<sub>8</sub>) resulted in the formation of corresponding *bis*-phosphonium dication salts **9a-e** (Scheme 3). All five derivatives were isolated and fully characterized. It is


curious to note that the salts **9b** and **9d**, featuring linkers with an even number of CH<sub>2</sub> units, are only sparingly soluble in CD<sub>2</sub>Cl<sub>2</sub> whereas those with an odd number of methylene carbons in the linker (**9a**, **9c**, **9e**) show good solubility. The respective NMR spectra of **9a-e** exhibit AA'XX' spin systems (see SI). Compounds **9a-c** show higher order effects due to coupling between the nuclei of the magnetically inequivalent phosphonium moieties and thus the NMR parameters were obtained via a full line shape iteration procedure.<sup>[15]</sup> Interestingly **9a** exhibits a large P–F coupling constant (<sup>1</sup>J<sub>AX</sub> = 1024 Hz) and significant coupling between both fluorine atoms (<sup>4</sup>J<sub>XX'</sub> = 12 Hz) suggesting enhanced Lewis acidity. Remarkable large P–P coupling constant was also observed for **9b** (<sup>3</sup>J<sub>AA'</sub> = 70 Hz) suggesting a weak through space interaction between the phosphorus centers (see SI).<sup>[36]</sup>

Lewis acidity tests of this series of bis-fluorophosphonium ion salts **9a-e** by the Gutmann-Beckett method<sup>[28]</sup> were performed (Scheme 4). Determination of an acceptor number for **9a** was hampered by deprotonation of the central CH<sub>2</sub> moiety, affording [Et<sub>3</sub>POH][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] (**10**) and [(CH)<sub>1</sub>(Ph<sub>2</sub>PF)<sub>2</sub>][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] (**11**). Compound (**10**) showed broad <sup>31</sup>P{<sup>1</sup>H} NMR resonance at 74.1 ppm, while (**11**) reveals an AA'XX' spin system in the <sup>31</sup>P{<sup>1</sup>H} (δ<sub>A</sub> = 81.2 ppm) and <sup>19</sup>F{<sup>1</sup>H} NMR spectrum (δ<sub>X</sub> = -99.4 ppm). The formation of **11** was further supported by its independent synthesis from **9a** and *t*-Bu<sub>3</sub>P.

44.1 ppm).<sup>[28]</sup> In contrast **12d-e** show <sup>31</sup>P{<sup>1</sup>H} NMR resonances of in the typical range for discrete phosphine oxide<sup>[37]</sup> and fluorophosphonium moieties.



Scheme 4. Reaction of **9a-e** with Et<sub>3</sub>PO.



Scheme 3. Synthetic route to **8a-e** and **9a-e**.

The cations of **9b-e** interact weakly with Et<sub>3</sub>PO resulting in acceptor numbers of ~5. However, in all cases slow conversion to the monocationic species [(CH<sub>2</sub>)<sub>n</sub>(Ph<sub>2</sub>PO)(Ph<sub>2</sub>PF)][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] (**12b-e**) and [Et<sub>3</sub>PF]<sup>+</sup>-salt **6** was observed.<sup>[9]</sup> The shorter linkers in the case of **12b** and **12c** result in a chelate interaction of the phosphine oxide and the electrophilic fluorophosphonium center as evidenced by the high field shift of the fluoro-substituted P atom (**12b**: 2.2 ppm, <sup>1</sup>J<sub>PF</sub> = 717 Hz, **12c**: 45.2 ppm, <sup>1</sup>J<sub>PF</sub> = 823 Hz). This notion is also supported by the observation of comparatively low field <sup>31</sup>P NMR resonances arising for the PO fragments (**12b**: 53.1 ppm, **12d**:


The utility of the dications **5**, **9a-e** in a variety of Lewis acid catalysed reactions was probed (Scheme 5). 1,1-Diphenylethylene in CD<sub>2</sub>Cl<sub>2</sub> was treated with 2 mol% of **5** or **9a-e**. In the case of **5** almost quantitative Friedel-Crafts dimerization was observed within 24 h at ambient temperature and the resulting product was isolated in 94% yield.<sup>[38]</sup> Complete conversion was also achieved using **9a-c**, however, **9a** required only one hour reaction time, while **9b,c** were complete in 24 h. In the case of **9d-e** under the same conditions after 24 h, the Friedel-Crafts product was formed in lesser yields (**9d**: 50%, **9e**: 25%). The corresponding reaction of 1,1-diphenylethylene with Et<sub>3</sub>SiH in the presence of these Lewis acid catalysts was assessed (Scheme 5).<sup>[9, 10a]</sup> Using **9a** as the catalyst gave 89% of the hydrosilylated product in one hour at ambient temperature whereas **9b** and **5** achieve comparable yields after 24 h at 50 °C. No conversion is observed in the case of less Lewis acidic catalysts **9c-e**.

In a similar fashion treatment of benzophenone with two equivalents of Et<sub>3</sub>SiH in the presence of these Lewis acid catalysts **5** and **9a,b** led to hydrodeoxygenation and complete conversion to diphenylmethane and Et<sub>3</sub>SiOSiEt<sub>3</sub> after one hour (**5**, **9a**) or three hours (**9b**) respectively (Scheme 5). Compounds **9c-e** required elevated temperatures and 36 h reaction time and afforded product in decreased yields (**9c**: 93%, **9d**: 72%, **9e**: 46%).

A fourth test involved the utility of these catalysts in the dehydrocoupling of Si–H bonds in silanes with phenol (Scheme 5).<sup>[10b]</sup> Compounds **5** and **9a** results in rapid dehydrogenative coupling which gives PhOSiEt<sub>3</sub> in high yields within one hour

at ambient temperature. In contrast, elevated temperature (50 °C) and 24 h reaction time were required to observe similar reactivity for catalyst **9b**. With an increase in methylene-linker length a decrease in conversion was observed (**9c**: 30%, **9d**: 19%) and only traces of the coupling product are detected using **9e**.

Finally, these EPCs were evaluated as catalysts for the hydrodefluorination of fluoropentane using Et<sub>3</sub>SiH as the hydride source (Scheme 5).<sup>[8, 9]</sup> Compounds **9a** and **5** catalyse the hydrodefluorination affording almost complete conversion after one hour at ambient temperature for (**9a**) or 24 h at 50 °C for (**5**). The less Lewis acidic species **9b-e** effect incomplete to trace conversions (**9b**: 18%, **9c**: 4%, **9d**: 2%, **9e**: 2%) after 24 h at 50 °C.



**Scheme 5.** Lewis acid catalysis by salt **5** and **9a-e** as catalysts for selected Lewis acid mediated transformations. Conversions were determined by <sup>1</sup>H NMR or <sup>19</sup>F spectroscopy and isolated yields are given in parenthesis. a: Ambient temperature and one hour reaction time (only for **9a**). b: Ambient temperature and one hour reaction time. c: Ambient temperature and one hour (**9a**, **5**) or three hours (**9b**) reaction time

The above results reveal that these dications are generally effective catalysts for a variety of Lewis acid mediated reactions. Moreover the proximity of two fluorophosphonium centers as in **5** and **9a-b** acts to enhance the Lewis acidity and the resulting reactivity by a distal P-F interaction. Indeed, the longer chain species (**9c-e**) show diminished reactivity behaving more like independent [Ph<sub>2</sub>PF(alkyl)]<sup>+</sup> cations. In this regard it is also noteworthy that in each of the above Lewis

acid-catalysed reactions the species [Ph<sub>3</sub>PF][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] was completely ineffective, which further supports the view that Lewis acidity of the present species is enhanced by spatial proximity.

## Conclusions

In summary, stepwise oxidations of 1,8-bis-(diphenylphosphino)naphthalene and a series of bidentate (oligo)methylene-linked phosphines with XeF<sub>2</sub> provide mono- and bisphosphorane species. Subsequent stepwise fluoride abstraction yielded phosphine / phosphonium, phosphonium / phosphorane and bisphosphonium species. The close spatial proximity in the naphthalene derived compound and the mono-methylene-linked species gives rise to Lewis acidity that is enhanced by the neighbouring fluorophosphonium moieties as evidenced by the reactivity of these species in the Lewis acid catalysed transformations including Friedel Crafts-type dimerization, hydrosilylation, dehydrocoupling, hydrodeoxygenation and hydrodefluorination. We are continuing to explore the facile and high yielding synthetic protocol that converts readily available bidentate donors providing Lewis acids of tuneable strength. New applications in Lewis acid catalysis and applications of these P(V) Lewis acids in frustrated Lewis pair (FLP) chemistry are the subject of ongoing efforts.

## Acknowledgements

D.W.S. gratefully acknowledges the financial support of the NSERC of Canada and is grateful for the award of a Canada Research Chair. M.H.H. thanks the Alexander von Humboldt Foundation for a Feodor Lynen Research Fellowship. R.R.H. is grateful for the support of an NSERC CGS-M scholarship.

## Notes and references

<sup>a</sup> Department of Chemistry, University of Toronto  
80 St. George St, Toronto Ontario M5S3H6 (Canada)

e-mail: dstephan@chem.utoronto.ca

<sup>b</sup> Chemistry Department-Faculty of Science,

King Abdulaziz University

Jeddah 21589, Saudi Arabia

Electronic Supplementary Information (ESI) available: Preparative, spectroscopic and analytical details for the experiments described herein are deposited in the supplementary information. Crystallographic data re deposited in CCDC# 1041558-1041561, See DOI: 10.1039/b000000x/

- (a) W. E. Piers and T. Chivers, *Chem. Soc. Rev.* 1997, 26, 345; (b) G. Erker, *Dalton Trans.* 2005, 1883; (c) T. Stahl, H. F. T. Klare and M. Oestreich, *ACS Catal.* 2013, 3, 1578; (d) F. J. Fernández-Alvarez, A. M. Aitani and L. A. Oro, *Catal. Sci. Technol.* 2014, 4, 611; (e) E. Dimitrijević and M. S. Taylor, *ACS Catal.* 2013, 3, 945; (f) K. Ishihara and H. Yamamoto, *Eur. J. Org. Chem.* 1999, 527; (g) F. Blank, C. Janiak, *Coord. Chem. Rev.* 2009, 253, 827; (h) S. V. Kostjuk and F. Ganachaud, *Acc. Chem. Res.* 2010, 43, 357; (i) W.

- Kaminsky, *Macromolecules* 2012, **45**, 3289. Ref for Lewis acid catalysis in general and polymerisation co-catalysts.
- 2 (a) B. Inés, S. Holle, R. Goddard and M. Alcarazo, *Angew. Chem. Int. Ed.* **2010**, *49*, 8389; (b) M. H. Holthausen, T. Mahdi, C. Schleppehorst, J. J. Weigand and D. W. Stephan, *Chem. Commun.* 2014, **50**, 10038; (c) E. R. Clark and M. J. Ingleson, *Organometallics* 2013, **32**, 6712; (d) E. R. Clark and M. J. Ingleson, *Angew. Chem. Int. Ed.* 2014, **53**, 11306; (e) M. P. Boone and D. W. Stephan, *Chem. Eur. J.* 2014, **20**, 3333.
- 3 (a) A. Schäfer, M. Reißmann, A. Schäfer, W. Saak, D. Haase and T. Müller, *Angew. Chem. Int. Ed.* 2011, **50**, 12636; (b) A. Schäfer, W. Saak, D. Haase and T. Müller, *Angew. Chem. Int. Ed.* 2012, **51**, 2981; (c) T. J. Herrington, B. J. Ward, L. R. Doyle, J. McDermott, A. J. P. White, P. A. Hunt and A. F. Ashley, *Chem. Commun.* 2014, **50**, 12753.
- 4 A. H. Cowley and R. A. Kemp, *Chem. Rev.* 1985, **85**, 367.
- 5 (a) C. A. Dyker and N. Burford, *Chem. Asian J.* 2008, **3**, 28; (b) M. H. Holthausen and J. J. Weigand, *Chem. Rev.* 2014, **43**, 15201.
- 6 (a) T. W. Hudnall, Y. M. Kim, M. W. P. Bebbington, D. Bourissou and F. P. Gabbaï, *J. Am. Chem. Soc.* 2008, **130**, 10890; (b) C. W. Chiu, Y. Kim and F. P. Gabbaï, *J. Am. Chem. Soc.* 2009, **131**, 60.
- 7 (a) A. Werner, *Adv. Synth. Catal.* 2009, **351**, 1469; (b) M. Terada and M. Kouchi, *Tetrahedron* 2006, **62**, 401.
- 8 C. B. Caputo, L. J. Hounjet, R. Dobrovetsky and D. W. Stephan, *Science* 2013, **341**, 1374.
- 9 M. H. Holthausen, M. Mehta and D. W. Stephan *Angew. Chem. Int. Ed.* 2014, **53**, 65387.
- 10 (a) M. Pérez, L. J. Hounjet, C. B. Caputo, R. Dobrovetsky and D. W. Stephan, *J. Am. Chem. Soc.* 2013, **135**, 18308; (b) M. Pérez, C. B. Caputo, R. Dobrovetsky and D. W. Stephan, *Proc. Natl. Acad. Sci. USA* 2014, **111**, 10917.
- 11 (a) C. Jiang, O. Blacque and H. Berke, *Chem. Commun.* 2009, 5518; (b) X. Zhao and D. W. Stephan, *Chem. Commun.* 2011, **47**, 1833; (c) M. J. Sgro, J. Dömer and D. W. Stephan, *Chem. Commun.* 2012, **48**, 7253; (d) W. E. Piers, G. J. Irvine and V. C. Williams, *Eur. J. Inorg. Chem.* 2000, 2131.
- 12 (a) T. J. Karol, J. P. Hutchinson, J. R. Hyde, H. G. Kuivilla and J. A. Zubieta, *Organometallics* 1983, **2**, 106; (b) H. G. Kuivilla, T. J. Karol and K. Swami, *Organometallics* 1983, **2**, 909; (c) M. Gielen and K. Jurkschat, *J. Organomet. Chem.* 1984, **273**, 303; (d) M. Austin, K. Gebreyes, H. G. Kivilla, K. Swami and J. A. Zubieta, *Organometallics* 1987, **6**, 834; (e) K. Jurkschat, F. Hesselbarth, M. Dagatz, J. Lehmann, E. Kleinpeter and A. Tzschach, *J. Organomet. Chem.* 1990, **388**, 259; (f) D. Dakternieks, K. Jurkschat, H. Zhu and E. R. T. Tiekink, *Organometallics* 1995, **14**, 251; (g) N. Asao, P. Liu and K. Maruoka, *Angew. Chem. Int. Ed.* 1997, **36**, 2507.
- 13 M. Hirai and F. P. Gabbaï, *Angew. Chem. Int. Ed.* 2015, **54**, 1205.
- 14 L. J. Hounjet, C. B. Caputo and D. W. Stephan, *Dalton Trans.* 2013, **42**, 2629.
- 15 P. H. M. Budzelaar, *gNMR for Windows (5.0.6.0) NMR Simulation Program*, IvorySoft **2006**.
- 16 (a) G. R. Miller, A. W. Yankowsky and S. O. Grim, *J. Chem. Phys.* 1969, **51**, 3185; (b) E. Fluck and D. Wachtler *Liebigs Ann. Chem.* 1979, 1125.
- 17 (a) R. R. Holmes, J. M. Holmes, R. O. Day, K. C. Kumara Swamy, V. Chandrasekhar, *Phosphorus, Sulfur and Silicon* 1995, **103**, 153; (b) G. M. Sheldrick, *Acta. Crystallogr., Sect. B: Struct. Crystallogr. Cryst. Chem.*, 1975, **31**, 1776.
- 18 R. D. Jackson, S. James, A. G. Orpen and P. G. Pringle, *J. Organomet. Chem.* 1993, **458**, C3.
- 19 A. Bondi, *J. Phys. Chem.* 1964, **68**, 441.
- 20 (a) T. Costa and H. Schmidbaur, *Chem. Ber.* 1982, **115**, 1374; (b) D. M. Upulani, K. Somisara, M. Bühl, T. Lebl, N. V. Richardson, A. M. Z. Slawin, J. D. Woollins and P. Kilian, *Chem. Eur. J.* 2011, **17**, 2666; (c) A. Karacar, M. Freytag, H. Thönnessen, J. Omelańczuk, P. G. Jones, R. Bartsch and R. Schmutzler, *Heteroatom Chem.* 2001, **12**, 102.
- 21 (a) M. Sanchez, R. Réau, F. Dahan, M. Regitz and G. Bertrand, *Angew. Chem. Int. Ed.* 1996, **35**, 2228; (b) M. Sanchez, R. Réau, H. Gornitzka, F. Dahan, M. Regitz and G. Bertrand, *J. Am. Chem. Soc.* 1997, **119**, 9720; (c) P. Kilian, D. Philp, A. M. Z. Slawin and J. D. Woollins, *Eur. J. Inorg. Chem.* 2003, 249.
- 22 (a) I. Krossing, *J. Am. Chem. Soc.* 2001, **123**, 4603; (b) H. Tai, I. Krossing, M. Seth and D. V. Deubel, *Organometallics* 2004, **23**, 2343.
- 23 K. Owsianik, L. Vendier, J. Błaszczuk and L. Sieroń, *Tetrahedron* 2013, **69**, 1628.
- 24 L. J. Hounjet, C. B. Caputo and D. W. Stephan, *Angew. Chem. Int. Ed.*, 2012, **51**, 4714.
- 25 A. Karaçar, V. Klaukien, M. Freytag, H. Thönnessen, J. Omelańczuk, P. G. Jones, R. Bartsch and R. Schmutzler, *Z. Anorg. Allg. Chem.* 2001, **627**, 2589.
- 26 (a) B. A. Surgenor, M. Bühl, A. M. Z. Slawin, J. D. Woollins and P. Kilian, *Angew. Chem. Int. Ed.* 2012, **51**, 10150; (b) M. J. Ray, A. M. Z. Slawin, M. Bühl and P. Kilian, *Organometallics* 2013, **32**, 3481.
- 27 (a) P. Kilian, A. M. Z. Slawin and J. D. Woollins, *Dalton Trans.* 2006, 2175; (b) K. Owsianik, R. Chauvin, A. Baliń, M. Wiczorek, M. Cypryk and M. Mikołajczyk *Organometallics* 2009, **28**, 4929.
- 28 V. Gutmann, *Coord. Chem. Rev.* 1976, **18**, 225.
- 29 I. Ruppert and V. Bastian, *Angew. Chem. Int. Ed. Engl.* 1977, **16**, 718.
- 30 E. Fluck and R. Braun, *Synth. React. Inorg. Met.-Org. Chem.* 1988, **18**, 727.
- 31 R. R. Holmes, J. M. Holmes, R. O. Day, K. C. Kumara Swamy and V. Chandrasekhar, *Phosphorus, Sulfur, and Silicon* 1995, **103**, 153.
- 32 I. V. Shevchenko, R. N. Mikolenko, E. Lork, G.-V. Rösenthaller, *Eur. J. Inorg. Chem.* 2001, **9**, 2377.
- 33 (a) O. D. Gupta and J. M. Shreeve, *J. Chem. Soc., Chem. Commun.* 1984, 416; (b) O. D. Gupta, R. L. Kirchmeier and J. M. Shreeve, *J. Fluorine Chem.* 1991, **52**, 1.
- 34 O. D. Gupta, R. L. Kirchmeier and J. M. Shreeve, *Inorg. Chem.* 1990, **29**, 573.
- 35 R. Appelt and I. Ruppert, *Chem. Ber.* 1975, **108**, 919.
- 36 (a) K.-O. Feldmann and J. J. Weigand, *J. Am. Chem. Soc.* 2012, **134**, 15443; (b) M. H. Holthausen, S. K. Surmiak, P. Jerabek, G. Frenking and J. J. Weigand, *Angew. Chem. Int. Ed.* 2013, **52**, 11078.
- 37 (a) W. Peng and J. M. Shreeve, *J. Fluorine Chem.* **2005**, *126*, 1054; (b) G. E. Maciel and R. V. James, *Inorg. Chem.* 1964, **3**, 1650.
- 38 (a) A. G. Evans and E. D. Owen, *J. Chem. Soc.* 1959, 4123; (b) L. J. Hounjet, C. Bannwarth, C. N. Garon, C. B. Caputo, S. Grimme and D. W. Stephan, *Angew. Chem. Int. Ed.* 2013, **52**, 7492.

TOC graphic

