


# Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

## Rh-Catalyzed Decarbonylation of Conjugated Ynonees via Carbon–Alkyne Bond Activation: Reaction Scope and Mechanistic Exploration via DFT Calculations

Alpay Dermenci<sup>†Δψ</sup>, Rachel E. Whittaker<sup>†Δ</sup>, Yang Gao<sup>‡§</sup>, Faben A. Cruz<sup>Δ</sup>, Zhi-Xiang Yu<sup>\*‡</sup>, and Guangbin Dong<sup>\*Δ</sup>

<sup>Δ</sup> The University of Texas at Austin, Department of Chemistry, Austin, TX 78712, United States


<sup>ψ</sup> Current Address: Pfizer Inc., Eastern Point Road, Groton, CT 06340, United States

<sup>†</sup> AD and REW contributed equally.

<sup>‡</sup> Beijing National Laboratory of Molecular Sciences (BNLMS), Key Laboratory of Bioorganic Chemistry and Molecular Engineering, College of Chemistry, Peking University, Beijing, 100871, China

<sup>§</sup> Key Laboratory of Pesticide & Chemical Biology, Ministry of Education, College of Chemistry, Central China Normal University, Hubei, Wuhan 430079, China


**Abstract:** In this full article, detailed development of a catalytic decarbonylation of conjugated monoynonees to synthesize disubstituted alkynes is described. The reaction scope and limitation has been thoroughly investigated, and a broad range of functional groups including heterocycles were compatible under the catalytic conditions. Mechanistic exploration via DFT calculations has also been executed. Through the computational study, a proposed catalytic mechanism has been carefully evaluated. These efforts are expected to serve as an important exploratory study for developing catalytic alkyne-transfer reactions via carbon–alkyne bond activation.




## INTRODUCTION

Transition metal-mediated carbon–carbon  $\sigma$  bond (C–C) activation offers a distinct strategy to construct or assemble organic molecules from unexpected, yet readily available starting materials.<sup>1,2</sup> Despite a number of C–C activation modes reported to date, limited catalytic approaches are available without relying on release of ring strain or use of an auxiliary directing group.<sup>3</sup> One important example that avoids these requirements is the catalytic activation of C–CN bonds, which has found broad usage in organic synthesis enabling “CN transfer” transformations (Scheme 1A).<sup>4</sup> Given that the cyano group can be readily converted to other functional groups, such as amides or amines, the C–CN activation approach can potentially be employed to streamline synthesis of nitrogen-containing molecules.<sup>5</sup>

## A. Carbon-Cyanide Bond Activation




## B. Carbon-Alkyne Bond Activation




**Scheme 1.** C–C Activation of Nitriles and Ynones

Considering that alkynes also have  $sp$  hybridized carbons like the cyano group, it would be impactful if the analogous activation of the carbon–alkyne bond could be realized (Scheme 1B). Alkynes have rich chemical reactivity and can serve as a latent functional group for alkenes, alkanes, ketones, diones, vicinal carbenes, etc.<sup>6</sup> Thus, transformations coupled with carbon–alkyne bond activation should be synthetically useful. However, in contrast to the C–CN bond, the carbon–alkyne bond is much less polarized. Consequently, only a few isolated cases on carbon–alkyne bond activation, i.e. oxidative addition of a transition metal into a carbon–alkyne bond, have been reported. One seminal example is C–C cleavage followed by decarbonylation of conjugated diyones with stoichiometric Wilkinson’s complex by Müller in 1969;<sup>7</sup> later, oxidative addition of rhodium(I) into a quinoline-derived acyl–alkyne bond was disclosed by Suggs in 1981.<sup>8</sup> Another example is photochemical cleavage of the aryl–alkyl bond in diarylalkynes with platinum(0) complexes.<sup>9</sup> To the best of our knowledge, it was not until our recent report that the catalytic transformation involving carbon–alkyne bond activation was realized.<sup>10</sup> Our laboratory has been particularly interested in developing catalytic transformations involving C–C activation of ketone compounds.<sup>11</sup> In the previous communication, we described an initial effort on catalytic decarbonylation of diyones to synthesize various 1,3-diyne (Scheme 2A).<sup>10</sup> Under the optimized conditions, both symmetrical and unsymmetrical diyones are suitable substrates, and a number of functional groups are tolerated. This C–C activation approach is complementary to transition metal-catalyzed cross couplings (e.g. compatibility with aryl bromides and iodides), and has been further applied to natural product derivatization.

## A. Previous Studies



## B. This Work

**Scheme 2.** Catalytic Decarbonylation of Conjugated Diynones and Monoynes via C–C Activation

With these preliminary results in hand, two key questions remained to be addressed: 1) *are both alkyne moieties required to maintain the catalytic activity for cleaving the carbon-alkyne bond*; 2) *if not (i.e. if only one alkynyl group is sufficient), in the absence of any auxiliary directing group, which C–C bond gets cleaved first for monoynes (Scheme 2B)?* Stimulated by these questions, we first describe a detailed development of a catalytic system that is effective for decarbonylation of conjugated monoynes, then disclose the reaction scope and limitation, and finally report our mechanistic exploration via DFT calculations. Through the computational efforts, we obtained a better understanding about the reaction mechanism, particularly about the rate-limiting step and which C–C bond is first activated. These efforts are expected to serve as an important exploratory study for developing catalytic alkyne-transfer reactions via carbon–alkyne bond activation.

## RESULTS AND DISCUSSION

## 1. Reaction Optimization

In 1969, Müller reported a single example that bisphenylynone **1a** reacted with one equivalent of Wilkinson's complex  $[\text{RhCl}(\text{PPh}_3)_3]$  in refluxing xylenes giving 8% yield of diphenylacetylene **2a**.<sup>7</sup> Although occurring with low efficiency, this seminal observation offered an opportunity to apply our knowledge of diyne activation into developing a catalytic decarbonylation of monoynes. However, under our previously optimized conditions (*vide supra*, Scheme 2A), 2.5 mol %  $[\text{Rh}(\text{COD})\text{Cl}]_2$  and 6 mol % dpf in refluxing chlorobenzene did not provide any decarbonylation product **2a**. This initial result was a clear indication of the difference in reactivity between diynones and monoynes for decarbonylation. The significantly reduced reactivity of monoynes, compared to diynones, can be possibly explained by the following: 1) the C–C bonds  $\alpha$  to the carbonyl of monoynes are more sterically demanding; and 2) the carbonyl group is also less electrophilic (reduced LUMO coefficient) than the one of diynones (both

factors would hinder oxidative addition). Clearly, to develop a catalytic decarbonylation of monoyrones, a more active catalyst system needed to be discovered.

**Table 1.** Selected Optimization Studies

Entry	Ligand (12 mol %)	Solvent	Bite Angle (°) <sup>b</sup>	Yield <sup>c</sup>
1	dppf	xylenes	96	24% (34%)
2	dppm	xylenes	72	<5%
3	dppe	xylenes	85	<5%
4	dppp	xylenes	91	13%
5	dppb	xylenes	98	29%
6	DPEphos	xylenes	104	11%
7	<i>t</i> -BuXphos	xylenes	-	43%
8	Xantphos	xylenes	111	85%
9	Xantphos	<i>m</i> -xylene	111	63% (71%)
10	Xantphos	<i>o</i> -xylene	111	29% (35%)
11	Xantphos	<i>p</i> -xylene	111	62%
12	Xantphos	ethylbenzene	111	91%

a) Conditions: ynone **1a** (0.20 mmol), [Rh]: Ligand=1:1.2, solvent (0.1 M). b) See ref 12 for bite-angle values. c) Isolated yields; number in parenthesis is percent conversion of starting material.

The optimization studies began with ynone **1a** as the model substrate (Table 1). Solvents with higher boiling points than chlorobenzene were examined first. When the reaction was run with 5 mol % [Rh(COD)Cl]<sub>2</sub> and 12 mol % dppf in refluxing xylenes (150-157 °C), we were pleased to find that the desired decarbonylation product **2a** was obtained in 24% yield (34% conversion of starting material, entry 1). With all other variables held constant, we surveyed a number of bidentate ligands with various bite angles, which were previously found to be important for decarbonylating diyrones.<sup>10,13</sup> Ligands, such as dppm, dppe, and dppp, with bite angles less than dppf (96°) showed trace or decreased yields (entries 2-4). On the other hand, bidentate ligands with larger bite angles or bulky monodentate ligands provided increased yields: while dppb slightly improved the yield (29%, entry 5), *t*-BuXphos and Xantphos<sup>13b</sup> gave improved yields (43% and 85%, entries 7 and 8, respectively). Unexpectedly, DPEphos gave a lower yield (11%, entry 6). Satisfied with Xantphos as the ligand, other reaction parameters were then explored. The commercially available xylenes contain a mixture of *m*-, *o*-, and *p*-isomers, as well as a small amount of ethylbenzene. Surprisingly, all *m*-, *o*-, and *p*-xylenes showed lower yields (29 – 63%, entries 9-11) than mixed xylenes; in contrast, ethylbenzene gave the highest yield (91%, entry 12). In addition, a series of Lewis acids, ruthenium co-catalysts and rhodium precatalysts were also examined, albeit with no improvement observed (for details, see Supporting Information Table S1).


## 2. Substrate Scope and Limitation

With a standard set of conditions in hand, the substrate scope of the reaction was explored. Keeping the alkyne moiety of the substrate fixed, a range of aryl substituted yrones were investigated under the decarbonylation conditions (Table 2). In general, good to high yields can be afforded with substrates (**2a-2k**) containing either electron-donating or withdrawing aryl groups, showing no obvious electronic bias. Interestingly, the 4-nitrophenyl substrate (**1h**) gave a higher yield (77%, **2h**) when using dppf as the ligand and xylenes as the solvent, compared to 40% yield (85% conversion of starting material) under the standard conditions. Functional groups, such as -F, -CN, -Cl, and -CO<sub>2</sub>Me, were also found compatible.

Substrates containing heterocyclic groups, such as furans (**1l**) and pyridines (**1m** and **1n**), also underwent decarbonylation smoothly, particularly the 3-pyridyl group which showed superior reactivity (**1n**).

An important observation is that this reaction is highly sensitive to the sterics around the carbonyl group. Substrates having substituents at the *ortho*-position (**2o-2r**) showed a dramatic decrease in yield, potentially hindering the substrate binding to the metal center. In addition, replacement of the aryl group with an alkenyl or alkyl substituent (**1s-1u**) resulted in no conversion to products (recovery of most of the starting materials).

**Table 2.** Substrate Scope Based on Ketone Substitution<sup>a</sup>




a) Reactions were run on a 0.20 mmol scale; all yields are isolated yields. b) Number in parenthesis is percent conversion of starting material. c) dppf and xylenes were used.

Substitution on the alkyne end of the substrates was also explored with the ketone end held constant as a phenyl group (Table 3). In general, both electron-donating and withdrawing aryl substituents were tolerated, giving synthetically useful yields (**4a-4d**, 60-73%). However, substrates containing a *para*-halogen substituent provided much lower yields (**4e-4g**), though the exact reason is unclear (*vide infra*, enhanced yields in Table 4). Furan (**3h**) and thiophene (**3i**) substrates also furnished the desired products, albeit in low yields. Certain alkyl substituents at the alkyne end were also tolerated (**3j**) and *vide infra*, ethynyl estradiol-derived ynone Scheme 3). However, *t*-Bu, trifluoromethyl, trimethylsilyl or linear alkyl substituents proved unreactive under the standard conditions. Under forcing conditions, i.e. in refluxing mesitylene (168-170 °C with all other parameters remaining the same), linear alkyl substrates (**3l** and **3o**) gave exclusive formation of the cycloisomerized furan products, which is likely though an alkyne-allene isomerization pathway (for details, see Supporting Information, Scheme


S1). Moreover, while cyclohexenyl ynone **3n** showed no reaction under the standard conditions, in refluxing mesitylene the decarbonylation product **4n** was able to form in 14% yield.<sup>14</sup>

**Table 3.** Substrate Scope Based on Phenyl Ynone<sup>a</sup>




a) Reactions were run on a 0.2 mmol scale; unless otherwise mentioned, all yields are isolated yields. b) Number in parenthesis is percent conversion of starting material. c) Product **4j** is slightly volatile. d) The reaction was run in mesitylene at 170°C. e) The yield is based on <sup>1</sup>H NMR using C<sub>2</sub>H<sub>2</sub>Cl<sub>4</sub> as the internal standard.

In contrast, when 3-pyridyl was used as the acyl substituent, the reactivity of the ynone substrates was greatly increased (Table 4). We were pleased to observe that a range of pyridine-containing disubstituted alkynes were isolated with enhanced conversions, and many functional groups were tolerated. Notably, the yields for the substrates containing halogen and heterocycles were significantly improved (12% for **4e** vs 56% for **6f**, and 24% for **4h** vs 38% for **6h**). In addition, the scope for the alkenyl and alkyl substituted substrates were expanded. Though straight alkyl ynone remain problematic giving allene isomerization, to our delight, branched alkyl substrates (**5i** – **k**) were found to be reactive and afforded products (**6i** – **k**) in modest to good yield (26 – 45%).

**Table 4.** Substrate Scope Based on 3-Pyridyl Ynone<sup>a</sup>

a) Reactions were run on a 0.2 mmol scale; all yields are isolated yields. b) Number in parenthesis is percent conversion of starting material.

The monoynone decarbonylation reaction has been further investigated in the derivatization of natural products (Scheme 3). For example, the ethynyl estradiol and myrtenal derived monoyones (**8** and **11**) smoothly gave the corresponding decarbonylated products **9** and **12** in 55% and 57% yields, respectively. Note that the aryl groups coupled with the natural products ultimately come from the corresponding carboxylic acids.


**Scheme 3.** Applications in Natural-Product Derivatization

With a thorough exploration of the reaction scope and a better understanding of substrate reactivity, we finally examined substrates that can undergo multiple decarbonylations. When terephthalic acid-derived di-ynone **13** was subjected to the standard conditions, the doubly decarbonylated product (**14**)




was obtained albeit in low yield along with severe decomposition to unidentified oligomers (Scheme 4). After further examining the reaction conditions, we found that use of lower concentrations can dramatically minimize the product decomposition to unidentified oligomers. Finally, with an increase of the catalyst loading at 0.05 M, the double decarbonylation product can be obtained in 94% yield.<sup>15</sup> Additionally, when trimesic acid-derived tri-ynone **15** was subjected to the above-optimized conditions, the tri-yne product **16** was isolated in 74% yield.

#### "Double" Decarbonylation




#### "Triple" Decarbonylation



**Scheme 4.** Multiple Decarbonylations

### 3. Mechanistic Studies via DFT Calculation

Our proposed mechanism of the Rh-catalyzed decarbonylation involves four steps: ligand substitution, oxidative addition, decarbonylation, and reductive elimination (Figure 1). The initial step involves substrate coordination to the Rh(I) through the alkynyl group, giving complex **I** (step 1). The second step is oxidative addition, leading to Rh(III) complex **IIA** (rhodium is inserted into **bond a** between the alkynyl and carbonyl groups of the substrate, pathway *a*) or **IIB** (rhodium is inserted into **bond b** between the aryl and carbonyl groups of the substrate, pathway *b*). Decarbonylation transforms **IIA** or **IIB** into intermediate **III**, which then undergoes reductive elimination to give the final product (step 4). Herein, we report density functional theory (DFT) calculations, to support this proposal and gain a better understanding of the mechanistic details.




**Figure 1.** The Proposed Mechanism of the Rh-catalyzed Decarbonylation of Monoyones.

DFT calculations were based on the model reaction of ynone **1a** to alkyne **2a**. The full model of the best ligand, Xantphos, was used for the DFT studies. The energy profiles of paths *a* and *b* were shown in Figure 2. The discussed energies here are the relative free energies in the gas phase, considering that the conclusions extracted from the gas phase and solvent are the same (see the DFT computed values in the parentheses in Figure 2 for the relative free energies of the reaction in ethylbenzene solvent).

First, we discuss the energy surface of pathway *a* (Figure 2). The catalytic cycle starts from ligand exchange reaction between **CAT-P** and substrate **1a**, giving catalyst-substrate complex **INT1** and releasing the decarbonylation product **2a**. Substrate **1a** could coordinate to the Rh center through either the alkyne group or the carbonyl group. DFT calculations indicate that the alkyne-coordinated complex is more stable than the carbonyl-coordinated complex by 7.1 kcal/mol and therefore formation of **INT1** is preferred. **INT1** then undergoes oxidative addition into **bond a** (pathway *a*) via **TS1-A**, requiring an activation free energy of 23.9 kcal/mol. This step is endergonic by 8.4 kcal/mol and generates **INT2-A**. A reversible decarbonylation via **TS2-A** subsequently transforms **INT2-A** to **INT3-A**, requiring an activation free energy of 14.4 kcal/mol. The decarbonylation step is endergonic by 6.6 kcal/mol. Subsequently, ligand reorganization converts **INT3-A** to **INT4-A**, which undergoes reductive elimination to give to **INT5** (via **TS3-A**).<sup>16</sup> The final reductive elimination step has an activation free energy of 10.5 kcal/mol and is irreversible (it is exergonic by 29.9 kcal/mol). Our calculations indicated that in pathway *a*, the rate-determining step of the catalytic cycle is the reductive elimination step and the overall activation free energy of the catalytic cycle is 28.8 kcal/mol in gas phase. Using ethylbenzene as the solvent, the computed overall activation free energy is 30.8 kcal/mol.<sup>17</sup> *The calculation results here reasonably explain why experimentally the decarbonylation reaction had to be carried out at 150 °C.*


An alternative pathway is rhodium insertion (from **INT1**) into **bond b** (**INT2-B**, between the carbonyl and aryl groups (pathway *b*)), which is disfavored by more than 20 kcal/mol compared to the insertion into **bond a** in pathway *a*. The computed activation energy barrier for this step is 45.7 kcal/mol, which is much higher than the total activation energy in pathway *a*. Consequently, pathway *b* can be excluded

from further consideration. To rationalize the above observation, we propose that the regioselectivity of the C–C bond cleavage can be controlled by a *trans* effect (TE), also known as *trans* influence when considering the ground state of the complex.<sup>18</sup> The intermediate (INT-2A or INT-2B) after the oxidative addition step should contain three X-ligands: the acyl, phenyl, and acetylide. Acyl and phenyl are very strong TE  $\sigma$ -donor ligands, while acetylide ligand is a weak TE ligand (weaker than phosphine). Cleavage *b* bond will generate two strong TE ligands: the acyl and phenyl ligands. In this case, the chloride ligand (a moderately strong TE ligand) has to be in a *trans* position to either the acyl ligand or phenyl ligand, which is not favored based on the TE.<sup>18</sup> In contrast, cleavage of the *a* bond will generate one strong TE ligand, the acyl ligand, and one weak TE ligand, the acetylide ligand. In this case, the strong TE ligand (i.e. the acyl group) can be arranged to a position that is *trans* to the oxygen of the Xantphos ligand to reduce the TE, while the weak TE ligand (i.e. the acetylide group) can be *trans* to the chloride (the geometry rearrangement is illustrated in TS1-A).




**Figure 2.** Comparison of the energy profiles ( $\Delta G$  in kcal/mol) of path *a* (black) and path *b* (red) for decarbonylation of ynone **1a**, and DFT optimized structures of transition states and key intermediates (distances in Å, hydrogen atoms and phenyl groups of Xantphos were omitted for clarity).

Experimentally, we found that replacement of the aryl group with an alkyl substituent (such as methyl group, **2t**) resulted in no conversion to product (Table 2). DFT studies on the substituent effect between phenyl substrate **1a** and methyl substrate **1t** have been performed (Figure 4). The rate-determining step of **1t** is also the reductive elimination step, but the overall activation free energy for the decarbonylation is 34.2 kcal/mol, which is 5.4 kcal/mol higher than that of **1a**. Due to this reason, the reaction of **1t** did not occur under the experimental conditions that are suitable for **1a**.



**Figure 3.** Comparison of the energy profiles ( $\Delta G$  of gas phase in kcal/mol) of **1a** (black) and **1t** (red).

The higher activation free energy of **1t** compared to that of **1a** is mainly caused by the more difficult reductive elimination step in the former case. In **1t**, the reductive elimination has an energy barrier of 16.7 kcal/mol, which is 6.2 kcal/mol higher than that of **1a** (10.5 kcal/mol). This result is consistent with our previously observed faster reductive elimination with a C(sp<sup>2</sup>) group than a C(sp<sup>3</sup>) group through DFT calculations.<sup>19</sup> What is the intrinsic reason for this difference? Here is our proposed explanation. Although the Rh-phenyl bond in **INT3-A** has a higher energy than the Rh-methyl bond in **INT3-1t** (our calculated results, Figure 4), in the transition state of the reductive elimination step the migrating carbon in the phenyl group is four-coordinated and the charge in this phenyl group can be well distributed into the aromatic ring (**TS3-A**). In contrast, the migrating carbon in the methyl group (**TS3-1t**) is energetically disfavored (five-coordinated), and this requires additional energy compared to the four-coordinated phenyl group in **TS3-A**.



**Figure 4.** The computed BDEs of **INT3-A** and **INT3-1t** and the comparison of two reductive elimination transition states. The ligand here is Xantphos.

Our DFT calculations also found that the dppp ligand is not effective for the present decarbonylation reaction. This is mainly due to a disfavored reductive elimination step (even though the C–C cleavage step is not difficult). For the computed energy surface, see the Supporting information.

## CONCLUSION

In conclusion, we have reported a detailed experimental and computational study of the Rh-catalyzed decarbonylation of conjugated monoyne, which should serve as an initiative towards developing catalytic alkyne transfer reactions via carbon–alkyne bond activation (a long-term goal). We have discovered an active catalytic system that is suitable for a range of substrates and functional groups stemming from readily available carboxylic acids. From the experimental study, the scope and limitation of this transformation have been thoroughly explored. From the computational study, a proposed catalytic mechanism has been carefully evaluated. Employing the theoretic models, we now have a better understanding about how the C–C bond in monoyne is activated, ruling out the pathway involving initial cleavage of the aryl-carbonyl bond and favoring cleavage of the alkynyl-carbonyl bond. In addition, the calculation results support that reductive elimination is the rate-determining step for the catalytic cycle. Furthermore, we obtained key information about why the aryl ketones are more reactive than the corresponding alkyl ketones.

With all the mechanistic information of the ynone decarbonylation in hand, further investigations to discover the alkyne-transfer transformations (analogous to the CN transfer reactions<sup>6</sup>) are currently underway in our laboratories.

## COMPUTATIONAL DETAILS

All calculations were performed with the Gaussian 09 program.<sup>20</sup> Density functional theory calculations using the B3LYP method were used to locate all the minima and transition points involved.<sup>21</sup> The 6-31G(d) basis set<sup>22</sup> was applied for all elements except for Rh, for which the LANL2DZ basis set and pseudopotential<sup>23</sup> were used. The key word “5D” in Gaussian 09 program was used. Frequency calculations at the same level had been performed to confirm each stationary point to be either a

minimum or a transition structure and to evaluate its zero-point energy and the thermal corrections at 298 K. Both single-point energies and solvation energies based on the geometry structures obtained at the B3LYP level were obtained by M06L method<sup>24</sup> using a higher level basis set, LANL2TZ(f) basis set and pseudopotential<sup>25</sup> for Rh and 6-311+G(d,p) basis set for all the other atoms in order to take the dispersion energies into consideration. Solvation energies in ethylbenzene were evaluated by a self-consistent reaction field (SCRF) using the SMD model with radii and non-electrostatic terms. Bond dissociation energies (BDE) are discussed as bond-homolysis into two radicals in B3LYP/6-31G(d) level in gas phase. In the paper and the Supporting Information, all discussed energies are Gibbs free energies in gas phase ( $\Delta G_{\text{gas}}$ ) at 298 K unless specified. We found that the conclusions in both the gas phase and ethylbenzene are the same. Computed structures are illustrated using CYLVIEW drawings.<sup>26</sup>

## ACKNOWLEDGMENT

We thank UT Austin and CPRIT for a startup fund, NIGMS (R01GM109054-01) and the Welch Foundation (F 1781) for research grants. GD is a Searle Scholar. ZXY thanks the Natural Science Foundation of China (21232001) for final support. We acknowledge Johnson Matthey for a generous donation of Rh salts.

## ABBREVIATIONS

COD: cyclooctadiene; COE: cyclooctene; dppf: 1,1'-bis(diphenylphosphino)ferrocene; dppp: 1,3-bis(diphenylphosphino)propane; dppe: 1,2-bis(diphenylphosphino)ethane; dppb: 1,3-bis(diphenylphosphino)butane; Xantphos: 4,5-Bis(diphenylphosphino)-9,9-dimethylxanthene; DPEphos: (oxydi-2,1-phenylene)bis(diphenylphosphine); dpmm: 1,1-bis(diphenylphosphino)methane; *t*-BuXphos: 2-di-*tert*-butylphosphino-2',4'6'-triisopropylbiphenyl.

## NOTES & REFERENCES

- <sup>1</sup> For selected reviews of C–C activation, see: (a) M. Murakami, Y. Ito, *Top Organomet. Chem.*, 1999, **3**, 97-129; (b) C.H. Jun, *Chem. Soc. Rev.*, 2004, **33**, 610-618; (c) T. Kondo, T.A. Mitsudo, *Chem. Lett.*, 2005, **34**, 1462-1467; (d) B. Rybtchiski, D. Milstein, *Angew. Chem., Int. Ed.*, 1999, **38**, 870-883; (e) M.E. van der Boom, D. Milstein, *Chem. Rev.*, 2003, **103**, 1759-1792; (f) T. Satoh, M. Miura, *Top. Organomet. Chem.*, 2005, **14**, 1-20; (g) C.H. Jun, J.W. Park, *Top. Organomet. Chem.*, 2007, **24**, 117-143; (h) D. Necas, M. Kotora, *Curr. Org. Chem.*, 2007, **11**, 1566-1591; (i) R.H. Crabtree, *Chem. Rev.*, 1985, **85**, 245-269; (j) M. Gozin, A. Weisman, Y. Ben-David, D. Milstein, *Nature*, 1993, **364**, 699-701; (k) K. Ruhland, *Eur. J. Org. Chem.*, 2012, 2683-2706; (l) A. Korotvicka, D. Necas, M. Kotora, *Curr. Org. Chem.*, 2012, **16**, 1170-1214; (m) T. Seiser, T. Saget, D.N. Tran, N. Cramer, *Angew Chem., Int. Ed.* 2011, **50**, 7740-7742; (n) L. Jiao, Z.-X. Yu, *J. Org. Chem.*, 2013, **78**, 6842-6848; (o) A. Dermenci, J.W. Coe, G. Dong, *Org. Chem. Frontiers*, 2014, **1**, 567-581; (p) A. Dermenci, G. Dong, *Sci. China, Chem.*, 2013, 685-701.
- <sup>2</sup> For a book volume, see: *C–C Bond Activation; Topics in Current Chemistry*, v. 346, ed. G. Dong, Springer, Berlin, Heidelberg, 2014.
- <sup>3</sup> For reviews of a novel strategy that uses a temporary directing group, see: refs 1b, 1g and C.H. Jun, J.W. Park, *Topics in Current Chemistry*, 2014, **346**, 59-83.
- <sup>4</sup> For recent reviews of C–CN activation, see: (a) Y. Nakao, *Topics in Current Chemistry*, 2014, **346**, 33-58; (b) Y. Nakao, T. Hiyama, *Pure Appl. Chem.*, 2008, **80**, 1097-1107; (c) M. Tobisu, N. Chatani, *Chem. Soc. Rev.*, 2008, **37**, 300-307; (d) S.M. Bonesi, M. Fagnoni, *Chem. Eur. J.*, 2010, **16**, 13572-13589.
- <sup>5</sup> (a) Y. Nakao, A. Yada, S. Ebata, T. Hiyama, *J. Am. Chem. Soc.*, 2007, **129**, 2428-2429; (b) Y. Nakao, T. Hiyama, *Pure Appl. Chem.*, 2008, **80**, 1097-1107; (c) Y. Hirata, T. Yukava, N. Kashihara, Y. Nakao, T. Hiyama, *J. Am. Chem. Soc.*, 2009, **131**, 10964-10973; (d) C. Najera, J.M. Sansano, *Angew. Chem., Int. Ed.*, 2009, **48**, 2452-2456; (e) Y. Nakao,

- A. Yada, S. Ebata, T. Hiyama, *J. Am. Chem. Soc.*, 2010, **132**, 10024-10026. For examples of arylcyanation, see: (f) M.P. Watson, E.N. Jacobsen, *J. Am. Chem. Soc.*, 2008, **130**, 12594-12595; (g) Y. Nakao, S. Ebata, A. Yada, T. Hiyama, M. Ikawa, S. Ogoshi, *J. Am. Chem. Soc.*, 2008, **130**, 12874-12875. For examples of cyanoamidation, see: (h) Y. Kobayashi, H. Kamiisaki, H. Takeda, Y. Yasui, R. Yanada, Y. Takemoto, *Tetrahedron*, 2007, **63**, 2978-2989; (i) Y. Kobayashi, H. Kamisaki, R. Yanada, Y. Takemoto, *Org. Lett.*, 2006, **8**, 2711-2713; (j) Y. Yasui, H. Kamisaki, T. Ishida, Y. Takemoto, *Tetrahedron*, 2010, **66**, 1980-1989; (k) Y. Yasui, H. Kamisaki, Y. Takemoto, *Org. Lett.*, 2008, **10**, 3303-3306; (l) V.J. Reddy, C.J. Douglas, *Tetrahedron*, 2010, **66**, 4719-4729; (m) V.J. Reddy, C.J. Douglas, *Org. Lett.*, 2010, **12**, 952-955. For an example of cyanoesterification, see: (n) N.R. Rondla, S.M. Levi, J.M. Ryss, R.A. Vanden Berg, C.J. Douglas, *Org. Lett.*, 2011, **13**, 1940-1943.
- <sup>6</sup> For representative reviews on alkyne transformations and importance, see: (a) *Acetylene Chemistry*, eds. F. Diederich, P.J. Stang, R.R. Tykwinski, Wiley-VCH, Weinheim, 2006; (b) *Chemistry of Acetylenes*, eds. P. Cadiot, W. Chodkiewicz, Marcel Dekker, New York, 1969; (c) R. Chinchilla, C. Najera, *Chem. Rev.*, 2014, **114**, 1783-1826. (d) R.S. Atkinson, *Aliphatic Chem.*, 1974, **2**, 3-127.
- <sup>7</sup> For stoichiometric decarbonylation of ynones, see: (a) E. Muller, A. Segnitz, E. Langer, *Tetrahedron Lett.*, 1969, 1129-1131; (b) E. Muller, A. Stegnitz, *Liebigs Ann. Chem.*, 1973, 1583-1591.
- <sup>8</sup> J.W. Suggs, S. D. Cox, *J. Organomet. Chem.*, 1981, **221**, 199-201.
- <sup>9</sup> (a) G. K. Anderson, G. J. Lumetta, J. W. Siria, *J. Organomet. Chem.*, 1992, **434**, 253-259; (b) C. Müller, C.N. Iverson, R. J. Lachicotte, W. D. Jones, *J. Am. Chem. Soc.*, 2001, **123**, 9718-9719.
- <sup>10</sup> A. Dermenci, R. E. Whittaker, G. Dong, *Org. Lett.*, 2013, **15**, 2242-2245.
- <sup>11</sup> For our recent catalytic activation of strained ketones, see: (a) T. Xu, G. Dong, *Angew. Chem., Int. Ed.*, 2012, **51**, 7567-7571; (b) T. Xu, H. M. Ko, N. A. Savage, G. Dong, *J. Am. Chem. Soc.*, 2012, **134**, 20005-20008; (c) H.M. Ko, G. Dong, *Nature Chem.*, 2014, **6**, 739-744; (d) T. Xu, G. Dong, *Angew. Chem. Int. Ed.*, 2014, **53**, 10733-10736; (e) T. Xu, N. A. Savage, G. Dong, *Angew. Chem. Int. Ed.*, 2014, **53**, 1891-1895; (f) P. Chen, T. Xu, G. Dong, *Angew. Chem. Int. Ed.*, 2014, **53**, 1674-1678.
- <sup>12</sup> For average ligand bite angles, see: (a) P. Dierkes, P. W. N. M. van Leeuwen, *J. Chem. Soc., Dalton Trans.*, 1999, 1519-1530; (b) P. W. N. M. van Leeuwen, P. D. J. Kamer, J. N. H. Reek, P. Dierkes, *Chem. Rev.*, 2000, **100**, 2741-2769.
- <sup>13</sup> (a) For seminal work of bidentate ligand-assisted decarbonylation of aldehydes, see: D. H. Dougherty, L. H. Pignolet, *J. Am. Chem. Soc.*, 1978, **100**, 7083-7085. (b) For a recent seminal work on Xantphos-Rh-catalyzed transfer hydroformylation of aldehydes, see: S. K. Murphy, J.-W. Park, F. A. Cruz, V. M. Dong, *Science*, 2015, **2**, 56-60.
- <sup>14</sup> Only NMR yield was obtained for substrate **3o** by comparing with an authentic sample, due to the difficulty of its purification.
- <sup>15</sup> For an example of increased catalyst loading enhancing multiple decarbonylations of phenylglyoxal hydrate, see: A. Modak, A. Deb, T. Patra, S. Rana, S. Maity, D. Maiti, *Chem. Commun.*, 2012, **48**, 4253-4255.
- <sup>16</sup> DFT calculations found that reductive elimination from **INT3-A** is not favored than that from intermediate **INT4-A** (35.7 vs. 28.8 kcal/mol in gas phase, please see Supporting Information). In addition, there is another possible pathway for CO dissociation first, followed by reductive elimination. This possibility can be ruled out because the activation barrier here is as high as 47.9 kcal/mol in the gas phase (please see Supporting Information).
- <sup>17</sup> The regeneration of **INT1-A** starts from loss of CO from **INT5-A**, giving **CAT-2a**, which is then transformed to **INT1-A** through ligand exchange with the incoming substrate for the next catalytic cycle. This CO release step can be considered irreversible since the CO gas is purged out during the reaction. Therefore, the required overall activation energy for the present reaction is the difference between **INT1-A** and **TS3-A**. This could explain why the target decarbonylation reaction cannot be performed under pure CO pressure. In this case, regeneration of **INT1-A** need additional energy, due to the added CO coordination to **CAT-2a**.
- <sup>18</sup> B. J. Coe, S. J. Glenwright, *Coord. Chem. Rev.*, 2000, **203**, 5-80.
- <sup>19</sup> (a) Z.-X. Yu, P. A. Wender, K. N. Houk, *J. Am. Chem. Soc.*, 2004, **126**, 9154-9155; (b) Y. Wang, J. Wang, J. Su, F. Huang, L. Jiao, Y. Liang, D. Yang, S. Zhang, P.A. Wender, Z.-X. Yu, *J. Am. Chem. Soc.*, 2007, **129**, 10060-10061; (c) Z.-K. Yao, J. Li, Z.-X. Yu, *Org. Lett.*, 2010, **13**, 134-137; (d) L. Jiao, M. Lin, Z.-X. Yu, *J. Am. Chem. Soc.*, 2011, **133**, 447-461; (e) W. Liao, Z.-X. Yu, *J. Org. Chem.*, 2014, **79**, 11949-11960.
- <sup>20</sup> Full citation of Gaussian 09: Gaussian 09, Revision D.01, M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X.

Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, O. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski, D. J. Fox, Gaussian, Inc., Wallingford Connecticut, 2013.

<sup>21</sup> (a) A. D. Becke, *J. Chem. Phys.*, 1993, **98**, 5648-5652; (b) C. Lee, W. Yang, R. G. Parr, *Phys. Rev. B*, 1988, **37**, 785-789.

<sup>22</sup> *Ab Initio Molecular Orbital Theory*, eds. W. J. Hehre, L. Radom, P. V. R. Schleyer, J. A. Pople, Wiley, New York, 1986.

<sup>23</sup> P. J. Hay, W. R. Wadt, *J. Chem. Phys.*, 1985, **82**, 299-316.

<sup>24</sup> Y. Zhao, D. G. Truhlar, *J. Chem. Phys.*, 2006, **125**, 194101-194119.

<sup>25</sup> (a) P. J. Hay, W. R. Wadt, *J. Chem. Phys.*, 1985, **82**, 270-283; (b) P. J. Hay, W. R. Wadt, *J. Chem. Phys.*, 1985, **82**, 284-298; (c) P. J. Hay, W. R. Wadt, *J. Chem. Phys.*, 1985, **82**, 299-316.

<sup>26</sup> C. Y. Legault, CYLView, version 1.0b; Université de Sherbrooke, Sherbrooke, Quebec, Canada, 2009; <http://www.cylview.org>.

#### TOC:

