

Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

ARTICLE

Access to Enantioenriched 2,3- and 2,5-Dihydrofurans with a Fully Substituted C2 Stereocenter by Pd-Catalyzed Asymmetric Intermolecular Heck Reaction

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

G. M. Borrajo Calleja,^a V. Bizet,^a T. Bürgi^b and C. Mazet^{*a}

A palladium catalyzed intermolecular asymmetric Heck reaction with dihydrofurans with a trisubstituted double bond is reported. The use of two different chiral ligands provides access to valuable 2,3- and 2,5-dihydrofurans with a fully substituted C2 stereocenter with high levels of regio- and enantiocontrol.

Introduction

During the past decades, building on the seminal contributions of Overman and Shibasaki, the asymmetric intramolecular Heck reaction has been successfully applied in natural product synthesis to install tertiary and quaternary stereocenters.¹⁻² In stark contrast, the *intermolecular* version of this asymmetric transformation has not reached the same level of development and its synthetic utility remains limited (Figure 1).³ A survey of the literature following Hayashi's pioneering contribution indicates that this transformation has been used essentially as a benchmark reaction to validate the design of novel homotopic and heterotopic chiral ligands.⁴⁻⁵ While variations of the electronic and steric nature of the aryl halides and pseudo-halides has been investigated in details, most of these studies have been focused on the use of dihydrofurans or *N*-protected dihydropyrroles to afford the corresponding coupling products with a C2 tertiary center. Systematic studies where emphasis has been placed on exploring the diversity of the olefinic coupling partner are scarce, in particular for the construction of congested quaternary centers or related fully saturated centers. Very recently, Sigman has reported a series of elegant contributions for the redox-relay Heck arylations of acyclic alkenyl alcohols enabling the installation of remote tertiary and quaternary stereogenic centers.⁶⁻⁷ Despite these recent breakthroughs, it is apparent that the development of an intermolecular asymmetric Heck reaction to enable the construction of *fully saturated stereocenters* in *cyclic* systems constitutes an important synthetic challenge.⁸

In this report, we disclose a methodology that gives access to chiral 2,3- and 2,5-dihydrofurans (dhf) with a fully substituted C2 stereocenter. Our study highlights the


complementarity in product selectivity between two distinct chiral ligands while it also delineates the scope of trisubstituted dihydrofurans compatible with the catalytic systems.

♦ Intramolecular (representative example)




- ♦ extensively investigated
- ♦ high ee
- ♦ numerous applications in synthesis

♦ Intermolecular: acyclic substrates (redox-relay)



- ♦ recently investigated
- ♦ high ee
- ♦ substrate diversity

♦ Intermolecular: cyclic substrates



- R = H ♦ benchmark reaction
- ♦ good regioselectivity
- ♦ high ee
- R ≠ H ♦ 1 isolated example
- ♦ high ee
- ♦ poor reactivity
- ♦ low regioselectivity

Fig. 1 Access to quaternary and related fully saturated stereocenters by asymmetric Heck reactions.

Results and discussion

The choice of our model reaction was inspired by a report from Pregosin and co-workers where 5-methyl-2,3-dihydrofuran **1a** was cross-coupled with phenyl triflate **2a** using (*R*)-DTB-MeO-BIPHEP (*i.e.* (*R*)-(+)-2,2'-bis[di(3,5-di-*t*-butylphenyl)phosphino]-6,6'-dimethoxy-1,1'-biphenyl).⁹ *To the best of our knowledge this represents the only precedent of an asymmetric intermolecular Heck reaction employing a cyclic substrate with a trisubstituted olefinic moiety.* Despite an excellent enantioselectivity (**4aa**: 98% ee), the low yield (38%), the moderate regioselectivity (1.0:6.0) and the impractical

reaction time (7 days) certainly precluded further developments. We began our investigations by evaluating an array of chiral (P,N) and (P,P) ligands as their complementarity in providing access to the product of direct cross-coupling **3** and the product of further isomerization **4** respectively is well documented for unsubstituted cyclic substrates.^{4-5,10}

Table 1 Reaction optimization^a

Entry ^b	2	Ligand	3:4 ^c	3		4	
				Yield (%) ^d	ee (%) ^e	Yield (%) ^d	ee (%) ^e
1	2a	L1	22:78	<5	nd ^f	5	9
2	2a	L3	99:1	12	76	–	–
3	2a	L4	99:1	16	94	–	–
4	2b	L4	99:1	11	92	–	–
5	2b	L4	99:1	64	93	–	–
6	2c	L4	–	nr ^g	–	nr	–
7	2b	L5	98:2	34	98	–	–
8	2c	L5	–	nr	–	nr	–
9	2b	L2	44:56	21	rac	28	13
10	2c	L2	5:95	–	–	55	93
11	2b	L6	5:95	–	–	30	56
12	2c	L6	5:95	–	–	32	62
13	2b	L7	5:95	–	–	39	45
14	2c	L7	5:95	–	–	34	51
15	2b	L8	57:43	<5	–	<5	–
16	2c	L8	5:95	–	–	51	97

^a Reaction conditions: **1a** (1 mmol), **2a-c** (0.2 mmol). ^b entry 1-4: Pd(OAc)₂ (3 mol%), ligand (6 mol%), toluene, 110 °C, 62 h. entry 5-16: Pd₂(dba)₃ (2.5 mol%), ligand (10 mol%), 2-Me-THF, 100 °C, 48 h. ^c Determined by ¹H NMR of the crude reaction mixture. ^d Yield of pure compound after chromatography. ^e Determined by GC or HPLC with a chiral column. ^f Not determined. ^g No reaction.

With an initial set of reaction conditions (Pd(OAc)₂, Toluene, *i*-Pr₂NEt, 110 °C, 62 h), only traces of products **3aa** and **4aa** were observed in the coupling between **1a** and **2a** using **L1** (Table 1, entry 1). Evaluation of two of our home-made chiral

(P,N) ligands did not significantly enhanced the reactivity, even though **3aa** was the only detectable product of cross-coupling.¹¹ While **3aa** was obtained in 76% *ee* with **L3**, **L4** gave an excellent 94% *ee* value (entry 2-3). A slightly improved reactivity along with similar regio- and enantioselectivities were obtained with the electron-rich 4-methoxy phenyl triflate **2b** (entry 4). An extensive optimization of all reaction parameters was conducted next. Multiple combinations of palladium sources, solvents and bases along with adjustment of the temperature, stoichiometry, concentration and reaction time led to the identification of a very efficient catalytic system (Pd₂(dba)₃, 2-Me-THF, *i*-Pr₂NEt, 100 °C, 48 h; see SI for details) affording **3ab** in 64% and 93% *ee* with **L4** (entry 5).

Table 2 Scope of aryl triflates^a

♦ Scope of 2,5-dhf - using L4

3aa	3ab	3ad	3ae
27%, 99:1 s 92% ee (R)	64%, 99:1 s 93% ee (R)	47%, 99:1 s 92% ee (R)	65%, 99:1 s 89% ee (R)
3af^b	3ag^b	3ah	3aj^b
48%, 99:1 s 88% ee (R)	37%, 98:2 s 91% ee (R)	56%, 99:1 s 90% ee (R)	54%, 99:1 s 88% ee (R)
3aj^{c,d}	3ak	3al	
50%, 99:1 s 88% ee (R)	53%, 99:1 s 90% ee (R)	25%, 99:1 s 92% ee (R)	

♦ Scope of 2,3-dhf - using L8

4ac	4am	4an	4ao
59%, 95:5 s 97% ee (R)	51%, 95:5 s 98% ee (R)	54%, 94:6 s 98% ee (R)	35%, 96:4 s 97% ee (R)
4ap	4aq^f	4ar	
40%, 73:27 s 92% ee (R)	45%, 86:14 s 72% ee (R)	39%, 91:9 s 93% ee (R)	


^a Reaction conditions: **1a** (1 mmol), **2a-c** (0.2 mmol). Absolute configurations determined by VCD. ^b 3 days. ^c 4 days. ^d 95% purity.

Of important note, while the catalyst loading, the relative stoichiometry between the coupling partners and the yields

obtained (*ca.* 40–65%) are typical of *intermolecular* asymmetric Heck reactions, 48 h is unusually short for such a process involving cyclic olefins. Under classical thermal conditions, reaction times often range from 4 to 7 days.¹² Unexpectedly, when the electron-deficient aryl triflate **2c** was employed, no product of cross-coupling was observed.¹³ The prototypical (P,N) ligand **L5** for intermolecular Heck reactions using unsubstituted dhf did not display better performances (entry 7-8). For sake of operational simplicity, the same protocol was used in the survey of several (P,P) ligands (entry 9-16). Consistent with literature precedents for the coupling of unsubstituted dhf, a marked regioselectivity switch in favor of the isomerized products **4** was observed with most ligands. Noticeably, while poor results were obtained in the coupling reactions with the electron-rich triflate **2b**, the electron-deficient aryl triflate **2c** permitted to reach excellent enantioselectivity levels with **L2**, and **L8** (entry 10, 16).¹⁴

This dichotomous situation was further confirmed when the scope or aryl triflates compatible with both **L4** and **L8** was delineated (Table 2). With the (P,N) ligand **L4**, a wide variety of electron-rich aryl triflates afforded quasi-exclusively 2,5-dihydrofurans (98:2-99:1 *s*) in consistently very high enantioselectivity (88-93% *ee*) and in practical yields. Remarkably, even the sterically demanding *o*-methoxy phenyl triflate **2k** was cross-coupled efficiently (**3ak**: 53%, 99:1, 90% *ee*). Reduced yields were obtained with **2a** and **2l**, albeit the regio- and enantioselectivities remained excellent. Attempts to couple electron-deficient aryl triflates were unsuccessful. In contrast, with the (P,P) ligand **L8**, only electron-deficient aryl triflates **2c**, **2m-r** were found to be compatible electrophilic coupling partners. This afforded 2,3-dihydrofurans with a saturated C2 stereocenter in excellent enantioselectivity and allowed functional groups such as cyano, nitro, trifluoromethyl, ester, aldehyde, aryl and alkyl ketones to be introduced.


Table 3 Scope of 2,2-disubstituted 2,5-dhf with **L4**^a



^a Reaction conditions: **1a** (1 mmol), **2a-c** (0.2 mmol). Absolute configurations determined by VCD. ^b 3 days. ^c 4 days. ^d 95% purity.

Seven different 5-substituted 2,3-dhf were found to be adequate substrates with both methodologies (Table 3 and 4). These compounds proved particularly unstable and presented challenges for study because of the difficulty associated with their preparation and purification (see SI).

Table 4 Scope of 2,2-disubstituted 2,3-dhf with **L8**^a



^a Reaction conditions: **1a** (1 mmol), **2a-c** (0.2 mmol). Absolute configurations determined by VCD. ^b 3 days. ^c 4 days. ^d 95% purity.

The C_1 -symmetric (P,N) ligand **L4** and the C_2 -symmetric (P,P) ligand **L8** gave access respectively to 2,2-disubstituted 2,5-dhf and 2,2-disubstituted 2,3-dhf in usually practical yields and with excellent regio- and enantioselectivity. Both catalytic systems behaved similarly and overall excellent results have been obtained. The yield range obtained for these reactions are consistent with literature precedents.^{2-3,6} They do not only reflect the entropic cost inherent to an intermolecular cross-coupling reaction but also the difficulty associated with creating saturated stereocenters. In addition to primary alkyl substituents, a methyl ether, a free alcohol, a remote olefin and a benzyl substituent were compatible. Whereas the yields were slightly reduced in the reactions with **3e**, it is remarkable that no competing Heck arylation was observed in the reactions with **1f** which bears a remote olefinic moiety.⁶ Finally, a large scale experiment (8.5 mmol) was conducted for the cross-coupling between **1a** and **2c** (Figure 2). Using (*R*)-**L2**, the isomerized product (*R*)-**4ac** was isolated with excellent regioselectivity, good enantioselectivity and in 46 % yield (0.725 g).¹⁵




Fig. 2 Large scale experiment for the intermolecular asymmetric Heck reaction.

The absolute configuration of the cross-coupling products was established by vibrational circular dichroism (VCD).¹⁶ The most stable conformer of **3ab** (obtained with (*R,R,R*)-**L4**) and **4ap** (obtained with (*R*)-**L8**) were compared to the experimental IR absorption and VCD spectra recorded in CD₂Cl₂ solutions (see SI for details). The good agreement between the predicted and measured data enabled the assignment of a (*R*) configuration to both compounds. The absolute configuration of all other cross-coupling products was assigned by analogy, on the basis of their optical activity.

Conclusions

In summary, we have developed an efficient protocol for the enantioselective intermolecular Heck reaction using a variety of 5-substituted 2,3-dihydrofurans. We found that electron-rich aryl triflates can be coupled with various trisubstituted olefinic substrates when a chiral (P,N) ligand is used. The corresponding 2,5-dihydrofurans with a fully saturated C2 stereocenters are obtained in excellent regio- and enantioselectivities. Electron-deficient aryl triflates are compatible with the method when a commercial chiral (P,P) ligands is employed, leading to 2,2-disubstituted 2,3-dihydrofurans with excellent stereocontrol. The diversity, functional group tolerance and scalability of the approach have been also demonstrated and provide access to enantioenriched dihydrofurans that would be difficult to prepare otherwise.¹⁷ The underlying mechanistic features responsible for the marked electronic dichotomy observed between the two ligand systems along with extension of the method to other substrate classes are currently being investigated in our laboratories.

Acknowledgements

This work was supported by the University of Geneva. Stéphane Rosset is acknowledged for his expertise on analytical instruments and for assistance in ligand synthesis. Johnson-Matthey is acknowledged for a generous gift of palladium precursors.

Notes and references

^a University of Geneva, Department of Organic Chemistry, 30 quai Ernest Ansermet, 1211 Geneva-4, Switzerland.

^b University of Geneva, Department of Physical Chemistry, 30 quai Ernest Ansermet, 1211 Geneva-4, Switzerland.

† Electronic Supplementary Information (ESI) available: Experimental procedures, characterization of all new compounds, VCD studies and spectral data. See DOI: 10.1039/b000000x/

- (a) N. E. Carpenter, D. J. Kucera and L. E. Overman, *J. Org. Chem.*, 1989, **54**, 5846; (b) Y. Sato, M. Sodeoka and M. Shibasaki, *J. Org. Chem.*, 1989, **54**, 4738.
- (a) M. Oestreich, *The Mizoroki–Heck Reaction*, Wiley: New York, 2009; (b) A. B. Dounay and L. E. Overman, *Chem. Rev.*, 2003, **103**, 2945; (c) M. Shibasaki, E. M. Vogl and T. Ohshima, *Adv. Synth. Catal.*, 2004, **346**, 1533; (d) L. F. Tietze, H. Ila and H. P. Bell, *Chem. Rev.*, 2004, **104**, 3453; (e) M. Diéguez and O. Pàmies, *Isr. J. Chem.*, 2012, **52**, 572; (f) M. Oestreich, *Angew. Chem. Int. Ed.*, 2014, **53**, 2282.
- For a specific review on the intermolecular Heck reaction, see: D. Mc Cartney and P. J. Guiry, *Chem. Soc. Rev.*, 2011, **40**, 5122.
- (a) F. Ozawa, A. Kubo and T. Hayashi, *J. Am. Chem. Soc.*, 1991, **113**, 1417; (b) F. Ozawa and T. Hayashi, *J. Organomet. Chem.*, 1992, **428**, 267.
- For selected leading examples, see: (a) F. Ozawa, Y. Kobatake and T. Hayashi, *Tetrahedron Lett.*, 1993, **34**, 2505; (b) Y. Kurihara, M. Sodeoka and M. Shibasaki, *Chem. Pharm. Bull.*, 1994, **42**, 2357; (c) O. Loiseleur, M. Hayashi, N. Schmees and A. Pfaltz, *Synthesis*, 1997, 1338; (d) A. J. Hennessy, D. J. Connolly, Y. M. Malone and P. J. Guiry, *Tetrahedron Lett.*, 2000, **41**, 7757; (e) S. T. Henriksen, P.-O. Norrby, P. Kaukoranta and P. G. Andersson, *J. Am. Chem. Soc.*, 2008, **130**, 10414; (f) T. H. Wöste, and M. Oestreich, *Chem.—Eur. J.*, 2011, **17**, 11914; (g) J. Hu, H. Hirao, Y. Li and J. Zhou, *Angew. Chem., Int. Ed.*, 2013, **52**, 8676; (h) S. Liu and J. Zhou, *Chem. Commun.*, 2013, **49**, 11758; (i) C. Wu and J. Zhou, *J. Am. Chem. Soc.*, 2014, **136**, 650.
- (a) E. W. Werner, T.-S. Mei, A. J. Burckle and M. S. Sigman, *Science*, 2012, **338**, 1455; (b) T.-S. Mei, E. W. Werner, A. J. Burckle and M. S. Sigman, *J. Am. Chem. Soc.*, 2013, **135**, 6830; (c) T.-S. Mei, H. H. Patel and M. S. Sigman, *Nature*, 2014, **508**, 340; (d) L. Xu, M. J. Hilton, X. Zhang, P.-O. Norrby, Y.-D. Wu, M. S. Sigman and O. Wiest, *J. Am. Chem. Soc.*, 2014, **136**, 1960; (e) H. H. Patel and M. S. Sigman, *J. Am. Chem. Soc.*, 2015, **137**, 3462.
- Correia and co-workers reported a related enantioselective intermolecular Heck-Matsuda reaction employing (*Z*)-buten-1,4-diols or 3-cyclopentenol. We note that only tertiary centers were generated by this approach. For relevant references, see: (a) C. R. D. Correia, C. C. Oliveira, A. G. Salles Jr. and E. A. F. Santos, *Tetrahedron Lett.*, 2012, **53**, 3325; (b) C. C. Oliveira, R. A. Angnes and C. R. D. Correia, *J. Org. Chem.*, 2013, **78**, 4373; (c) R. A. Angnes, J. M. Oliveira, C. C. Oliveira, N. C. Martins and C. R. D. Correia, *Chem. Eur. J.*, 2014, **20**, 13117.
- In the strict sense, the stereogenic centers generated in this study are not quaternary, even though they are very congested. We therefore use the terms ‘fully saturated stereocenters’ to highlight this notion. A fully saturated center is defined by a tetrasubstituted C atom with at least one substituent different from H or C. For relevant reviews on the challenges associated with installing congested stereocenters, see: (a) J. Christoffers and A. Baro, *Quaternary Stereocentres: Challenges and Solutions for Organic Synthesis*, Wiley-VCH: Weinheim, 2005; (b) K. Fuji, *Chem. Rev.*, 1993, **93**, 2037; (c) M. Bella and T. Gasperi *Synthesis*, 2009, **10**, 1583; (d) I. Marek and G. Sklute, *Chem. Commun.*, 2007, 1683; (e) C. Hawner and A. Alexakis, *Chem. Commun.*, 2010, **46**, 7295.
- M. Tschoerner, P. S. Pregosin and A. Albinati, *Organometallics*, 1999, **18**, 670.
- (a) R. Grigg, V. Loganathan, V. Santhakumar, V. Sridharan and A. Teasdale, *Tetrahedron Lett.*, 1991, **32**, 687; (b) Y. Fall, F. Berthiol, H. Doucet, M. Santelli, *Synthesis*, 2007, 1683; (c) B. M. M. Wheatley and B. A. Keay, *J. Org. Chem.*, 2007, **72**, 7253; (d) M. G. Lauer, M. K. Thompson and K. H. Shaughnessy, *J. Org. Chem.*, 2014, **79**, 10837.

- 11 (a) P. Nareddy, L. Mantilli, L. Guénée and C. Mazet, *Angew. Chem. Int. Ed.*, 2012, **51**, 3826; (b) N. Humbert, E. Larionov, L. Mantilli, P. Nareddy, C. Besnard, L. Guénée and C. Mazet, *Chem. Eur. J.*, 2014, **20**, 745.
- 12 Asymmetric intermolecular Heck reactions can be accelerated using microwave irradiation. However, in some cases a drop in enantioselectivity has been observed. When the reaction reported in Table 1 (entry 5) was conducted under microwave irradiation (150 °C, 2 h), **3ab** was isolated in 38% yield and 87% *ee*. For relevant examples see: (a) P. Nilsson, H. Gold, M. Larhed and A. Hallberg, *Synthesis*, 2002, 1611; (b) P. Kaukoranta, K. Källstrom and P. G. Andersson, *Adv. Synth. Catal.*, 2007, **349**, 2595; (c) Y. Mata, O. Pàmies and M. Diéguez, *Chem. Eur. J.*, 2007, **13**, 3296; (d) J. Mazuela, O. Pàmies and M. Diéguez, *Chem. Eur. J.*, 2010, **16**, 3434.
- 13 Analysis of the crude reaction mixture by ¹⁹F NMR indicates deflation of **2c**, suggesting that oxidative addition into the C(sp²)-O bond of the aryl triflate is not problematic.
- 14 J. P. Genet, T. Ayad and V. Ratovelomanana-Vidal, *Chem. Rev.*, 2014, **114**, 2824.
- 15 Ligand **L2** is ca. 8 fold less-expensive than **L8**. This aspect and the relatively similar performances reported in Table 1 (entry 10 and 16) justify the use of **L2** for large scale experiments.
- 16 L. A. Nafie, T. A. Keiderling and P. J. Stephens, *J. Am. Chem. Soc.*, 1976, **98**, 2715; (b) G. Holzwarth, E. C. Hsu, H. S. Mosher, T. R. Faulkner and A. Moscovitz, *J. Am. Chem. Soc.*, 1974, **96**, 251; (c) T. B. Freedman, X. Cao, R. K. Dukor and L. A. Nafie, *Chirality*, 2003, **15**, 743.
- 17 For potential use and derivatization of the products obtained in this work, see: J. Cornella and R. Martin, *Org. Lett.*, 2013, **15**, 6298.