

# Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Chemical Science

## ARTICLE

## RNA G-Quadruplex Formation in Defined Sequence in Living Cells Detected by Bimolecular Fluorescence Complementation

Hong-he Liu,<sup>a</sup> Ke-wei Zheng,<sup>a</sup> Yi-de He,<sup>a</sup> Quan Chen,<sup>a</sup> Yu-hua Hao,<sup>a</sup> Zheng Tan<sup>a\*</sup>

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

G-quadruplexes are implicated in many essential cellular processes and sequences with potential to form G-quadruplex are widely present in DNA and RNA. However, it is difficult to know whether a sequence of interest naturally forms G-quadruplex in living cells. Here we report the detection of G-quadruplex in defined RNA sequences in living cells in a natural intracellular environment. A G-quadruplex forming sequence in RNA transcript is tagged at proximity with an aptamer. The two structures are recognized respectively by two probe proteins each of which is fused with a split half of an enhanced green fluorescent protein (eGFP). Simultaneous binding of the two proteins to the RNA brings the two halves of the eGFP into proximity, permitting them to reconstitute into a functional eGFP that emits fluorescence to signal the formation of G-quadruplex in the RNA. We show that G-quadruplex can form in RNA and can be detected with sequence and structure specificity under both in vitro and in vivo conditions. The results, therefore, provides a direct evidence for the formation of RNA G-quadruplexes in live cells and the method provides a useful tool to validate G-quadruplex formation in a specific sequence under a natural cellular condition.

### Introduction

G-quadruplexes are four-stranded structures formed in guanine-rich nucleic acids. Because of their implication in physiological and pathological processes as well as their potential use as therapeutic targets,<sup>1-5</sup> G-quadruplexes are gaining increasing attention from researchers in different disciplines in recent years. Sequences with potential to form intramolecular G-quadruplexes are extremely abundant in genomes and genes. For example, ~370,000 such sequences are present in the human genome.<sup>6</sup> These numbers are expanded because certain non-canonical sequences are also capable of forming G-quadruplexes. For example, >700,000 G-quadruplex-forming sequences have been identified in human genome by high-throughput sequencing.<sup>7</sup> Recently, we found that G-quadruplexes with an incomplete G-quartet layer can form in DNA, resulting in structures that response to guanine derivatives in environment.<sup>8</sup> The amount of motifs with potential to form such structures is in the same order of those for the canonical G-quadruplexes in human genome. In our earlier studies, we also found that a hybrid type of G-quadruplexes can form during transcription when a non-template DNA strand bears two or more guanine tracts (G-tract) downstream of the transcription start sites (TSS).<sup>9-12</sup> This finding raised the number of potential G-quadruplex forming sites to >1.5 million in the human genome.<sup>9</sup>

Besides DNA, RNA also forms G-quadruplexes, for example, in the human telomere sequence,<sup>13-17</sup> a most studied sequence in the G-quadruplex field. In human, there are >100,000 potential G-quadruplex-forming motifs in RNAs.<sup>9</sup> RNA G-quadruplexes forms during transcription to stimulate mitochondrial transcription termination and primer formation.<sup>18</sup> Their biological implications can be found in gene expression and telomere maintenance.<sup>19, 20</sup>

Detection of G-quadruplex formation in cells is necessary for determining the function of G-quadruplexes. The progresses made in the past few years on G-quadruplex-recognizing antibodies and ligands have resulted in the establishment of several detecting methods. In an early study, G-quadruplexes were detected in the telomeres of ciliated protozoa *Stylonychia*. The huge numbers of telomeres and their particular organization pattern in the macronucleus facilitated a direct visualization of G-quadruplexes by antibodies generated from ribosome display.<sup>21</sup> More recently, G-quadruplex structures have been visualized by antibody,<sup>22, 23</sup> and captured by chemical ligands<sup>24</sup> in human cells, providing evidence for the existence of G-quadruplexes in cells. A common feature of these methods is that they detect G-quadruplexes in cells in a large set of different sequences without sequence identities and most of them involve disruption or fixation of cells.<sup>21-25</sup> Therefore, they have an advantage of viewing an overall picture of G-quadruplex formation in cells. A potential G-quadruplex forming sequence may not necessarily form G-quadruplex in cells. For a specific G-quadruplex forming sequence, it is still difficult to know if a G-quadruplex can form inside cells in the sequence. In this case, the detection of G-quadruplex formation in the sequence

<sup>a</sup>State Key Laboratory of Membrane Biology, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, P. R. China.

Electronic Supplementary Information (ESI) available: Figs. S1-S7. See DOI: 10.1039/x0xx00000x

is essential to establish a connection between the structure and a cellular process.


In this work, we attempted to detect G-quadruplex structures in defined G-quadruplex forming sequences in RNA transcripts in living bacteria cells under a natural cellular condition by adapting the bimolecular fluorescence complementation (BiFC) technique.<sup>26</sup> By transformation with appropriate plasmids, RNAs containing a G-quadruplex-forming sequence and an aptamer tag side-by-side were expressed in *E. coli* cells. In the meantime, two probe proteins that were fused respectively with two split halves of an enhanced green fluorescent protein (eGFP) were also expressed by transformation with another plasmid. When the two fusion proteins bound simultaneously to the aptamer and G-quadruplex on the RNA, the formation of G-quadruplex in the RNA was reported by the fluorescence generated by the complementation of the two split halves of the eGFP brought together by the two probe proteins. By this method, we demonstrate that G-quadruplexes can form in RNA in living cells which can be detected with both sequence and structure specificity.

## Results


### Design of BiFC for detection of RNA G-quadruplex

We constructed several plasmids (Supporting Information, Fig. S1A) that could express RNAs containing a *REX(A)* aptamer at the 5' side of three G-quadruplex forming motifs, *TERC(Q)*, *G4T(Q)*, and *PITX(Q)*, respectively (Fig. 1) (RNA motifs are presented in italic and probe proteins in normal font here and thereafter). We examined G-quadruplex formation in the RNA sequences, *TERC(Q)*, *G4T(Q)*, or *PITX(Q)*, using CD spectroscopy. Their CD spectra all displayed a positive peak at 265 and a negative peak at 240 nm (Supporting Information, Fig. S2) which are characteristic of parallel G-quadruplexes.

Another plasmid (Supporting Information, Fig. S1B) was constructed to express two probe proteins,  $G_N$ -*REX(A)* and *RHAU(Q)*- $G_C$  (Supporting Information, Fig. S3), that could bind the RNA aptamer and G-quadruplexes, respectively. The *REX(A)* aptamer in the RNA was recognized by  $G_N$ -*REX(A)*,<sup>27</sup> a *REX* protein fused with the N-terminal half of the eGFP ( $G_N$ ).



**Fig. 1.** Schematic illustration of detection of RNA G-quadruplex by the BiFC technique. An aptamer-binding protein *REX(A)* is fused with the N-terminal half ( $G_N$ ) and a G-quadruplex-binding protein *RHAU(Q)* is fused with the C-terminal half ( $G_C$ ) of an eGFP, respectively. When the  $G_N$ -*REX(A)* and *RHAU(Q)*- $G_C$  bind to their targets *REX(A)* and G-quadruplex (GQ) on a same RNA, the two split halves of the eGFP are brought to close vicinity with each other such that they can reconstitute into an active eGFP, resulting in fluorescence emission. "A" and "Q" in parentheses indicates that a probe fusion protein is intended to target aptamer or G-quadruplex, or that a RNA motif forms aptamer or G-quadruplex, respectively.




**Fig. 2.** Specific probe protein binding to intended RNA target revealed by electrophoretic mobility shift assay (EMSA). (A) *REX(A)*-*TERC(Q)*, (B) *REX(A)*-*G4T(Q)*, (C) *REX(A)*-*PITX(Q)* RNA and their G-quadruplex mutants, *REX(A)*-*TERC(Qm)*, *REX(A)*-*G4T(Qm)*, and *REX(A)*-*PITX(Qm)* were incubated with either protein  $G_N$ -*REX(A)* or *RHAU(Q)*- $G_C$  or both in  $K^+$  solution before being resolved by native gel electrophoresis. The -, +, ++, +++ indicate protein concentrations of 0, 125, 500, and 1250 nM, respectively, for  $G_N$ -*REX(A)* and of 0, 250, 1000, and 2500 nM, respectively, for *RHAU(Q)*- $G_C$ . "Qm" in parenthesis indicates mutated G-quadruplex sequence in this and the rest Figs..

*REX* is a human T-cell leukemia virus type 1 (HTLV-1) encoded protein with high binding affinity to the *REX(A)* aptamer.<sup>27</sup> On the other hand, the G-quadruplexes were recognized by *RHAU(Q)*- $G_C$ , a *RHAU* protein fused with the C-terminal half of the eGFP ( $G_C$ ). *RHAU* is a human DEAH-box RNA helicase that has been reported to bind intramolecular RNA G-quadruplex in *TERC* and *PITX* in vitro and in cells.<sup>28-30</sup> When both probe fusion proteins bound to their targets on a same RNA, the two eGFP fragments would be brought into close proximity to form a functional eGFP that could emit fluorescence to signal the presence of G-quadruplex in the sequence (Fig. 1). If G-quadruplex was not present on the RNA, its recognizing protein would not bind the RNA. In this case, the RNA-mediated complementation would not take place.

### In vitro RNA target recognition by probe proteins

We first used electrophoretic mobility shift assay (EMSA) to test if the probe proteins could recognize their intended targets. RNAs containing *TERC(Q)*, *PITX(Q)*, *G4T(Q)*, and their mutants, *TERC(Qm)*, *PITX(Qm)*, *G4T(Qm)*, were incubated with



**Fig. 3.** Fluorescence emission as a result of complementation between probe proteins upon binding to their RNA targets. Two probe proteins,  $G_N$ -REX(A) and RHAU(Q)- $G_C$  were incubated individually or together in (A) the absence or (B-D) together in the presence of (B) *REX(A)-TERC(Q)* and *REX(A)-TERC(Qm)*, or (C) *REX(A)-G4T(Q)* and *REX(A)-G4T(Qm)*, or (D) *REX(A)-PITX(Q)* and *REX(A)-PITX(Qm)* RNAs. Fluorescence was monitored over time after sample mixing.

each individual probe protein and the RNA/protein complexes were resolved on a native gel. The results in Fig. 2A show that both the  $G_N$ -REX(A) (lanes 2-4, 9-11) and RHAU(Q)- $G_C$  (lanes 5-7) protein bound their targets in a concentration-dependent manner. The binding of RHAU(Q)- $G_C$  to the *TERC(Q)* motif was specific to G-quadruplex because when the *TERC(Q)* motif in the RNA was mutated to *TERC(Qm)* to disable the formation of G-quadruplex, no binding was detected (lanes 12-14). When the wild type *REX(A)-TERC(Q)* RNA was incubated with both  $G_N$ -REX(A) and RHAU(Q)- $G_C$ , an additional band appeared (lane 15), demonstrating that the two proteins could simultaneously bind to the RNA. The simultaneous binding also relied on the presence of G-quadruplex because the extra band disappeared when the mutated *TERC(Qm)* motif was used (lane 16). For the *REX(Q)-G4T(Q)* and *REX(Q)-PITX(Q)* RNAs, similar results were also obtained. The binding of RHAU(A)- $G_C$  to the *G4T(Q)* (Fig. 2B, lanes 5-7) and *PITX(Q)* (Fig. 2C, lanes 5-7) also depended on the presence of G-quadruplex structure and mutation in the *REX(Q)-G4T(Qm)* and *REX(Q)-PITX(Qm)* abolished the binding (Fig. 2B and 2C, lanes 12-14).

G-quadruplex is stabilized by monovalent cations in the order of  $K^+ > Na^+ > Li^+$ .<sup>31</sup> Our aforementioned EMSA was carried out in  $K^+$  solution to promote formation of G-quadruplex in the RNAs. To further verify the structural specificity of RHAU(Q)- $G_C$ , we performed the same experiments with  $K^+$  being replaced by  $Li^+$ . As is shown in Fig. S5, this substitution dramatically reduced the binding of RHAU(Q)- $G_C$  to the G-quadruplex targets. For example, a majority of the *REX(A)-TERC(Q)* RNAs remained free in  $Li^+$  solution whereas they became fully shifted in  $K^+$  solution in the presence of  $>1 \mu M$  of RHAU(Q)- $G_C$  (Fig. 2A, lanes 6-7 versus Fig. S5A, lanes 6-7). This phenomenon also hold true for the *REX(A)-G4T(Q)* RNA (lanes 6-7 in Fig. 2B versus in Supporting Information, Fig. S5B). Collectively, the results obtained with mutations in the RNAs (Fig. 2) and wild-type RNA in  $Li^+$  solution


(Fig. S5) confirmed that the binding of RHAU(Q)- $G_C$  to the RNAs were G-quadruplex specific.

### In vitro fluorescence generation upon target binding

The capability of the  $G_N$ -REX(A) and RHAU(Q)- $G_C$  to simultaneously bind their targets in a RNA molecule provides a basis for the two proteins to form a BiFC complex. The two proteins did not show fluorescence emission when they were mixed together in the absence of RNA target (Fig. 3A). When the RNA carrying a *REX(A)* aptamer and a *TERC(Q)* G-quadruplex was added, fluorescence was detected that increased over time (Fig. 3B). Mutation in the *TERC(Qm)* prevented the generation of fluorescence because of the lack of G-quadruplex. Similar results were obtained using the other two aptamer/G-quadruplex pairs (Fig. 3C and 3D). These results indicated that complementation took place when both  $G_N$ -REX(A) and RHAU(Q)- $G_C$  bound to their targets in the RNA. They also demonstrated that the fluorescence signal could serve as an indication of G-quadruplex formation in the RNA.

### Detection of G-quadruplexes in living *E. coli*

Using the fluorescence signal of the  $G_N$ -REX(A) and RHAU(Q)- $G_C$  proteins generated upon binding to their targets, we detected G-quadruplex formation in RNA in living *E. coli* cells. After the expression of the two proteins and RNA targets (Fig. S1A, 1B) were induced by an addition of IPTG, the *E. coli* cells were analyzed by flow cytometry (Fig. 4). When the cells expressed the *REX(A)-TERC(Qm)* RNA in which the mutation prevented G-quadruplex formation, they showed certain level of background fluorescence (Fig. 4A, green curve), probably as a result of self-assembly of the two split halves of the eGFP.<sup>32</sup> When the *REX(A)-TERC(Q)* RNA was expressed, however, a




**Fig. 4.** Detection of BiFC fluorescence in *E. coli* cells by flow cytometry. Distribution of (A-C) eGFP fluorescence and (D-F) forward scattering intensity was collected of cells expressing  $G_N$ -REX(A) and RHAU(Q)- $G_C$  probe proteins plus the indicated RNA. (G-I) Relative RNA expression (mean  $\pm$  SD of three experiments) assayed by qPCR.


new population of cells with much higher fluorescence intensities emerged, resulting in an additional peak at the right hand side of the histogram (Fig. 4A, red curve). Again, expression of the other two RNAs carrying the *REX(A)-G4T(Q)* and *REX(A)-PITX(Q)*, respectively, also showed increased fluorescence in the cells comparing with those expressing the corresponding RNA mutants (Fig. 4B and 4C, red versus green curve). The cells expressing the wild and mutated RNAs displayed little difference in their forward scatter (FSC) signal (Fig. 4D-4F). The RNA expression levels were also similar between the two species (Fig. 4G-4I). These two results supported that the skewed distribution in eGFP fluorescence in the cells expressing the G-quadruplex-bearing RNAs was the result of G-quadruplex formation, rather than possible variation in cell size or RNA level.

To test if the enhancement in fluorescence involved target RNA, we replaced the RNA expressing plasmid (Fig. S1A) with a new one (Fig. S4A) by which the expression of *REX(A)-TERC(Q)* and *REX(A)-TERC(Qm)* RNA can be independently induced by anhydrotetracycline (ATc). After testing several induction intervals, we found that fluorescence showed greater difference between cells expressing *REX(A)-TERC(Q)* and *REX(A)-TERC(Qm)* when they were induced about at the same time as the probe proteins. As is shown in Fig. 5A, the enhancement of fluorescence increased over time after the induction of probe proteins in the absence of ATc, which might result from a background expression of target RNAs. When the RNAs were simultaneously induced with the proteins, the fluorescence enhancement was initially reduced (red versus green curve) probably because of a reduction in the protein expression due to the competition for resources from the RNA expression. At a later time, however, the fluorescence enhancement became significantly greater than that in the cells without ATc treatment. This increment in fluorescence enhancement upon RNA induction clearly showed that the greater fluorescence in the *REX(A)-TERC(Q)*-expressing cells was an RNA- and G-quadruplex-dependent event.

To verify the structural specificity of the BiFC fluorescence towards G-quadruplex, we replaced the G-quadruplex forming motif with a set of four sequences, in which the number of  $G_3$  tracts increased from 1 to 4 (Supporting Information, Fig. S6,




**Fig. 5.** Dependence of BiFC fluorescence enhancement on target RNA expression. (A) Time course of difference in mean fluorescence between cells expressing RNA *REX(A)-TERC(Q)* and *REX(A)-TERC(Qm)* in the presence and absence of RNA induction by ATc. (B) RNA level in cells before and after induction with ATc. Data are given in mean  $\pm$  SD of three experiments. IPTG was added at 0 hr with or without ATc. Representative distribution of eGFP fluorescence of corresponding cells is given in Fig. S4B.



**Fig. 6.** Dependence of BiFC fluorescence on the number of  $G_3$  tracts in the RNA target expressed in *E. coli*. Distribution of (A) eGFP fluorescence or (B) forward scattering intensity or (C) relative RNA expression (mean  $\pm$  SD of three experiments) was assayed and processed as in Fig. 4.

1G3-4G3(Q)). The fluorescence of the corresponding cell populations showed little change with a sequential addition of  $G_3$  tract till their number reached 3 when these changes in the sequence did not bring a possibility of G-quadruplex formation (Fig. 6A, colored curves). However, a further addition of a single  $G_3$  tract resulted in an increase in fluorescence once the sequence became capable of forming G-quadruplex (Fig. 6A, black curve). Since the forward scatter (Fig. 6B) and RNA expression (Fig. 6C) was similar among the plasmids, this result further demonstrated that the recognition of the RHAU(Q)- $G_C$  was specific to G-quadruplex.

The complementation efficiency of the two eGFP halves is dependent of the distance between the two partners. To further confirm the target specificity of the fluorescence signal, we added an additional 20 nt sequence between the aptamer *REX(A)* and G-quadruplex *TERC(Q)* and *G4T(Q)*, respectively, in the RNAs (Supporting Information, Fig. S7). This modification would either push the aptamer and G-quadruplexes further apart or made the linkage between them more rigid by folding into a double-stranded stem, which were expected to reduce the complementation and, as a result, BiFC fluorescence. Indeed, for the two RNAs, the fluorescence was significantly reduced (Fig. 7, blue curve) compared to that of



**Fig. 7.** Dependence of BiFC fluorescence on the distance between the aptamer and G-quadruplex targets. Distribution of (A and B) eGFP fluorescence or (C and D) forward scattering intensity was collected by flow cytometry and processed as in Fig. 4. Cells expressed  $G_N$ -*REX(A)* and RHAU(Q)- $G_C$  plus the indicated RNA without (red, and green curves) or with a 20 bp stem spacer (blue curve) between the aptamer and G-quadruplex motif.

the cells expressing the corresponding RNAs without the spacer (Fig. 7, red curve).

### Immuno-detection of probe protein-G-quadruplex interaction in living *E. coli*

We verified the interaction between RHAU(Q)-G<sub>C</sub> and RNA G-quadruplexes in cells by a pull-down experiment. The target RNA was pulled down by an immobilized DNA oligonucleotide that was complementary to the 22 nt region at the 5' end of the RNAs. The protein associated with the RNA was then detected by a monoclonal antibody against eGFP. While information regarding the amino acid sequence recognized by the antibody is not available from the manufacturer, the antibody appeared to recognize both fragments of the eGFP (Fig. 8). When IPTG was not added to induce the expression of the probe proteins and target RNA or when RNA-expressing plasmid was omitted, only a weak background was detected (Fig. 8A, lanes 7-10, top gel). When *REX(A)-TERC(Q)* was expressed, the two protein bands showed dramatically enhanced intensity at similar magnitudes (lane 11). Mutation at the G-quadruplex moiety significantly reduced the intensity of RHAU(Q)-G<sub>C</sub> band as expected (lane 12). In the same lane, the G<sub>N</sub>-*REX(A)* displayed a smaller decrease in the intensity, which might be resulted from the loss of the RHAU(Q)-G<sub>C</sub> bound to the RNA. A fraction of the G<sub>N</sub>-*REX(A)* proteins on the RNA might be simply associated with RHAU(Q)-G<sub>C</sub> by self-assembly without itself binding the *REX(A)* aptamer such that they were lost when the binding of RHAU(Q)-G<sub>C</sub> on the RNA


was reduced. Therefore the reduction in the signal for both proteins signified that the binding of the RHAU(Q)-G<sub>C</sub> to the RNA was G-quadruplex specific. As expected, hydrolysis of the RNA with RNase A (lanes 13-14) significantly reduced the intensity of the two probe proteins, indicating that the proteins were pulled down by the target RNA via a protein-RNA interaction. Similar results were also obtained with the *REX(A)-G4T(Q)* RNA and its mutant (Fig. 8B, lanes 7-14, top gel). The results for both RNAs therefore confirmed the interaction between the probe proteins and their target RNAs in the cells.

### Discussion

In summary, we successfully detected G-quadruplex formation in living bacterial cells with sequence and structure specificity using plasmids expressing RNA and target-recognizing fusion probe proteins. Our results provide evidence that G-quadruplex can form in living cells under a truly natural condition. By the *in vitro* EMSA assays, we proved that the probe proteins can specifically recognize and simultaneously bind to their intended targets on the RNAs (Fig. 2). As a result, bimolecular complementation took place as expected between the two proteins, giving rise to fluorescence to signal the presence of G-quadruplex structure (Fig. 3). The binding was suppressed in solution of Li<sup>+</sup> that does not stabilize G-quadruplex (Fig. S5). In addition, mutation in the G-quadruplex forming motifs to prevent G-quadruplex formation abolished the binding of the G-quadruplex probe protein to the RNAs (Figs. 2 and 3). Our *in vivo* assays further showed that these properties led to a successful application of the BiFC system in living cells (Figs. 4-6, and S4). The correlation between the increase in fluorescence and capability of the sequences to form G-quadruplex confirmed the structure specificity (Figs. 4, 5 and S4). The dependence of fluorescence emission of the eGFP on the distance between the two probe proteins supported the complementation interaction between the two split halves of the eGFP (Fig. 7).

The BiFC events mediated by RHAU indicate not only a formation of G-quadruplex, but also an interaction between the target G-quadruplex and RHAU in cells. The RHAU is able to bind both intra- and intermolecular G-quadruplexes.<sup>30</sup> There were two or three G-tracts in our RNA mutants *REX(A)-TERC(Qm)*, *REX(A)-G4T(Qm)*, *REX(A)-PITX(Qm)*, *REX(A)-3G3*, and *REX(A)-2G3*. They are capable of forming intermolecular G-quadruplexes and, therefore, might also lead to BiFC. Because these RNAs were used as controls, the fluorescence enhancement associated with the wild RNAs was an indication of intramolecular G-quadruplex formation. A recent study shows that the RHAU binds parallel quadruplexes, but not non-parallel ones,<sup>33</sup> suggesting that the RHAU only binds the folded form. This property eases the concern on a possibility that the protein may fold a sequence into a G-quadruplex.

The bimolecular complementation strategy in the BiFC and the choice of plasmid-generated RNA transcript offer several advantages. Firstly, the detection can be conducted in living



**Fig. 8.** Pull-down of G-quadruplex-recognizing proteins by RNA from *E. coli*. Cells were lysed in the presence of RNase inhibitor or RNase A. Probe proteins associated with the indicated target RNAs were pulled down with an immobilized DNA oligomer complementary to the 5' end of the RNA and detected by an antibody against eGFP. "NS" indicates a non-specifically stained band that can serve as a loading control.

cells in a natural cellular environment. Secondly, negative controls can be arranged in BiFC to strictly evaluate the structural specificity. When G-quadruplexes in endogenous nucleic acid molecules are to be examined, it is technically difficult to have a negative control. Thirdly, the construction of a plasmid made it possible to study any G-quadruplex-forming RNA sequences by simply insert a corresponding DNA between the promoter and terminator (Fig. S1). Lastly, BiFC is normally irreversible.<sup>32</sup> This is a desired property that should enhance the detection sensitivity because BiFC is expected to accumulate even when interacting G-quadruplex dissociates or unfolds.

Since the introduction of an aptamer at one side of the G-quadruplex-forming motif, the native RNA sequence is altered. This may influence but is unlikely to prevent the formation of G-quadruplexes because G-quadruplexes are stable structures. On the other hand, the formation of aptamer reduces its interaction with the rest part of a RNA and such interaction may be further minimized when the aptamer is bound by a probe protein. RNA-protein interaction constantly takes place in cells. The disturbance brought by the aptamer and its interaction with a probe protein may not be different from those by the other RNA-protein interactions naturally occurring in cells. Therefore, once G-quadruplexes are detected, it is most likely that G-quadruplexes would also form in the endogenous native sequence. In future development of the method, the aptamer may be removed if DNA G-quadruplexes are detected using the ZFN, TALEN, and CRISPR/Cas technique in which recognition of virtually any sequence is possible.<sup>34</sup>

Our data demonstrated that certain factors affect the detection of G-quadruplexes. The detection relies on a proper G-quadruplex-binding protein. Similar to antibodies and chemical ligands, a protein may better recognize some type of G-quadruplexes than others. This might be why the BiFC fluorescence varied when different G-quadruplex targets were used (Fig. 3). For the same reason, when a G-quadruplex-forming sequence has more than four G-tracts and therefore has chance to adopt different forms of G-quadruplexes, some may be better captured than others depending on the selectivity of the probe protein. Therefore, testing different binding proteins may be required when positive signal is not seen with a G-quadruplex forming sequence. When a protein is able to bind a G-quadruplex target, the distance between the G-quadruplex and aptamer was critical for a given fusion probe protein pair (Fig. 7). The optimal distance may also vary with different probe protein pairs depending on their size and spatial orientation when bound to their targets. Our method also inherited an issue from the BiFC in that self-assembly can occur between the two halves of an eGFP,<sup>32</sup> resulting in background fluorescence especially when the two proteins are expressed at high level. Given this, the timing of RNA and probe protein expression becomes important. If the concentration of RNA overweighs that of the probe proteins, the two halves of the protein tend to bind to different RNA molecules, reducing their chance to meet with each other. On the other hand, if the RNA expression lags behind that of the

proteins, the level of proteins may become too high leading to high fluorescence background as a result of self-assembly. It seems that inducing both RNA and protein expression at the same time is a practical option to produce optimal results and simplify the expressing system. While a thorough exploration on these technical issues is not within the scope of this work, our study demonstrates that the BiFC-based technique should benefit studies towards specific G-quadruplex in a sequence of interest in a natural cellular environment, thus, providing a complementation to the existing methods.

## Experimental Section

### RNA- and protein-expressing plasmids

An IPTG-inducible RNA-expressing plasmid was constructed base on the pET28b (Novagen) backbone (Fig. S1A) to which an anti-REX RNA aptamer, *REX(A)*, was inserted between the Xba I and Xho I sites and a G-quadruplex forming sequence, *GQ*, or mutant was inserted between the Xho I and Bpu1102 I sites downstream of the T7 promoter.

An ATc-inducible RNA-expressing plasmid was constructed base on the pETDuet-1 (Novagen) backbone (Fig. S4A) to which 2×tetO, a *REX(A)-TERC(Q)* or *REX(A)-TERC(Qm)* sequence, and a T7 terminator was inserted between the Xba I and Hind III site downstream of the first T7 promoter. A TetR protein coding sequence was amplified from the pLenti6/TR vector (Invitrogen) and inserted between the Bgl II and Kpn I site downstream of the second T7 promoter of the modified plasmid. Then the LacI promoter and part of the LacI coding sequence were cut out at the Sph I and Mlu I site. Finally, the second T7 promoter was replaced by a lacI promoter by insertion at Not I and Bgl II site.

The protein-expressing plasmid for BiFC was constructed based on the pACYCDuet1 (Novagen) to express two fusion proteins simultaneously (Fig. S1, right).<sup>35</sup> The aptamer-binding protein, *G<sub>N</sub>-REX(A)*, was a REX protein fused to the C-terminus of a N-terminal fragment (aa 1-158) of an eGFP. The G-quadruplex-binding protein, *RHAU(Q)-G<sub>C</sub>*, consisted of a RHAU protein fragment (aa 53-105) fused to the N-terminus of a C-terminal fragment (aa 159-238) of an eGFP. The sequence of the two components in each of the fusion protein was separated by a same flexible 10-mer oligopeptide. For protein purification, the coding sequences of *G<sub>N</sub>-REX(A)* and *RHAU(Q)-G<sub>C</sub>* were amplified from the pACYCDuet1 plasmid by PCR and inserted into the pET28b plasmid between the Nde I and Xho I sites downstream of a T7 promoter.

### RNA preparation

To prepare RNA for EMSA and *in vitro* BiFC assays, the RNA-expressing plasmid was linearized by cutting at the Bpu1102 I site of the pET28b. The resulting DNA (50 ng/μl) was transcribed at 37 °C for 3 hr in 80 mM HEPES-LiOH buffer (pH 7.5), 10 mM MgCl<sub>2</sub>, 50 mM LiCl, 10 mM DTT, 1 mM spermidine, 4 U/μl T7 RNA Polymerase (Fermentas, USA), 2 mM rNTPs, 1 U/μl RiboLock RNase Inhibitor (Fermentas, USA), and 0.025 U/μl Pyrophosphatase, Inorganic (Fermentas, USA).

The samples were then treated at 37 °C for 30 min with DNase I at a final concentration of 0.04 U/ $\mu$ l to remove DNA, followed by an addition of 20 mM EDTA final concentration to stop the reaction. RNA transcript was purified using the RNAClean Kit (TIANGEN, China) and stored at -70 °C in RNase-Free water containing 1 U/ $\mu$ l RiboLock RNase Inhibitor (Fermentas, USA). The preparation of RNA of the quadruplex core or the corresponding mutant sequence for circular dichroism (CD) spectroscopy was carried out in the same way except that synthesized DNA was used as template at a final concentration of 0.1  $\mu$ M.

#### Protein expression and purification

Protein-expressing plasmid was transformed into the BL21 (DE3) *E. coli* strain (TransGen Biotech) and the expressed proteins in inclusion body was purified as described<sup>36</sup> with modification. Briefly, transformed *E. coli* cells were grown at 37 °C to OD<sub>600</sub>  $\approx$  0.5, then protein expression was induced by adding 0.5 mM IPTG and lowering the temperature to 25 °C. After an overnight growth, cells were collected by centrifugation and resuspended in 50 mM Tris-HCl (pH 8.5), 25% sucrose, 1 mM EDTA, and 10 mM DTT. They were then disrupted by freezing at -70 °C followed by sonication with a Vibra-Cell (Sonics, USA) in ice-water bath for 30 min in cycles of 3 sec on and 7 sec off. After centrifugation, the pellet was resuspended in washing buffer (50 mM Tris-HCl, pH 8.5, 0.5% TritonX-100, 100 mM NaCl, 1 mM DTT, and 1 mM NaEDTA) by a brief sonication. The sample was centrifuged and the pellet was solubilized in a buffer of 25 mM MES (pH 8.5), 8 M urea, 10 mM NaEDTA, 0.1 mM DTT. The solution was added slowly drop by drop to a refolding buffer (50 mM Tris-HCl, pH 8.5, 500 mM NaCl, 1 mM DTT) to a 1:150 fold dilution and kept at 4 °C overnight. The refolded His-tagged protein was purified with the His GraviTrap column (GE Healthcare) and concentrated by ultrafiltration with the Ultra-4 Centrifugal Filter Units (Amicon, Millipore). Purified protein (Fig. S3) was stored in 20 mM Tris-HCl, pH 7.4, 200 mM NaCl, 1 mM EDTA; 1 mM DTT, 50% (v/v) glycerol at -70 °C until use.

#### Circular dichroism (CD) spectroscopy

Half  $\mu$ M RNA obtained by *in vitro* transcription was denatured at 95 °C for 5 min in 10 mM Tris-HCl (pH 7.4), 50 mM KCl, 1 mM EDTA, and quickly cooled on ice. CD spectra were measured at 25 °C on a Chirascan-plus CD Spectrometer (Applied Photophysics, United Kingdom) with a 10 mm pathlength quartz cuvette.

#### Electrophoretic mobility shift assay (EMSA)

EMSA was carried out as described.<sup>37</sup> RNA (0.4  $\mu$ M) was mixed with a synthetic 3' 6-FAM labeled DNA oligomer (0.8  $\mu$ M) complementary to a 22 nt<sup>32</sup> region at the 5' end of the RNA in 10 mM Tris-HCl (pH 7.4), 50 mM KCl, and 1 mM EDTA. The mixture was heated to 95 °C and slowly cooled down to 25 °C. The RNA (0.04  $\mu$ M) was then incubated with G<sub>N</sub>-REX(A) or RHAU(Q)-G<sub>C</sub> or both at the indicated concentration in buffer of 10 mM Tris-HCl, 100 mM KCl, 1 mM EDTA, 1 mM DTT, 5% (v/v)

glycerol, 0.4 U/ $\mu$ l RiboLock RNase Inhibitor (Fermentas, USA), and 110 ng/ $\mu$ l yeast tRNA. RNA-protein complexes were resolved on a 8% non-denaturing polyacrylamide gel (37.5:1 acrylamide:bis) at 4 °C for 90 min in 1 $\times$  TBE buffer containing 75 mM KCl.

#### Spectroscopy of real-time BiFC *in vitro*

*In vitro* real-time monitoring of BiFC by eGFP fluorescence was performed in a 96-Well Black Solid Plate (Costar, Corning). First, 1  $\mu$ M RNA in 10 mM Tris-HCl (pH 7.4), 50 mM KCl, and 1 mM EDTA was heated to 95 °C and then slowly cooled down to 25 °C. Then 10  $\mu$ l RNA was added to the 96-Well plate containing 100  $\mu$ l of 20 mM Tris-HCl (pH 7.4), 100 mM KCl, 5 mM MgCl<sub>2</sub>, 1 mM DTT, 0.4 U/ $\mu$ l RiboLock RNase Inhibitor (Fermentas, USA), and 0.2  $\mu$ M fusion protein each or both. Fluorescence was recorded at 507 nm with an excitation at 480 nm at a 2 min interval in a Synergy 4 Hybrid Microplate Reader (BioTek, USA) at 25 °C.

#### Flow cytometry of BiFC in cells

For simultaneous expression of RNA and probe proteins, the RNA and protein expressing plasmids (Fig. S1) were transformed into the BL21 (DE3) (TransGen Biotech) *E. coli* strain. Cells were cultured at 37 °C in 5 ml LB medium on a shaking incubator overnight. The culture was then diluted 100 times in fresh LB media and maintained at 37 °C for about 2–3 hrs until OD<sub>600</sub>  $\approx$  0.5. Then 0.1 mM IPTG was added to induce the expression of RNA and probe proteins overnight at 25 °C. The cells were collected by centrifugation at 4000 $\times$  g for 2 min and washed twice with PBS by filtration on a 0.22- $\mu$ m filter. They were then diluted in PBS to an OD<sub>600</sub>  $\approx$  0.05. Fluorescence was measured on a FACSCalibur flow cytometer (Becton-Dickinson, USA) using the FL1 channel (excitation at 488 nm, emission filter at 530/30 nm).

To monitor the BiFC with an independent induction of target RNA, the RNA (Fig. S4A) and probe protein (Fig. S1B) expressing plasmids were transformed into the BL21 (DE3). Colony was selected to start the culturing in a LB medium with 100 mg/ml of ampicillin and 34 mg/ml of chloramphenicol at 37 °C overnight. The cultures were then diluted 100 times into fresh LB media and maintained at 37°C for about 3 hrs until OD<sub>600</sub>  $\approx$  0.4. Then 0.1 mM IPTG was added together with or without 2  $\mu$ g/ml anhydrotetracycline (ATc) to induce the expression of probe proteins/RNA or protein only. Cells were collected at different time points and analyzed by flow cytometry as aforementioned. This RNA expressing plasmid was only used in the experiments for Fig. S4.

#### Quantitative polymerase chain reaction (qPCR)

Total RNAs were extracted from the *E. coli* cells using the RNAprep Pure Cell/Bacteria Kit (TIANGEN, China). The RNAs were then polyadenylated by *E. coli* Poly(A) Polymerase (M0276S, NEB) in a 50  $\mu$ l volume containing 50 mM Tris-HCl (pH 7.9), 250 mM NaCl, 10 mM MgCl<sub>2</sub>, 1.5  $\mu$ g RNA, 2.5 U poly(A) Polymerase, and 1 mM ATP. cDNA was generated from the purified tailed RNA using M-MLV Reverse Transcriptase



(M368B, Promega). Briefly, 1  $\mu\text{g}$  tailed RNA and 0.05 nmol Oligo(dT)<sub>18</sub> in a total volume of 15  $\mu\text{l}$  in RNase free water were heated at 70 °C for 5 min and then cooled immediately on ice for 5 min. They were then mixed in a total volume of 50  $\mu\text{l}$  50 mM Tris-HCl (pH 8.3) buffer with 4 U/ $\mu\text{l}$  M-MLV RT (H-) Point Mutant, 0.25 mM dNTP each, 75 mM LiCl, 3 mM MgCl<sub>2</sub> and 10mM DTT. After a reaction with 3 cycles of 45 °C for 5 min, 48 °C for 5 min, 52 °C for 5 min, and then followed by 52 °C for 30 min, qPCR reaction was carried out in 20  $\mu\text{l}$  mixture containing 10  $\mu\text{l}$  GoTaq qPCR Master Mix (A6001, Promega), 1  $\mu\text{l}$  1/5 diluted cDNA, 0.2  $\mu\text{M}$  forward primer and 0.2  $\mu\text{M}$  reverse primer on the real-time cycler qTower (Analytik Jena AG, Jena, Germany) with the following setting: 95 °C for 2 min, followed by 40 cycles of 95 °C for 15 sec, 60 °C for 60 sec. The forward and reverse primer sequences for the target RNA were 5'-TTAGCGCAGGTACGCAAGT-3' and 5'-TCGAGAAATAGCCGGCG-3', respectively. For the IPTG-induced RNA expression (Fig. S1A), an internal control was amplified from KanR using a primer pair of 5'-GCAATCAGGTGCGACAATCT-3' (forward) and 5'-TCATTGGCAACGCTACCTTT-3' (reverse). For the ATc-induced RNA expression (Fig. S4A), an internal control was amplified from CmR using a primer pair of 5'-CACATTCTTGCCCGCTGAT-3' (forward) and 5'-TCACCGTCTTCATTGCCATAC-3' (reverse).

#### Immunoblotting of RNA-Protein interaction

Cells from 4 ml of culture were pelleted and resuspended in 300  $\mu\text{l}$  ice-cold lysis buffer containing 50 mM Tris-HCl, pH 7.4, 150 mM KCl, 1 mM EDTA, 1 mM DTT, 0.1 mM AEBSF, 1 U/ $\mu\text{l}$  RiboLock RNase Inhibitor (Fermentas, USA) or 10  $\mu\text{g}/\text{ml}$  RNase A (Fermentas, USA), followed by sonication for 15 min with 5 sec on and 5 sec off cycles in ice-water bath (Bioruptor-Plus, Diagenode). The lysate was centrifuged at 16000 $\times$  g for 20 min and the supernatant was collected and incubated at room temperature under constant rotation for 1 hr with 20  $\mu\text{l}$  streptavidin-agarose beads (S1638, Sigma) that was pre-loaded with a biotinylated DNA oligomer complementary to the 22 nt region at the 5' end of the target RNA. After centrifugation at 2000 $\times$  g for 1 min, resin-bound RNA-protein complexes were washed three times with 10 resin bed volume of washing buffer (50 mM Tris-HCl, pH 7.4, 150 mM KCl, 1mM EDTA, 1 mM DTT, 0.1 mM AEBSF). The RNA-protein complexes were then released by digestion with RNase H (Fermentas, USA) in 20 mM Tris-HCl (pH 7.8), 40 mM KCl, 10 mM MgCl<sub>2</sub>, and 1 mM DTT. After a centrifugation at 12000 $\times$  g for 1 min, the supernatant was collected and resolved on a 15% Tris-glycine PAGE gel. The proteins were transferred to a Protran BA 85 nitrocellulose membrane (Whatman-Schlicher and Schuell) using a semi-dry electroblotting system (Pyxis Protein Transfer/Staining System). The membrane was blocked by 5% fat-free milk in TBST buffer (50 mM Tris-HCl, PH 7.6, 150 mM NaCl, 0.1% Tween 20) for 1 hr at room temperature and washed three times by TBST buffer each for 5 min at room temperature. The membrane was then incubated overnight at 4 °C with anti-GFP antibody (E385, abcam) at 1:2000 dilution in TBST buffer containing 5% fat-free milk. After three times wash with TBST buffer, the membrane was incubated with

1:10000 diluted HRP Goat anti-Rabbit IgG Antibody (LP1001a, abcam) for 1 hr at room temperature. After another three times of wash, the eGFP was detected using the eECL Western blotting kit (CW0049, CWbiotech) according to the manufacturer's instruction.

#### Acknowledgements

This work was supported by the Ministry of Science and Technology of China, grant # 2013CB530802 and 2012CB720601 and the National Science Foundation of China, grant # 21432008 and 31470783.

#### Notes and references

- 1 L. A. Cahoon and H. S. Seifert, *Science*, 2009, **325**, 764-767.
- 2 R. Rodriguez, K. M. Miller, J. V. Forment, C. R. Bradshaw, M. Nikan, S. Britton, T. Oelschlaegel, B. Xhemalce, S. Balasubramanian and S. P. Jackson, *Nat. Chem. Biol.*, 2012, **8**, 301-310.
- 3 L. T. Gray, A. C. Vallur, J. Eddy and N. Maizels, *Nat. Chem. Biol.*, 2014, **10**, 313-318.
- 4 G. H. Nguyen, W. Tang, A. I. Robles, R. P. Beyer, L. T. Gray, J. A. Welsh, A. J. Schetter, K. Kumamoto, X. W. Wang, I. D. Hickson, N. Maizels, R. J. Monnat, Jr. and C. C. Harris, *Proc. Natl. Acad. Sci. USA*, 2014, **111**, 9905-9910.
- 5 N. Maizels, *EMBO Rep.*, 2015, **16**, 910-922.
- 6 J. L. Huppert and S. Balasubramanian, *Nucleic Acids Res.*, 2005, **33**, 2908-2916.
- 7 V. S. Chambers, G. Marsico, J. M. Boutell, M. Di Antonio, G. P. Smith and S. Balasubramanian, *Nat. Biotechnol.*, 2015, **33**, 877-881.
- 8 X. M. Li, K. W. Zheng, J. Y. Zhang, H. H. Liu, Y. D. He, B. F. Yuan, Y. H. Hao and Z. Tan, *Proc. Natl. Acad. Sci. USA*, 2015, **112**, 14581-14586.
- 9 K. W. Zheng, S. Xiao, J. Q. Liu, J. Y. Zhang, Y. H. Hao and Z. Tan, *Nucleic Acids Res.*, 2013, **41**, 5533-5541.
- 10 R. Y. Wu, K. W. Zheng, J. Y. Zhang, Y. H. Hao and Z. Tan, *Angew. Chem. Int. Ed. Engl.*, 2015, **54**, 2447-2451.
- 11 K. W. Zheng, R. Y. Wu, Y. D. He, S. Xiao, J. Y. Zhang, J. Q. Liu, Y. H. Hao and Z. Tan, *Nucleic Acids Res.*, 2014, **42**, 10832-10844.
- 12 S. Xiao, J. Y. Zhang, J. Wu, R. Y. Wu, Y. Xia, K. W. Zheng, Y. H. Hao, X. Zhou and Z. Tan, *Angew. Chem. Int. Ed. Engl.*, 2014, **53**, 13110-13114.
- 13 Y. Xu, Y. Suzuki, K. Ito and M. Komiyama, *Proc. Natl. Acad. Sci. USA*, 2010, **107**, 14579-14584.
- 14 G. W. Collie, S. M. Haider, S. Neidle and G. N. Parkinson, *Nucleic Acids Res.*, 2010, **38**, 5569-5580.
- 15 A. Randall and J. D. Griffith, *J. Biol. Chem.*, 2009, **284**, 13980-13986.
- 16 H. Martadinata and A. T. Phan, *J. Am. Chem. Soc.*, 2009, **131**, 2570-2578.
- 17 Y. Xu, K. Kaminaga and M. Komiyama, *J. Am. Chem. Soc.*, 2008, **130**, 11179-11184.

- 18 P. H. Wanrooij, J. P. Uhler, T. Simonsson, M. Falkenberg and C. M. Gustafsson, *Proc. Natl. Acad. Sci. USA*, 2010, **107**, 16072-16077.
- 19 A. Bugaut and S. Balasubramanian, *Nucleic Acids Res.*, 2012, **40**, 4727-4741.
- 20 S. Millevoi, H. Moine and S. Vagner, *Wiley Interdiscip Rev RNA*, 2012, **3**, 495-507.
- 21 C. Schaffitzel, I. Berger, J. Postberg, J. Hanes, H. J. Lipps and A. Pluckthun, *Proc. Natl. Acad. Sci. USA*, 2001, **98**, 8572-8577.
- 22 G. Biffi, D. Tannahill, J. McCafferty and S. Balasubramanian, *Nat. Chem.*, 2013, **5**, 182-186.
- 23 G. Biffi, M. Di Antonio, D. Tannahill and S. Balasubramanian, *Nat. Chem.*, 2014, **6**, 75-80.
- 24 Y. P. Xing, C. Liu, X. H. Zhou and H. C. Shi, *Sci Rep*, 2015, **5**, 8125.
- 25 A. Laguerre, K. Hukezalie, P. Winckler, F. Katranji, G. Chanteloup, M. Pirrotta, J. M. Perrier-Cornet, J. M. Wong and D. Monchaud, *J. Am. Chem. Soc.*, 2015, **137**, 8521-8525.
- 26 T. K. Kerppola, *Nat. Protoc.*, 2006, **1**, 1278-1286.
- 27 S. Baskerville, M. Zapp and A. D. Ellington, *J. Virol.*, 1999, **73**, 4962-4971.
- 28 S. Lattmann, B. Giri, J. P. Vaughn, S. A. Akman and Y. Nagamine, *Nucleic Acids Res.*, 2010, **38**, 6219-6233.
- 29 S. Lattmann, M. B. Stadler, J. P. Vaughn, S. A. Akman and Y. Nagamine, *Nucleic Acids Res.*, 2011, **39**, 9390-9404.
- 30 E. P. Booy, R. Howard, O. Marushchak, E. O. Ariyo, M. Meier, S. K. Novakowski, S. R. Deo, E. Dzananovic, J. Stetefeld and S. A. McKenna, *Nucleic Acids Res.*, 2014, **42**, 3346-3361.
- 31 N. V. Hud, F. W. Smith, F. A. Anet and J. Feigon, *Biochemistry*, 1996, **35**, 15383-15390.
- 32 Y. J. Shyu and C. D. Hu, *Trends Biotechnol.*, 2008, **26**, 622-630.
- 33 B. Heddi, V. V. Cheong, H. Martadinata and A. T. Phan, *Proc. Natl. Acad. Sci. USA*, 2015, **112**, 9608-9613.
- 34 T. Gaj, C. A. Gersbach and C. F. Barbas, 3rd, *Trends Biotechnol.*, 2013, **31**, 397-405.
- 35 M. Valencia-Burton, R. M. McCullough, C. R. Cantor and N. E. Broude, *Nat. Methods*, 2007, **4**, 421-427.
- 36 V. V. Demidov and N. E. Broude, *Nat. Protoc.*, 2006, **1**, 714-719.
- 37 G. J. Smith, T. R. Sosnick, N. F. Scherer and T. Pan, *RNA*, 2005, **11**, 234-239.

Graphical and textual abstract for

## RNA G-Quadruplex Formation in Defined Sequence in Living Cells Detected by Bimolecular Fluorescence Complementation

Hong-he Liu, Ke-wei Zheng, Yi-de He, Yu-hua Hao, Quan Chen, Zheng Tan\*

State Key Laboratory of Membrane Biology, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, P. R. China

RNA G-quadruplex formation in defined sequence in living bacterial cells is detected by bimolecular fluorescence complementation with sequence and structure specificity

