

# Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



## Dehydrocoupling of Phosphine-Boranes Using the $[\text{RhCp}^*\text{Me}(\text{PMe}_3)(\text{CH}_2\text{Cl}_2)][\text{BAR}^{\text{F}}_4]$ Precatalyst: Stoichiometric and Catalytic Studies

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x


www.rsc.org/

Thomas N. Hooper<sup>a</sup>, Andrew S. Weller<sup>a\*</sup>, Nicholas A. Beattie<sup>b</sup> and Stuart A. Macgregor<sup>b\*</sup>

We report a detailed, combined experimental and computational study on the fundamental B–H and P–H bond activation steps involved in dehydrocoupling/dehydropolymerization of primary and secondary phosphine-boranes,  $\text{H}_3\text{B}\cdot\text{PPhR}'\text{H}$  ( $\text{R} = \text{Ph}, \text{H}$ ), using  $[\text{RhCp}^*(\text{PMe}_3)\text{Me}(\text{ClCH}_2\text{Cl})][\text{BAR}^{\text{F}}_4]$ , to either form polyphosphino-boranes  $[\text{H}_2\text{B}\cdot\text{PPhH}]_n$ ,  $M_n \sim 15\,000\text{ g mol}^{-1}$ ,  $\text{PDI} = 2.2$ ) or the linear diboraphosphine  $\text{H}_3\text{B}\cdot\text{PPh}_2\text{BH}_2\cdot\text{PPh}_2\text{H}$ . A likely polymer-growth pathway of reversible chain transfer step-growth is suggested for  $\text{H}_3\text{B}\cdot\text{PPh}_2\text{H}$ . Using secondary phosphine-boranes as model substrates a combined synthesis, structural (X-ray crystallography), labelling and computational approach reveals: initial bond activation pathways (B–H activation precedes P–H activation); key intermediates (phosphido-boranes,  $\alpha$ -B–agostic base-stabilized boryls); and a catalytic route to the primary diboraphosphine ( $\text{H}_3\text{B}\cdot\text{PPhHBH}_2\cdot\text{PPh}_2\text{H}$ ). It is also shown that by changing the substituent at phosphorus (Ph or Cy versus <sup>t</sup>Bu) different final products result (phosphido-borane or base stabilized phosphino-borane respectively). These studies provide detailed insight into the pathways that are operating during dehydropolymerization.

### Introduction

The polymerization of alkenes using transition metal-based catalysts to afford societally and technologically ubiquitous polyolefins is well-established, yet equivalent catalytic routes to polymeric materials containing main-group elements is considerably less developed.<sup>[1, 2]</sup> In particular, the group 13/15 mixed polymers provide one example that promises to lead to significant scientific and technological opportunities, given that polyphosphino-boranes, along with polyamino-boranes,<sup>[3]</sup> are (valence) isoelectronic with polyolefins and are finding uses in a variety of applications from lithography to pre-ceramics.<sup>[4, 5]</sup> Ill-defined polyphosphino-boranes were first synthesised in 1959 through thermal dehydrocoupling of primary phosphine-boranes,<sup>[6]</sup> but a faster and more selective dehydrocoupling/dehydropolymerisation process was reported by Manners and co-workers in the early 2000's using transition metal pre-catalysts primarily based upon  $[\text{Rh}(\text{COD})\text{Cl}]_2$  and  $[\text{Rh}(\text{COD})_2][\text{OTf}]$ , operating under melt conditions.<sup>[7–10]</sup> Others have since used similar catalyst systems to prepare related polyphosphino-boranes, or elegant demonstrations of highly selective cross-dehydrocouplings.<sup>[11, 12]</sup> For primary phosphine-boranes,  $\text{H}_3\text{B}\cdot\text{PRH}_2$ , polyphosphino-boranes are formed, whereas for secondary phosphine-




**Scheme 1.** Rh-catalyzed dehydrocoupling of primary and secondary phosphineboranes

boranes,  $\text{H}_3\text{B}\cdot\text{PR}_2\text{H}$ , linear diboraphosphines or cyclic oligomers form (Scheme 1). Although catalysis has been shown to be homogenous rather than heterogeneous,<sup>[13, 14]</sup> the melt conditions required for effective dehydrocoupling meant that resolving intermediates/resting states or kinetics was challenging. In contrast, the mechanism of amine-borane dehydrocoupling using transition metal catalysts is much better understood as catalysis can be performed in solution at room temperature.<sup>[15]</sup> Very recently the non-metal-catalyzed addition polymerization of *in situ* generated phosphino-boranes, such as  $[\text{H}_2\text{BP}^t\text{Bu}_2]$ , has been described,<sup>[16]</sup> that avoids the use of melt conditions.

Recently, *in situ* sampling using ESI-MS (electrospray ionisation mass spectrometry) led to identification of a  $[\text{Rh}(\text{PHR}_2)_2]^+$  fragment as an active species in the dehydrocoupling of secondary phosphine-boranes under melt conditions to form  $\text{H}_3\text{B}\cdot\text{PR}_2\text{H}_2\text{B}\cdot\text{PHR}_2$  when using  $[\text{Rh}(\text{COD})_2][\text{BAR}^{\text{F}}_4]$  as the precatalyst, ( $\text{R} = \text{Ph}, {}^t\text{Bu}$ ;  $\text{Ar}^{\text{F}} = 3,5\text{-(CF}_3)_2\text{C}_6\text{H}_3$ ).<sup>[17]</sup> This arises from cleavage of the relatively weak

<sup>a</sup> Department of Chemistry, Chemistry Research Laboratories, University of Oxford, Mansfield Road, Oxford, OX1 3TA, UK.


<sup>b</sup> Institute of Chemical Sciences, Heriot Watt University, Edinburgh, EH14 4AS, UK  
Electronic Supplementary Information (ESI) available: Synthesis and characterisation data, computational details. See DOI: 10.1039/x0xx00000x



**Scheme 2.** Intermediates observed in the dehydrocoupling of  $\text{H}_3\text{B}\cdot\text{PPh}_2\text{H}$  using the  $[\text{Rh}(\text{Ph}_2\text{P}(\text{CH}_2)_3\text{PPh}_2)]^+$  fragment.  $[\text{BAR}^F_4]^-$  anions not shown.

P–B bond in the substrate.<sup>[18]</sup> Simple replacement of the monodentate phosphine ligands with a bidentate phosphine produced a metal fragment, i.e.  $[\text{Rh}(\text{Ph}_2\text{P}(\text{CH}_2)_3\text{PPh}_2)]^+$ , which did not suffer from ligand redistribution, allowing for a detailed study of the mechanism, including isolation of intermediates, isotopic labelling studies and determination of activation parameters.<sup>[19, 20]</sup> Thus intermediate complexes that relate to overall P–H activation of  $\text{H}_3\text{B}\cdot\text{PPh}_2\text{H}$  at a Rh(I) center (**A** Scheme 2), and subsequent P–B bond formation (**B**), were isolated, while B–H activation of the second phosphine-borane to form a boryl intermediate was proposed to be involved in the rate-determining step that follows from **A**. However, because of relatively rapid H/D exchange between P and B the elementary P–H/B–H activation steps could not be delineated using labelling studies. In addition, although this dehydrocoupling occurred at room temperature, melt conditions were required for turnover. This same fragment was also found to dehydrocouple primary phosphine-boranes under melt conditions to produce ill-defined low molecular weight polymer. The mechanism was proposed to be the same as with secondary phosphine-boranes, but with the added complexity of diastereomer formation caused by P–H activation of the prochiral phosphorus centre.<sup>[20]</sup>

A catalytic system which does not require melt conditions, produces well-defined, high molecular weight polyphosphino-borane ( $M_n = 59\,000\text{ g mol}^{-1}$ , PDI = 1.6) and operates via a chain growth process was reported in 2015 by Manners *et al.* using the  $\text{FeCp}(\text{CO})_2(\text{OTf})$  catalyst.<sup>[5]</sup> Heating (toluene, 100°C) in the presence of phosphine-borane was required to promote CO and  $[\text{OTf}]^-$  loss and the formation of an initial phosphido-borane complex (**C**, Scheme 3, isolated for the  $\text{H}_3\text{B}\cdot\text{PPh}_2$



**Scheme 3.** (Top) Intermediates (isolated and suggested) in the dehydrocoupling of  $\text{H}_3\text{B}\cdot\text{PPh}_2$  as catalysed by  $\text{FeCp}(\text{CO})_2\text{OTf}$ . (Bottom) Relationship between  $\text{FeCp}(\text{CO})_2\text{Y}$  and  $[\text{RhCp}^*(\text{PMe}_3)\text{Y}]^+$  (Y = anionic ligand)

analogue). In the mechanism it was suggested that the Fe centre adopts a constant oxidation state with B–H/P–H activation and P–B coupling proposed (**D** and **E**), using DFT calculations, to proceed via multiple sigma-complex assisted metathesis steps.<sup>[21, 22]</sup>

Central to control of the dehydrocoupling process is a detailed understanding of the fundamental, elementary, steps that are occurring. Inspired by this recent report by Manners on the  $\text{FeCp}(\text{CO})_2(\text{OTf})$  system, and also aware that this system still required heating to promote CO loss, we turned to  $[\text{RhCp}^*\text{Me}(\text{PMe}_3)(\text{CH}_2\text{Cl}_2)]^+[\text{BAR}^F_4]^-$  (**1**, Scheme 3,  $\text{Cp}^* = \eta^5\text{-C}_5\text{Me}_5$ )<sup>[23, 24]</sup> as an alternative entry point (cf. structures **F** and **G**), proposing that B–H/P–H activation may be studied at ambient temperature under solution conditions. This complex provides a latent vacant site through  $\text{CH}_2\text{Cl}_2$  dissociation and also a methyl group that is well set up for loss as methane after B–H or P–H transfer. It is also well-established to mediate bond activation processes via sigma-bond metathesis, and related, processes,<sup>[23, 24]</sup> while the  $\{\text{RhCp}^*\}$  fragment more generally catalyzes C–H, B–H, and P–H activation and bond coupling.<sup>[25–28]</sup>


We report here that complex **1** is an effective catalyst for the dehydrocoupling of  $\text{H}_3\text{B}\cdot\text{PPh}_2$ , and also allows for a study of the elementary B–H/P–H activation processes occurring via a combined experimental and computational approach. In particular the order of B–H/P–H activation is determined in these systems, as well as a subsequent isomerisation and P–B bond forming events. This provides insight into both the order of events and the likely intermediates involved in dehydrocoupling of phosphine-boranes.

## Results and Discussion

### Catalysis: Dehydrocoupling of $\text{H}_3\text{B}\cdot\text{PPh}_2$

Initial catalytic screening showed that complex **1** was an active precatalyst (1 mol%, 0.01 M, toluene, 100 °C, 72 h, system open to Ar) for the dehydrocoupling of  $\text{H}_3\text{B}\cdot\text{PPh}_2$ . After work-up by precipitation into hexanes, the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of the resulting solid shows a well-defined peak at  $\delta -49.5$ , while in the  $^{11}\text{B}$  NMR spectrum a broad peak at  $\delta -34.0$  is observed ( $\text{CDCl}_3$ ), in good agreement with that reported by Manners *et al.* for polymer formed using the  $\text{FeCp}(\text{CO})_2(\text{OTf})$ <sup>[5]</sup> and  $[\text{Rh}(\text{COD})_2][\text{OTf}]$ <sup>[8]</sup> catalysts. The simple doublet observed in the  $^{31}\text{P}$  NMR spectrum suggests a linear  $[\text{H}_2\text{BPPH}]_n$  structure to the polymer, rather than a branched structure that would invoke a quaternary phosphorus;<sup>[8, 29]</sup> although a low intensity ill-defined broad shoulder is observed between  $\delta -50$  to  $-60$  that is suggestive of a small proportion of shorter chain oligomers or some branching. Consistent with this NMR data, the isolated polymer was shown by GPC to consist of a moderate molecular weight fraction ( $M_n = 15\,000\text{ g mol}^{-1}$ , PDI = 2.2) alongside lower molecular weight material (less than  $1\,000\text{ g mol}^{-1}$ ). Although similar to that reported for the  $[\text{Rh}(\text{COD})_2][\text{OTf}]$  catalyst ( $M_w = 30\,000\text{ g mol}^{-1}$ )<sup>[7, 8]</sup> it falls short of the  $\text{FeCp}(\text{CO})_2(\text{OTf})$  system at 1 mol% ( $M_n = 59\,000\text{ g mol}^{-1}$ ,

PDI = 1.6).<sup>[5]</sup> The organometallic species in the catalytic mixture could not be identified. However, a signal corresponding to  $\text{H}_3\text{B}\cdot\text{PMe}_3$  was observed,<sup>[30]</sup> suggesting dissociation (or substitution) of  $\text{PMe}_3$  in complex **1** during catalysis. If dehydropolymerization is carried out at a higher catalyst loading (5 mol%, 0.05 M, 72 hours) moderate molecular weight polymer is also formed as measured by GPC of hexane-precipitated material ( $M_n = 13\,000\text{ gmol}^{-1}$ , PDI = 1.5), and low molecular weight polyphosphino-borane is again present (less than  $1\,000\text{ gmol}^{-1}$ ). The isolated polymer was also analysed by ESI-MS with a broad range of molecular weight chains  $[\text{H}\{\text{PPhBH}_2\}_n\text{PPhH}_2]^+$  and clear repeat units of  $\{\text{PPhBH}_2\}$  ( $m/z = 122$ ) observed. The highest molecular weight polymer measured by this technique was  $n = 20$ ,  $m/z = 2551.9$ .




**Scheme 4.** Purified  $[\text{H}_2\text{BPPH}]_n$  from the dehydrocoupling of  $\text{H}_3\text{B}\cdot\text{PPhH}_2$  catalysed by **1**, 1 mol%. Inset (A) purified **2**; (B)  $^{11}\text{B}\{^1\text{H}\}$  NMR after 1 h: \*  $\text{H}_3\text{B}\cdot\text{PPhHBH}_2\cdot\text{PPhH}_2$  **2**, †  $\text{H}_3\text{B}\cdot\text{PPhH}_2$ , +  $\text{H}_3\text{B}\cdot\text{PMe}_3$ , # short chain oligomers.

Monitoring this reaction by  $^{11}\text{B}$  NMR spectroscopy shows that the  $\text{H}_3\text{B}\cdot\text{PPhH}_2$  monomer is consumed after only four hours, suggesting its relatively rapid oligomerization, but the slower formation of higher molecular weight polymer. If dehydropolymerization is stopped after only 1 hour the  $^{11}\text{B}\{^1\text{H}\}$  NMR spectrum now shows signals due to  $\text{H}_3\text{B}\cdot\text{PPhH}_2$ , a broad signal at  $\delta -33.6$  assigned to oligomer/polymer,  $\text{H}_3\text{B}\cdot\text{PMe}_3$  and significant amounts of a new compound assigned to the primary diboraphosphine  $\text{H}_3\text{B}\cdot\text{PPhHBH}_2\cdot\text{PPhH}_2$  **2** (Scheme 4). Compound **2** is present in significantly greater amounts at 5 mol% loading [ $\text{H}_3\text{B}\cdot\text{PPhH}_2$  : **2**; 1:1, 5 mol%; 6:1, 1 mol%], and could be isolated in 25% yield by removing the toluene *in vacuo* and extracting with hexane to give a very pale yellow oil that could be fully characterized by NMR spectroscopy [e.g.  $^{11}\text{B}\{^1\text{H}\}$   $\delta -36.5$  vt,  $J(\text{PB}) \sim 70\text{Hz}$ ;  $-38.9$  (d,  $J(\text{PB}) \sim 50\text{Hz}$ )], with data similar to both the secondary diboraphosphine

$\text{H}_3\text{B}\cdot\text{PPh}_2\text{BH}_2\cdot\text{PPh}_2\text{H}$ ,<sup>[8]</sup> and the primary analogue,  $\text{H}_3\text{B}\cdot\text{PCyHBH}_2\cdot\text{PCyH}_2$ .<sup>[20]</sup> The thermal dehydrogenation of  $\text{H}_3\text{B}\cdot\text{PPhH}_2$  in the absence of **1** (toluene, 0.625 M) produces **2** only slowly ( $\sim 50\%$  conversion after 16 hrs) alongside a small amount of oligomeric product and unreacted  $\text{H}_3\text{B}\cdot\text{PPhH}_2$ .


The lack of significant change in  $M_n$  on increasing the catalyst loading from 1 to 5 mol% suggests that a coordination chain-growth type mechanism is not operating, in which the polymer chain grows on the metal centre by successive monomer insertion events, as suggested for  $\text{FeCp}(\text{CO})_2(\text{OTf})$  system for phosphine borane and  $[\text{Rh}(\text{Xanthphos})]^+$  for amine-borane dehydropolymerization.<sup>[5, 31]</sup> Under this mechanistic model lower catalyst loadings would be expected to lead to higher molecular weight polymer, although such an analysis can be complicated by the fact that the metal has to both dehydrogenate and couple the reactive monomers.<sup>[32]</sup> Instead, that at short reaction times **2** is observed in significant quantities, especially at higher catalyst loadings, and  $\text{H}_3\text{B}\cdot\text{PPhH}_2$  is completely consumed after only 4 hours hints at a step-growth-type mechanism, as suggested for  $[\text{Rh}(\text{COD})\text{Cl}]_2$ -catalysed systems.<sup>[29]</sup> Under this regime, a greater catalyst loading might be expected to increase the molecular weight of the resulting polymer.<sup>[29, 32]</sup> However the analysis of the mechanism of polymer growth is further complicated by the fact that both isolated **2** and higher  $M_n$  polymer undergo P-B bond cleavage in the presence of **1**. For example, heating **2** in the presence of 5 mol% **1** for 1 hour (100°C, toluene) resulted in a mixture of **2**,  $\text{H}_3\text{B}\cdot\text{PPhH}_2$  (approx. 3:1 ratio by  $^{11}\text{B}\{^1\text{H}\}$  NMR spectroscopy) and signals assigned to oligomers. Further heating overnight resulted in complete consumption of **2** and  $\text{H}_3\text{B}\cdot\text{PPhH}_2$  to reveal signals in the  $^{11}\text{B}$  NMR spectrum consistent with low molecular weight polymer, Scheme 5. Heating a sample of high molecular weight polymer (100°C, toluene) with 5 mol% **1** also resulted in P-B cleavage events, with lower molecular weight species observed by  $^{31}\text{P}$  NMR spectroscopy. Linear diborazanes have also been observed to undergo B-N bond cleavage and product redistribution processes through both thermal and metal catalysed pathways, with a mixture of monomeric amine-borane and oligomeric products generated.<sup>[33]</sup>



**Scheme 5:** P-B bond cleavage and polymerisation of **2** as catalysed by **1**

On balance we thus suggest that a process in which reversible chain transfer between an oligomer (polymer) sigma-bound to a metal centre and free  $\text{H}_3\text{B}\cdot\text{PPhH}_2$ , either initially present or generated by P-B bond cleavage, accounts best for these observations. We have previously demonstrated similar behaviour (as monitored by ESI-MS) using  $\text{H}_3\text{B}\cdot\text{NH}_3$  and a  $[\text{Ir}(\text{PCy}_3)_2(\text{H})_2]^+$  fragment.<sup>[34]</sup>

#### Catalysis: Dehydrocoupling of $\text{H}_3\text{B}\cdot\text{PPhH}_2$



Scheme 6. The dehydrocoupling of  $\text{H}_3\text{B-PPh}_2\text{H}$  as catalysed by **1** and **5** to form **3**

To further probe the mechanism of dehydrocoupling using **1** the secondary phosphine-borane  $\text{H}_3\text{B-PPh}_2\text{H}$  was used, which has been shown to afford the diboraphosphine  $\text{H}_3\text{B-PPh}_2\text{BH}_2\text{-PPh}_2\text{H}$  **3** or cyclic species depending on dehydrocoupling conditions.<sup>[8]</sup> Treatment of pre-catalyst **1** (5 mol%, 0.0313 M,  $100^\circ\text{C}$ , toluene, 16 h) with  $\text{H}_3\text{B-PPh}_2\text{H}$  resulted in almost full conversion to **3** (95% by  $^{31}\text{P}$  and  $^{11}\text{B}$  NMR spectroscopy), Scheme 6. Analysis of the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum post-catalysis showed one dominant phosphine-containing organometallic species, as a doublet at  $\delta$  26.7 [ $J(\text{RhP})$  139 Hz] which splits into a doublet of doublets in the  $^{31}\text{P}$  NMR spectrum [ $J(\text{PH})$  391 Hz], demonstrating a direct P-H bond.  $\text{H}_3\text{B-PMe}_3$  was also observed to be formed. The  $^1\text{H}$  NMR spectrum of the reaction mixture showed a doublet of triplets at  $\delta$  -11.36 which simplified to a doublet upon  $^{31}\text{P}$  decoupling, suggesting a rhodium-bound hydride coupling to two phosphorus centres. ESI-MS showed one dominant peak at  $m/z$  = 611.15, with an isotope pattern that corresponds to the cation  $[\text{RhCp}^*(\text{H})(\text{PPh}_2)_2]^+$ , **4**<sup>+</sup>, fully consistent with the NMR data. Species closely related to cationic **4**<sup>+</sup> have been previously structurally characterised.<sup>[35, 36]</sup> Addition of Hg to the catalytic mixture after 4 hours resulted in no significant change to the overall conversion or rate, suggesting that the catalyst is not colloidal.<sup>[14]</sup>


The diphenylphosphine ligands required for the formation of cation **4**<sup>+</sup> likely result from P-B cleavage of the starting material  $\text{H}_3\text{B-PPh}_2\text{H}$  and resulting exchange at the metal centre to release  $\text{PMe}_3$ , which is trapped as  $\text{H}_3\text{B-PMe}_3$ . Following the temporal evolution of catalysis using  $^{31}\text{P}\{^1\text{H}\}$  NMR spectroscopy and ESI-MS<sup>[37]</sup> showed that after 1 hour **4**<sup>+</sup> was present, but also a pair of doublet of doublet resonances at  $\delta$  19.2 and 2.3 were observed, that correlate with signals in the ESI-MS spectrum assigned to the cation  $[\text{RhCp}^*(\text{H})(\text{PMe}_3)(\text{PPh}_2)_2]^+$  (**5**<sup>+</sup>). After 4 hours at  $100^\circ\text{C}$  complex **4**<sup>+</sup> was dominant, suggesting that the cation **5**<sup>+</sup> evolves to give **4**<sup>+</sup> during catalysis. The ESI-MS also revealed signals with isotopic patterns which correspond to  $[\text{RhCp}^*(\text{PPh}_2\text{-BH}_3)(\text{PPh}_2)_2]^+$  (at  $m/z$  = 809.23) and  $[\text{RhCp}^*(\text{PPh}_2\text{-BH}_2\text{PPh}_2\text{-BH}_3)(\text{PPh}_2)_2]^+$  ( $m/z$  = 1007.31) which we assume are Rh-P bound (*vide infra*). Phosphido-borane species have been detected and proposed as catalytic intermediates in phosphine-borane dehydrocoupling in systems based on the  $\{\text{Rh}(\text{Ph}_2\text{P}(\text{CH}_2)_3\text{PPh}_2)\}^+$  and  $\{\text{FeCp}(\text{CO})\}^+$

fragments.<sup>[5, 19, 20]</sup> Addition of a further 20 equivalents of  $\text{H}_3\text{B-PPh}_2\text{H}$  to this reaction mixture post catalysis and heating to  $100^\circ\text{C}$  resulted in complete conversion to diboraphosphine **3** after 22 hrs, suggesting that cation **4**<sup>+</sup> is active in catalysis. Further evidence for complexes of general formula  $[\text{RhCp}^*(\text{H})(\text{PR}_3)_2]^+$  being the active species comes from the isolation of **5** as pure material as the  $[\text{BAR}_4^{\text{F}}]^-$  salt (*vide infra*). Complex **5** is also a competent pre-catalyst for the dehydrocoupling of  $\text{H}_3\text{B-PPh}_2\text{H}$  (5 mol%,  $100^\circ\text{C}$ ) reaching completion within 22 hours. Again, cation **4**<sup>+</sup> is observed to be formed in the reaction mixture by  $^{31}\text{P}$  NMR spectroscopy, and the associated release of  $\text{PMe}_3$  was confirmed by the detection of  $\text{H}_3\text{B-PMe}_3$ . Addition of  $\text{PPh}_3$  (10 equivalents) to complex **5** and monitoring by ESI-MS shows, after 2 hours at 298 K, the formation of  $[\text{RhCp}^*(\text{H})(\text{PMe}_3)(\text{PPh}_3)]^+$  ( $m/z$  = 577.17) showing that phosphine exchange also occurs at 298 K. At room temperature, neither *in situ* generated **4**, or pure **5**, displayed any reactivity towards one equivalent of  $\text{H}_3\text{B-PPh}_2\text{H}$ . This suggests that under these conditions phosphine-borane is not a competitive ligand with phosphine, requiring higher temperatures and a large excess to promote reactivity at the metal center when there are two phosphines bound. The generation of vacant sites has been suggested to be important in the mode of action of  $\text{FeCp}(\text{CO})_2(\text{OTf})$  in dehydrocoupling.<sup>[5]</sup> Consistent this we show next that **1**, which is a masked source of  $\{\text{RhCp}^*\text{Me}(\text{PMe}_3)\}^+$  and thus does not require phosphine dissociation, reacts very rapidly with  $\text{H}_3\text{B-PPh}_2\text{H}$ .

Overall these data show that the  $\{\text{RhCp}^*\text{Me}(\text{PMe}_3)\}^+$  pre-catalyst, and related species formed during catalysis such as cation **4**<sup>+</sup>, are implicated in the dehydrocoupling/dehydropolymerization of both primary and secondary phosphine-boranes. In order to determine the role the metal fragment plays in this stoichiometric reactivity was studied, as is described next.

#### Stoichiometric Reactivity with $\text{H}_3\text{B-PPh}_2\text{H}$


Reaction of 1 equivalent of  $\text{H}_3\text{B-PPh}_2\text{H}$  with **1** at room temperature in  $\text{CD}_2\text{Cl}_2$  solution resulted in rapid effervescence and a colour change from orange to yellow.  $^{31}\text{P}\{^1\text{H}\}$  NMR spectroscopy of the resulting solution showed one sharp doublet of doublets at  $\delta$  -6.6 [ $J(\text{RhP})$  = 139 Hz,  $J(\text{PP})$  = 22 Hz] assigned to the  $\text{PMe}_3$  ligand and one broad peak at  $\delta$  6.9 [ $fwhm$  = 222 Hz] assigned to a phosphine-borane moiety, which was essentially unchanged in line shape in the  $^{31}\text{P}$  NMR spectrum. The  $^1\text{H}$  NMR spectrum demonstrated a lack of P-H and Rh-Me signals, and dissolved  $\text{CH}_4$  was detected [ $\delta$  0.21<sup>[38]</sup>]. A very broad peak was observed at  $\delta$  0.3 (relative integral 2H) which sharpens on  $^{11}\text{B}$  decoupling and splits into



Scheme 7. Complex **6**.  $[\text{BAR}_4^{\text{F}}]^-$  anions are not shown.


two distinct resonances at  $\delta$  0.49 and  $\delta$  -0.03 in a 1:1 ratio. A broad peak is observed at  $\delta$  -10.81 that also sharpens on decoupling  $^{11}\text{B}$ , under which conditions it also resolves into a broad doublet of doublets. These 3 upfield resonances are assigned to a  $\text{BH}_3$  unit binding to the metal centre through one  $\text{Rh}-\text{H}-\text{B}$  3 centre-2 electron bond that is not undergoing exchange on the NMR timescale between terminal and bridging environments. In the  $^{11}\text{B}$  NMR spectrum a signal at  $\delta$  -45.5 was observed, shifted slightly upfield from  $\text{H}_3\text{B}-\text{PPh}_2\text{H}$  [ $\delta$  -40.1]. Overall, these data are consistent with the formation of a phosphido-borane complex which also has a rather tight  $\beta$ -B-agostic interaction:  $[\text{RhCp}^*(\text{PPh}_2\text{BH}_3)(\text{PMe}_3)][\text{BAR}^{\text{F}}_4]$  (**6**), Scheme 7.

Yellow crystals were grown from the reaction mixture and isolated in good yield (76%). A resulting single-crystal X-ray diffraction study (Figure 1) confirmed the structure as a phosphido-borane species with a  $\beta$ -B-agostic interaction. Although the B-H hydrogen atoms were located in the difference map, in the final refinement they were placed at



**Figure 1:** X-ray molecular structure of  $[\text{RhCp}^*(\text{PPh}_2\text{BH}_3)(\text{PMe}_3)][\text{BAR}^{\text{F}}_4]$  **6**.  $[\text{BAR}^{\text{F}}_4]$  anion and selected hydrogen atoms omitted for clarity. Ellipsoids shown at 50% probability. Selected bond lengths (Å) and angles ( $^\circ$ ): P(1)-B(1) 1.896(4), Rh(1)-P(1) 2.302(1), Rh(1)-B(1) 2.464(4), Rh(1)-P(2) 2.3241(10), Rh(1)-Cp\*(centroid) 1.859, P(1)-Rh(1)-P(2) 95.35(3), Rh(1)-P(1)-B(1) 71.13(13), B(1)-P(1)-C(11) 116.09(19), B(1)-P(1)-C(17) 119.55(19), C(11)-P(1)-C(17) 103.34(15).

fixed positions. The P-B distance in **6** [1.896(4) Å] is slightly shorter than the reported P-B bonds in  $\text{H}_3\text{B}-\text{P}(\text{Mes})_2\text{H}$  [1.938 Å]<sup>[39]</sup> and in  $\text{H}_3\text{B}-\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_2\text{H}$  [1.917(2) Å]<sup>[10]</sup> (the structure of  $\text{H}_3\text{B}-\text{PPh}_2$  has not been reported) but longer than most of the crystallographically characterised monomeric phosphino-boranes, which usually bear bulky substituents to prevent oligomerisation (1.76–1.88 Å).<sup>[40, 41]</sup> The NMR data are also characteristic of a four-coordinate boron, indicating a  $\beta$ -B-agostic structure rather a phosphino-borane complex with



**Scheme 8.** Reactivity of complex **6**.  $[\text{BAR}^{\text{F}}_4]$  anions are not shown


concomitant hydride transfer to Rh. Further evidence for a  $\beta$ -B-agostic structure was obtained from DFT calculations<sup>[42]</sup> which revealed a significant lengthening of the agostic B(1)-H(1A) bond (1.39 Å) compared to the non-agostic B(1)-H(1B)/H(1C) bonds (both 1.21 Å), as well as a short  $\text{Rh}(1)\cdots\text{H}(1\text{A})$  contact of 1.72 Å. Other heavy atom bond metrics were in good agreement with experiment, including a computed P(1)-B(1) distance of 1.92 Å (see ESI for full details).  $\beta$ -B-agostic interactions of this type have been previously observed in phosphido-borane complexes with Mo,<sup>[43, 44]</sup> Fe,<sup>[45]</sup> Ti,<sup>[46]</sup> Rh<sup>[20]</sup> and alkaline earth metals,<sup>[47-49]</sup> but the structure of **6**, and the salient NMR data, most closely resemble the neutral compound  $[\text{FeCp}(\text{PPh}_2\text{BH}_3)(\text{CO})]$ .<sup>[5]</sup> Finally, the  $\beta$ -B-agostic interaction observed in **6** is in contrast with valence isoelectronic  $[\text{RhCp}^*(\text{H})(\text{H}_2\text{C}=\text{CH}_2)\text{P}(\text{OMe})_3][\text{BF}_4]$  that although in equilibrium with the corresponding  $\beta$ -agostic complex, favours the former.<sup>[50]</sup> Complex **6** is stable in  $\text{CD}_2\text{Cl}_2$  solution for at least 2 weeks.

The  $\beta$ -B-agostic interaction in **6** could be viewed as a source of masked highly reactive, phosphino-borane i.e.  $\{\text{H}_2\text{BPPh}_2\} / \{\text{Cp}^*\text{RhH}(\text{PMe}_3)\}^+$  in which Rh-H acts as a Lewis base to boron and phosphorus a Lewis base to the Rh-center. The parent  $\text{H}_2\text{BPPh}_2$  has been shown to oligomerise at [Ti] centres,<sup>[51-53]</sup> or form polymeric materials when generated *in situ*.<sup>[16]</sup> To explore whether phosphino-borane  $\text{H}_2\text{BPPh}_2$  could be liberated, as signalled by the formation of  $[\text{Ph}_2\text{PBH}_2]_n$  ( $n = 3$  or 4),<sup>[8, 16]</sup> complex **6** was heated to 100 °C in toluene for 4 hours. However, the only product that could be observed by NMR spectroscopy was the P-B cleavage product **5**, while the fate of the remaining  $\{\text{BH}\}$  is unclear (Scheme 8). This process is therefore the likely route to formation of **5** from **1** under catalytic conditions. Complex **5** could also be formed cleanly by pressurising a 1,2-difluorobenzene solution of **6** with  $\text{H}_2$  (~ 4 atm) at room temperature for 16 hours. In this case the boron-containing by-product of P-B cleavage was determined to be  $\text{B}_2\text{H}_6$  by  $^{11}\text{B}$  NMR spectroscopy.<sup>[54]</sup> Complex **6** does not react with  $\text{H}_3\text{B}-\text{PPh}_2\text{H}$  at 298 K, reflecting the strong  $\text{Rh}\cdots\text{H}-\text{B}$  interaction.

### Stoichiometric Reactivity with $\text{H}_3\text{B}-\text{PCy}_2\text{H}$

Reaction of one equivalent of  $\text{H}_3\text{B}-\text{PCy}_2\text{H}$  with **1** in  $\text{CD}_2\text{Cl}_2$  resulted in rapid effervescence (methane). Analysis by NMR spectroscopy after 5 minutes indicated the formation of a complex very similar to **6**:  $[\text{RhCp}^*(\text{PCy}_2\text{BH}_3)(\text{PMe}_3)][\text{BAR}^{\text{F}}_4]$ , **7**, in particular an upfield signal in the  $^1\text{H}$  NMR spectrum is observed at  $\delta$  -11.42, assigned to the  $\beta$ -B-agostic interaction. Single crystals of **7** suitable for X-ray diffraction were grown from a cooled  $\text{CH}_2\text{Cl}_2$ /pentane solution, and the solid state structure confirms a  $\beta$ -B-agostic phosphidoborane ligand chelating with the rhodium centre (Figure 2). The bond lengths and angles in the structure were broadly similar to those found in **6**, and this was also borne out when comparing the DFT-optimised structures (ESI). In contrast to complex **6**, **7** is not stable in  $\text{CD}_2\text{Cl}_2$  solution, decomposing fully after approximately 24 hours to form a mixture, from which the major component could be characterised spectroscopically as  $[\text{RhCp}^*(\text{H})(\text{PCy}_2\text{H})(\text{PMe}_3)][\text{BAR}^{\text{F}}_4]$  **8**, i.e. the analogue of **5**. This


low temperature instability to P–B cleavage can be contrasted with **6**, that only decomposes upon heating. P–B bond cleavage in phosphine–borane complexes has previously been noted to be a function of both the electron withdrawing nature and the steric bulk of the P–substituents, the latter suggested to be dominating here.<sup>[3, 20]</sup>



**Figure 2:** X-ray molecular structure of  $[\text{RhCp}^*(\text{PCy}_2\text{BH}_3)(\text{PMe}_3)][\text{BAR}_4^{\text{F}}]$  (**7**).  $[\text{BAR}_4^{\text{F}}]$  anion and selected hydrogen atoms omitted for clarity. Ellipsoids shown at 50% probability. Selected bond lengths (Å) and angles ( $^\circ$ ): P(1)–B(1) 1.910(7), Rh(1)–P(1) 2.3425(14), Rh(1)–B(1) 2.468(7), Rh(1)–P(2) 2.2878(16), Rh(1)–Cp\*(centroid) 1.875, P(1)–Rh(1)–P(2) 93.94(6), Rh(1)–P(1)–B(1) 70.1(2), B(1)–P(1)–C(11) 109.3(3), B(1)–P(1)–C(17) 118.3(3), C(11)–P(1)–C(17) 110.0(2).

### Stoichiometric reactivity of $\text{H}_3\text{B}\cdot\text{P}^t\text{Bu}_2\text{H}$


One equivalent of  $\text{H}_3\text{B}\cdot\text{P}^t\text{Bu}_2\text{H}$  was added to complex **1** to explore further the effect of increasing the steric bulk at the phosphorus center. After mixing, the yellow solution rapidly turned dark red and effervescence was observed. Over the course of two hours at 298 K this intense colour was lost to give a yellow/orange solution. Analysis by  $^{31}\text{P}\{^1\text{H}\}$  NMR spectroscopy of this final solution showed two broad peaks at  $\delta$  54.8 and  $-7.8$ , alongside minor unidentified species. The  $^1\text{H}$  NMR spectrum showed two resonances in the hydride region at  $\delta$   $-10.79$  and  $-13.76$  (the former being considerably broader but sharpened on decoupling  $^{11}\text{B}$ ) which, in contrast to **6** and **7**, suggest the presence of *both* Rh–H–B and Rh–H groups respectively. A broad peak at  $\delta$  0.50 (BH, integral 1 H) was also observed, in addition to phosphine and Cp\* resonances. Moreover the  $^{11}\text{B}\{^1\text{H}\}$  NMR spectrum revealed a broad virtual triplet at  $\delta$   $-45.4$  [ $J(\text{BP}) \approx 95$  Hz] suggestive of coupling to two phosphorus centres. The structure of this new species was resolved by a single-crystal X-ray diffraction study (Figure 3) to be  $[\text{RhCp}^*(\text{H})(\text{P}^t\text{Bu}_2\text{BH}_2\cdot\text{PMe}_3)][\text{BAR}_4^{\text{F}}]$  **9**, in which the  $\text{PMe}_3$  ligand has migrated to the boron centre to afford a Lewis base stabilised phosphino-borane, chelating to the rhodium centre through  $\text{P}^t\text{Bu}_2$  and a  $\beta$ -B-agostic interaction. The  $\text{P}^t\text{Bu}_2$  unit is disordered over two sites meaning that the P–B bond metrics cannot be discussed in detail, but it is similar to those observed in the phosphido-borane species **6** and **7**, suggesting a single P–B bond. DFT calculations on **9** provide optimised P(1A)–B(1) and P(2)–B(1) distances of 1.95 Å and 1.96 Å, respectively, consistent with single bond character. Lewis–base stabilised phosphino-boranes were first synthesised by Burg in 1978,<sup>[55]</sup> and have recently been used by Scheer and coworkers to form metal complexes<sup>[16, 51, 53]</sup> that can also undergo P–B coupling reactions.<sup>[52]</sup> Similar phosphine ligand migration to a boron centre in a transient phosphino-borane has been previously proposed in the formation of  $[\text{Rh}(\text{PPh}_3)_2(\text{PPh}_2\text{BH}_2\cdot\text{PPh}_3)][\text{BAR}_4^{\text{F}}]$ <sup>[56]</sup> which also has a Lewis base-stabilised phosphino-borane with a  $\beta$ -B-agostic



**Figure 3:** X-ray molecular structure of (**9**). The  $\text{P}^t\text{Bu}_2$  unit is disordered over 2 sites, only the major component labelled, i.e. P(1A), C(11A), is shown.  $[\text{BAR}_4^{\text{F}}]$  anion and selected hydrogen atoms omitted for clarity. Selected bond lengths (Å) and angles ( $^\circ$ ): P(1A)–B(1) 1.99(2), P(1B)–B(1) 1.901(14), B(1)–P(2) 1.918(5), Rh(1)–P(1A) 2.30(3), Rh(1)–P(1B) 2.258(14), Rh(1)–B(1) 2.431(5), Rh(1)–Cp\*(centroid) 1.870; P(1A)–B(1)–P(2) 126.2(7).

interaction to the Rh(I) centre [Rh–B: 2.407(5); B–P: 1.915(5), 1.945(5) Å].

A low temperature NMR spectroscopy study was performed to help elucidate the mechanism by which **9** is formed, and in particular the identity of the observed dark red intermediate.  $\text{CD}_2\text{Cl}_2$  solutions of  $\text{H}_3\text{B}\cdot\text{P}^t\text{Bu}_2\text{H}$  and **1** were combined at  $-78^\circ\text{C}$  to form a yellow solution after mixing. After loading into a precooled NMR spectrometer the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum at  $-80^\circ\text{C}$  showed a new species by a sharp doublet  $\delta$  8.3 and a broad signal  $\delta$  35.4, consistent with Rh– $\text{PMe}_3$  and  $\text{H}_3\text{B}\cdot\text{PR}_2\text{H}$  environments respectively. The  $^1\text{H}$  NMR spectrum was more revealing with a very broad upfield peak observed at  $\delta$   $-4.01$  (3 H relative integral) consistent with a Rh $\cdots\text{H}_3\text{B}$  unit. A broad signal was also observed at  $\delta$  0.79 (3 H relative integral), assigned to Rh–Me. The P–H bond is still intact, as shown by a doublet at  $\delta$  4.08 [ $J(\text{HP})$  363 Hz] which collapsed to a singlet on  $^{31}\text{P}$  decoupling. These data suggest that this species is an  $\eta^1$ -sigma complex with the bound dichloromethane molecule of **1** replaced by the phosphino-borane to form  $[\text{RhCp}^*\text{Me}(\text{PMe}_3)(\eta^1\text{-H}_3\text{B}\cdot\text{P}^t\text{Bu}_2\text{H})][\text{BAR}_4^{\text{F}}]$ , **10**, Scheme 9 That only one B–H environment is observed, even at  $-80^\circ\text{C}$ , suggests rapid terminal/bridging B–H exchange on the NMR timescale.  $\eta^1$ -sigma binding with a variety of metal-ligand fragments has been observed for both phosphine- and amine-boranes, with low-energy exchange between bridging and terminal B–H sites observed on the NMR timescale.<sup>[57–60]</sup> The  $^{11}\text{B}\{^1\text{H}\}$  NMR spectrum shows a chemical shift at  $\delta$   $-44.8$ , characteristic<sup>[61]</sup> of an  $\eta^1$ -M $\cdots\text{H}_3\text{B}\cdot\text{PR}_3$  interaction, being barely shifted from free phosphine-borane ( $\delta$   $-42.9$ ).




**Scheme 9.** Formation of complex (9). Observed and proposed intermediates.  $[\text{BAR}^{\text{F}}_4]^-$  anions are not shown.

When this solution was warmed to  $-40^\circ\text{C}$  inside the spectrometer after approximately one hour a new species, **11**, was formed at the expense of complex **10**. The  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum showed two new resonances at  $\delta$  25.1 and  $-1.9$ , as a broad peak and a sharp doublet respectively. The  $^1\text{H}$  NMR spectrum revealed the disappearance of the Rh-Me signal with concomitant appearance of dissolved  $\text{CH}_4$  ( $\delta$  0.15).<sup>[38]</sup> Two broad peaks (both 1 H relative integral) at  $\delta$  7.1 and  $\delta$   $-12.76$  [ $d, J(\text{RhH}) = 38$  Hz] were observed, both of which sharpen on decoupling  $^{11}\text{B}$ , and a doublet of multiplets at  $\delta$  4.68 [ $J(\text{RhP})$  380 Hz], consistent with a P-H group. In the  $^{11}\text{B}\{^1\text{H}\}$  NMR spectrum there is a peak at  $\delta$  47.6, downfield shifted by 92.4 ppm compared to **10**. These data suggest that **11** corresponds to a base-stabilised boryl complex,  $[\text{RhCp}^*(\text{PMe}_3)(\text{H}_2\text{B-P}^t\text{Bu}_2\text{H})][\text{BAR}^{\text{F}}_4]$ , featuring a strong  $\alpha$ -B-agostic interaction, as the two, now diastereotopic, B-H groups do not undergo exchange.

As far as we are aware there is only one other reported base-stabilised  $\alpha$ -B-agostic boryl complex albeit a dimeric motif,<sup>[62]</sup> although examples that may be described as having  $\alpha$ -B-agostic amino-boryl limiting structures have been discussed.<sup>[63, 64]</sup> DFT calculations on the dehydrogenation of  $\text{H}_3\text{B-NMe}_2\text{H}$  using the  $\{\text{Ir}(\text{PCy}_3)_2(\text{H})_2\}^+$  fragment suggest intermediates with structures closely related to **11**.<sup>[65]</sup> Similar B-H activation and elimination of methane (under photolytic conditions) has been reported by Shimoi and co-workers to form  $\text{M}(\eta^5\text{-C}_5\text{R}_5)(\text{CO})_n(\text{BH}_2\text{-PMe}_3)$  [ $n = 2, \text{M} = \text{Mn}; n = 3, \text{M} = \text{W}, \text{Mo}, \text{R} = \text{H}, \text{Me}$ ] from the corresponding metal methyl precursors.<sup>[66, 67]</sup> Interestingly these, and other closely related complexes,<sup>[68, 69]</sup> only show small (ca. 13 ppm) downfield shifts, when compared to free  $\text{H}_3\text{B-PMe}_3$ , on formation of the boryl moiety, in contrast to the ca. 92 ppm shift observed between **10** and **11**. In fact the  $^{11}\text{B}$  chemical shift is more similar to complexes featuring 3-coordinate boron (e.g.  $\delta$  30–50).<sup>[63, 70–72]</sup> The  $^1\text{H}$  NMR spectrum of **11** shows a large  $J(\text{RhH})$  coupling in the low field hydride-like signal [ $J(\text{RhH})$  38 Hz], whereas in complexes **6** and **7** no such coupling is observed. Moreover the other BH group resonates at rather low field ( $\delta$  7.11), compared with **6** ( $\delta$  0.49 and  $-0.03$ ). In comparison, Shimoi's  $\text{M}(\eta^5\text{-$

$\text{C}_5\text{R}_5)(\text{CO})_n(\text{BH}_2\text{-PMe}_3)$  species (which do not feature an  $\alpha$ -B-agostic interaction) exhibit BH chemical shifts around 1.5,<sup>[66]</sup> whereas hydrido-amino-boryls  $\text{Ir}(\text{PMe}_3)_3(\text{H})\text{Cl}\{\text{B}(\text{H})(\text{NCy}_2)\}^{[70]}$  and  $[\text{Rh}(\kappa^3\text{-P}_{\text{O},\text{P}}\text{-Xantphos})(\text{H})\{\text{B}(\text{H})(\text{N}^i\text{Pr}_2)\}(\text{NCMe})][\text{BAR}^{\text{F}}_4]^{[31]}$  (featuring 3-coordinate boron) show B-H and  $^{11}\text{B}$  chemical shifts more like **11** [ $\delta^{11}\text{B}$  43, 49 respectively]. These data suggest that complex **11** could also be described as a hydrido base-stabilised borylene complex, at least in a limiting form. However, it is also possible that a tight  $\alpha$ -B-agostic interaction could induce a downfield shift in the  $^{11}\text{B}$  NMR spectrum, similar to  $\alpha$ -C-agostic interactions probed by  $^{13}\text{C}$  NMR spectroscopy.<sup>[73]</sup>


In an attempt to resolve this structural ambiguity, dark red single crystals of **11** were grown at  $-20^\circ\text{C}$ , however the resulting structure was of poor quality and only showed the connectivity of the heavy atoms that demonstrate a Rh-B interaction (see ESI). Instead both limiting forms were characterized via DFT calculations which revealed the  $\alpha$ -B-agostic boryl (**11**) to lie 2.1 kcal/mol below the hydrido base-stabilised borylene complex (**11'**, see Figure 4).<sup>[74, 75]</sup> This preference was reproduced with a range of other functionals. A third form, **11''**, featuring an agostic interaction with one  $^t\text{Bu}$  C-H bond was also located and was 5.4 kcal/mol above **11** (see ESI). Computed barriers suggest rapid interconversion between all three species, with **11** being the dominant species in solution. The computed structure of **11** exhibits a strong  $\alpha$ -B-agostic interaction, with a short  $\text{Rh}\cdots\text{H}^1$  contact of 1.79 Å and significant elongation of the  $\text{B}^1\text{-H}^1$  bond (1.35 Å) compared to the terminal  $\text{B}^1\text{-H}^2$  bond (1.22 Å). Further support for the  $\alpha$ -B-agostic assignment was seen in the computed  $^{11}\text{B}$  chemical shifts, the value for **11** ( $\delta$  53.7 ppm) being both in good absolute agreement with experiment ( $\delta$  47.6) and significantly better than that computed for **11'** ( $\delta$



**Figure 4:** (a) Computed isomers and interconversions of  $[\text{RhCp}^*(\text{PMe}_3)(\text{H}_2\text{B-PH}^t\text{Bu}_2)]^+$ ; (b) Computed structures of  $\alpha$ -B-agostic boryl complex **11** and hydrido base-stabilised borylene complex, **11'**. Selected distances are in Å and C-bound H atoms are omitted for clarity. Free energies are quoted relative to **11** set to 0.0 kcal/mol and are at the BP86-D3( $\text{CH}_2\text{Cl}_2$ ) level; computed  $^{11}\text{B}$  chemical shifts are at the B3LYP(BS2)//BP86(BS1) level (see ESI for full details).

119.3 ppm).

Removal of the NMR tube from the spectrometer while at low temperature showed complex **11** to be responsible for the intermediate deep red colour observed. Warming to room temperature over two hours produced the yellow/orange solution in which **9** was the major product (Scheme 9). The formation of complex **9** was signalled in the  $^{11}\text{B}$  NMR spectrum by a dramatic upfield shift to  $\delta -45.4$  (computed value =  $-49.1$ ). Complex **9** forms from **11** by P–H activation and migration of the  $\text{PMe}_3$  ligand to the boron centre. We suggest that this may occur via a phosphino-borane intermediate (**H**, Scheme 9) that then undergoes intramolecular attack by  $\text{PMe}_3$ . A structural analogue of **H** has been reported by Bourissou and co-workers in  $[\text{Cy}_2\text{PB}(\text{C}_6\text{F}_5)_2\text{Pt}(\text{PMe}_3)_2]$ .<sup>[76]</sup>




**Figure 5:** Computed free energy reaction profile (kcal/mol, BP86-D3( $\text{CH}_2\text{Cl}_2$ ) level) for formation of **9** from **10** ( $\text{R} = \text{tBu}$ ) with equivalent data for  $\text{R} = \text{Ph}$  provided in italics. All free energies are quoted relative to **10** + free  $\text{H}_3\text{B-PHR}_2$  at 0.0 kcal/mol; see Figure 4 for details of species **11**, **11'** and **11''** when  $\text{R} = \text{tBu}$ .

DFT calculations were employed to assess this proposed mechanism and the results are summarised in Figure 5 (which also presents data for the analogous reaction of  $\text{H}_3\text{B-PPH}_2$  that will be discussed below). Starting from species **10** (set to 0.0 kcal/mol) B–H activation involves a sigma-CAM process<sup>[21]</sup> via **TS(10-11'')** ( $G = +14.1$  kcal/mol) to generate intermediate **Int(10-11'')** ( $G = +6.9$  kcal/mol) featuring both phosphine-stabilised boryl and methane ligands. **TS(10-11'')** exhibits a short  $\text{Rh-H}^3$  distance of 1.61 Å, indicative of significant  $\text{Rh(V)}$  character at this point (see Figure 6(a) which also gives the labelling scheme employed). Facile loss of  $\text{CH}_4$  initially yields the C–H agostic species **11''** ( $G = -1.6$  kcal/mol) which readily isomerises to **11** at  $-7.0$  kcal/mol.

The onward reaction of **11** requires an initial rearrangement back to **11''**. This proves to be necessary as it swaps the strong  $\alpha$ -B-agostic interaction in **11** for a weak C–H agostic in **11''** which then allows the transfer of  $\text{H}^4$  from  $\text{P}^1$  to Rh via **TS(11''-9')1** ( $G = +17.2$  kcal/mol). The intermediate generated, **Int(11''-9')** ( $G = -4.0$  kcal/mol, Figure 6(b)), features a  $\{\text{tBu}_2\text{PBH}_2\}$  phosphino-borane moiety and is equivalent to the postulated intermediate **H** of Scheme 8. **Int(11''-9')** exhibits a  $\text{P}^1\text{-B}^1$  distance of 1.87 Å, lying between the computed B–P distances of  $\text{H}_3\text{B-P}^t\text{Bu}_2\text{H}$  (1.96 Å) and  $\text{H}_2\text{B=P}^t\text{Bu}_2$  (1.83 Å), see ESI. This suggests a degree of back-bonding from the metal to

the phosphino-borane, but perhaps less than is implied in




**Figure 6:** Computed structures and free energies (BP86-D3( $\text{CH}_2\text{Cl}_2$ )) for (a) **TS(10-11'')** and (b) **Int(11''-9')**; selected distances are in Å and C-bound H atoms are omitted for clarity.

$[\text{Cy}_2\text{PB}(\text{C}_6\text{F}_5)_2\text{Pt}(\text{PMe}_3)_2]$ ,<sup>[76]</sup> for which a  $\text{P-B}$  distance of 1.917(3) Å has been determined crystallographically. It is also notable that the hydride and  $\{\text{BH}_2\}$  unit in **Int(11''-9')** are orientated *trans*, while the  $\text{PMe}_3$  and  $\text{BH}_2$  are *cis*. Thus  $\text{B}^1\text{-P}^2$  coupling can occur via **TS(11''-9')2** with a modest barrier of only  $+11.7$  kcal/mol to give **9'**, which is related to the observed species **9** ( $G = -16.9$  kcal/mol) via rotation about the new B– $\text{PMe}_3$  bond. The overall barrier for the formation of **9** from **11** is 24.2 kcal/mol, and so is somewhat higher than that for the formation of **11** from **10** (14.1 kcal/mol). These relative barriers are qualitatively consistent with the rapid formation of **11** at low temperature, compared to the onwards slower generation of **9** (room temperature, 2 hours). The higher barrier for P–H activation (from **11**), compared to the initial B–H activation (from **10**) is also consistent with previous experimental and computational studies on related amine-borane chemistry,<sup>[65, 77]</sup> and for  $\text{H}_3\text{B-P}^t\text{Bu}_2\text{H}$  dehydrocoupling using the  $[\text{Rh}(\text{Ph}_2\text{P}(\text{CH}_2)_3\text{PPh}_2)]^+$  fragment.<sup>[19]</sup>

### Reactions with $\text{H}_3\text{BPCy}_3$

In an attempt to produce a stable boryl complex,  $\text{H}_3\text{B-PCy}_3$  was reacted with **1**, in the anticipation that the lack of a P–H group would stop onward reactivity. Reaction formed a deep red phosphine-boryl complex which was characterised




**Scheme 10.** Spectroscopically observed boryl complex and reactivity with  $\text{H}_2$ .  $[\text{BAR}_4^-]$  anions are not shown.

spectroscopically as  $[\text{RhCp}^*(\text{PMe}_3)(\text{H}_2\text{B-PCy}_3)][\text{BAR}^{\text{F}}_4]$ , **12**, which was stable at room temperature for 4 hours before any decomposition (to unidentified products) was observed (Scheme 10). The NMR spectra of complex **12** are very similar to **11**. In particular in the  $^1\text{H}$  NMR spectrum a broad upfield peak at  $\delta -13.57$  is observed,<sup>[78]</sup> along with the characteristic downfield shift of the  $^{11}\text{B}$  NMR resonance ( $\delta 53.0$ ). Attempts to crystallise **12** resulted in intractable oils. Addition of  $\text{H}_2$  (4 atm) to **12** resulted in loss of the deep red colour to form an orange/brown solution, which was characterised spectroscopically as  $[\text{RhCp}^*\text{H}(\text{PMe}_3)(\text{H}_3\text{B-PCy}_3)][\text{BAR}^{\text{F}}_4]$ , **13**.  $^{11}\text{B}$  NMR spectroscopy at room temperature revealed a considerable upfield shift in the  $^{11}\text{B}$  NMR shift in which the boryl signal had been replaced by one at  $\delta -45.6$ , characteristic of a  $\sigma$ -phosphine-borane. In the  $^1\text{H}$  NMR spectrum (under a  $\text{H}_2$  atmosphere) one very broad upfield signal was observed at  $\delta -4.14$ . Cooling to  $-60^\circ\text{C}$  resolved this into a quadrupolar broadened peak at  $\delta -4.07$  (relative integral 3 H), assigned to a  $\text{Rh}\cdots\text{H}_3\text{B}$  unit, and a sharp doublet of doublets at  $\delta -11.53$  (integral 1 H), assigned to  $\text{Rh}-\text{H}$ . These are exchanging at room temperature, and we suggest that the mechanism for this is likely be through a boryl-dihydrogen complex  $[\text{RhCp}^*(\text{PMe}_3)(\text{H}_2\text{B-PCy}_3)(\text{H}_2)][\text{BAR}^{\text{F}}_4]$ , operating via a sigma-CAM mechanism.<sup>[21]</sup> Addition of  $\text{PPh}_3$  to **12** results in a loss of the high-field signal, and the appearance of two signals at  $\delta 2.42$  and  $0.23$  in the  $^1\text{H}\{^{11}\text{B}\}$  NMR spectrum assigned to  $\text{RhBH}_2\text{PCy}_3$ . Furthermore the  $^{11}\text{B}$  NMR spectrum shows a significant upfield shift to  $\delta -39.5$ , consistent with previously reported, non- $\alpha$ -B-agostic, base-stabilised boryls.<sup>[66-69]</sup> These, and associated  $^{31}\text{P}\{^1\text{H}\}$  NMR data, signal the formation of complex **14**:  $[\text{RhCp}^*(\text{PMe}_3)(\text{PPh}_3)(\text{H}_2\text{B-PCy}_3)][\text{BAR}^{\text{F}}_4]$ .

### D-Labelling Experiments.

The observation of the  $\alpha$ -B-agostic boryl intermediate **11** en route to complex **9** led us to speculate upon the mechanism of formation of the phosphido-borane species **6** (and **7**), and whether Ph- and Cy-analogues of **11** are intermediates in the formation of these species from **1** and the corresponding phosphine-borane. To probe this  $\text{D}_3\text{B-PPh}_2$  was added to **1**. Two scenarios follow: (i) B-D activation followed by P-H activation would lead to a  $\{\text{HD}_2\text{BPR}_2\}$  unit in the final product and the release of  $\text{CH}_3\text{D}$ , or (ii) initial P-H activation would result in liberation of  $\text{CH}_4$  and no incorporation of  $^1\text{H}$  into the borane (Scheme 11).  $^{31}\text{P}$  and  $^{11}\text{B}$  NMR spectroscopy confirmed clean formation of the phosphido-borane product; while  $^1\text{H}$



Scheme 11. D-labelling experiments

and  $^2\text{H}$  NMR spectroscopy (ESI) showed H and D in all positions of the  $\beta$ -B-agostic borane, with an overall relative integral of 1-H measured from the  $^1\text{H}$  NMR spectrum indicating a H:D ratio of 1:2. This suggests route (i) is operating, as observed spectroscopically for complex **11**. That  $^1\text{H}$  signals are observed in all 3 B-H positions of the final product **d-6** suggests slow exchange between terminal and bridging positions which was confirmed by a spin saturation  $^1\text{H}$  NMR exchange experiment.<sup>[79]</sup>  $\text{CH}_3\text{D}$  is observed [ $\delta 0.19$ , t,  $J(\text{HD}) 2.0$  Hz,  $\text{CD}_2\text{Cl}_2$ ], that disappears on degassing the solution.




Figure 7. Computed free energy reaction profiles (kcal/mol, BP86-D3(DCM) level) for formation of **6<sub>R</sub>** and **9<sub>R</sub>** from phosphino-borane adducts **Int(11'-9')**<sub>R</sub>: (R = <sup>t</sup>Bu and Ph). All free energies are quoted relative to **10** set to 0.0 kcal/mol. <sup>a</sup>An intermediate corresponding to a ca. 90° rotation of the phosphino-borane ligand was located between **Int(11'-9')**<sub>R</sub> and **6<sub>R</sub>** and only the energy of the higher-lying transition state is indicated. See text and ESI for full details.

The observation of a phosphido-borane complex  $[\text{RhCp}^*(\text{PR}_2\text{-BH}_3)(\text{PMe}_3)]^+$  when R = Ph (**6**) and Cy (**7**) is in sharp contrast to the formation of  $[\text{RhCp}^*(\text{H})(\text{PR}_2\text{-BH}_2\text{-PMe}_3)]^+$  when R = <sup>t</sup>Bu (**9**). The above labelling studies (R = Ph) and calculations (R = <sup>t</sup>Bu and Ph, Figure 5) are all consistent with initial B-H activation to form  $[\text{RhCp}^*(\text{H}_2\text{B-PHR}_2)(\text{PMe}_3)]^+$ , **11<sub>R</sub>**, as a common intermediate. Figure 5 also indicates that the reaction profile for the formation of **9<sub>Ph</sub>** from **11<sub>Ph</sub>** would follow a similar course to the <sup>t</sup>Bu system, although significantly different energetics are seen around the  $\beta$ -H transfer step from **11'<sub>R</sub>**, which has a much lower barrier and is far more exergonic when R = Ph. The onward reactivities of the resultant phosphino-borane intermediates **Int(11'-9')**<sub>R</sub> are compared in Figure 7. The stability of **Int(11'-9')**<sub>Ph</sub> ( $G = -28.0$  kcal/mol) means the subsequent P-B coupling step towards **9<sub>Ph</sub>** encounters a significant barrier of 22.5 kcal/mol via **TS(11'-9')**<sub>2Ph</sub> at -5.5 kcal/mol. Alternatively, we found that the phosphino-borane ligand in **Int(11'-9')**<sub>Ph</sub> can undergo a two-step rotation that leads directly to **6<sub>Ph</sub>**. This process involves first a transition state **TS(11'-6)**<sub>2Ph</sub> at -12.7 kcal/mol which leads to an intermediate in which the phosphino-borane ligand lies parallel to the Rh-Cp\*(centroid) direction with the  $\{\text{BH}_2\}$  moiety adjacent to the Cp\* ring (**Int(11'-6)**<sub>2Ph</sub>,  $G = -17.4$

kcal/mol). The rotation is completed via a transition state at -15.9 kcal/mol and this second step was also found to be coupled to B-H bond formation involving the Rh-H ligand, resulting in the formation of **6<sub>Ph</sub>**. Note that for clarity only the energy of **TS(11''-6)<sub>Ph</sub>** (the highest point in the rotation process) is indicated in Figure 7; full details are provided in the ESI. Overall this rotation process is kinetically favoured over P-B bond coupling towards **9<sub>Ph</sub>** by 7.2 kcal/mol; moreover the formation of **6<sub>Ph</sub>** is also thermodynamically favoured over **9<sub>Ph</sub>** by 6.5 kcal/mol.

In the light of these results phosphino-borane rotation in **Int(11''-9)<sub>tBu</sub>** was also assessed and was found to proceed with a low overall barrier of 5.8 kcal/mol. This also involves two steps, although in this case the rotated phosphino-borane intermediate has the {P<sup>t</sup>Bu<sub>2</sub>} moiety adjacent to the Cp\* ring. The resultant phosphido-borane, **6<sub>tBu</sub>**, is located at -4.5 kcal/mol and so can readily revert to **Int(11''-9)<sub>tBu</sub>** with a barrier of only 6.3 kcal/mol, from which it can access the competing P-B bond coupling via **TS(11''-9)<sub>tBu</sub>**. The overall barrier for this (from **6<sub>tBu</sub>**) is therefore only 12.2 kcal/mol and leads to first **9'<sub>tBu</sub>** and then **9<sub>tBu</sub>** in processes that are both significantly exergonic. The calculations therefore suggest rapid, but reversible formation of **6<sub>tBu</sub>** before the thermodynamically favoured pathway to **9<sub>tBu</sub>** takes over.<sup>[80]</sup>


The differences in the reaction profiles when R = <sup>t</sup>Bu and Ph in Figure 7 can be attributed to the greater steric encumbrance of the <sup>t</sup>Bu system. This is particularly apparent for **6<sub>tBu</sub>**, the formation of which is 31 kcal/mol less accessible than **6<sub>Ph</sub>**. The combination of the steric bulk derived from both the <sup>t</sup>Bu substituents and the Cp\* ligands is important in this: thus with H<sub>3</sub>B-PMe<sub>2</sub>H (i.e. exchanging Me for <sup>t</sup>Bu) the formation of **6<sub>Me</sub>** becomes exergonic by 17.5 kcal/mol, while the equivalent reaction of [RhCp(Me)(H<sub>3</sub>B-P<sup>t</sup>Bu<sub>2</sub>H)(PMe<sub>3</sub>)]<sup>+</sup> (i.e. retaining the <sup>t</sup>Bu substituents but exchanging Cp for Cp\*) is downhill by 27.6 kcal/mol. Similar arguments explain the greater relative stability of **9<sub>Ph</sub>** over **9<sub>tBu</sub>**. In these systems, however, a PMe<sub>3</sub> ligand has migrated from Rh onto B to be replaced by a much smaller hydride. The accumulative steric effect around the metal is therefore much less significant meaning that **9<sub>tBu</sub>** is only 11.9 kcal/mol less accessible than **9<sub>Ph</sub>**; moreover, the formation of **9<sub>tBu</sub>** becomes thermodynamically viable. Calculations also show that H<sub>3</sub>B-PHCy<sub>2</sub> follows the pattern of behaviour computed for H<sub>3</sub>B-PPhH<sub>2</sub>, consistent with the observed formation of **7** in this case (see ESI for full details).

### Comments on Mechanism of Dehydropolymerization of H<sub>3</sub>B-PRH<sub>2</sub>

These studies suggest that the two likely limiting mechanisms for dehydropolymerization of H<sub>3</sub>B-PPhH<sub>2</sub>, step-growth-like via reversible chain transfer or coordination chain-growth, both likely flow from a common phosphido-borane intermediate (**I**, Scheme 12) that is an analogue of complex **6**. Stoichiometric, labelling and computational studies on secondary phosphine-borane systems suggest that such a species is likely formed from initial B-H activation of a phosphine-borane, followed by P-H transfer and

rearrangement of a resultant hydrido phosphino-borane intermediate, modelled in this study as **Int(11''-9')**.


The observation of significant amounts of oligomer **2** at



**Scheme 12.** Suggested mechanisms for dehydropolymerization. [Rh] = Rh(PR<sub>3</sub>)Cp\* (PR<sub>3</sub> = PMe<sub>3</sub> or PPhH<sub>2</sub>).

short reaction times, alongside the rapid consumption of H<sub>3</sub>B-PPhH<sub>2</sub>, points to reversible chain transfer (B, Scheme 12) as a likely mechanism. That M<sub>n</sub> is essentially unchanged with catalyst loading suggests this mechanism could be further modified by (observed) increasingly more P-B cleavage of the polymer at higher catalyst loadings. Based on our observations a coordination chain growth mechanism (C, Scheme 12) appears less likely; as H<sub>3</sub>B-PPhH<sub>2</sub> would be expected to be consumed gradually throughout the whole polymerization, **2** should not form in significant quantities, and M<sub>n</sub> should increase with decreased catalyst loadings. If chain growth was occurring, slow propagation and faster termination/chain transfer steps would be required to account for our observations. We cannot discount a scenario where both mechanisms operate in ensemble, or there is a change from reversible chain transfer (step growth) to chain growth at lower [H<sub>3</sub>B-PPhH<sub>2</sub>] / higher [oligomer]. Related dual mechanisms have been discussed before with regard to polymer growth kinetics.<sup>[81, 82]</sup>

The contrast with Manners' FeCp(CO)<sub>2</sub>(OTf) system is interesting,<sup>[5]</sup> as this shows coordination chain-growth-type polymerisation kinetics. We currently do not have a clear reason why this would be, although cationic Rh versus neutral Fe, and PR<sub>3</sub> versus CO ligands, are obvious electronic differences. Common to both Rh and Fe systems is the implication of β-B-agostic phosphido-borane complexes of the type [MCp(L)(PRHBH<sub>3</sub>)]<sup>n+</sup>, and we thus suggest that such species, as well as precursor metal-bound phosphino-boranes



**Scheme 13.** Manners' and co-workers observations on polymer molecular weight and P-R substituent.

such as  $[\text{MCp(L)(H)(PRHBH}_2\text{)]}^{n+}$ , play a role in dehydropolymerization. As shown here the reactivity of such phosphino-borane intermediates is dependent on the steric bulk at phosphorus: for  $\text{R} = \text{Ph}$  phosphido-boranes are favoured thermodynamically, whereas for bulkier  $\text{R} = \text{tBu}$  this is the kinetic product, and the thermodynamic product arises from transfer of a metal bound ancillary ligand ( $\text{PMe}_3$ ) to the phosphino-borane. In this regard it is interesting to compare the differences in reported dehydropolymerization efficacy for  $\text{FeCp(CO)}_2(\text{OTf})$ .<sup>[5, 16]</sup> For  $\text{H}_3\text{B-PhH}_2$  high molecular weight polymer is formed ( $M_n$  59 000  $\text{g mol}^{-1}$  in 24 hrs), whereas for  $\text{H}_3\text{B-tBuH}_2$  only short chain oligomers  $[\text{H}_2\text{BP}^t\text{BuH}]_x$  ( $x < 10$ ) are formed after 172 hrs. Given our observations presented here we speculate that this may be due to deactivation routes that are modelled by complexes such as **9** when  $\text{R} = \text{tBu}$  (Scheme 13), that in turn arise from differing reactivity pathways of the corresponding phosphino-boranes.

### Conclusions

By choosing a system that can produce well-defined, moderate molecular weight, poly- $[\text{H}_2\text{BPPH}]_n$ , and is also designed to be latent low-coordinate, the intimate details of initial phosphine-borane activation in dehydropolymerization can be studied. Studies on model systems with secondary phosphine-boranes show that B-H activation precedes P-H activation, to give the kinetic product of a base-stabilised  $\alpha$ -B-agostic boryl complex, subsequent P-H transfer, that operates via a hydrido-phosphino-borane species, leads to the observed phosphido-borane as the thermodynamic product. Together these three species offer many possibilities for pathways operating during dehydropolymerization.

Given the ambiguity related to the mechanism of dehydropolymerisation (step or chain growth-like) in this system we are reluctant to say definitively which mechanism is operating, but our general observations are consistent with those recently proposed mechanisms operating for  $\text{FeCp(CO)}_2(\text{OTf})$ ,  $[\text{Rh(Ph}_2\text{P(CH}_2\text{)}_3\text{PPh}_2)]^+$  and  $\{\text{Fe(N-N)(P-P)}\}$  systems (N-N, P-P = chelating ligands),<sup>[5, 19, 83]</sup> in as much that the proposed species that undergo the P-B bond forming event have M-P bonds (i.e. phosphido-boranes). Moreover, given that boryl, phosphino-borane and phosphido-boranes are all accessible they should all be considered as viable intermediates in catalytic dehydrocoupling and dehydropolymerization processes. This work also lends insight into related amine-borane dehydropolymerization in which amido-boranes, structurally related to **6** have been proposed as actual catalysts, and proposed to form via a N-H activation from a sigma-amine borane precursor,<sup>[31, 83]</sup> similar to that described in detail here for phosphido-boranes. The ubiquity of B-agostic interactions in the systems discussed here, whether  $\alpha$ - or  $\beta$ -, also shows that such interactions also need to be explicitly considered when discussing the mechanism of dehydropolymerization. This mirrors olefin polymerisation, in which  $\alpha$ - and  $\beta$ -agostic interactions play key roles in migratory insertion and polymerization processes.<sup>[84, 85]</sup> Such detail makes a further step towards fully understanding the mechanisms of group 13/15 dehydropolymerizations, and thus

the further development of catalysts that can deliver tailored new polymeric materials.<sup>[2]</sup>

### Acknowledgements

The EPSRC for funding through EP/J02127X (ASW, TNH) and with Heriot-Watt University, for a DTP studentship (NAB). Professor Ian Manners and Dr Titel Jurca (University of Bristol) for GPC analysis and useful discussions.

### Notes and references

- [1] T. Chivers and I. Manners, *Inorganic Rings and Polymers of the p-Block Elements: From Fundamentals to Applications*, Royal Soc. Chem., Cambridge, **2009**.
- [2] E. M. Leitao, T. Jurca and I. Manners, *Nat. Chem.* **2013**, *5*, 817.
- [3] A. Staubitz, A. P. M. Robertson, M. E. Sloan and I. Manners, *Chem. Rev.* **2010**, *110*, 4023.
- [4] T. J. Clark, K. Lee and I. Manners, *Chem. Eur. J.* **2006**, *12*, 8634.
- [5] A. Schäfer, T. Jurca, J. Turner, J. R. Vance, K. Lee, V. A. Du, M. F. Haddow, G. R. Whittell and I. Manners, *Angew. Chem. Int. Ed.* **2015**, *54*, 4836.
- [6] A. B. Burg and R. I. Wagner, *J. Am. Chem. Soc.* **1953**, *75*, 3872.
- [7] H. Dorn, R. A. Singh, J. A. Massey, A. J. Lough and I. Manners, *Angew. Chem. Int. Ed.* **1999**, *38*, 3321.
- [8] H. Dorn, R. A. Singh, J. A. Massey, J. M. Nelson, C. A. Jaska, A. J. Lough and I. Manners, *J. Am. Chem. Soc.* **2000**, *122*, 6669.
- [9] H. Dorn, E. Vejzovic, A. J. Lough and I. Manners, *Inorg. Chem.* **2001**, *40*, 4327.
- [10] T. L. Clark, J. M. Rodezno, S. B. Clendinning, S. Aouba, P. M. Brodersen, A. J. Lough, H. E. Ruda and I. Manners, *Chem. Eur. J.* **2005**, *11*, 4526.
- [11] S. Pandey, P. Lönnecke and E. Hey-Hawkins, *Inorg. Chem.* **2014**, *53*, 8242.
- [12] S. Pandey, P. Lönnecke and E. Hey-Hawkins, *Eur. J. Inorg. Chem.* **2014**, *2014*, 2456.
- [13] C. A. Jaska and I. Manners, *J. Am. Chem. Soc.* **2004**, *126*, 1334.
- [14] C. A. Jaska and I. Manners, *J. Am. Chem. Soc.* **2004**, *126*, 9776.
- [15] H. C. Johnson, T. N. Hooper and A. S. Weller in *The Catalytic Dehydrocoupling of Amine-Boranes and Phosphine-Boranes*, Vol. 49 Eds.: E. Fernández and A. Whiting), Springer International Publishing, **2015**.
- [16] C. Marquardt, T. Jurca, K.-C. Schwan, A. Stauber, A. V. Virovets, G. R. Whittell, I. Manners and M. Scheer, *Angew. Chem. Int. Ed.* **2015**, *54*, 13782.
- [17] M. A. Huertos and A. S. Weller, *Chem. Commun.* **2012**, *48*, 7185.
- [18] D. J. Grant and D. A. Dixon, *J. Phys. Chem. A.* **2006**, *110*, 12955.
- [19] M. A. Huertos and A. S. Weller, *Chem. Sci.* **2013**, *4*, 1881.
- [20] T. N. Hooper, M. A. Huertos, T. Jurca, S. D. Pike, A. S. Weller and I. Manners, *Inorg. Chem.* **2014**, *53*, 3716.
- [21] R. N. Perutz and S. Sabo-Etienne, *Angew. Chem. Int. Ed.* **2007**, *46*, 2578.
- [22] R. Waterman, *Organometallics* **2013**, *32*, 7249.
- [23] F. L. Taw, H. Mellows, P. S. White, F. J. Hollander, R. G. Bergman, M. Brookhart and D. M. Heinekey, *J. Am. Chem. Soc.* **2002**, *124*, 5100.
- [24] B. K. Corkey, F. L. Taw, R. G. Bergman and M. Brookhart, *Polyhedron* **2004**, *23*, 2943.

- [25] A. H. Roy, C. P. Lenges and M. Brookhart, *J. Am. Chem. Soc.* **2007**, *129*, 2082.
- [26] J. F. Hartwig, K. S. Cook, M. Hapke, C. D. Incarvito, Y. Fan, C. E. Webster and M. B. Hall, *J. Am. Chem. Soc.* **2005**, *127*, 2538.
- [27] V. P. W. Böhm and M. Brookhart, *Angew. Chem. Int. Ed.* **2001**, *40*, 4694.
- [28] F. Wang and X. Li, *Chem. Soc. Rev.* **2012**, *41*, 3651.
- [29] H. Dorn, J. M. Rodezno, B. Brunnhöfer, E. Rivard, J. A. Massey and I. Manners, *Macromolecules* **2003**, *36*, 291.
- [30] A. H. Cowley and M. C. Damasco, *J. Am. Chem. Soc.* **1971**, *93*, 6815.
- [31] H. C. Johnson, E. M. Leitao, G. R. Whittell, I. Manners, G. C. Lloyd-Jones and A. S. Weller, *J. Am. Chem. Soc.* **2014**, *136*, 9078.
- [32] A. Staubitz, M. E. Sloan, A. P. M. Robertson, A. Friedrich, S. Schneider, P. J. Gates, J. r. S. a. d. Günne and I. Manners, *J. Am. Chem. Soc.* **2010**, *132*, 13332.
- [33] A. P. M. Robertson, E. M. Leitao, T. Jurca, M. F. Haddow, H. Helten, G. C. Lloyd-Jones and I. Manners, *J. Am. Chem. Soc.* **2013**, *135*, 12670.
- [34] A. Kumar, H. C. Johnson, T. N. Hooper, A. S. Weller, A. G. Algarra and S. A. Macgregor, *Chem. Sci.* **2014**, *5*, 2546.
- [35] D. Michael, P. Mingos, P. C. Minshall, M. B. Hursthouse, K. M. A. Malik and S. D. Willoughby, *J. Organomet. Chem.* **1979**, *181*, 169.
- [36] Z.-J. Yao, X.-K. Huo and G.-X. Jin, *Chem. Commun.* **2012**, *48*, 6714.
- [37] A. T. Lubben, J. S. McIndoe and A. S. Weller, *Organometallics* **2008**, *27*, 3303.
- [38] G. R. Fulmer, A. J. M. Miller, N. H. Sherden, H. E. Gottlieb, A. Nudelman, B. M. Stoltz, J. E. Bercaw and K. I. Goldberg, *Organometallics* **2010**, *29*, 2176.
- [39] E. M. Pelczar, E. A. Nytko, M. A. Zhuravel, J. M. Smith, D. S. Glueck, R. Sommer, C. D. Incarvito and A. L. Rheingold, *Polyhedron* **2002**, *21*, 2409.
- [40] R. T. Paine and H. Noeth, *Chem. Rev.* **1995**, *95*, 343.
- [41] J. A. Bailey and P. G. Pringle, *Coord. Chem. Rev.*
- [42] DFT calculations were performed with the Gaussian 03 and 09 suites and optimisations employed the BP86 functional. See Supporting Information for full details.
- [43] W. F. McNamara, E. N. Duesler, R. T. Paine, J. V. Ortiz, P. Koelle and H. Noeth, *Organometallics* **1986**, *5*, 380.
- [44] I. Amor, D. García-Vivó, M. E. García, M. A. Ruiz, D. Sáez, H. Hamidov and J. C. Jeffery, *Organometallics* **2007**, *26*, 466.
- [45] M. A. Alvarez, M. E. Garcia, R. Gonzalez and M. A. Ruiz, *Dalton Trans.* **2012**, *41*, 14498.
- [46] H. Helten, B. Dutta, J. R. Vance, M. E. Sloan, M. F. Haddow, S. Sproules, D. Collison, G. R. Whittell, G. C. Lloyd-Jones and I. Manners, *Angew. Chem. Int. Ed.* **2013**, *52*, 437.
- [47] M. S. Hill, M. Hodgson, D. J. Liptrot and M. F. Mahon, *Dalton Trans.* **2011**, *40*, 7783.
- [48] D. J. Liptrot, M. S. Hill, M. F. Mahon and D. J. MacDougall, *Chem. Eur. J.* **2010**, *16*, 8508.
- [49] J. Spielmann, D. F. J. Piesik and S. Harder, *Chem. Eur. J.* **2010**, *16*, 8307.
- [50] M. Brookhart and D. M. Lincoln, *J. Am. Chem. Soc.* **1988**, *110*, 8719.
- [51] U. Vogel, P. Hoemensch, K.-C. Schwan, A. Y. Timoshkin and M. Scheer, *Chem. Eur. J.* **2003**, *9*, 515.
- [52] C. Thoms, C. Marquardt, A. Y. Timoshkin, M. Bodensteiner and M. Scheer, *Angew. Chem. Int. Ed.* **2013**, *52*, 5150.
- [53] K.-C. Schwan, A. Y. Timoshkin, M. Zabel and M. Scheer, *Chem. Eur. J.* **2006**, *12*, 4900.
- [54] B. Wrackmeyer, *Z.Naturforsch.(B)* **2004**, *59*, 1192.
- [55] A. B. Burg, *Inorg. Chem.* **1978**, *17*, 593.
- [56] T. A. Shuttleworth, M. A. Huertos, I. Pernik, R. D. Young and A. S. Weller, *Dalton Trans.* **2013**, *42*, 12917.
- [57] Y. Kawano, T. Yasue and M. Shimoi, *J. Am. Chem. Soc.* **1999**, *121*, 11744.
- [58] Y. Kawano, M. Hashiva and M. Shimoi, *Organometallics* **2006**, *25*, 4420.
- [59] H. C. Johnson, R. Torry--Harris, L. Ortega, R. Theron, J. S. McIndoe and A. Weller, *Cat. Sci. Tech.* **2014**, *4*, 3486
- [60] H. C. Johnson, C. L. McMullin, S. D. Pike, S. A. Macgregor and A. S. Weller, *Angew. Chem. Int. Ed.* **2013**, *52*, 9776.
- [61] N. Merle, G. Koicok-Köhn, M. F. Mahon, C. G. Frost, G. D. Ruggerio, A. S. Weller and M. C. Willis, *Dalton Trans.* **2004**, 3883.
- [62] A. B. Chaplin and A. S. Weller, *Angew. Chem. Int. Ed.* **2010**, *49*, 581.
- [63] C. Y. Tang, A. L. Thompson and S. Aldridge, *J. Am. Chem. Soc.* **2010**, *132*, 10578.
- [64] G. Bénac-Lestrille, U. Helmstedt, L. Vendier, G. Alcaraz, E. Clot and S. Sabo-Etienne, *Inorg. Chem.* **2011**, *50*, 11039.
- [65] C. J. Stevens, R. Dallanegra, A. B. Chaplin, A. S. Weller, S. A. Macgregor, B. Ward, D. Mckay, G. Alcaraz and S. Sabo-Etienne, *Chem. Eur. J.* **2011**, *17*, 3011.
- [66] T. Yasue, Y. Kawano and M. Shimoi, *Angew. Chem. Int. Ed.* **2003**, *42*, 1727.
- [67] Y. Kawano, T. Yasue and M. Shimoi, *J. Am. Chem. Soc.* **1999**, *121*, 1.
- [68] T. Yasue, Y. Kawano and M. Shimoi, *Chem. Lett.* **2000**, 58.
- [69] H. Nakazawa, M. Ohba and M. Itazaki, *Organometallics* **2006**, *25*, 2903.
- [70] M. O'Neill, D. A. Addy, I. Riddlestone, M. Kelly, N. Phillips and S. Aldridge, *J. Am. Chem. Soc.* **2011**, *133*, 11500.
- [71] H. Braunschweig, K. Radacki, F. Seeler and G. R. Whittell, *Organometallics* **2004**, *23*, 4178.
- [72] G. Alcaraz and S. Sabo-Etienne, *Coord. Chem. Rev.* **2008**, *252*, 2395.
- [73] M. Etienne, *Organometallics* **1994**, *13*, 410.
- [74] H. Braunschweig, R. D. Dewhurst and V. H. Gessner, *Chem. Soc. Rev.* **2013**, *42*, 3197.
- [75] H. Braunschweig, K. Radacki, D. Rais and D. Scheschkewitz, *Angew. Chem. Int. Ed.* **2005**, *44*, 5651.
- [76] A. Amgoune, S. Ladeira, K. Miqueu and D. Bourissou, *J. Am. Chem. Soc.* **2012**, *134*, 6560.
- [77] T. M. Douglas, A. B. Chaplin, A. S. Weller, X. Yang and M. B. Hall, *J. Am. Chem. Soc.* **2009**, *131*, 15440.
- [78] We could not observe the corresponding B–H (terminal) signal for (**12**), and it may be obscured by the anion resonances, as the equivalent signal in (**11**) comes at  $\delta$  7.11.
- [79] The  $^1\text{H}\{^{11}\text{B}\}$  spectra of complex **6/d-6** are suggestive of an isotopic perturbation of equilibrium. Relative integrals of the BH signals at 0.50 (BH), –0.02 (BH) and –10.80 (BH...Rh) ppm: **6**, 0.30:0.31:0.39; **d-6**, 0.26:0.25:0.49. This suggests a preference for Rh...H–B rather than Rh...D–B in **d-6**.
- [80] A process involving P–H activation with direct H-transfer onto B was also characterised but exhibited significantly higher barriers (see ESI).
- [81] M. Mizutani, K. Satoh and M. Kamigaito, *J. Am. Chem. Soc.* **2010**, *132*, 7498.
- [82] L. You and J. Ling, *Macromolecules* **2014**, *47*, 2219.
- [83] R. T. Baker, J. C. Gordon, C. W. Hamilton, N. J. Henson, P.-H. Lin, S. Maguire, M. Murugesu, B. L. Scott and N. C. Smythe, *J. Am. Chem. Soc.* **2012**, *134*, 5598.

[84] M. Brookhart, M. L. H. Green and G. Parkin, *Proc. Nat. Acad. Sci USA* **2007**, *104*, 6908.

[85] M. Brookhart, A. F. Volpe, D. M. Lincoln, I. T. Horvath and J. M. Millar, *J. Am. Chem. Soc.* **1990**, *112*, 5634.