

Soft Matter

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Formation of adhesion domains in stressed and confined membranes

Nadiv Dharan¹ and Oded Farago^{1,2}

¹Department of Biomedical Engineering, ²Ilse Katz Institute
for Nanoscale Science and Technology, Ben Gurion University,
Be'er Sheva 84105, Israel

The adhesion bonds connecting a lipid bilayer to an underlying surface may undergo a condensation transition resulting from an interplay between a short range attractive potential between them, and a long range fluctuation-induced potential of mean force. Here, we use computer simulations of a coarse-grained molecular model of supported lipid bilayers to study this transition in confined membranes, and in membranes subjected to a non-vanishing surface tension. Our results show that confinement may alter significantly the condensation transition of the adhesion bonds, whereas the application of surface tension has a very minor effect on it. We also investigate domain formation in membranes under negative tension which, in free membranes, causes enhancement of the amplitude of the membrane thermal undulations. Our results indicate that in supported membranes, this effect of a negative surface tension on the fluctuation spectrum is largely eliminated by the pressure resulting from the mixing entropy of the adhesion bonds.

PACS numbers:

I. INTRODUCTION

Lipid bilayers serve as a physical barrier separating the content of the cell from its surroundings. They are embedded with different types of proteins responsible for numerous biological functions. An important class of membrane proteins are cell adhesion molecules that form specific (ligand-receptor) bonds with different biological elements, e.g., the extracellular matrix, the cytoskeleton, and neighboring cell membranes [1]. Cell adhesion plays a central role in many biological processes, such as T-cell activation as part of the immune response [2], cell migration [3] and tissue formation [4].

When membranes adhere to another surface, the adhesion bonds may condense to form large adhesion domains [5–7], which not only provide mechanical stability but also promote cellular signaling pathways necessary for many biological processes [8, 9]. The aggregation of adhesion bonds is driven by attractive forces, including electrostatic and Van der Waals forces [10], interactions resulting from cytoskeleton remodeling [11], and membrane-mediated interactions [12]. The latter mechanism stems from the suppression of membrane thermal fluctuation caused by the adhesion. The associated decrease in fluctuation entropy is minimized when the bonds cluster together into a single adhesion domain.

Over the past two decades, considerable efforts have been made to better understand the role of membrane mediated interactions in the formation of adhesion domains. To this end, a common theoretical approach is the use of discrete lattice models, where the occupied sites represent membrane adhesion bonds, while empty sites stand for free fluctuating membrane segments [13–19]. Computer simulations of such models have shown that membrane fluctuations can facilitate conditions for condensation of adhesion bonds. Nevertheless, it has been also concluded that the fluctuation induced interactions

alone are too weak to promote aggregation and, hence, other attractive interactions between the bonds must also exist. A similar conclusion was recently reached in simulation studies employing coarse-grained (CG) molecular models, which provide a more physical description of the system. In one such study, we simulated a bilayer membrane with a small fraction of lipids from the lower monolayer connected to an underlying surface, and additionally introduced a short range attractive potential of depth ϵ between the adhesive lipids [20]. Our simulations revealed that the adhesive lipids underwent a first order condensation transition when the strength of the short range potential between them exceeds a threshold value $\epsilon_c > 0$. The fact that adhesion domains do not form for $\epsilon = 0$ implies that the fluctuation entropy gained by the aggregation of the adhesive lipids only partially compensates for their loss of mixing entropy. In another CG simulation study, Noguchi demonstrated that cell junctions connecting *more than two* membranes can aggregate without an additional attractive potential between them [21]. This observation can be understood because the total entropy of thermal undulations increases linearly with the number of membranes, N_m , while the mixing entropy of the cell junctions is independent of N_m .

Previous simulation studies have focused on the aggregation behavior of adhesion bonds in *tensionless* membranes. Here, we wish extend our investigations to lipid bilayers subjected to surface tension and confinement. Under such conditions, the membrane long wavelength thermal undulations are suppressed, which weakens the fluctuation induced attraction. Specifically, a harmonic confining potential of strength $\gamma > 0$ suppresses thermal fluctuations at length scales larger than $\xi_\gamma \sim (\kappa/\gamma)^{1/4}$, where κ is the bending modulus of the membrane. Likewise, surface tension $\tau > 0$ affects bending modes with wavelengths greater than $\xi_\tau \sim (\kappa/\tau)^{1/2}$. The influence of tension and confinement on the fluctuation mediated potential of mean force (PMF) was analyzed for

the case of two adhesion bonds [22]. It was shown that the PMF depends logarithmically on the pair distance: $\phi(r) = Ck_B T \ln(r)$, where $C = 2$ at short distances, $C = 1$ at long distances ($r \gg \xi_\tau$) for stressed membranes, and $C = 0$ for long distances ($r \gg \xi_\gamma$) for confined membranes. These findings suggest that tension and confinement may hinder the formation of adhesion domains, and that the phase transition into the condensed state requires a stronger short range attraction between the bonds. Therefore, one should expect the transition threshold value ϵ_c to grow with τ and γ ; however, the magnitude of the increase in ϵ_c remains hard to assess due to the many body nature of the PMF [19]. One may also consider the case of a bilayer membrane under negative tension. Negative tensions amplify the amplitude of the long wavelength undulations (and, for very large negative values, even lead to instability), and this should cause a reduction in ϵ_c . It is especially interesting to examine whether ϵ_c can decrease to zero, in which case the adhesion domains will form without an additional short range potential, i.e., on purely entropic grounds. As will be shown further on, the results of the current study suggest that confinement might have a very strong impact on ϵ_c , while the application of a positive surface tension hardly influences the transition into an aggregated phase. In cases where the membrane is subjected to a negative surface tension, we observe the formation of elongated adhesion domains close to the transition point.

II. MONTE CARLO SIMULATIONS

In order to address the above issues, we conducted Monte Carlo simulations of a lipid bilayer using the Cooke-Deserno implicit-solvent CG model [23]. The details of the simulations can be found in ref. [20]. Briefly, we simulated bilayers of $2N = 2000$ lipids (1000 lipids per monolayer), where each lipid consists of one hydrophilic (head) bead and two hydrophobic (tail) beads of size σ . A flat impenetrable surface was placed below the membrane at $z = 0$, and adhesion was established by connecting N_b head beads from the lower monolayer to it. A Lennard-Jones type potential with a tunable depth ϵ was introduced between the adhesive head beads. We simulated systems with various densities of adhesive lipids, $\phi = N_b/N$, and for different values of the interaction potential depth ϵ . Our simulations for confined membranes included an additional impermeable surface placed above the upper monolayer at $z = z_{\text{conf}}$. Simulations of membranes under constant mechanical surface tension τ were carried out according to the method described in ref. [24].

Typical equilibrium configurations of tensionless membranes with $\phi = 0.2$ are shown in Fig. 1 for $\epsilon = 0.2k_B T$ (A) and $\epsilon = 1.4k_B T$ (B). In the former, the adhesive lipids (whose head beads are shown in the figure in red) are scattered across the surface, while in the latter they condense into a large cluster. Fig. 1(C) shows the average energy E of pair interactions between adhesive




FIG. 1: Bottom view of equilibrium snapshots of a tensionless bilayer with $\phi = 0.2$ showing (A) a scattered distribution of adhesion bonds for $\epsilon = 0.2k_B T$, and (B) a condensed phase for $\epsilon = 1.4k_B T$. The head and tail beads are colored in white and blue, respectively, while the adhesive beads are colored in red. (C) The average number of contacts between the adhesion bonds as function of ϵ . The condensation transition value ϵ_c is empirically defined via the equality $\langle N_C \rangle \simeq 1.5$. The line serves as a guide to the reader's eye.

beads, normalized per bond and expressed in dimensionless units by dividing by the potential strength ϵ . This quantity, which is a measure for the typical number of contacts between the adhesive beads $\langle N_C \rangle$, increases sharply when the condensation transition takes place. From the results of Fig. 1(C), we empirically determine the transition value as the one for which $\langle N_C \rangle \simeq 1.5$.

III. RESULTS

A. Membranes under confinement

We first start with our simulation results for non stressed membranes confined between the supporting underlying surface (on which the adhesive beads reside) and a second upper surface. The former is located at $z = 0$, just underneath the tips of the head beads of the lower leaflet, while the latter is placed at $z = z_{\text{conf}} \geq 6\sigma$. The degree of confinement increases when z_{conf} decreases which, in turn, would lead to stronger suppression of the membrane thermal fluctuations and a shift in the transition threshold ϵ_c to larger values. This trend is demonstrated in Fig. 2, showing our simulation results for




FIG. 2: The average number of contacts between the adhesion bonds in membranes confined by a surface located at $z_{\text{conf}} = 9\sigma$ (green diamonds), $z_{\text{conf}} = 7.5\sigma$ (blue triangles) and $z_{\text{conf}} = 6\sigma$ (red squares), as a function of ϵ , for (A) $\phi = 0.1$, and (B) $\phi = 0.2$. Results for non-confined membranes are denoted by black circles. The lines serve as a guide to the reader's eye. The statistical errors are comparable to the size of the symbols.

$z_{\text{conf}} = 6\sigma$, 7.5σ , and 9σ in supported membranes with $\phi = 0.1$ (A) and $\phi = 0.2$ (B). For $z_{\text{conf}} = 9\sigma$, our results (green diamonds) match perfectly with the results obtained for non-confined membranes (black circles). This means that the rate of collisions between the membrane and the upper surface is negligibly small and, therefore, it does not affect the fluctuation spectrum. Lowering the confining surface by a distance equal to the size of a bead and a half to $z_{\text{conf}} = 7.5\sigma$ has a more noticeable effect on membrane thermal undulations, which leads to an increase in the condensation transition value from $\epsilon_c \simeq 0.65k_B T$ for non-confined membranes at $\phi = 0.1$ to $\epsilon_c \simeq 0.8k_B T$. For $\phi = 0.2$, the shift is smaller, from $\epsilon_c \simeq 0.6k_B T$ to $\epsilon_c \simeq 0.7k_B T$. When the upper surface is further lowered to $z_{\text{conf}} = 6\sigma$, it touches the tips of the head beads in the upper leaflet, as the thickness of the bilayer is equal to the size of six beads. A confining surface located at $z_{\text{conf}} = 6\sigma$ completely suppresses thermal undulations, and eliminates the fluctuation mediated interactions between the adhesion bonds. Under these conditions, the threshold value for aggregation increases to




FIG. 3: The average number of contacts between the adhesion bonds in membranes under mechanical surface tension $\tau = 0.48k_B T/\sigma^2$, as a function of ϵ for (A) $\phi = 0.1$, and (B) $\phi = 0.2$. The results are plotted in red squares, and are compared with the results for tensionless membranes ($\tau = 0$) that are depicted by black open circles. The lines serve as guides to the eye. The statistical errors are comparable to the size of the symbols.

$\epsilon_c \simeq 1.2k_B T$ at $\phi = 0.1$, and $\epsilon_c \simeq 1k_B T$ for $\phi = 0.2$. These values are approximately twice larger than the corresponding values found when no upper plate exists ($\epsilon_c \simeq 0.65k_B T$ and $\epsilon_c \simeq 0.6k_B T$, respectively), which is in accord with the conclusions of refs. [19, 21], that the entropic gain of aggregation compensates for, roughly, half of the loss in mixing entropy of the adhesion bonds.

B. Membranes under surface tension

We now proceed to our simulation results for non-confined membranes subjected to a lateral surface tension $\tau > 0$. Since, just like steric confinement, surface tension also suppresses the thermal undulations of the membrane, one may expect to also find an increase in the value of ϵ_c . However, our computational results that are depicted in Fig 3, demonstrate that the $\langle N_C \rangle$ vs. ϵ curve remains almost unchanged when the tension increases from $\tau = 0$ (black open circles) to $\tau = 0.48k_B T/\sigma^2$ (red squares) for both $\phi = 0.1$ (A) and $\phi = 0.2$ (B). This ob-




FIG. 4: The average number of contacts between the adhesion bonds in membranes under surface tension $\tau = -0.24k_B T/\sigma^2$, as a function of ϵ for (A) $\phi = 0.1$, and (B) $\phi = 0.2$. The results are plotted in red squares, and are compared with the results for tensionless membranes ($\tau = 0$) that are depicted by black open circles. (C) The mean projected area per lipid as a function of ϵ , for $\phi = 0.1$ (black) and $\phi = 0.2$ (red). Circles denote the results for tensionless membranes, while the results for $\tau = -0.24k_B T/\sigma^2$ are shown in squares. The lines serve as guides to the eye.

servation suggests that the fluctuation spectrum of the simulated membranes is very little affected by the tension, which is expected given that the bending modulus of the model membrane has been previously estimated to be $\kappa \simeq 8k_B T$ [25]. For $\tau = 0.48k_B T/\sigma^2$, the wavelength of the modes that are suppressed are larger than

$\xi_\tau = 2\pi\sqrt{\kappa/\sigma} \simeq 25\sigma$, which is much larger than the typical distance $\bar{d} = \sqrt{A_P/N_b} \simeq 4\sigma$ between the adhesion bonds when the latter are uniformly distributed on the surface (where A_P is the projected area of the membrane). In order for the effect of tension to be appreciable, one must apply stronger tensions; however, the application of a stronger tension would open pores in the membrane and tear it apart. In simulations with a surface tension of $\tau = 0.64k_B T/\sigma^2$, we observed pores in most of the simulated membranes, and in the few that had managed to survive the tension, we still did not observe any considerable changes in the strength of membrane mediated interactions.

We next aimed to address the implications of applying a negative surface tension. In general, a negative tension imposed on the membrane leads to a reduction in its projected area, A_P , and amplifies the long wavelength bending modes. Hence, the fluctuation-induced attraction between the adhesion bonds is expected to be stronger which, in turn, implies that the threshold for condensation ϵ_c should become smaller than in tensionless membranes. To test this hypothesis, we simulated the membrane under a negative tension of $\tau = -0.24k_B T/\sigma^2$. We performed two sets of independent MC simulations, one starting from a random distribution of adhesion bonds, and another where initially the adhesion bonds were organized in one large cluster. The system was equilibrated until configurations originating from these two distinct initial conditions achieved similar characteristics. Fig. 4 shows our results for $\langle N_C \rangle$ as a function of ϵ for $\phi = 0.1$ (A) and $\phi = 0.2$ (B). Contrary to our expectation to observe a reduction in ϵ_c , the data for $\tau = -0.24k_B T/\sigma^2$ appears almost identical to the results of the tensionless case, with $\epsilon_c \simeq 0.6k_B T$ for both values of ϕ . Taken together with the data for $\tau = +0.48k_B T/\sigma^2$ (see Fig. 3), we conclude that application of surface tension has a negligible effect on the condensation transition, within the range of tensions where the supported membrane is mechanically stable.

The negative tension, however, does have an impact on the shape of membranes. Freely fluctuating bilayers assume buckled configurations at negative tensions larger (in absolute value) than $\tau_c \simeq -4\pi^2\kappa/A_P$ [26, 27]. In this study, $A_P/N \simeq 1.33\sigma^2$ [see Fig. 4(C)], which gives $\tau_c \simeq -0.24k_B T/\sigma^2$. In ref. [28], a similar model membrane consisting of the same number of lipids was simulated and, indeed, for $\tau = \tau_c$ the membrane appeared quite buckled. In supported membranes, however, the emergence of buckled configurations occurs only in membranes with large adhesion domains (i.e., for $\epsilon \gtrsim \epsilon_c$). Fig. 5 shows typical equilibrium configuration for $\phi = 0.1$ with $\epsilon = 0$ (A), $0.6k_B T$ (B), and $1.0k_B T$ (C). Each configuration is shown both in side and bottom views. When the attractive potential is set to $\epsilon = 0$, the distribution of the adhesion bonds is scattered and the membrane remains fairly flat. This indicates that the mixing entropy of the bonds dominates the fluctuation entropy of the bilayer, despite the imposed negative ten-




FIG. 5: Typical equilibrium configurations of membranes under surface tension $\tau = -0.24k_B T/\sigma^2$ with density of adhesion bonds $\phi = 0.1$ for (A) $\epsilon = 0$, (B) $\epsilon = 0.6k_B T$ and (C) $\epsilon = 1.0k_B T$. The figures in the upper and lower rows display bottom and side views of the system, respectively. Color coding is the same as in Fig. 1.

sion [see Fig. 5(A)]. For $\epsilon = 1.0k_B T$, the short range pair interactions between the bonds lead to their aggregation. Once the bonds condense, their influence on the thermal behavior of the membrane is greatly weakened, and strong bending undulations appear [see Fig. 5(C)]. Close to the condensation transition, at $\epsilon = 0.6k_B T$, the system exhibits some interesting features: The amplitude of one of the two longest wavelength bending modes [with wavevector $\vec{q}_{(1,0)} = (2\pi/\sqrt{A_P})(1,0)$, or $\vec{q}_{(0,1)} = (2\pi/\sqrt{A_P})(0,1)$] grows considerably, and the membrane assumes an anisotropic buckled configuration. The adhesion bonds are concentrated throughout the minimum of the dominating bending mode, forming an elongated domain (“stripe”) [see Fig. 5(B)]. These observed characteristics represent an intricate balance between the driving forces that govern the thermodynamic behavior of the system. Under negative tension, the system benefits from a reduction in the projected area, leading to a decrease in the Gibbs free energy. The membrane, however, is quite incompressible and, thus, the reduction in A_P must be accompanied by an increase in the area stored in thermal fluctuations whose amplitudes grow. The modes that experience the largest increase in amplitude are the softest ones, corresponding to $\vec{q}_{(1,0)}$ and $\vec{q}_{(0,1)}$ [26] (see footnote ref. [29]). In our simulations, we seldom observed situations where both these modes were simultaneously excited [30], which can be linked to the mixing entropy of the adhesion bonds. When only one of the long modes is dominant, the contact area between the membrane and the surface, which is available for the presence of the adhesion bonds, is larger than in configurations where both modes are excited.

The interplay between the mixing entropy of the adhesion bonds and the contribution of the negative tension to the free energy is further demonstrated in Fig. 4(C), depicting the mean projected area $\langle A_P \rangle$ (normalized by the




FIG. 6: Configurations of membranes with $\phi = 0.1$ and $\epsilon = 1.0k_B T$ under a strong negative tension $\tau = -0.32k_B T/\sigma^2$, showing a spherical protrusion (A) and a tubular one (B). The upper and lower rows display top and side views of the membrane respectively. The tail and adhesive beads are colored in blue and red, respectively. The head beads are colored in grayscale to reflect their height above the surface, with lighter colors representing a higher bead.

number of lipids per monolayer $N = 1000$) as a function of ϵ for $\tau = -0.24k_B T/\sigma^2$ (squares) and $\tau = 0$ (circles). In the tensionless case, we observe a very mild decrease in $\langle A_P \rangle$ as ϵ increases, occurring mainly around ϵ_c . For $\tau = -0.24k_B T/\sigma^2$, $\langle A_P \rangle$ maintains a value close to the tensionless case for $\epsilon < \epsilon_c$, and drops significantly for $\epsilon > \epsilon_c$. In the latter regime, we also observe an increase in the area fluctuations, resulting in larger uncertainties (error bars) in our estimates of $\langle A_P \rangle$. The sharp decrease in $\langle A_P \rangle$ and the concurrent increase in the area fluctuations are anticipated outcomes of a negative surface tension [28]. The fact that they are observed only above the condensation transition is consistent with the picture discussed in the previous paragraph that, below ϵ_c , the effect of the negative tension is largely eliminated by the pressure resulting from the mixing entropy of the adhesion bonds. Notice that the sharp decrease in the mean projected area and the increase in the area fluctuations are much more noticeable for $\phi = 0.1$ than for $\phi = 0.2$. This is to be expected because the smaller ϕ , the smaller the restrictions imposed by the adhesion bonds on large thermal undulations which are directly coupled to the projected area by the highly incompressible character of the membrane.

Applying an even stronger negative tension causes the

membrane to lose its mechanical stability. This is demonstrated in Fig. 6, showing snapshots in top and side views of membranes with $\phi = 0.1$ and $\epsilon = 1.0k_B T$ subjected to $\tau = -0.32k_B T/\sigma^2$. In these snapshots, the head beads are colored in grayscale, with lighter colors indicating beads located higher above the underlying surface. The application of a strong negative tension causes the supported membrane to develop large protrusions with either spherical [Fig. 6(A)] or tubular [Fig. 6(B)] shapes. The adhesion bonds (which are colored in red, and are only partially visible) are concentrated in the periphery of the protrusion, where the membrane is in contact with the underlying surface. We note that the observed protrusions tend to evolve slowly and, therefore, the snapshots shown in Fig. 6 may not represent true equilibrium structures. On the other hand, we also note that very similar equilibrium structures, featuring spherical and tubular protrusions, have been recently observed in an experimental study where supported lipid bilayers were subjected to compression [31].

IV. SUMMARY

The thermal fluctuations of a supported membrane induce an attractive PMF between the adhesion bonds, that promotes the formation of adhesion domains. In previous studies of systems where the adhesion bonds interact with each other via a pairwise short range attractive potential, it has been concluded that the fluctuation-induced PMF significantly decreases the condensation transition point, ϵ_c , of the adhesion bonds. In this work, we investigate adhesion domain formation in stressed and

confined membranes by using MC simulations of a CG supported bilayer model. Our simulations reveal that placing a plate above the membrane (in addition to the surface supporting the membrane from below), significantly alters the condensation point. In the most extreme case, where the membrane is completely flattened between the two plates, the value of ϵ_c increases by almost a factor of two. In contrast to the impact of steric confinement, the application of a positive surface tension, which also acts to suppress membrane undulations, hardly affects the transition within the range of tensions where the supported membrane is mechanically stable. Similarly, we find that a negative tension also has a very minor effect on the condensation transition. Nevertheless, once the adhesion bonds are aggregated into a large domain (i.e., for $\epsilon > \epsilon_c$), the negative tension affects the shape of the membrane and causes it to buckle. Below ϵ_c the adhesion bonds are scattered across the membrane, which prevents the formation of buckled configurations. From the fact that ϵ_c is hardly affected by the application of a negative tension, we conclude that the mixing entropy of the adhesion bonds dominates the demixing entropy of thermal fluctuations. Close to ϵ_c , we observed the formation of elongated adhesion stripes. Such configurations emerge from an interplay between the mixing entropy of the adhesion bonds, the short range residual potential, and the applied negative tension. Finally, under a very strong negative tension, we observe tubular and spherical structures protruding out of the membrane's plane, which indicates that the system is at the onset of mechanical instability.

This work was supported by the Israel Science Foundation (ISF) through grant No. 1087/13.

-
- [1] B. Alberts, D. Bray, J. Lewis, M. Raff, K. Roberts, and J. D. Watson, *Molecular Biology of the Cell* (Garland, New York, 1994).
 - [2] T. A. Springer, *Nature* **340**, 425 (1990).
 - [3] A. Huttenlocher, R. R. Sandborg, and A. F. Horwitz, *Curr. Opin. Cell. Biol.* **7**, 697 (1995).
 - [4] K. Vlemminckx and R. Kemler, *BioEssays* **21**, 211 (1999).
 - [5] Y. Wang and B. Rose, *J. Cell Sci.* **108**, 3501 (1995).
 - [6] C. R. F. Monks *et al.*, *Nature* **395**, 82 (1998).
 - [7] C. Selhuber-Unkel, M. López-García, H. Kessler, and J. P. Spatz, *Biophys. J.* **95**, 5424 (2008).
 - [8] K. Burridge and M. Chrzanowska-Wodnicka, *Annu. Rev. Cell Dev. Biol.* **12**, 463 (1996).
 - [9] F. G. Giancotti and E. Ruoslahti, *Science* **285**, 1028 (1999).
 - [10] J. N. Israelachvili, *Intermolecular and Surface Forces* (Academic Press, San Diego, 1992).
 - [11] C. Wulffing and M. M. Davis, *Science* **282**, 2266 (1998).
 - [12] R. Bruinsma and P. Pincus, *Curr. Opin. Solid St. M.* **1**, 401 (1996).
 - [13] R. Lipowsky, *Phys. Rev. Lett.* **77**, 1652 (1996).
 - [14] T. Weikl *et al.*, *Soft Matter* **5**, 3213 (2009).
 - [15] T. R. Weikl and R. Lipowsky, *Phys. Rev. E* **64**, 011903 (2001).
 - [16] T. Speck and R. L. C. Vink, *Phys. Rev. E* **86**, 031923 (2012).
 - [17] R. L. C. Vink and T. Speck, *Soft Matter* **9**, 11197 (2013).
 - [18] T. Bihr, U. Seifert, and A.S. Smith, *Phys. Rev. Lett.* **109**, 258101 (2012).
 - [19] N. Weil and O. Farago, *Eur. Phys. J. E* **33**, 81 (2010).
 - [20] N. Dharan and O. Farago, *J. Chem. Phys.* **141**, 024903 (2014).
 - [21] H. Noguchi, *Europhys. Lett.* **102**, 68001 (2013).
 - [22] O. Farago, *Phys. Rev. E* **81**, 050902R (2010).
 - [23] I. R. Cooke, K. Kremer, and M. Deserno, *Phys. Rev. E* **72**, 011506 (2005).
 - [24] O. Farago and N. Grønbech-Jensen, *Biophys. J.* **92**, 3228 (2007).
 - [25] O. Farago, *J. Chem. Phys.* **128**, 184105 (2008).
 - [26] H. Noguchi, *Phys. Rev. E* **83**, 061919 (2011).
 - [27] W. K. den Otter, *J. Chem. Phys.* **123**, 214906 (2005).
 - [28] Y. Avital and O. Farago, To appear in *J. Chem. Phys.* Also see arXiv:1503.02818[cond-mat.soft].
 - [29] Notice that in ref. [26], the membrane is simulated in the fixed-area ensemble, $A_P = L_x L_y$, with $L_x \neq L_y$ which generates buckled configurations with anisotropic surface

tension. In this study, we simulate the fixed (isotropic) tension ensemble, where the projected area A_P is allowed to fluctuate, but with $L_x = L_y$.

[30] Out of 15 independent realizations of the system corresponding to $\epsilon = 0.6k_B T$, in more than half we observed one dominant long Fourier mode. For larger values of ϵ ,

realizations where both modes were simultaneously excited occurred more frequently.

[31] M. Staykova, M. Arroyo, M. Rahimi, and H. A. Stone, Phys. Rev. Lett. **110**, 028101 (2013).