

# PCCP

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE TYPE

Cite this: DOI: 10.1039/xxxxxxxxxx

## Comparing molecules and solids across structural and alchemical space<sup>†</sup>

Sandip De,<sup>a,b</sup> Albert P. Bartók,<sup>c</sup> Gábor Csányi,<sup>c</sup> and Michele Ceriotti <sup>\*a,b</sup>Received Date  
Accepted Date

DOI: 10.1039/xxxxxxxxxx

www.rsc.org/journalname

Evaluating the (dis)similarity of crystalline, disordered and molecular compounds is a critical step in the development of algorithms to navigate automatically the configuration space of complex materials. For instance, a structural similarity metric is crucial for classifying structures, searching chemical space for better compounds and materials, and driving the next generation of machine-learning techniques for predicting the stability and properties of molecules and materials. In the last few years several strategies have been designed to compare atomic coordination environments. In particular, the Smooth Overlap of Atomic Positions (SOAP) has emerged as an elegant framework to obtain translation, rotation and permutation-invariant descriptors of groups of atoms driven by the design of various classes of machine-learned inter-atomic potentials. Here we discuss how one can combine such local descriptors using a Regularized Entropy Match (REMatch) approach to describe the similarity of both whole molecular and bulk periodic structures, introducing powerful metrics that enable the navigation of alchemical and structural complexity within a unified framework. Furthermore, using this kernel and a ridge regression method we can predict atomization energies for a database of small organic molecules with a mean absolute error below 1 kcal/mol, reaching an important milestone in the application of machine-learning techniques to the evaluation of molecular properties.

### 1 Introduction

The increase of available computational power, together with the development of more accurate and efficient simulation algorithms, have made it possible to reliably predict the properties of materials and molecules of increasing levels of complexity. Furthermore, high-throughput computational screening of existing and hypothetical compounds promises to dramatically accelerate the development of materials with the better performances or custom-tailored properties<sup>1–6</sup>.

These developments have made even more urgent the need for

automated tools to analyze, classify<sup>7–11</sup> and represent<sup>12–16</sup> large amounts of structural data, as well as techniques to leverage this wealth of information to estimate inexpensively the properties of materials using machine-learning techniques, circumventing the need for computationally demanding quantum mechanical calculations<sup>17–28</sup>.

At the most fundamental level, the crucial ingredient for all these techniques is a mathematical formulation of the concept of (dis)similarity between atomic configurations, that can take the form of a distance - that can be used for dimensionality reduction or clustering - or of a kernel function, that could be used for ridge regression or automated classification.<sup>29–32</sup> The most obvious choice for a metric to compare atomic structures would involve the Euclidean distance between the Cartesian coordinates of the atoms, commonly known as root mean square displacement (RMSD) distance, that can be easily made invariant to relative translations and rotations. It is however highly non-trivial to extend the RMSD to deal with situations in which atoms in the two structures cannot be mapped unequivocally onto each other. The

<sup>a</sup> National Center for Computational Design and Discovery of Novel Materials (MARVEL)

<sup>b</sup> Laboratory of Computational Science and Modelling, Institute of Materials, Ecole Polytechnique Fédérale de Lausanne, Lausanne, Switzerland; E-mail: michele.ceriotti@epfl.ch

<sup>c</sup> Engineering Laboratory, University of Cambridge, Trumpington Street, Cambridge CB2 1PZ, United Kingdom

<sup>†</sup> Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/b000000x/

deterministic evaluation of a “permutationally invariant” RMSD scales combinatorially with the size of the molecules to be compared<sup>33</sup>, and introduces cusps at locations where the mapping of atom identities changes. Furthermore, as we will discuss later on, the RMSD is perhaps the most straightforward, but not necessarily the most flexible or effective strategy to compare molecular and condensed-phase configurations.

In the last few years, a large number of “fingerprint” functions have been developed to represent the state of structures, or of groups of atoms within a structure. Structural descriptors have been developed based on graph-theoretic procedures (e.g. SPRINTS<sup>34</sup>), as well as on analogies with electronic structure methods (e.g. Hamiltonian matrix, Hessian matrix, Overlap matrix of Gaussian type Orbitals (GTO) or even Kohn-Sham eigenvalues fingerprints<sup>33</sup>). Most of these approaches have been introduced to provide a fast and reliable estimate of the dissimilarity between structures. Several other descriptors have been also used in machine learning, to predict properties of materials and molecules circumventing the need for an expensive electronic structure calculation. A non-comprehensive list of such methods include Coulomb matrices<sup>17</sup>, bags of bonds<sup>28</sup>, “symmetry functions”<sup>35</sup>, scattering transformation applied on a linear superposition of atomic densities<sup>23</sup>.

A particularly promising approach to compare structures in a way that is invariant to rotations, translations, and permutations of equivalent atoms, is to start from descriptors designed to represent *local* atomic environments and that fulfill these requirements, and combine them to yield a *global* measure of similarity between structures. This idea typically relies on finding the best match between pairs of environments in the two configurations<sup>22,33,36</sup>, and can also be traced back to methods developed to compare images based on the matching of local features<sup>37</sup>.

In the present work we start from a recently-developed strategy to define a similarity kernel between local environments – the smooth overlap of atomic positions (SOAP)<sup>38</sup> – and discuss the different ways one can process the set of all possible matchings between atomic environments to generate a global kernel to compare two structures. In particular, we introduce a regularized entropy match (REMatch) strategy that is based on techniques in optimal-transport theory<sup>39</sup>, and that is both more efficient and tunable than previously-applied methods. We discuss the relative merits of different approaches, and generalize this strategy to the comparison between structures with different numbers and kinds of atoms. We demonstrate the behavior of the different global kernels when applied to completely different classes of problems, ranging from elemental clusters, to bulk structures, to the conformers of oligopeptides and to a heterogeneous database of small organic molecules. We visualize the behavior of the distance associated with these kernels using sketch-map<sup>13</sup>, a non-linear dimensionality reduction technique,

and demonstrate the great promise shown by the straightforward application of the REMatch-SOAP kernel to the machine-learning of molecular properties. Finally, we present our conclusions.

## 2 Theory

Let us start by introducing the notation we will employ in the rest of the paper. We will label structures to be compared by capital letters, use a lowercase Latin letter to indicate the index of an atom, and when necessary use a Greek lowercase letter to mark its chemical identity. For instance, the position of the  $i$ -th atom within the structure  $A$  will be labeled as  $\mathbf{x}_i^A$ . The *environment* of that atom, i.e. the abstract descriptor of the arrangement of atoms in its vicinity will be labelled with a calligraphic upper case letter, e.g.  $\mathcal{X}_i^A$ , and the sub-set of such environment that singles out atoms of species  $\alpha$  will be indicated as  $\mathcal{X}_i^{A,\alpha}$ .

Among the many descriptors of local environments that have been developed in the recent years<sup>1–3,5,6,17–22,24–28,33,36</sup>, we will refer in particular to the SOAP fingerprints<sup>38</sup>, that have been proven to be a very elegant and robust strategy to describe coordination environments in a way that is naturally invariant with respect to translations, rotations and permutations of atoms. We will use the notation  $k(\mathcal{X}, \mathcal{X}')$  to indicate the similarity kernel (normalized to one) between two environments – which one would use in a kernel ridge regression method<sup>31,32,40</sup> – and  $d(\mathcal{X}, \mathcal{X}')^2 = 2 - 2k(\mathcal{X}, \mathcal{X}')$  to indicate the (squared) kernel distance between the environments – which one would use in a dimensionality reduction method<sup>13,16</sup>. In what follows we will discuss different ways by which environment kernels can be combined to yield a *global* similarity kernel between two structures  $K(A, B)$ , and the associated squared distance  $D(A, B)^2 = 2 - 2K(A, B)$ .

### 2.1 SOAP similarity kernels and local environment distance

We will first focus on the comparison between the environment of two atoms in a pure compound made up of a single atomic species  $\alpha$ . The crucial ingredient in making the comparison is a kernel function based on the distribution of atoms in the two environments. In the context of SOAP kernels one represents the local density of atoms within the environment  $\mathcal{X}$  as a sum of Gaussian functions with variance  $\sigma^2$ , centered on each of the neighbors of the central atom, as well as on the central atom itself:

$$\rho_{\mathcal{X}}(\mathbf{r}) = \sum_{i \in \mathcal{X}} \exp\left(-\frac{(\mathbf{x}_i - \mathbf{r})^2}{2\sigma^2}\right). \quad (1)$$

The SOAP kernel is then defined as the overlap of the two local atomic neighbour densities, integrated over all three-dimensional rotations  $\hat{R}$ ,

$$\tilde{k}(\mathcal{X}, \mathcal{X}') = \int d\hat{R} \left| \int \rho_{\mathcal{X}}(\mathbf{r}) \rho_{\mathcal{X}'}(\hat{R}\mathbf{r}) d\mathbf{r} \right|^n. \quad (2)$$

Note that in the  $n = 1$  case the two integrals can be switched, and therefore the kernel loses all angular information, so we focus on the  $n = 2$  case exclusively. For most applications it is helpful to normalise the kernel so that the self-similarity of any environment is unity, giving the final kernel

$$k(\mathcal{X}, \mathcal{X}') = \tilde{k}(\mathcal{X}, \mathcal{X}') / \sqrt{\tilde{k}(\mathcal{X}, \mathcal{X})\tilde{k}(\mathcal{X}', \mathcal{X}')} \quad (3)$$

It is a remarkable property of the SOAP kernel that the integration over all rotations can be carried out analytically. First, the atomic neighbour density is expanded in a basis composed of spherical harmonics and a set of orthogonal radial basis functions  $\{g_b(r)\}$ ,

$$\rho_{\mathcal{X}}(\mathbf{r}) = \sum_{blm} c_{blm} g_b(|\mathbf{r}|) Y_{lm}(\hat{\mathbf{r}}), \quad (4)$$

then the rotationally invariant *power spectrum* is given by

$$p(\mathcal{X})_{b_1 b_2 l} = \sum_m (c_{b_1 l m})^\dagger c_{b_2 l m}. \quad (5)$$

Collecting the elements of the power spectrum into a unit-length vector  $\hat{\mathbf{p}}(\mathcal{X})$ , the SOAP kernel is shown<sup>38</sup> to be given by

$$k(\mathcal{X}, \mathcal{X}') = \hat{\mathbf{p}}(\mathcal{X}) \cdot \hat{\mathbf{p}}(\mathcal{X}') \quad (6)$$

eventually leaving a definition of the distance as

$$d(\mathcal{X}, \mathcal{X}') = \sqrt{2 - 2\hat{\mathbf{p}}(\mathcal{X}) \cdot \hat{\mathbf{p}}(\mathcal{X}')} \quad (7)$$

The SOAP kernel can be written in the form of a dot product, therefore it is manifestly positive definite, which implies that the distance function (7) is a proper metric.

## 2.2 From local descriptors to structure matching

The vectors that enter the definition of the environments are defined in such a way that their dot product is the overlap of (smoothed) atomic distributions. Given two structures with the same number  $N$  of atoms, we can compute an *environment covariance matrix* that contains all the possible pairings of environments

$$C_{ij}(A, B) = k(\mathcal{X}_i^A, \mathcal{X}_j^B), \quad (8)$$

This matrix contains the complete information on the pair-wise similarity of all the environments between the two systems. Based on it, one can introduce a global kernel to compare two structures or molecules. We will discuss and compare four different approaches. All of them are meant to be normalized, i.e. the given expressions for  $K(A, B)$  are to be divided by  $\sqrt{K(A, A)K(B, B)}$  whenever the kernel is not normalized to one by construction.

**Average structural kernel** A first possibility to compare two structures involves computing an *average kernel*

$$\begin{aligned} \bar{K}(A, B) &= \frac{1}{N^2} \sum_{ij} C_{ij}(A, B) = \\ &= \left[ \frac{1}{N} \sum_i \mathbf{p}(\mathcal{X}_i^A) \right] \cdot \left[ \frac{1}{N} \sum_j \mathbf{p}(\mathcal{X}_j^B) \right]. \end{aligned} \quad (9)$$

One sees that  $\bar{K}$  can be computed inexpensively by just storing the average SOAP fingerprint between all environments of the two structures. This kernel is also positive-definite, being based on a scalar product<sup>41</sup>, and therefore induces a metric  $\bar{D}(A, B) = \sqrt{2 - 2\bar{K}(A, B)}$ . On the other hand, it is not a very sensitive metric: two very different structures can appear to be the same if they are composed of environments that give the same fingerprint upon averaging.

**Best-match structural kernel** Another possibility, that has been used previously with different kinds of structural fingerprints<sup>22,33,42,43</sup> is to identify the best match between the environments of the two structures,

$$\hat{K}(A, B) = \frac{1}{N} \max_i \sum_j C_{ij}(A, B). \quad (10)$$

which can be accomplished with an  $\mathcal{O}(N^3)$  effort using the Munkres algorithm<sup>44</sup>. The corresponding distance has the properties of a metric, which means it can still be safely used to assess similarity between structures and molecules. Unfortunately, this “best-match” kernel is not guaranteed to be positive-definite, which makes it less than ideal for use in machine-learning applications. Furthermore, the distance obtained by a best-match strategy is continuous, but has discontinuous derivatives whenever the matching of environments changes. These problems can be solved or alleviated by matching the environments based on a different strategy, that combines features of the average and the best-match kernels.

**Regularized entropy match kernel** The best match problem can be also stated in an alternative form, namely

$$\hat{K}(A, B) = \max_{\mathbf{P} \in \mathcal{Z}(N, N)} \sum_{ij} C_{ij}(A, B) P_{ij}. \quad (11)$$

where  $\mathcal{Z}(N, N)$  is the set of  $N \times N$  (scaled) doubly stochastic matrices, whose rows and columns sum to  $1/N$ , i.e.  $\sum_i P_{ij} = \sum_j P_{ij} = 1/N$ . We can then borrow an idea that was recently introduced in the field of optimal transport<sup>39</sup> to *regularize* this problem, adding a penalty that instead aims at maximizing the information entropy for the matrix  $\mathbf{P}$  subject to the aforementioned constraints on its marginals. Such “regularized-entropy match” (REMatch)

kernel is defined as

$$\begin{aligned} \hat{K}^\gamma(A, B) &= \text{Tr} \mathbf{P}^\gamma \mathbf{C}(A, B), \\ \mathbf{P}^\gamma &= \underset{\mathbf{P} \in \mathcal{U}(N, N)}{\text{argmin}} \sum_{ij} P_{ij} (1 - C_{ij} + \gamma \ln P_{ij}), \end{aligned} \quad (12)$$

where the regularization is given by an entropy term  $E(\mathbf{P}) = -\sum_{ij} P_{ij} \ln P_{ij}$ .  $\mathbf{P}^\gamma$  can be computed very efficiently, with  $\mathcal{O}(N^2)$  effort, by the Sinkhorn algorithm<sup>39</sup> (see Appendix C). For  $\gamma \rightarrow 0$ , the entropic penalty becomes negligible, and  $\hat{K}^\gamma(A, B) \rightarrow \hat{K}(A, B)$ . For  $\gamma \rightarrow \infty$ , one selects the  $\mathbf{P}$  with the least information content, that is one with constant  $P_{ij} = 1/N^2$ . Hence, in this limit  $\hat{K}^\gamma(A, B) \rightarrow \bar{K}(A, B)$ .

**Permutation structural kernel** For the sake of completeness, we also discuss a fourth option: rather than summing over all possible pairs of environments, one can consider each pairing of environments separately, and sum over all the  $N!$  possible permutations that define the pairings. In order to kill off more rapidly the combinations of environments that contain bad matches, one can *multiply* the kernels that appear in each pairing, and define a *permutation kernel*

$$\check{K}(A, B) = \frac{1}{N!} \sum_{\pi} \prod_i C_{i\pi_i}(A, B) = \text{perm} \mathbf{C}(A, B). \quad (13)$$

This choice corresponds to the evaluation of the permanent of the environment kernel matrix, and has some appeal as it is guaranteed to yield a positive-definite kernel<sup>45</sup>. The evaluation of the permanent of a matrix, however, has combinatorial computational complexity\*. Its application is limited to small molecules, and we will not discuss it further in the present work.

### 2.3 Matching structures containing multiple species

When comparing structures that contain different atomic species, the first problem that has to be addressed is that of extending the local environment metric so that the presence of multiple elements is properly accounted for.

SOAP descriptors provide a straightforward way to do this: a separate density can be built for each atomic species

$$\rho_{\mathcal{X}}^\alpha(\mathbf{r}) = \sum_{i \in \mathcal{X}_\alpha} \exp\left(-\frac{(\mathbf{x}_i - \mathbf{r})^2}{2\sigma^2}\right), \quad (14)$$

and a (non-normalized) kernel be defined by matching separately

the different species:

$$\begin{aligned} \tilde{k}(\mathcal{X}, \mathcal{X}') &= \int d\hat{\mathbf{R}} \left| \int \sum_{\alpha} \rho_{\mathcal{X}}^\alpha(\mathbf{r}) \rho_{\mathcal{X}'}^\alpha(\hat{\mathbf{R}}\mathbf{r}) d\mathbf{r} \right|^2 \\ &= \sum_{\alpha\beta} \mathbf{p}_{\alpha\beta}(\mathcal{X}) \cdot \mathbf{p}_{\alpha\beta}(\mathcal{X}'). \end{aligned} \quad (15)$$

Here we have introduced “partial” power spectra  $\mathbf{p}_{\alpha\beta}$  that encode information on the relative arrangement of pairs of species, and can be written as

$$p(\mathcal{X})_{b_1 b_2 l}^{\alpha\beta} = \frac{8\pi^2}{2l+1} \sum_m (c_{b_1 l m}^\alpha)^\dagger c_{b_2 l m}^\beta, \quad (16)$$

where we built in the angular channel dependent weights into the elements of the power spectrum. The expansion coefficients describe the atomic density of species  $\alpha$

$$\rho_{\mathcal{X}}^\alpha(\mathbf{r}) = \sum_{blm} c_{blm}^\alpha g_b(|\mathbf{r}|) Y_{lm}(\hat{\mathbf{r}}) \quad (17)$$

in terms of a basis set, which is a combination of spherical harmonics and orthogonal radial functions. The kernel in Eq. (15) can then be normalized as in Eq. (3).

Note that, even though the overlap between the environments of the different species is considered to be zero, the kernel is sensitive the relative correlations of different species. This is because, due to the squaring of the density overlap within the rotational average, the SO(3) power spectrum vectors contain mixed-species components. One could also introduce a notion of “alchemical similarity” between different species. For instance, when comparing structures of III-V semiconductors one could disregard the chemical information on the identity of an atom as long as it belongs to the same column of the periodic table. Such a notion can be readily implemented, defining an alchemical similarity kernel  $\kappa_{\alpha\beta}$  which is one for pairs that should be considered interchangeable, and tend to zero for pairs that one wants to consider as completely unrelated. The expression then becomes

$$\begin{aligned} \tilde{k}(\mathcal{X}, \mathcal{X}') &= \int d\hat{\mathbf{R}} \left| \int \sum_{\alpha\alpha'} \kappa_{\alpha\alpha'} \rho_{\mathcal{X}}^\alpha(\mathbf{r}) \rho_{\mathcal{X}'}^{\alpha'}(\hat{\mathbf{R}}\mathbf{r}) d\mathbf{r} \right|^2 \\ &= \sum_{\alpha\beta\alpha'\beta'} \mathbf{p}_{\alpha\beta}(\mathcal{X}) \cdot \mathbf{p}_{\alpha'\beta'}(\mathcal{X}') \kappa_{\alpha\alpha'} \kappa_{\beta\beta'}. \end{aligned} \quad (18)$$

The original expression (15) can be recovered by setting  $\kappa_{\alpha\beta} = \delta_{\alpha\beta}$ . Global similarity kernels can then be transparently introduced to compare structures composed of different atomic species, with geometry and alchemical composition treated on the same footings and the possibility of adapting the definition of similarity to the system and application.

\*Although stochastic algorithms do exist to compute it to a desired precision in polynomial time<sup>46</sup>

## 2.4 Matching structures with different numbers of atoms

The definitions above can be readily extended to compare structures containing different numbers of atoms  $N_A$  and  $N_B$ . We discuss two possible strategies. When comparing crystalline, periodic structures, it may be the case that one of the structures corresponds to a slight distortion of the other, that needs a larger unit cell for a proper representation. Comparing the structures using the average kernel (9) does automatically the “right thing”, that is performing the comparison in a way that is independent of the number of times the two structures have to be replicated to match atom counts. In the case of the permutation kernel and of the best-match kernel, the most effective way to perform the comparison is to evaluate the least common multiple  $N$  of  $N_A$  and  $N_B$ , and replicate the environment similarity matrix to form a square matrix. One can then proceed to compute the permanent, or the linear assignment problem, based on such replicated matrix. The advantage of this procedure is that one does not need to explicitly find the relation between the shape of the two unit cells and replicate them to perform the comparison: the environment similarities can be evaluated including periodic replicas, and the minimum number of comparisons will be naturally performed among any pairs of structures. However, the least common multiple can become very large, making even the best-match kernel (10) impractically demanding, although the cost can be reduced by exploiting the redundancy in the extended environment covariance matrix. As shown in the Appendix, the REMatch kernel (12) can be computed easily also for a rectangular matrix, which constitutes an additional advantage of formulating the environment matching problem in terms of a regularized transport optimization.

When comparing molecules or molecular fragments, it may be advisable to proceed differently – since in that case the chemical composition might differ, and it may not make sense to compare molecules as if they were part of an infinite periodic assembly. A possible strategy is then to consider, given a molecular database, the smallest pool (“kit”) of atoms from which every molecule in the set can be constructed. Then, when comparing each pair of structures, the atoms that are not needed to form either of the two molecules would still be part of the comparison, in the form of idealized “isolated” species. Alternatively, for instance when the full database is not known a priori, such “reference kit” could be chosen dynamically for each pair of molecules. Since the SO(3) fingerprints that underlie the definition of the SOAP kernel can also be evaluated for isolated atoms<sup>†</sup>, it is then possible to introduce a natural definition of the covariance between an environment and an isolated atom. One of the advantages of such approach is that the global kernels will then vary smoothly if a

molecule is continuously broken up into its constituent atoms, which lends itself to a very effective description of atomization processes.

## 2.5 Representing (al)chemical landscapes

In this work we will demonstrate the flexibility, transferability and effectiveness of the framework we have just introduced to compare molecular and condensed-phase structures. To this aim, we will build two dimensional maps that represent proximity relations between the structures – as assessed by the kernel-induced metric – using sketch-map<sup>13</sup>, a non-linear dimensionality reduction (NLDR) scheme specifically designed to deal with atomistic simulation data. As we will demonstrate, the combination of SOAP-based structural metrics and NLDR representation provides a broadly applicable protocol to generate an insightful representation of the structural and alchemical landscape of complex molecular and condensed-phase systems. Of course, one could use the SOAP-based global kernels, or the corresponding distances, as the basis of other non-linear dimensionality reduction techniques, such as multi-dimensional scaling<sup>47</sup> or diffusion maps<sup>12,16,48</sup>.

We refer the reader to the relevant literature for a detailed explanation of the sketch-map algorithm<sup>13–15</sup>. The main idea derives from multi-dimensional scaling, and is based on optimizing a non-linear objective function


$$S^2 = \sum_{ij} [F[D(X_i, X_j)] - f[d(x_i, x_j)]]^2 \quad (19)$$

where  $\{X_i\}$  and  $\{x_i\}$  correspond respectively to high-dimensional reference structures and to vectors in a low-dimensional space. The metric  $d$  in low dimension is typically taken to be the Euclidean distance, whereas the metric in high dimension could be more complex. In this case,  $X_i$  can be regarded as an abstract descriptor of a structure or molecule, and  $D$  is one of the kernel-based distance metrics discussed above.  $F$  and  $f$  are non-linear sigmoid functions of the form

$$F(r) = 1 - (1 + (2^{a/b} - 1)(r/\sigma)^a)^{-b/a}, \quad (20)$$

which serve to focus the optimization of (19) on the most significant, intermediate distances, disregarding local distortion (e.g. induced by thermal fluctuations) and the relation between completely unrelated portions of configurational landscape. The choice of the parameters in the sigmoid functions is discussed in Ref.<sup>15</sup>. Here we will label synthetically each sketch-map representation using the notation  $\sigma_{-A\_B-a\_b}$  where  $A$  and  $B$  denote the exponents used for the high-dimensional function  $F$ ,  $a$  and  $b$  denote the exponents for the low-dimensional function  $f$ , and  $\sigma$  the threshold for the switching function. Open-source software to perform the dimensionality-reduction step, as well as to compute the different similarity kernels we have introduced, is available

<sup>†</sup> The density of atoms defined in equation (1) contains the central atom.



**Fig. 1** The figure compares the value of global structural similarities for different pairs of structures taken from the 80 local energy minima of  $C_{60}$  discussed in Ref. <sup>42</sup>. The structural similarities considered include the absolute difference in energy per atom, the (permutation invariant<sup>33</sup>) RMSD per atom, and the best-match combination of SOAP kernels computed with different cutoff distances (2Å, 3.5Å, 7Å). The correlation between RMSD and  $\hat{D}$  based on 2Å-cutoff SOAP is enlarged, color-coded based on energy differences and annotated with selected pairs of structures corresponding to different distances.

from the authors upon request. Interactive versions of the structural maps discussed in the text are provided in the supporting information (SI), and are available as an on-line repository<sup>49</sup>.

### 3 Examples and applications

After having described the theoretical and algorithmic background of our strategy to define a structural similarity kernel, let us present a series of applications. In order to demonstrate that our approach can be seamlessly applied to the most diverse atomistic simulation problems, we have chosen examples of increasing complexity, from clusters, to crystalline and amorphous solids, to biological molecules and a database of small organic compounds, containing varying number of both atoms and atomic species. As we will discuss, SOAP-based structural kernels contain several adjustable hyperparameters, that can be regulated to focus the dissimilarity measure onto the desired features. Unless otherwise specified, however, we have not explored fully this possibility, and we have simply chosen reasonable values of the parameters without much fine-tuning.

#### 3.1 The energy landscape of C<sub>60</sub> clusters

Let us start with a relatively simple test case. We consider the same set of 80 local minima for C<sub>60</sub> discussed in Ref.<sup>42</sup>, which were obtained by exploring the Density Functional Theory energy landscape of C<sub>60</sub> using the Minima Hopping<sup>50</sup> global structure search algorithm. Figure: 1 contrasts different similarity matrices: the permutation-invariant RMSD<sup>33</sup>, the absolute difference between the potential energy, and the best-match distances obtained from SOAP descriptors computed with different environment cutoff. RMSD distance does not correlate very well with SOAP-based metrics, particularly for the smaller cutoff value. The  $\bar{D}(2\text{\AA})$ -RMSD correlation plot is enlarged, and allows us to discuss the source of this discrepancy. Hollow fullerene-like structures (A, with reference to the labeling in the figure) and compact structures containing internal connections (A,G) are extremely different from the point of view of the short-range connectivity, but differ comparatively less in terms of RMSD, since they are both fairly compact. On the other hand, flake-like structures based on a honeycomb motif (F,E) have the same basic first-neighbor connectivity as the defective fullerene structures (C,D) but have much different spatial extent. Then, one sees that the discrepancy between RMSD and small-cutoff  $\hat{D}$  indicates just the focus on different structural features: the global arrangement of atoms in the first case, and the local connectivity in the latter. In the case of SOAP-based metrics, however it is easy to extend the sensitivity of the metric to longer distances just by increasing the cutoff: by going from 2Å to 3.5 and 7, one sees that  $\hat{D}$  and RMSD become progressively more correlated, as the focus shifts from the nearest-neighbor coordination to the overall geometry of the cluster.


It is worth stressing that the RMSD, albeit a very natural measure of structural similarity, is not necessarily the best metric to compare configurations. To see why, consider the absolute energy difference as a measure of similarity: even though one can obviously have configurations with very different geometry and similar energies, in general one would expect that on the contrary large energy differences should be associated with highly dissimilar structures in a given system – which is not the case for RMSD. One sees that the intermediate-cutoff  $\bar{D}(3.5\text{\AA})$  shows a nice correlation between energetic and structural differences.

These considerations underline a theme that will recur in other examples: SOAP-based structural metrics offer a mathematically sound framework that can be transparently adapted to focus on the aspects that are most relevant to a given application. For instance, power-spectrum based environment kernels are invariant to mirror symmetry, and therefore the derived metrics cannot distinguish enantiomers. If one needed to do so, however, it would be sufficient to use a bispectrum-based SOAP kernel<sup>38</sup> – which corresponds to  $n = 3$  in eq. (2) and is invariant to rotations but not to mirror symmetry operations – as the basis for obtaining a global comparison that is sensitive to chirality.


Having established a connection between traditional structural similarity metrics and the best-match SOAP kernel, let us use the example of C<sub>60</sub> to compare the three main strategies we propose to build a global kernel: the average kernel  $\bar{K}$ , the best-match kernel  $\hat{K}$ , and the regularized entropy match kernel  $\hat{K}^\gamma$  with an intermediate regularization parameter  $\gamma = 0.1$ .

The distance-distance correlation plot for each pair of structures, that compares the distances induced by the three kernels, is reported in Fig.2. The  $\bar{D} - \hat{D}$  plot shows overall linear correlation except for very small values of  $\bar{D}$ . This is expected as the average kernel is under-determined, and could in principle label two structures as identical even though they might be composed of different environments. The best-match kernel, therefore, provides better resolving power. As we will discuss in more detail later on, the regularized best-match kernel  $\hat{D}^\gamma$  can be tuned to interpolate between these two extremes. As an example, we chose here an intermediate value  $\gamma = 0.1$ : as shown in Fig. 2, the resulting distance correlates strongly with both  $\bar{D}$  and the conventional best-match distance  $\hat{D}$ .

Fig.2 also shows annotated sketch-maps obtained based on the three metrics. Once the sketch-map parameters have been adjusted following the guidelines in Ref.<sup>15</sup>, the three maps are effectively equivalent – indicating that the three kernels give similar qualitative information on the similarity between different structures. Given the much lower computational cost associated with the evaluation of the average kernel, this observation suggests it might conveniently be used to preliminarily screen a dataset before proceeding to a more accurate comparison of similar structures based on the best-match, or REMatch distance.



**Fig. 2** The figure compares the value of global structural distances induced by the average, best-match, and REMatch kernels discussed in Section 2.2, for 80 local-minimum structures of  $C_{60}$ . On the diagonal we report the sketch-map projections of the structural landscape based on the three metrics, colored according to the energy of each structure, as obtained by Sandip et al.<sup>42</sup>. Eight representative structures and their positions on the Sketch-maps have been indicated with letters on color coded disks. The numeric value on the top of each structure represents their energy in eV, relative to the global minimum. SOAP descriptors were computed using a cutoff of 3.5Å and the Sketch-map parameters are indicated on the map according to the syntax described in the text.



**Fig. 3** Sketch-map of 1274 crystalline and amorphous silicon structures obtained by sampling different phases from the phase diagram (disks), polymorphs obtained by ab initio random structure search<sup>51</sup> (+ signs) and by minima hopping<sup>52</sup> (x signs). The color and size of the points varies according to their atomic energy and atomic volumes respectively. Regions of the plot which represents different phases have been outlined with dotted contours.

### 3.2 Natural and hypothetical polymorphs of silicon

As a second example, let us move on to a condensed-phase application. Here we start from a database of 1274 bulk silicon structures containing ideal and distorted configurations from the phase diagram (e.g. diamond, simple hexagonal,  $\beta$ -tin, liquid and quenched amorphous structures). SOAP environment kernels with a 5 Å cutoff distance were used, and combined with a best-match strategy to obtain the (dis)similarity matrix ‡ We selected 100 landmark configurations out of this data set (using farthest point sampling based on kernel distance) and built a sketch-map, on which the rest of configurations were projected. The outcome of such mapping procedure is shown in Fig. 3, where points are colored according to the DFT atomic energy, and point sizes are scaled to a size proportional to volume per atom. As seen in the Fig. 3 the map is extremely well correlated with both atomic energy and density. Furthermore, structures that were obtained by distorting and heating up structures coming from different portions of the phase diagram are clustered together: rough outlines have been drawn on the map to indicate different phases.

Although the map has been built using only reference configurations from a few of the conventional Si phases, we have also projected on it (using out-of-sample embedding) two sets of hypothetical configurations obtained by minima hopping<sup>52</sup> and by ab initio random structure search (AIRSS)<sup>51,53</sup>. These structures were not included in the landmarks selection phase. Still, the out-of-sample embedding procedure correctly identifies not only that in most cases AIRSS structures differ significantly from stable phases of silicon, but also clusters together hypothetical polymorphs that share common features. For instance, the AIRSS structures outlined in the lower portion of the map are all taken from Ref.<sup>53</sup>. The structures were proposed as possible metastable polymorphs arising as a result of a microexplosion (induced by powerful, ultrashort and tightly focused laser pulses) in crystalline cubic diamond silicon phase, hence their structural motif naturally carries resemblance with silicon diamond phase. It is interesting to see that they indeed are projected close to the diamond phase on the map. All of the minima hopping low-density Si polymorphs are also clustered together, which is consistent with the fact that they are all based on combinations of a few base motifs. Thus, Figure 3 shows not only that SOAP-based structural similarity distances can be very effective in the study of bulk crystalline structures, but also testifies the extrapolative power of a sketch-map representation based on such a metric.

‡ Some of the structures in the overall data set have numbers of atoms in the unit cell that would lead to a large least common multiple when repeating the environment similarity matrix to form a square matrix. One can keep the cost of computing the similarity matrix low by exploiting the redundancy in the similarity matrix, or by approximating  $\hat{K}$  using a REMatch kernel with a small entropy regularization.

### 3.3 Arginine Dipeptide

Having shown that SOAP-based structural similarity kernels are equally effective for clusters and for bulk configurations of elemental materials, let us consider a case of a multi-species chemical compound. We selected a library of 5062 locally stable conformers of arginine dipeptide (845 with and 4217 without a  $\text{Ca}^{2+}$  counterion) from a public database of oligopeptides structures developed by Ropo et al<sup>54</sup>. We used a cut-off of 3.5Å in the definition of environment SOAP kernels, and combined them using a best-match strategy. Since H atoms stay at almost fixed positions relative to their neighboring atoms, we decided to include them in the environment descriptors of other atoms, but did not include them explicitly as centers of atomic environments. This is another example of how SOAP-based structural metrics are effective in a broad variety of contexts, but at the same time can be easily and transparently refined based on intuition, prior experience, or a clear understanding of the objectives of the structural comparison.

In Fig. 4 we show the sketch-map representation for these two sets of structures, highlighting the correlation between the location on the map and structural and energetic properties of the conformers. In the absence of a complexing cation, the dipeptide can exist in a very large number of local minima, spanning a relatively narrow range of energies. The map shows very clearly partitioning of configuration space in four disconnected regions. Conventional wisdom<sup>55</sup> assumes that the  $\text{C}_\alpha$  dihedral angles  $\phi$  and  $\psi$  are the most important descriptors of oligopeptide structure. One quickly realizes, however, that the order parameters corresponding to the four lobes are connected to the cis-trans isomerization of the two peptide bonds. Within each of the lobes, configurations with different  $\phi$ - $\psi$  dihedral angles are clearly clustered together, but in this case they constitute features of secondary importance. This observation demonstrates the advantages of using a general-purpose descriptor, that does not rely on pre-conceived assumptions on the behavior of the molecule being studied, but instead captures automatically the intrinsic structural hierarchy of minima in the configuration landscape.

The presence of a  $\text{Ca}^{2+}$  cation has a dramatic impact on the landscape for the dipeptide. The distribution of configurations becomes considerably more sparse and spans a broader range of energies. The strong electrostatic interaction with the cation means that there is not a clear separation anymore between the energy scale for  $\phi$ - $\psi$  flexibility of the backbone and the isomerization of the peptide bonds.

A remarkable observation in this analysis is the realization that the presence of the cation catalyzed unexpected proton transfer reactions, that change the chemical structure of the molecule. Configurations that underwent a chemical reaction are clustered on one side of the map (Fig. 5), with further internal structure reflecting the fact that SOAP-based structural metrics treat on the





Fig. 4 Energy landscape representation of locally stable arginine dipeptide conformers, without (top) and with (bottom) a  $\text{Ca}^{2+}$  ion. Left-hand panels are colored according to the energy relative to the minimum energy form, while the smaller maps on the right are colored according to the values of different dihedral angles, as indicated in the legend.




**Fig. 5** Sketch-map representation of stable configurations of Arginine dipeptide complexed with a  $\text{Ca}^{2+}$  ion. The structures that have undergone a proton transfer reaction relative to the neutral molecule have been highlighted, and a few representative snapshots of the molecular structure are also reported.

same footing information on the chemical bonding and on the conformational variability of the molecule. It is again worth noting that by changing the cut-off value for the SOAP descriptors, one can “focus” the structural metric on different molecular features. A short cutoff of  $2\text{\AA}$  makes the chemically different structures stand out more as outliers – which would for instance be useful to detect automatically this kind of unwanted transitions in an automatically-generated data set – while on the contrary a longer cutoff would give more importance to the difference between collapsed and extended molecular conformers.


### 3.4 Mapping (al)chemical space

As a final example of the evaluation of a structural and alchemical similarity metric, and its use to represent complex ensembles of compounds, let us consider the QM7b database<sup>24</sup>. This set of compounds contains 7211 minimum-energy structures for small organic compounds containing up to seven non-hydrogen atoms (C, N, O, S, Cl), saturated with H to different degrees. This database constitutes a small fraction of a larger chemical library that contains millions of hypothetical structures screened for accessible synthetic pathways<sup>56</sup>.

This is an extremely challenging data set to benchmark a structural similarity metric: molecules differ by number of atoms, chemical composition, bonding and conformation. To simplify the description, we decided to use SOAP descriptors with a cut-



**Fig. 6** Correlations between structural similarity distances induced by the average kernel  $\bar{K}$ , the best-match kernel  $\hat{K}$ , and *regularized* best-match kernels  $\hat{K}^\gamma$  with different regularization parameters  $\gamma$ . Distances are computed between pairs of 200 structures, randomly selected from the QM7b database<sup>24,56</sup>.



**Fig. 7** Sketch-map representation of minimum-energy structures from a database of molecules containing up to 7 non-hydrogen atoms (C O N S Cl), and saturated with hydrogen to varying degrees<sup>24</sup>. Left-hand panels show the map colored according to the atomization energy as computed by DFT. In the right-hand images, the points are colored according to the number of constituent C, O, N, S atoms. The top row corresponds to an alchemical kernel that treats all species as different, the middle row treats all the non-H atoms as the same species, whereas the bottom row introduces an alchemical kernel that depends on the difference in electronegativity between species.

off of 3Å, and to include H atoms in the environments but not as environment centers, to simplify the description – considering also that in the case of arginine dipeptide this choice did not prevent clear identification of isomers that only differed by a proton transfer reaction. We used a best-match strategy to compare configurations, and topped them up with isolated atoms up to the maximum number of each species that is present in the database. This effectively corresponds to choosing a “kit” (in other terms, a fully atomized reference state) starting from which all of the compounds can be assembled.

This is a fairly extreme case for the application of our idea of compounding local structure matching to obtain a global structural metric, so it is worth returning on a comparison of the different strategies we proposed. Fig. 6 compares average and best-match distances to REMatch kernels using different regularization parameters. Despite the very different context, the outcome is similar to what we observed in Figure 2 for  $C_{60}$  clusters. The average kernel is reasonably well correlated with the more demanding best-match kernel, although in most cases it has poorer resolution. By varying  $\gamma$ , the regularized match distance  $\hat{D}^\gamma$  varies between these two extremes, and for  $\gamma < 1$  provides a smooth, inexpensive approximation to the best-match distance.

For the sake of simplicity (and given we reduced the size of the environment covariance matrix  $C$  not considering H atoms as environment centers) we used the conventional best-match distance for the rest of our analyses. As shown in Fig. 7, the SOAP-based metric nicely separates out “islands” with homogeneous composition in terms of the number of non-H atoms. Within each group of atoms, one can recognize some sub-structure, with configurations roughly arranged in terms of the atomization energy – which in turns strongly correlates with the degree of H saturation. As it can be seen from inspection of the database (see the SI) in many cases one can notice that structures with similar chemical skeleton (presence of cycles, chemical groups, etc.) are clustered close to each other in the map. However, it is of course very difficult to quantitatively assess how well the map corresponds to chemical intuition, and how much departures from it are to be considered a failure of the metric, of the sketch-map procedure or of the notion of “chemical intuition”.

Our objective here is more to demonstrate how the fingerprint-based structural metric we introduced can cope with widely different classes of problems, and how it can treat on the same footings alchemical and structural variability. As an example we have also computed the similarity matrix and mapped the QM7b landscape using a modified alchemical similarity metric between the non-H atoms (we always take  $\kappa_{\alpha H} = \delta_{\alpha H}$ ). First, we set  $\kappa_{\alpha\beta} = 1$  (which means we are treating species  $\alpha$  and  $\beta$  as the same species) for all of atoms except H. The clear separation of the map into islands with the same stoichiometry is lost. However, there is now near-perfect correlation between position on

the map and atomization energy, and at the same time one can see some residual clustering of molecules with similar composition. This can be explained because information on the alchemical identity of the atoms is encoded in their atomic coordination and bond lengths. This is for instance evident for sulfur, that has considerably larger bond lengths, leading to better clustering of sulfur-containing compounds than in the case of oxygen or nitrogen.

Obviously, assuming that all atom kinds are interchangeable is an extreme choice, and it is hard to imagine circumstances in which this “element agnostic” metric would be advantageous over one that exploited knowledge of the chemical identity of atoms. On the other hand, one could foresee to encode information on the “alchemical similarity” using one of the many quantities chemists have used historically to rationalize trends in reactivity across the periodic table. As an example, we used the electronegativity  $E_\alpha$  to define

$$\kappa_{\alpha\beta} = e^{-(E_\alpha - E_\beta)^2 / 2\Delta^2} \quad (21)$$

where  $\Delta$  is a parameter that determines how sensitive is the alchemical kernel to differences in electronegativities. We used  $\Delta = 1$  to generate the last set of maps in Fig. 7. The map now separates out quite accurately regions with homogeneous stoichiometry. Whereas in the  $\kappa_{\alpha\beta} = \delta_{\alpha\beta}$  the different “islands” were roughly arranged according to a square grid pattern corresponding to  $n_O$  and  $n_N$  along two orthogonal directions, now stripe-shaped islands are arranged in 1D, following numbers of  $n_O$  and  $n_C$ , with the number of nitrogen atoms coming out clustered in adjacent “stripes”, but less clear-cut partitioning than for the other two elements. This is perhaps unsurprising given that nitrogen has an intermediate electronegativity between that of oxygen and carbon, and the metric tries to separate most efficiently the elements that differ most based on the alchemical similarity kernel.

This last example gives perhaps the most compelling demonstration of how a structural similarity metric based on a combination of SOAP kernels gives an effective, broadly applicable and easily customizable strategy to assess the similarity of materials and molecules, and how a sketch-map construction based on such metric provides an insightful representation of structural and alchemical landscapes.

### 3.5 Learning molecular properties

In this paper we focused mainly on the definition of a compound structural similarity kernel, and on characterizing its behavior by means of sketch-map representations. It is however important to keep in mind that an effective tool to compare atomic structures can find application to a broad range of problems - one of the most intriguing being the inexpensive prediction of physical-chemical properties of materials and molecules. To demonstrate

the great promise of REMatch-SOAP kernels for machine-learning of molecular properties, we used a standard kernel-ridge regression (KRR) method<sup>32</sup> to reproduce the 14 properties that had been reported in Ref.<sup>24</sup> for the 7211 molecules we described in the previous paragraph.

We randomly selected 5000 training structures, and used the remainder as an out-of-sample validation set. After having computed the REMatch-SOAP kernel matrix  $\mathbf{K}$  between all the structures, using a cutoff of 3 Å and a regularization parameter  $\gamma = 0.5$  – in this case including also H atoms in the list of environments – we computed the KRR weights vector

$$\mathbf{w} = \left( \mathbf{K}_{\text{train}}^{\xi} + \sigma \mathbf{1} \right)^{-1} \mathbf{y}_{\text{train}}. \quad (22)$$

Here  $\mathbf{K}_{\text{train}}$  and  $\mathbf{y}_{\text{train}}$  are the kernel matrix and property values restricted to the training set,  $\xi$  indicates entry-wise exponentiation to tune the spatial range of the kernel, and  $\sigma$  is a regularization hyperparameter. The prediction of the properties for the test set can then be obtained as  $\mathbf{y}_{\text{test}} = \mathbf{K}_{\text{test}}^{\xi} \mathbf{w}$ , where  $\mathbf{K}_{\text{test}}$  is the matrix containing the REMatch-SOAP kernels between the test points and the training points. The procedure was repeated 10 times, and the average mean absolute error (MAE) and root mean square error (RMSE) on the test set were computed.

We optimized the  $\xi$  and  $\sigma$  hyperparameters by minimising the MAE on the atomization energy, and then used the same values to perform a KRR for all the other molecular properties. Since we did not further adjust the choice of kernel and the  $\xi$  exponent, all the properties could be estimated at the same time, as discussed e.g. in Ref.<sup>57</sup>. The results of this procedure are reported in Table 1, and demonstrate the extraordinary performance of REMatch-SOAP for machine-learning applications. For the atomization energy we can obtain a MAE of less than 1 kcal/mol – a four-fold improvement relative to previous results that were based on a Coulomb matrix representation of structures and a deep-neural-network learning strategy. What is more, even without separately tuning the KRR hyperparameters, we can improve or match the performance of prior methods for almost all of the properties, the only exceptions being some of the properties computed with semi-empirical methods. The fact we can obtain such a dramatic improvement using a standard regression technique is a testament to the effectiveness of our kernel. The crucial importance of the choice of descriptors is also apparent by noting that a MAE of about 1.5 kcal/mol was recently obtained by regression based on a “bag of bonds” description of molecules, coupled with a Laplacian kernel<sup>28</sup>.

Reaching chemical accuracy in the automated prediction of atomization energies is an important milestone, and the fact that we could achieve that without fully exploring the flexibility of the REMatch-SOAP framework (e.g. by optimizing the entropy regularization parameter, the environment cutoff, eliminating the

Property	SD	MAE	RMSE	MAE <sup>24</sup>	RMSE <sup>24</sup>
$E$ (PBE0)	9.70	0.04	0.07	0.16	0.36
$\alpha$ (PBE0)	1.34	0.05	0.07	0.11	0.18
$\alpha$ (SCS)	1.47	0.02	0.04	0.08	0.12
HOMO (GW)	0.70	0.12	0.17	0.16	0.22
HOMO (PBE0)	0.63	0.11	0.15	0.15	0.21
HOMO (ZINDO)	0.96	0.13	0.18	0.15	0.22
LUMO (GW)	0.48	0.12	0.17	0.13	0.21
LUMO (PBE0)	0.68	0.08	0.12	0.12	0.20
LUMO (ZINDO)	1.31	0.10	0.15	0.11	0.18
IP (ZINDO)	0.96	0.19	0.28	0.17	0.26
EA (ZINDO)	1.41	0.13	0.18	0.11	0.18
$E_{1st}^*$ (ZINDO)	1.87	0.18	0.41	0.13	0.31
$E_{max}^*$ (ZINDO)	2.82	1.56	2.16	1.06	1.76
$I_{max}$ (ZINDO)	0.22	0.08	0.12	0.07	0.12

**Table 1** Mean absolute errors (MAEs) and root mean square errors (RMSE) for the KRR estimation of 14 molecular properties, together with previously published estimation<sup>24</sup> for the same data set. The standard deviation of the values of the properties across all 7211 molecules in the database is shown in the second column. Errors in the KRR estimation refer to a test set of 2200 randomly selected configurations, while the remaining structures were used for training. Property labels refer to the level of theory and molecular property, i.e. atomization energy ( $E$ ), averaged molecular polarizability ( $\alpha$ ), HOMO and LUMO eigenvalues, ionization potential (IP), electron affinity (EA), first excitation energy ( $E_{1st}^*$ ), excitation frequency of maximal absorption ( $E_{max}^*$ ) and the corresponding maximal absorption intensity ( $I_{max}$ ). Energies, polarizabilities and intensities are in eV, Å<sup>3</sup> and arbitrary units, respectively.

outliers, combining multiple layers of description or using a non-diagonal alchemical similarity matrix) highlights the potential of our approach. Future work will be devoted to analyzing the performance, convergence and limits of machine-learning of molecular and materials’ properties using our SOAP-based structural similarity kernel.

## 4 Conclusions

Distances between atomic structures based on combinations of local similarity kernels provide a flexible framework to define a metric in structural and alchemical space. Atom-centered environment information can be combined to provide a global measure of (dis)similarity. An average kernel  $\bar{K}$  provides an inexpensive strategy to do so, with a cost that scales linearly with the size of the structures to be compared, but might under-estimate the difference between two configurations – since in principle two different structures might yield zero  $\bar{D}$ . Alternatively, one can compute the local kernel between every possible pair of environments (which itself involves a cost scaling with the square of the number of environments), and then build a compound kernel  $\hat{K}$  by finding the best-match permutation of the environments – which gives a metric with better resolving power, but entails solving a cubic-

scaling linear assignment problem. Introducing an entropy regularization makes it possible at the same time to reduce the size-scaling to quadratic, and to obtain a better behaved, smoothly varying metric, that interpolates - depending on the regularization parameter - between the average and best-match limit.

This strategy to compare atomic configurations builds on the very general notion that complex bulk and molecular structures arise from the combination of local building blocks, and can be applied seamlessly to systems as diverse as clusters, bulk phases of an element, conformation of a biomolecules and an assembly of small chemical compounds with varying atom kinds and number. At the same time, the structure of the underlying SOAP kernels allows for very effective fine-tuning. For instance, by choosing the cutoff radius over which atomic densities are compared between environments, one can make the metric more sensitive to the first-neighbor chemical connectivity, or vice versa, include information on the long-range conformation of flexible molecules. What is more, it is possible to treat structural and alchemical complexity on the same footing, by introducing an alchemical similarity kernel that makes it possible to specify whether atoms of different species should be considered completely separate, or whether a notion of chemical distance (based e.g. on the difference in electronegativity) should be introduced to give different weights to substitutions between elements with similar reactivity.

We also demonstrate that straightforward application of the REMatch-SOAP kernel to the ridge-regression evaluation of molecular properties matches or outperforms all previously-presented approaches, reaching chemical accuracy in the prediction of the atomization energies of a set of small organic molecules. We believe that in this respect we are only scratching the surface of the potential applications to machine-learning of our kernels, since these results were obtained without using any of the more sophisticated techniques (e.g. introducing a hierarchy of models to capture the variance of properties at different structural scales<sup>23</sup>) that have been shown to significantly improve this kind of procedures when using other structural descriptors.

The similarity metric we introduce could find application as the workhorse of a number of simulation protocols, machine-learning algorithms and data mining strategies. For instance, it could be used to detect outliers in automated high-throughput screenings of materials, to cluster similar configurations together, to accelerate the exploration of chemical and conformational space of materials and molecules. Here, we show in particular how it can be combined with a non-linear dimensionality reduction technique such as sketch-map, to give simple and insightful two-dimensional representation of a given molecular or structural data set. As atomistic modelling adventures into larger-scale structures, and unsupervised exploration of materials space, maps such as these can provide a valuable tool to convey intuitive information on complex structural and alchemical landscapes, to ratio-

nalize structure-property relations, and to predict physical observables of novel compounds by training machine-learning models to libraries of known materials.

## 5 Acknowledgments

S.D and M.C would like to acknowledge support from the NCCR MARVEL. A.P.B. was supported by a Leverhulme Early Career Fellowship with joint funding from the Isaac Newton Trust. We would like to thank M. Cuturi and C. Ortner for insightful discussion. We thank C. Pickard for providing silicon structures found with AIRSS, and S. Goedecker and M. Amsler for sharing with us the crystal structures of low-density silicon polymorphs.

### A SOAP kernel for multi-species environments

### B Derivation of the multi-species kernel

Let us show how the alchemical kernel in Eq. (18) can be derived. The overlap kernel

$$\tilde{k}(\mathcal{X}, \mathcal{X}') = \int d\hat{R} \left| \int d\mathbf{r} \sum_{\alpha\alpha'} \kappa_{\alpha\alpha'} \rho_{\mathcal{X}}^{\alpha}(\mathbf{r}) \rho_{\mathcal{X}'}^{\alpha'}(\hat{R}\mathbf{r}) \right|^2 \quad (23)$$

is first rewritten in terms of the expansion of the atomic density functions (17)

$$\tilde{k}(\mathcal{X}, \mathcal{X}') = \int d\hat{R} \left| \sum_{\substack{\alpha\alpha'n \\ lmm'}} \kappa_{\alpha\alpha'} [c_{nlm}^{\alpha}(\mathcal{X})]^{\dagger} \times \right. \\ \left. D_{mm'}^l(\hat{R}) c_{nlm'}^{\alpha'}(\mathcal{X}') \right|^2, \quad (24)$$

where we carried out the spatial integration. Expanding this result, we obtain

$$\tilde{k}(\mathcal{X}, \mathcal{X}') = \int d\hat{R} \sum_{\substack{\alpha\alpha' \\ nlm}} \kappa_{\alpha\alpha'} [c_{nlm}^{\alpha}(\mathcal{X})]^{\dagger} D_{mm'}^l(\hat{R}) c_{nlm'}^{\alpha'}(\mathcal{X}') \\ \sum_{\substack{\beta\beta' \\ n'l'm''}} \kappa_{\beta\beta'} c_{n'l'm''}^{\beta}(\mathcal{X}) [D_{m''m'''}^{\beta'}(\hat{R}) c_{n'l'm'''}^{\beta'}(\mathcal{X}')]^{\dagger}, \quad (25)$$

which allows us to integrate analytically over all possible rotations  $\hat{R}$  and exploit the orthogonality relations of the Wigner ro-

tation matrices.

$$\begin{aligned} \tilde{k}(\mathcal{X}, \mathcal{X}') = & \sum_{\substack{\alpha\alpha'nlmm' \\ \beta\beta'n'l'm''}} \frac{8\pi^2}{2l+1} \delta_{ll'} \delta_{mm''} \delta_{m'm''} \times \\ & \kappa_{\alpha\alpha'} [c_{nlm}^{\alpha}(\mathcal{X})]^{\dagger} c_{nlm'}^{\alpha'}(\mathcal{X}') \times \\ & \kappa_{\beta\beta'} c_{n'l'm''}^{\beta}(\mathcal{X}) [c_{n'l'm''}^{\beta'}(\mathcal{X}')]^{\dagger} \end{aligned} \quad (26)$$

The final formula of the overlap kernel couples the radial, angular, and species channels of the expansion coefficients while being rotationally invariant

$$\begin{aligned} \tilde{k}(\mathcal{X}, \mathcal{X}') = & \sum_{\substack{\alpha\alpha'\beta\beta' \\ m'l'mm'}} \frac{8\pi^2}{2l+1} \kappa_{\alpha\alpha'} \kappa_{\beta\beta'} \times \\ & [c_{nlm}^{\alpha}(\mathcal{X})]^{\dagger} c_{n'l'm}^{\beta}(\mathcal{X}) c_{nlm'}^{\alpha'}(\mathcal{X}') [c_{n'l'm'}^{\beta'}(\mathcal{X}')]^{\dagger}. \end{aligned} \quad (27)$$

In terms of the power spectrum (16) the kernel may be regarded as a dot-product kernel

$$\tilde{k}(\mathcal{X}, \mathcal{X}') = \sum_{\substack{\alpha\alpha'\beta\beta' \\ m'l}} P_{m'l}^{\alpha\beta}(\mathcal{X}) P_{m'l}^{\alpha'\beta'}(\mathcal{X}') \kappa_{\alpha\alpha'} \kappa_{\beta\beta'}. \quad (28)$$

Finally, it is easy to see that one can recover Eq. (15) by setting the alchemical kernel to the Kronecker-delta  $\kappa_{\alpha\beta} = \delta_{\alpha\beta}$ :

$$\tilde{k}(\mathcal{X}, \mathcal{X}') = \sum_{\substack{\alpha\beta \\ m'l}} P_{m'l}^{\alpha\beta}(\mathcal{X}) P_{m'l}^{\alpha\beta}(\mathcal{X}'). \quad (29)$$

## C Sinkhorn distance for structural similarity

Let us discuss briefly how the REMatch procedure can be implemented in practice. Consider for generality the  $N \times M$  environment similarity matrix  $\mathbf{C}(A, B)$  between two structures with  $N$  and  $M$  atoms respectively. The expression (12) given in Section 2 for the optimal-transport-inspired definition of  $\hat{K}$  generalizes straightforwardly to non-square matrices<sup>39</sup>:

$$\begin{aligned} \hat{K}^{\gamma}(A, B) = & \text{Tr} \mathbf{P}^{\gamma T} \mathbf{C}(A, B) \\ \mathbf{P}^{\gamma} = & \underset{\mathbf{P} \in \mathcal{U}(M, N)}{\text{argmin}} \sum_{ij} P_{ij} (1 - C_{ij} + \gamma \ln P_{ij}), \end{aligned} \quad (30)$$

where  $\mathbf{P} \in \mathcal{U}(N, M)$  is a (scaled) doubly-stochastic  $N \times M$  matrix for which  $\sum_i P_{ij} = 1/M$  and  $\sum_j P_{ij} = 1/N$ .

The Sinkhorn algorithm finds the optimal  $\mathbf{P}^{\gamma}$  by the decomposition  $\mathbf{P}^{\gamma} = \text{diag} \mathbf{u} \mathbf{K} \text{diag} \mathbf{v} = \mathbf{K} \circ \mathbf{u} \mathbf{v}^T$ , where  $\circ$  indicates the Hadamard product, and  $\mathbf{K}$  is the entry-wise exponential of  $(\mathbf{C} -$

$1)/\gamma$ , i.e.

$$P_{ij}^{\gamma} = u_i v_j \exp[(C_{ij} - 1)/\gamma]. \quad (31)$$

The balancing vectors  $\mathbf{u}$  and  $\mathbf{v}$  can be obtained by the iteration

$$\begin{aligned} \mathbf{u} & \leftarrow \mathbf{e}_N / \mathbf{K} \mathbf{v} \\ \mathbf{v} & \leftarrow \mathbf{e}_M / \mathbf{K}^T \mathbf{u}. \end{aligned} \quad (32)$$

where  $(\mathbf{e}_N)_i = 1/N$  are scaled stochastic vectors, and the iteration can be initialized by setting  $\mathbf{v} = \mathbf{e}_M$ .

One of the advantages of a regularized match strategy is that the kernel becomes a smooth function of the environment kernels. Computing its derivative  $\partial_{\alpha} K^{\gamma}$  with respect to a parameter  $\alpha$  (a Cartesian coordinate, for instance), is however not completely trivial. Such a derivative is in fact composed of two terms

$$\partial_{\alpha} \hat{K}^{\gamma}(A, B) = \text{Tr} \mathbf{P}^{\gamma} \partial_{\alpha} \mathbf{C} + \text{Tr} \partial_{\alpha} \mathbf{P}^{\gamma} \mathbf{C}. \quad (33)$$

The first term is easy to compute – provided that one can obtain  $\partial_{\alpha} \mathbf{C}$ , the derivative of all environment kernels with respect to  $\alpha$ . The second term can be further broken down based on the Sinkhorn decomposition of  $\mathbf{P}^{\gamma}$ :

$$\partial_{\alpha} \mathbf{P}^{\gamma} = \partial_{\alpha} \mathbf{K} \circ \mathbf{u} \mathbf{v}^T + \mathbf{K} \circ \partial_{\alpha} (\mathbf{u} \mathbf{v}^T) \quad (34)$$

The critical issue here is that direct evaluation of  $\partial_{\alpha} (\mathbf{u} \mathbf{v}^T)$  would involve performing a separate calculation for each derivative  $\alpha$ , which could make the approach prohibitively expensive when, for instance, one would want to compute derivatives with respect to the coordinates of each atom.

By straightforward albeit tedious algebra, one can reformulate the problem in such a way that the derivative can be computed cheaply for any variational parameter, given  $\partial_{\alpha} \mathbf{C}$ :

$$\partial_{\alpha} \hat{K}^{\gamma}(A, B) = \text{Tr} \mathbf{Q}^T \partial_{\alpha} \mathbf{C}, \quad (35)$$

with

$$Q_{ij} = u_i K_{ij} v_j \left[ 1 + \frac{1}{\gamma} (C_{ij} + a_j - N u_i b_i + M v_j c_j) \right]. \quad (36)$$

The  $\mathbf{Q}$  matrix can be fully evaluated based on intermediate terms that do not depend on  $\delta_{\alpha} \mathbf{C}$ :

$$\begin{aligned} \mathbf{a} & = -M \mathbf{v} \circ (\mathbf{K} \circ \mathbf{C})^T \mathbf{u} \\ \mathbf{b} & = (\mathbf{1} - \mathbf{W})^{-T} [(\mathbf{K} \circ \mathbf{C}) \mathbf{v} + \mathbf{K} (\mathbf{v} \circ \mathbf{a})] \\ \mathbf{c} & = N (\mathbf{b} \circ \mathbf{u}^2) \mathbf{K} \\ \mathbf{W} & = \text{diag} (N \mathbf{u}^2) \mathbf{K} \text{diag} (M \mathbf{v}^2) \mathbf{K}^T, \end{aligned} \quad (37)$$

were with  $\mathbf{u}^2 = \mathbf{u} \circ \mathbf{u}$  we indicate the entry-wise square. The only caveat here is that  $(\mathbf{1} - \mathbf{W})$  is singular, and so it cannot be straight-

forwardly inverted. Nevertheless,  $\mathbf{b}$  can be computed by the fixed-point iteration  $\mathbf{b} \leftarrow \mathbf{W}^T \mathbf{b} + \mathbf{y}$  with  $\mathbf{y} = [(\mathbf{K} \circ \mathbf{C}) \mathbf{v} + \mathbf{K}(\mathbf{v} \circ \mathbf{a})]$ . Due to the potential instability of the procedure, it is crucial however to check the convergence on the overall value of  $\partial_\alpha K^\gamma$ , and not to push the convergence to higher relative accuracy levels than those used for the original solution to the Sinkhorn balancing problem.

## References

- 1 L. M. Ghiringhelli, J. Vybiral, S. V. Levchenko, C. Draxl and M. Scheffler, *Phys. Rev. Lett.*, 2015, **114**, 105503.
- 2 T. D. Huan, A. Mannodi-Kanakkithodi and R. Ramprasad, *Phys. Rev. B*, 2015, **92**, 014106.
- 3 V. Botu and R. Ramprasad, *Phys. Rev. B*, 2015, **92**, 094306.
- 4 A. G. Kusne, T. Gao, A. Mehta, L. Ke, M. C. Nguyen, K.-M. Ho, V. Antropov, C.-Z. Wang, M. J. Kramer, C. Long and I. Takeuchi, *Scientific Reports*, 2014, **4**, 6367 EP –.
- 5 R. Ramakrishnan, P. O. Dral, M. Rupp and O. A. von Lilienfeld, *Scientific Data*, 2014, **1**, 140022 EP –.
- 6 L.-F. m. c. Arsenault, A. Lopez-Bezanilla, O. A. von Lilienfeld and A. J. Millis, *Phys. Rev. B*, 2014, **90**, 155136.
- 7 A. Rodriguez and A. Laio, *Science*, 2014, **344**, 1492–6.
- 8 R. Xu and I. Wunsch, D., *Neural Networks, IEEE Transactions on*, 2005, **16**, 645–678.
- 9 G. Yu, J. Chen and L. Zhu, Knowledge Acquisition and Modeling, 2009. KAM '09. Second International Symposium on, 2009, pp. 189–192.
- 10 O. Isayev, D. Fourches, E. N. Muratov, C. Oses, K. Rasch, A. Tropsha and S. Curtarolo, *Chemistry of Materials*, 2015, **27**, 735–743.
- 11 P. V. Balachandran, J. Theiler, J. M. Rondinelli and T. Lookman, *Scientific Reports*, 2015, **5**, 13285 EP –.
- 12 A. L. Ferguson, A. Z. Panagiotopoulos, P. G. Debenedetti and I. G. Kevrekidis, *Proc. Natl. Acad. Sci. USA*, 2010, **107**, 13597–602.
- 13 M. Ceriotti, G. a. Tribello and M. Parrinello, *Proc. Natl. Acad. Sci. USA*, 2011, **108**, 13023–8.
- 14 G. A. Tribello, M. Ceriotti and M. Parrinello, *Proc. Natl. Acad. Sci. USA*, 2012, **109**, 5196–201.
- 15 M. Ceriotti, G. A. Tribello and M. Parrinello, *J. Chem. Theory Comput.*, 2013, **9**, 1521–1532.
- 16 M. A. Rohrdanz, W. Zheng and C. Clementi, *Annu. Rev. Phys. Chem.*, 2013, **64**, 295–316.
- 17 M. Rupp, A. Tkatchenko, K.-R. Müller and O. A. von Lilienfeld, *Phys. Rev. Lett.*, 2012, **108**, 058301.
- 18 W. J. Szlachta, A. P. Bartók and G. Csányi, *Phys. Rev. B*, 2014, **90**, 104108.
- 19 A. Lopez-Bezanilla and O. A. von Lilienfeld, *Phys. Rev. B*, 2014, **89**, 235411.
- 20 G. Paliana, C. Wang, X. Jiang, S. Rajasekaran and R. Ramprasad, *Scientific Reports*, 2013, **3**, 2810 EP –.
- 21 A. P. Bartók, M. J. Gillan, F. R. Manby and G. Csányi, *Phys. Rev. B*, 2013, **88**, 054104.
- 22 M. Rupp, E. Proschak and G. Schneider, *J. Chem. Inf. Model.*, 2007, **47**, 2280–2286.
- 23 M. Hirn, N. Poilvert and S. Mallat, *arXiv Prepr. arXiv1502.02077*, 2015.
- 24 G. Montavon, M. Rupp, V. Gobre, A. Vazquez-Mayagoitia, K. Hansen, A. Tkatchenko, K.-R. MÅijller and O. A. von Lilienfeld, *New Journal of Physics*, 2013, **15**, 095003.
- 25 J. C. Snyder, M. Rupp, K. Hansen, K.-R. Müller and K. Burke, *Phys. Rev. Lett.*, 2012, **108**, 253002.
- 26 S. A. Ghasemi, A. Hofstetter, S. Saha and S. Goedecker, *Phys. Rev. B*, 2015, **92**, 045131.
- 27 O. A. von Lilienfeld, *International Journal of Quantum Chemistry*, 2013, **113**, 1676–1689.
- 28 K. Hansen, F. Biegler, R. Ramakrishnan, W. Pronobis, O. A. von Lilienfeld, K.-R. Müller and A. Tkatchenko, *The Journal of Physical Chemistry Letters*, 2015, **6**, 2326–2331.
- 29 J. B. Endelman, 2011, **4**, year.
- 30 S. An, W. Liu and S. Venkatesh, Computer Vision and Pattern Recognition, 2007. CVPR '07. IEEE Conference on, 2007, pp. 1–7.
- 31 C. E. Rasmussen and C. K. I. Williams, *Gaussian Processes for Machine Learning (Adaptive Computation and Machine Learning)*, The MIT Press, 2005.
- 32 T. Hastie, R. Tibshirani and J. Friedman, *The Elements of Statistical Learning*, Springer New York, New York, NY, 2009.
- 33 A. Sadeghi, S. A. Ghasemi, B. Schaefer, S. Mohr, M. A. Lill and S. Goedecker, *J. Chem. Phys.*, 2013, **139**, 184118.
- 34 F. Pietrucci and W. Andreoni, *Phys. Rev. Lett.*, 2011, **107**, 085504.
- 35 J. Behler, *The Journal of Chemical Physics*, 2011, **134**, year.
- 36 L. Zhu, M. Amsler, T. Fuhrer, B. Schaefer, S. Faraji, S. Rostami, S. A. Ghasemi, A. Sadeghi, M. Grauzinyte, C. Wolverson and S. Goedecker, *The Journal of Chemical Physics*, 2016, **144**,.
- 37 K. Grauman and T. Darrell, *Proc. IEEE Int. Conf. Comput. Vis.*, 2005, **II**, 1458–1465.
- 38 A. P. Bartók, R. Kondor and G. Csányi, *Phys. Rev. B*, 2013, **87**, 184115.
- 39 M. Cuturi, *Adv. Neural Inf. Process. Syst. 26*, Curran Associates, Inc., 2013, pp. 2292–2300.
- 40 B. Schölkopf, A. Smola and K.-R. Müller, *Neural Comput.*, 1998, **10**, 1299–1319.
- 41 C. Berg, J. P. R. Christensen and P. Ressel, *Harmonic Analysis on Semigroups*, Springer New York, New York, NY, 1984, vol. 100.
- 42 S. De, A. Willand, M. Amsler, P. Pochet, L. Genovese and

- S. Goedecker, *Physical Review Letters*, 2011, **106**, 225502.
- 43 S. De, B. Schaefer, A. Sadeghi, M. Sicher, D. G. Kanhere and S. Goedecker, *Physical Review Letters*, 2014, **112**, 083401.
- 44 H. W. Kuhn, *Nav. Res. Logist. Q.*, 1955, **2**, 83–97.
- 45 M. Cuturi, *Int. Jt. Conf. Artif. Intell. IJCAI*, 2007, 732–737.
- 46 M. Jerrum, A. Sinclair and E. Vigoda, *J. ACM*, 2004, **51**, 671–697.
- 47 T. F. Cox and M. A. A. Cox, *Multidimensional scaling*, CRC Press, 2010.
- 48 R. R. Coifman, S. Lafon, A. B. Lee, M. Maggioni, B. Nadler, F. Warner and S. W. Zucker, *Proc. Natl. Acad. Sci. USA*, 2005, **102**, 7426–31.
- 49 M. Ceriotti, G. A. Tribello and S. De, *sketchmap.org*, <http://sketchmap.org>, 2016.
- 50 S. Goedecker, *The Journal of chemical physics*, 2004, **120**, 9911–9917.
- 51 C. J. Pickard and R. J. Needs, *Journal of Physics: Condensed Matter*, 2011, **23**, 053201.
- 52 M. Amsler, S. Botti, M. A. L. Marques, T. J. Lenosky and S. Goedecker, *Phys. Rev. B*, 2015, **92**, 014101.
- 53 L. Rapp, B. Haberl, C. J. Pickard, J. E. Bradby, E. G. Gamaly, J. S. Williams and A. V. Rode, *Nature Communications*, 2015, **6**, year.
- 54 M. Ropo, M. Schneider, C. Baldauf and V. Blum, *Sci. Data*, 2016, **3**, 160009.
- 55 G. Ramachandran, C. Ramakrishnan and V. Sasisekharan, *Journal of Molecular Biology*, 1963, **7**, 95 – 99.
- 56 T. Fink and J.-L. Reymond, *Journal of Chemical Information and Modeling*, 2007, **47**, 342–353.
- 57 R. Ramakrishnan and O. A. von Lilienfeld, *Chimia. Int. J. Chem.*, 2015, **69**, 182–186.