



Surface-active ionic liquids in micellar catalysis: Impact of anion selection on reaction rates in nucleophilic substitutions

Journal:	<i>Physical Chemistry Chemical Physics</i>
Manuscript ID	CP-ART-01-2016-000493.R1
Article Type:	Paper
Date Submitted by the Author:	30-Mar-2016
Complete List of Authors:	Cognigni, Alice; Vienna University of Technology, Institute of Applied Synthetic Chemistry Gärtner, Peter; Vienna University of Technology, Institute of Applied Synthetic Chemistry Zirbs, Ronald; University of Natural Resources and Life Sciences Vienna, Department of nanobiotechnology Peterlik, Herwig; University of Vienna, Faculty of Physics Prochazka, Katharina; University of Vienna, Dynamics of Condensed Systems Schröder, Christian; University Vienna, Computational Biological Chemistry Bica, Katharina; Vienna University of Technology, Institute of Applied and Synthetic Chemistry



Journal Name

ARTICLE

Surface-active ionic liquids in micellar catalysis: Impact of anion selection on reaction rates in nucleophilic substitutions

Alice Cognigni,^a Peter Gaertner,^a Ronald Zirbs,^b Herwig Peterlik,^c Katharina Prochazka,^c Christian Schröder,^d and Katharina Bica^{*a}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

A series of surface-active ionic liquids based on the 1-dodecyl-3-methylimidazolium cation and different anions such as halides and alkylsulfates was synthesized. The aggregation behavior of these ionic liquids in water was characterized by surface tension, conductivity measurements and UV-Vis spectroscopy in order to determine the critical micelle concentration (CMC) and to provide aggregation parameters. The determination of surface activity and aggregation properties of amphiphilic ionic liquids was accompanied by SAXS studies on selected surface-active ionic liquids. The application of these surface-active ionic liquids with different anions was tested in nucleophilic substitution reactions for the degradation of organophosphorus compounds. Kinetic studies via UV-Vis spectrophotometry showed a strong acceleration of the reaction in the micellar system compared to pure water. In addition, an influence of the anion was observed, resulting in a correlation between the anion binding to the micelle and the reaction rate constants, indicating that the careful choice of the surface-active ionic liquid can considerably affect the outcome of reactions.

Introduction

In recent years the scope of ionic liquids as functional fluids has been expanded to include their mixtures with water for multiple applications.¹ The microstructure and -heterogeneity in liquid state, a fundamental aspect of pure ionic liquids continues to exist in aqueous solution, and several reviews have been published focusing structure and behaviour of ionic liquid clusters in water.² The self-assembly of ionic liquids in aqueous solution is particularly well investigated for long alkyl chain ionic liquids that have been found to display amphiphilic character analogous to conventional cationic surfactants.³ The co-existence of an ionic hydrophilic head group and a long hydrophobic tail results in the formation of aggregates in water; however, as general rule the alkyl chain must contain at least eight carbon atoms for aggregation in water.⁴ A number of computational and experimental techniques, including molecular dynamics simulations, surface tension and

conductivity measurements, potentiometry, UV-Vis spectroscopy, fluorescence probes, NMR spectroscopy, mass spectrometry, isothermal titration calorimetry, light scattering and small-angle X-ray and neutron scattering (SAXS and SANS) have been applied to understand self-aggregation of surface-active ionic liquids in water.^{4,5,6,7,8,9,10,11}

While water traces in ionic liquids are often considered as problematic due to their difficult removal and their large impact on physical properties, the deliberate combination of these two media in well-defined mixtures may result in unique properties that can be fully exploited by varying their composition.¹² For example, the limited solubility of many organic compounds in water could be enhanced with ionic liquids, whereas price or viscosity issues of ionic liquids could be balanced when used in aqueous solution. At the same time the specific properties of the highly tunable ionic liquids and the strong hydrogen bond of water can be combined to afford a reaction environment with new properties. Applications of well-defined ionic liquid/water mixtures have already been identified and range from biomolecule stabilization towards their application in catalysis or in novel separation processes. The impact of self-assembly of ionic liquids in aqueous solution has also been exploited in our group, showing that surface-active ionic liquids can be efficiently applied for micellar catalysis of Diels Alder-reactions, for asymmetric transfer hydrogenations in water, but also for natural products extraction.^{13,14}

Studies on the self-assembly and structural organization of ionic liquids in water so far deal mainly with monocationic alkyl imidazolium based ionic liquids, however also other structures have already been investigated. Full Critical Micellar Concentration (CMC) characterization and extensive

^a Institute of Applied Synthetic Chemistry, Vienna University of Technology, Getreidemarkt 9/163, 1060 Vienna, Austria.

^b Group for Biologically Inspired Materials, Institute of Nanobiotechnology (DNBT), University of Natural Resources and Life Sciences, Muthgasse 11, 1190 Vienna, Austria.

^c Dynamics of Condensed Systems, University of Vienna, Boltzmanngasse 5, 1090 Vienna, Austria.

^d Institute of Computational Biological Chemistry, University of Vienna, Währingerstrasse 17, 1090 Vienna, Austria

*Corresponding author Katharina Schröder (née Katharina Bica). Email: katharina.schroeder@tuwien.ac.at; Fax: +43 1 58801 16360; Tel: +43 1 58801 163601.

Electronic Supplementary Information (ESI) available: Formula for calculating aggregation parameters and fitting of kinetic constants, copies of NMR spectra. See DOI: 10.1039/x0xx00000x

thermodynamic studies allow identifying specific features that influence the properties of these ionic liquids. Several studies showed that the CMC is strongly influenced by the structure of the ionic liquid, in particular in terms of the alkyl chain length of the hydrophobic tail unit. $3'4'5'6'$ ^{15,16,17,18,19} As for traditional cationic surfactants, the Klevens equation was found to be valid.²⁰ A linear relationship between the number of carbon atoms on the alkyl chain and the logarithm of the CMC has been verified¹⁹ which can be applied for predicting the CMC for a homologous series of linear-single chain amphiphiles at a fixed temperature. Cations with different structures have also been reported.^{5'6'}²¹ Blesic *et al.*²¹ investigated the effect of different cationic rings by comparing alkylmethylimidazolium with alkylmethylpyridinium, alkylmethylpiperidinium and alkylmethylpyrrolidinium core groups. The results show that pyrrolidinium has the lowest CMC while the others present an almost identical surface tension profile, however due to the very complex interactions a univocal explanation based on geometry and/or electronic distribution could not be provided. This is in accordance with a study from Cornellas *et al.*⁵ where no substantial difference was identified in the behaviour for surface-active ionic liquids with imidazolium and pyridinium head groups. The introduction of an ester or amide functionality in the side chain of these head groups resulted in improved surface activity according to the studies of Comelles *et al.*^{22,23}; moreover biodegradability was improved by the ester functionality, while the amide group provided a higher thermal stability. On the other hand El Seoud *et al.*²⁴ showed that the introduction of a vinyl moiety on the imidazolium cation resulted in higher CMC values compared to the corresponding ethyl analogues due to the less hydrophobic character of the vinyl group. Baltazar *et al.*²⁵ also investigated the aggregation behaviour of dicationic alkylimidazolium ionic liquids, and CMC values were found to depend not only on the chain length of the imidazolium cation, but also on the length of the linkage chain between the two head groups. Similar results were also obtained by Anderson *et al.*²⁶ who investigated tricationic surface active ionic liquids. Beside showing a better surface activity compared to dicationic ones, this could be further improved by increasing the size and aromaticity of the internal linker. In addition, surface-active ionic liquids with different anions have been investigated, but studies concerning the role of the anion are less systematic. Blesic *et al.*⁴ compared the aggregation behaviour of 1-dodecyl-3-methylimidazolium ($[C_{12}mim]^+$) based ionic liquids with Cl^- , $[PF_6]^-$, $[NTf_2]^-$ as counterion. Aggregates in aqueous solution were found only in case of chloride anions, whereas the low solubility of the ionic liquids containing the $[PF_6]^-$ and $[NTf_2]^-$ anions resulted in phase separation prior to aggregates formation. In a different study on different halides counterions Kim *et al.* reported higher surface activity for $[C_{12}mim]I$ than for the same ionic liquids containing Br^- or Cl^- as counter anion; this was explained with the higher polarizability and binding to the micelles of iodide ions.²⁷ In a study conducted by Frizzo *et al.*²⁸ concerning dicationic imidazolium based surface-active ionic liquids dissolved in 4.75% ethanol-water solution, a correlation between the

aggregation and the anion hydrophobicity was identified. Recently Nowicki *et al.*²⁹ reported a series of alkylimidazolium surface-active ionic liquids with HSO_4^- as counterion. These ionic liquids had a key role as co-catalyst in an oxirane ring opening reaction of epoxidized methyl oleate, and their performance was dependent on the length of the alkyl chain attached to the imidazolium cation. Using a bulky amphiphilic anion such as ilbuprofenate allows to detect a critical aggregation concentration even when coupled with $[C_4mim]^+$ or $[C_6mim]^+$ cations; moreover it was suggested that ionic liquids containing an active pharmaceutical ingredient can be incorporated in ionogels and used for drug delivery applications.³⁰

In contrast, fewer studies exist for ionic liquids with surface-active anions in literature. Early examples report surface-activity of octylsulfate combined with $[C_4mim]^+$, which also contributes to decrease the toxicity of the ionic liquid.^{10,31} Comprehensive work was conducted by Blesic *et al.*^{32,33} on alkylimidazolium alkylsulfonate ionic liquids. A matrix of ionic liquids with variable chain length in both cation and anion was synthesized and physico-chemical properties were determined. Again, a strong dependence of the CMC on the chain length of the cation was identified, but surface activity could be considerably improved by simultaneously introducing a long alkyl chain in the anion. A series of protic ionic liquids based on long-alkyl chain carboxylates was investigated with different cations such as imidazolium and pyrrolidinium. These ionic liquids not only showed a higher surface activity than cationic surfactants, but performed also better than the corresponding surfactants with inorganic cations.³⁴ When diisopropylethylammonium was used as cation, the formation of bilayer vesicle-like aggregates was also demonstrated.³⁵


Eventually, fluorinated surface-active ionic liquids composed of perfluorinated anions have only recently been investigated.^{36,37} An extensive study by Pereiro *et al.* showed that perfluorinated alkylsulfonates strongly decrease the CMC in comparison to the corresponding halide-based ionic liquids, and even for $[C_2mim][C_4F_9SO_3]$ and $[C_2mim][C_8F_{17}SO_3]$ a critical aggregation concentration could be detected.

In here, we intend to present further insight into the nature of self-organization of ionic liquids in water and its impact on micellar catalysis of nucleophilic substitution reactions in water. In particular, we focused on the role of the counteranion and aimed for a correlation between anion binding to the micelle and reaction rate constants, suggesting that the careful choice of the surface-active ionic liquid can considerably affect the outcome of reactions.

Results and discussion

Synthesis of surface-active ionic liquids

We focused on 1-dodecyl-3-methylimidazolium-based ionic liquids $[C_{12}mim]X$, as this is a known and already characterized surface-active cation and we functionalized it with different anions. Our synthetic route is based on two different approaches (Scheme 1). In the case of the surface active $[C_{12}mim]Cl$ **1**, $[C_{12}mim]Br$ **2** and $[C_{12}mim]I$ **3** the classical alkylation of *N*-methylimidazole with the corresponding dodecyl halides was performed. In contrast, surface-active ionic liquids $[C_{12}mim]OMs$ **4**, $[C_{12}mim]OTs$ **5** and $[C_{12}mim]OTf$ **6** were prepared in a two steps synthesis route, including the initial synthesis of *N*-dodecylimidazole by alkylation of imidazole. Afterwards the final ionic liquids were obtained via direct alkylation of *N*-dodecylimidazole with the methyl esters of the corresponding anions. This second route offers a halide free pathway to the synthesis of ionic liquids. In any case, good yields between 83 and 98% were obtained. The final products were purified by repeated crystallization from organic solvents to isolate all ionic liquids as colourless crystals in high purity. In contrast, crystallization was not possible for $[C_{12}mim]OTf$ **6**; however, the highly volatile anion precursors could be easily removed under reduced pressure, allowing to isolate the ionic liquid as low-melting colourless solid.



Scheme 1: Synthesis with surface-active ionic liquids based on the 1-dodecyl-3-methylimidazolium cation with variable anions.

Characterization of surface-activity

Prior to their application in micellar catalysis all ionic liquids were characterized by different experimental techniques in order to investigate the formation of aggregates in water. Several techniques can and have been used for the determination of the CMC of alkyl methyl imidazolium-based ionic liquids, and the reported literature values for a specific ionic liquid vary considerably according to the different techniques as can be seen from the ionic liquid $[C_{14}mim]Cl$, where CMC values ranging from 2.98-4.6 mM have been reported. In here, we selected three different techniques, including surface tension, conductivity and UV-Vis spectroscopy. The results are summarized in Table 1 and will be discussed in the following paragraphs in detail.

Conductivity measurements

Conductivity measurements can be utilized for the characterization of ionic surfactants due to the different

mobility of the individual ions and their aggregates. Curves of the specific conductivity (κ) versus the ionic liquid concentration are reported in Fig. 1 (left). Two different linear regimes could be typically identified that allowed calculating the CMC values via their breaking point. In principle, two phenomena are responsible for this curve shape: formed micelles have a lower mobility than single ions. Moreover, parts of the anions are adsorbed on the surface of the aggregates, decreasing the total species contributing to the solution conductivity upon micelles formation. The degree of counterion binding (β) can be estimated from the ratio of the slopes²⁷ and indicates the amount of anions on the surface of the micelles. The observed CMC values for ionic liquids already reported are in agreement with the results obtained from other research groups.^{27,32} A clear trend can be seen in the halide series, and CMC values decrease in the order $Cl > Br > I$. Similarly, a trend is visible for sulfonates where tosylate (OTs^-) and triflate (OTf^-) based ionic liquids have a lower CMC than $[C_{12}mim]OMs$ **4**. A larger and more hydrophobic anion such as OTs^- tends to adsorb on the micellar surface instead of being hydrated, and consequently favours the aggregates formation by shielding the electrostatic repulsion forces of the charged headgroups.³⁸ This behaviour is also supported by the degree of counterion binding (β): a higher value corresponds to a lower CMC due to the higher amount of anion bound to the micellar surface. Additionally, the Gibbs energy of micellization (ΔG_{mic}^0) could be calculated from the conductivity data, proving that micelles formation is a spontaneous process that is accompanied by a negative free energy change (see ESI for details).³⁹



Journal Name

ARTICLE




Figure 1. Surface tension (left), conductivity (center) and UV-Vis profiles (right) of the investigated ionic liquids at 25 °C.

Table 1. Physical-chemical characterization of the surface-active ionic liquids

Ionic Liquid ^a	Surface tension ^b				Conductivity ^c			UV-Vis ^d
	CMC (mM)	Π_{CMC} (mN/m)	A_{min} (nm ²)	P	CMC (mM)	β	ΔG_{mic}^0 (kJ/mol)	
[C ₁₂ mim]Cl 1	13.25 (Lit. 13.17) ⁶	28.6	0.72	0.29	14.53	0.43	-29.1	-
[C ₁₂ mim]Br 2	9.29	29.3	0.71	0.30	10.29	0.70	-36.2	10.24
[C ₁₂ mim]I 3	4.76	37.7	0.62	0.34	5.19	0.84	-42.3	4.12
[C ₁₂ mim]OMs 4	12.50	29.4	1.13	0.18	14.71	0.45	-29.5	15.26
[C ₁₂ mim]OTf 5	2.37	40.2	0.55	0.38	3.31	0.73	-41.7	2.69
[C ₁₂ mim]OTs 6	2.25	38.3	0.65	0.32	3.44	0.89	-45.4	-

^aSolutions were prepared with doubly-distilled Millipore Milli-Q water and left under shaking with 360 min⁻¹ for 24 h to equilibrate. Samples were equilibrated at 25.0 ± 0.1 °C with a HAAKE K15 thermostat before measurements. ^bSurface tension was determined with the Du Noüy ring method on a Krüss tensiometer. ^cConductivity measurements were performed at 25 °C with a Mettler Toledo SevenExcellence InLAB® 741-ISM electrode. ^dUV-Vis measurements were performed at 25 °C, following the absorbance at 312 nm using benzoylacetone. Π_{CMC} is the effectiveness of the surface tension reduction, A_{min} is the area per molecule residing at the surface, P is the packing parameter, β is the degree of counterion binding, ΔG_{mic}^0 is Gibbs energy of micellization (for details see ESI).

Surface-tension measurements

Figure 1 (center) displays the surface tension (γ) curves as a function of ionic liquids concentration. As a consequence of ionic liquid adsorption at the surface, the surface tension decreases with increasing ionic liquid concentration, until a stable value is obtained after surface saturation. The breaking point of the curve can be used to determine the CMC value of the respective surfactant (see ESI for details). This characteristic behaviour allowed calculating CMC values for all ionic liquids. In general, the obtained CMC values were systematically slightly lower than those measured via conductivity, however the same trend can be observed for the anions. According to the effectiveness of the surface tension reduction (Π_{CMC}) all investigated ionic liquids reduce the water surface tension of 30-40 mN/m. The effectiveness can also be

related to the degree of packing of adsorbed surfactant molecules at the interface. The Gibbs adsorption isotherm equation can be used to estimate the average area per molecule residing at the surface (A_{min}), assuming that the structure pattern corresponds to a monolayer.⁴⁰ A lower A_{min} value corresponds to a closer packing of monomers at the interface. This value could be used to estimate the shape of the formed aggregates according to the packing parameter⁴¹ (P) which includes the volume (v) of the hydrocarbon chain imbedded in the hydrocarbon core of the aggregate and the maximum effective length (l_c) that the chain can assume, calculated according to Tanford equation (see ESI for details)⁴². In general, a packing parameter $P < 0.33$ corresponds to spherical micelles, an intermediate value $0.33 < P < 0.5$ corresponds to non-spherical, ellipsoidal micelles, while higher

values correspond to rod-like micelles and various interconnected structures.⁴⁴ Results in table 1 show that both spherical and ellipsoidal aggregates are presumably formed and depend on the anion size. According to this methodology, smaller anions such as bromide or mesylate form spherical micelles, while bigger anions, e.g. iodide or tosylate favour the formation of ellipsoidal aggregates. This is to some extent in agreement with a study on the shape of alkyl imidazolium bromides by Goodchild *et al.*⁹, who performed SANS experiments and found that $[C_{12}mim]Br$ forms spherical particles at low concentration near the CMC, while the shape tends to turn into elongated micelles at higher concentrations. Assuming that the structure pattern corresponds to a monolayer.⁴³ A lower A_{min} value, i.e. the average area per molecule residing at the surface corresponds to a closer packing of monomers at the interface. This value could be used to estimate the shape of the formed aggregates according to the packing parameter⁴⁴ (P) which includes the volume (v) of the hydrocarbon chain imbedded in the hydrocarbon core of the aggregate and the maximum effective length (l_c) that the chain can assume, calculated according to Tanford equation (see ESI for details)⁴⁵. In general, a packing parameter $P < 0.33$ corresponds to spherical micelles, an intermediate value $0.33 < P < 0.5$ corresponds to non-spherical, ellipsoidal micelles, while higher values correspond to rod-like micelles and various interconnected structures.⁴⁴ Results in table 1 show that both spherical and ellipsoidal aggregates are presumably formed and depend on the anion size. According to this methodology, smaller anions such as bromide or mesylate form spherical micelles, while bigger anions, e.g. iodide or tosylate favour the formation of ellipsoidal aggregates. This is to some extent in agreement with a study on the shape of alkyl imidazolium bromides by Goodchild *et al.*⁹, who performed SANS experiments and found that $[C_{12}mim]Br$ forms spherical particles at low concentration near the CMC, while the shape tends to turn into elongated micelles at higher concentrations.

UV-vis measurements

The determination of the CMC of surfactants with UV-Vis is based on the keto-enolic equilibrium of benzoylacetone. The enolic form can be stabilized by intramolecular hydrogen bonding forming a 6-membered ring. This structure is favoured inside the micelles where no competition for hydrogen bonding with the solvent is present. Consequently the formation of micelles with increasing ionic liquid concentration will shift the keto-enolic equilibrium towards the enol form. This results in an increase of the absorbance at 312 nm which can be used to calculate the CMC value from the breaking point of the curves.^{46,47} The values depicted in Fig 1 (right) are in good agreement with those obtained with the previous techniques, with the exception of $[C_{12}mim]Cl$ **1**, where we were not able to find an unambiguous break point. However, we could not perform measurements for $[C_{12}mim]OTs$ due to overlap of the absorption spectrum of the ionic liquids with benzoylacetone.

Small-angle X-ray scattering

The determination of surface activity and aggregation properties of amphiphilic ionic liquids was accompanied by SAXS studies. As the SAXS measurements and the evaluation was very time consuming, we decided to select three representative ionic liquids **1**, **2** and **6**, which cover the observed range of different conductivities (high, medium, and low conductivity in Figure 1, center).




Figure 2. Small-angle X-ray intensities for the different ionic liquids. The experimental intensities are shown as filled circles, the fit from the core-shell model with grey fit lines and the inserts show the scattering intensities towards very low q -values.

The SAXS intensities (Fig. 2) were fitted with a model describing spherical micelles with a radius of the core R_c , a Schulz distribution for the radii and a shell thickness δ . The core radius is about 1 nm and obtains a more uniform and

narrow size distribution with increasing concentration (see the Figs.S1a to S1c in the supporting information). The shell is of a similar size, about 1 nm for [C₁₂mim]Cl **1** and a little bit higher for [C₁₂mim]Br **2** and [C₁₂mim]OTs **6**. The most important information is the parameter γ , which describes the electron differences between core, shell and surrounding medium⁴⁸ and is defined in the supporting information: It decreases with increasing concentration as well as it is significantly lower for [C₁₂mim]Cl (even close to zero for [C₁₂mim]Cl with 100 mM concentration) than for both other ionic liquids. As a value of $\gamma=0$ means that the scatterings only arises from the core⁴⁸ and the density of the shell is the same as the density of the surrounding solution, one may conclude that Cl is mobile and distributed in the solution, whereas OTs is only very weakly affected by different concentrations. The insert in Fig. 2 shows the scattering intensities towards very low q-values. There is also a weak maximum visible, which shifts towards smaller q-values for higher concentrations. This can be interpreted as a weak tendency for agglomeration of micelles with a cluster size of approximately 15 nm for the 25 mM concentration and considerably larger the higher the concentration.


Table 2. Fit parameters for spheres with core shell structure and a Schulz distribution of the radius of the core (parameters core radius R_c , Z). γ is the scaled medium contrast ($\gamma=1$, the scatter origins only from the shell, $\gamma=0$, only from the core).³⁵

Ionic Liquid	Concentration	Core radius R_c / nm	Z	Shell thickness δ / nm	γ
[C ₁₂ mim]OTs	25 mM	0.95	15	1.23	0.16
[C ₁₂ mim]OTs	50 mM	0.97	28	1.26	0.13
[C ₁₂ mim]OTs	100 mM	1.02	84	1.22	0.12
[C ₁₂ mim]Br	25 mM	0.97	8.3	1.27	0.20
[C ₁₂ mim]Br	50 mM	0.97	13	1.39	0.13
[C ₁₂ mim]Br	100 mM	0.98	25	1.59	0.09
[C ₁₂ mim]Cl	25 mM	1.14	52	0.95	0.07
[C ₁₂ mim]Cl	50 mM	1.18	250	0.94	0.06
[C ₁₂ mim]Cl	100 mM	1.14	51	0.94	0.007

Application in nucleophilic substitutions

Micellar catalysis has already been proven to be a very powerful tool to overcome issues of conventional synthesis in both pure organic solvents and water, and recent developments in the fields have been thoroughly reviewed.^{49,50,51} The problems created by the massive use of volatile, toxic and flammable organic solvents in industrial processes have increased the demand for more benign reaction media, and water is a prime candidate here. However, issues related to the poor solubility of organic compounds in water are well known, a problem that can be overcome via the addition of surfactants. It has already been reported that surface active ionic liquids aggregates can increase the solubility of apolar organic compound that exhibit very poor solubility in water.⁵² Furthermore micelles have the capability to drastically alter reaction rates of organic reactions, as the local concentration experienced by the reactants at/in the micellar zone is higher than in a bulk solution. Surfactants

based on ionic liquids can also be finely tuned by adjustments on the head group, alkyl chain or counterion structure in order to offer an optimal reaction environment increasing parameters such as yield, rate and selectivity. Successful results were already obtained in our previous research on Diels-Alder reactions.¹⁴ The highest reaction rate occurred above the CMC, clearly showing the positive effect of ionic liquid-derived micelles.



Scheme 2: Nucleophilic substitution of 4-nitrophenyl diphenyl phosphate (PNPDPP) with acetaldoxime

Nucleophilic substitutions of organophosphorus compounds can be accelerated by cationic surfactants and are strongly influenced by different anions present in the reaction mixture.⁵³ Optimal conditions for the substitution of organophosphorus compounds are obtained with α -nucleophiles, i.e. nucleophiles possessing a heteroatom with an unshared electron pair adjacent to the nucleophilic centre that exhibit a higher nucleophilic reactivity compared with common nucleophiles of similar basicity, a concept that is also known under the term “ α -effect”. More specifically, oximates have been shown to efficiently promote the hydrolysis of phosphate triesters as they prevent the competition with other nucleophiles such as OH⁻ or halides.^{54,55,56} Studies concerning the effect of several oximes⁵⁷ and substrate features^{54,56} have been reported.

In this study, we investigated the reaction between 4-nitrophenyl diphenyl phosphate (PNPDPP) and acetaldoxime (pKa=11.82) in different surface-active ionic liquid/water mixtures. PNPDPP is a highly lipophilic substrate which tends to interact strongly with micellar aggregates in aqueous solution. Moreover, it has an analogue structure to pesticides such as paraoxon or fenitron, both toxic organophosphorous compounds that are widely used as pesticides in agriculture for field crop and fruit protection against a variety of insect and show considerable toxicity towards human and other mammals.⁵⁷ In literature second order kinetic constants of the reaction occurring in the micellar phase and in the bulk water are reported to be very similar, consequently, we were only interested in the observed pseudo first order kinetic constant that shows the overall reaction rate acceleration.^{56,57} The substitution reaction was conducted under pseudo-first order conditions where the oxime was used in 100-fold excess - an oxime concentration high enough to act as a buffer itself. Different possible reaction pathways have been reported for nucleophilic substitutions of organophosphorus compounds. In our case the only possible pathway is the attack of the

nucleophile at the phosphorous centre, which follows a $S_N2(P)$ mechanism,⁵⁷ while the attack on the aromatic carbons is very unlikely.^{56,58,59} In principle, two leaving groups are possible, either the favoured *p*-nitrophenolate, whose formation can be followed at 402.5 nm or phenolate, which cannot be seen due to spectral overlap with other components in the reaction mixture. We observed that the final absorbance reached during the kinetic runs did not correspond to the maximum theoretical one, suggesting the formation of a by-product. This was verified through ^{31}P experiments, where the formation of both *p*-nitrophenyl phenyl phosphate and diphenyl phosphate as by-product was observed, although the formation of the latter one was considerably slower. Eventually, rate constants for both the desired reaction, and the by-product formation were calculated through least-squares exponential fit of the experimental data: results reported here as observed reaction rates k_{obs} refer to the desired nucleophilic reaction only (see ESI for details).

In order to investigate the effect of our ILs anions we initially investigated the reaction in all halide based ionic liquids $[C_{12}mim]X$ with $X = Cl, Br$ and I in a concentration range from 0 to 100 mM. The observed reaction rates are shown in Fig. 3, and three interesting factors can be identified: In all cases a strong increase in the reaction rate was observed with increasing concentration of ionic liquid, resulting in strongly enhanced reaction rates compared to pure water. This rate enhancement in the micellar medium arises from two effects: on one side the lipophilic substrate is accumulating in the micelles, and at the same time the electrostatic attraction of the cationic head groups to the negatively charged nucleophile leads to the increment of its concentration at the micellar interface where the reaction is taking place.




Figure 3. Pseudo first order reaction rate constant of the nucleophilic substitution of nitrophenyl diphenyl phosphate (PNPDPP) with acetaldoxime as a function of $[C_{12}mim]$ with $X = Cl, Br$ and I in water.

Moreover, it can be observed that in all three cases the maximum in acceleration for each ionic liquid corresponds to their CMC. Below the CMC the formation of aggregates induced from the solute presence causes the rate enhancement compared to water, while above most probably both a dilution effect over more micelles and a higher competition with the higher amount of inert anions takes place. This effect is in accordance with the concept of micellar catalysis and has also been observed in our previous work on Diels Alder reaction in surface-active ionic liquid/water mixtures.¹⁴

Most importantly, the highest enhancement of the reaction rate was observed with chloride-based ionic liquids, suggesting that the choice of anion plays a fundamental role. It has been reported that the presence of the inert anions influences the reaction rate due to a competition at the micellar interface with the reactive oximate.^{56, 60} As a consequence different rate enhancement will take place depending on the extent at which the inert anions are bound to the micellar surface.^{53,61} The observed reaction rates are therefore in accordance with physico-chemical data of the respective ionic liquid, as the chloride anion is less bound to the micellar surface and hence more readily exchangeable compared to bromide or iodide as it can also be seen from the β value calculated from the conductivity data.

This trend also exists when kinetic studies were extended to all surface-active ionic liquids involved in this study. Table 3 reports the maximum rate constants for all ionic liquids $[C_{12}mim]X$ 1-6 at a concentration close to their respective CMC, as this should correspond to the highest rate constants.

Table 3. Comparison of pseudo first order reaction rate constant of the nucleophilic substitution of nitrophenyl diphenyl phosphate (PNPDPP) with acetaldoxime at the CMC of different surface-active ionic liquids

Ionic Liquid	Concentration (mM)	Rate constant $10^3 k_{obs}$ (s^{-1})
$[C_{12}mim]Cl$ 1	15	2.74 ± 0.46
$[C_{12}mim]Br$ 2	10	1.30 ± 0.23
$[C_{12}mim]I$ 3	5	0.51 ± 0.08
$[C_{12}mim]OMs$ 4	15	1.68 ± 0.29
$[C_{12}mim]OTs$ 5	3	0.42 ± 0.07
$[C_{12}mim]OTf$ 6	2.5	0.18 ± 0.03
water	-	0.08 ± 0.01

Kinetics experiments were done using UV-visible spectroscopy on a Shimadzu UV1800 spectrometer at 25 °C. All kinetic runs were performed under pseudo first order conditions, the initial concentration of PNPDPP and oxime in the cuvette were 2.3×10^{-5} M and 2.3×10^{-3} M respectively. The appearance of the product *p*-nitrophenolate was monitored at 402.5 nm.

The results reported in Table 2 show again a dependence between the CMC and the rate enhancement, as ionic liquids with lower CMC show poorer performance than those with higher CMC, due to ability of the anion to exchange with the nucleophile. Moreover, the correlation between the degree of anion binding to the micelle and the reaction rate constants exists also in the sulfonate-based series with $[C_{12}mim]X$ 4-6. The highest rate among these surface-active ionic liquids was observed for $[C_{12}mim]OMs$, indicating again that the careful

choice of the surface-active ionic liquid can considerably affect the outcome of reactions. In fact, this suggests that it would be possible to govern the rate of a desired reaction by applying definite micellar system with targeted surface-active ionic liquids, and various projects in this regard are currently investigated in our lab.

Experimental

Materials and methods

Commercially available reagents and solvents were used as received from Sigma Aldrich unless otherwise specified. Doubly-distilled deionized water was obtained from a Millipore Milli-Q water purification system (Millipore, USA). ^1H and ^{13}C NMR spectra were recorded on a Bruker AC 200 at 200 MHz, using the solvent peak as reference. *N*-Dodecylimidazole was synthesized according to a procedure already reported in literature and distilled before use.⁶² All ionic liquids were dried for at least 48 h at room temperature and 0.01 mbar before use and were stored under Argon atmosphere.

Synthetic procedures

Synthesis of $[\text{C}_{12}\text{mim}]\text{X}$, $\text{X}=\text{Cl}^-, \text{Br}^-, \text{I}^-$

The halide-based imidazolium salts were synthesized according to standard methodologies⁶³, which include the alkylation of *N*-methylimidazole with the appropriate alkyl halide to afford the corresponding imidazolium halide. The ILs were repeatedly crystallized in order to obtain colorless solids.

1-Dodecyl-3-methylimidazolium chloride (1)

After following the general procedure the product was recrystallized from THF and obtained in 92% yield. ^1H NMR (200 Hz, CDCl_3): δ (ppm) = 0.81 (3H, t, $J = 6.36$ Hz, $-\text{C}_{11}\text{H}_{22}-\text{CH}_3$), 1.18 (18H, m, $-\text{C}_2\text{H}_4-\text{C}_9\text{H}_{18}-\text{CH}_3$), 1.84 (2H, t, $J = 6.26$ Hz $-\text{CH}_2-\text{CH}_2-\text{C}_{10}\text{H}_{21}$), 4.06 (3H, s, $\text{N}-\text{CH}_3$), 4.25 (2H, t, $J = 7.35$ Hz, $-\text{CH}_2-\text{C}_{11}\text{H}_{23}$), 7.29 (1H, s, $-\text{N}-\text{CH}-\text{CH}-$), 7.45 (1H, s, $-\text{N}-\text{CH}-\text{CH}-$), 10.54 (1H, s, $-\text{N}-\text{CH}-\text{N}-$). The data were in accordance with the literature.⁶⁴

1-Dodecyl-3-methylimidazolium bromide (2)

After following the general procedure the product was recrystallized from THF and obtained in 92% yield. ^1H NMR (200 Hz, CDCl_3): δ (ppm) = 0.81 (3H, t, $J = 7.41$ Hz, $-\text{C}_{11}\text{H}_{22}-\text{CH}_3$), 1.18 (18H, m, $-\text{C}_2\text{H}_4-\text{C}_9\text{H}_{18}-\text{CH}_3$), 1.85 (2H, t, $J = 6.95$ Hz $-\text{CH}_2-\text{CH}_2-\text{C}_{10}\text{H}_{21}$), 4.07 (3H, s, $\text{N}-\text{CH}_3$), 4.25 (2H, t, $J = 7.41$ Hz, $-\text{CH}_2-\text{C}_{11}\text{H}_{23}$), 7.31 (1H, s, $-\text{N}-\text{CH}-\text{CH}-$), 7.46 (1H, s, $-\text{N}-\text{CH}-\text{CH}-$), 10.35 (1H, s, $-\text{N}-\text{CH}-\text{N}-$). The data were in accordance with the literature.⁵

1-Dodecyl-3-methylimidazolium iodide (3)

After following the general procedure the product was crystallized from toluene and further washed with *n*-hexane and obtained in 95% yield. mp: 39°C. $\text{C}_{16}\text{H}_{31}\text{IN}_2$ (378.34): calcd. C 50.79, H 8.26, N 7.40%; calcd. (including $1.2 \times \text{H}_2\text{O}$) C 48.05, H 8.42, N 7.00 % found C 48.08, H 8.06, N 6.90. ν_{max} (cm^{-1}): 3415 (O-H), 2916 (C-H), 1565(C=C), 1163 (C-N). ^1H NMR (200

Hz, CDCl_3): δ (ppm) = 0.82 (3H, m, $-\text{C}_{11}\text{H}_{22}-\text{CH}_3$), 1.19 (18H, m, $-\text{C}_2\text{H}_4-\text{C}_9\text{H}_{18}-\text{CH}_3$), 1.88 (2H, m, $-\text{CH}_2-\text{CH}_2-\text{C}_{10}\text{H}_{21}$), 4.08 (3H, s, $\text{N}-\text{CH}_3$), 4.27 (2H, t, $J = 7.36$ Hz, $-\text{CH}_2-\text{C}_{11}\text{H}_{23}$), 7.45 (1H, s, $-\text{N}-\text{CH}-\text{CH}-$), 7.59 (1H, s, $-\text{N}-\text{CH}-\text{CH}-$), 9.94 (1H, s, $-\text{N}-\text{CH}-\text{N}-$). ^{13}C NMR (200 Hz, CDCl_3): δ (ppm) = 136.59, 123.84, 122.18, 50.20, 37.07, 31.82, 30.23, 29.52 (2C), 29.44, 29.32, 29.25, 28.93, 26.18, 22.60, 14.06.

1-Dodecyl-3-methylimidazolium methanesulfonate (4)

N-Dodecylimidazole (27.06 g, 30 mmol) was dissolved in 8 ml ethyl acetate, and a solution of methyl methanesulfonate (3.47 g, 31.5 mmol) in 8 ml ethyl acetate was added dropwise at RT. The reaction mixture was then stirred for 6 h at 50 °C. The product precipitated upon cooling and was recrystallized two more times to obtain the product in high purity. After removal of solvent traces under reduced pressure for at least 2 d at 0.01 mbar, 1-dodecyl-3-methylimidazolium methanesulfonate was obtained as colorless powder in 83% yield. ^1H NMR (200 Hz, CDCl_3): δ (ppm) = 0.88 (3H, t, $J = 6.17$ Hz, $-\text{C}_{11}\text{H}_{22}-\text{CH}_3$), 1.25 (18H, m, $-\text{C}_2\text{H}_4-\text{C}_9\text{H}_{18}-\text{CH}_3$), 1.88 (2H, m, $-\text{CH}_2-\text{CH}_2-\text{C}_{10}\text{H}_{21}$), 2.83 (3H, s, $-\text{S}-\text{CH}_3$), 4.06 (3H, s, $\text{N}-\text{CH}_3$), 4.26 (2H, t, $J = 7.33$ Hz, $-\text{CH}_2-\text{C}_{11}\text{H}_{23}$), 7.21 (1H, s, $-\text{N}-\text{CH}-\text{CH}-$), 7.27 (1H, s, $-\text{N}-\text{CH}-\text{CH}-$), 10.10 (1H, s, $-\text{N}-\text{CH}-\text{N}-$). The data were in accordance with the literature.³²

1-Dodecyl-3-methylimidazolium *p*-toluenesulfonate (5)

N-Dodecylimidazole (2.85 g, 12 mmol) was dissolved in 8 ml ethyl acetate, and a solution of methyl *p*-toluenesulfonate (2.37 g, 12.6 mmol) in 8 ml ethyl acetate was added dropwise at RT. The reaction mixture was then stirred for 3 h at 50 °C. The product precipitated upon cooling and was recrystallized two more times to obtain the product in high purity. After removal of solvent traces under reduced pressure for at least 2 d at 0.01 mbar, 1-dodecyl-3-methylimidazolium *p*-toluenesulfonate was obtained as colorless crystal in 85% yield. mp: 97°C. $\text{C}_{23}\text{H}_{38}\text{N}_2\text{O}_3\text{S}$ (422.63): calcd. C 65.37, H 9.06, N 6.63%; found C 65.42, H 9.20, N 6.66. ν_{max} (cm^{-1}): 2919 (C-H), 1575 (C=C), 1498 (S=O), 1194(Ar-C-C), 1034 (C-N). ^1H NMR (200 Hz, CDCl_3): δ (ppm) = 0.81 (3H, t, $J = 12.8$ Hz, $-\text{C}_{11}\text{H}_{22}-\text{CH}_3$), 1.16 (18H, m, $-\text{C}_2\text{H}_4-\text{C}_9\text{H}_{18}-\text{CH}_3$), 1.71 (2H, t, $J = 6.55$ Hz, $-\text{CH}_2-\text{CH}_2-\text{C}_{10}\text{H}_{21}$), 2.27 (3H, s, $-\text{C}_6\text{H}_6-\text{CH}_3$), 3.91 (3H, s, $\text{N}-\text{CH}_3$), 4.09 (2H, t, $J = 14.87$ Hz, $-\text{CH}_2-\text{C}_{11}\text{H}_{23}$), 7.07 (2H, d, $J = 7.98$ Hz, aromatic proton), 7.18 (1H, s, $-\text{N}-\text{CH}-\text{CH}-$), 7.32 (1H, s, $-\text{N}-\text{CH}-\text{CH}-$), 7.71 (2H, d, $J = 8.12$ Hz, aromatic proton), 9.78 (1H, s, $-\text{N}-\text{CH}-\text{N}-$). ^{13}C NMR (50 Hz, CDCl_3): δ (ppm) = 143.91, 139.25, 137.6, 137.44, 128.60, 125.73, 123.79, 122.01, 49.74, 36.15, 31.83, 30.87, 30.14, 29.55-29.48 (3C), 29.34-29.27 (2C), 28.93, 26.13, 22.61, 21.21, 14.06.

1-Dodecyl-3-methylimidazolium trifluoromethane sulfonate (6)

N-Dodecylimidazole (5.16 g, 21.8 mmol) was dissolved in 25 ml anhydrous DCM under argon and freshly distilled methyl trifluoromethane sulfonate (4.3 g, 3 ml, 26.2 mmol) was added dropwise while cooling in an ice bath. The reaction mixture was then stirred for 3 h at RT until NMR showed complete conversion. The volatile materials were removed under reduced pressure, and the product was repeatedly washed

with *n*-hexane. 1-Dodecyl-3-methylimidazolium trifluoromethane sulfonate was obtained via filtration as pale solid in 97% yield. ¹NMR (CDCl₃): δ(ppm) = 0.87 (3H, t, J = 6.36 Hz, -C₁₁H₂₂-CH₃), 1.25 (18H, m, -C₂H₄-C₉H₁₈-CH₃), 1.87 (2H, br, -CH₂-CH₂-C₁₀H₂₁), 3.98 (3H, s, -N-CH₃), 4.18 (2H, t, J = 7.24 Hz, -CH₂-C₁₁H₂₃), 7.31 (1H, s, -N-CH-CH-), 7.38 (1H, s, -N-CH-CH-), 9.13 (1H, s, -N-CH-N-). The data were in accordance with the literature.⁶⁵

Characterization of surface-activity

Solutions at various concentration of the ionic liquids were prepared with doubly-distilled Millipore Milli-Q water and left under shaking at 360 min⁻¹ for 24 h at RT to equilibrate.

Conductivity measurements

The samples were previously equilibrated at (25 ± 0.1) °C in a HAAKE K15 thermostat, then conductivity measurement were performed on a Mettler Toledo Seven Excellence system, equipped with a InLAB[®] 741-ISM electrode (cell constant *k* = 0.105). The conductimeter was calibrated with a standard KCl solution and two independent measurements were performed. The CMC was calculated as the intersection point of the two linear regimes in the conductivity/concentration graph. The degree of counter ion binding was calculated with the following equation β = 1 - α, where α is the degree of ionization and corresponds to the ratio of the slopes between the two linear fragments of the conductivity curves.

Surface tension measurements

Surface tension was determined with the Du Noüy ring method on a Krüss manual tensiometer K6 at RT. Each measurement was repeated 5 times.

UV-Vis measurements

For the UV-Vis measurements a stock solution of benzoylacetone (31 mM) in 1,4-dioxane was prepared, this was used to prepare a diluted solution in water (0.5 mM). For UV-Vis measurements 0.2 ml of benzoylacetone solution in water were added to the ionic liquid solution to a final volume of 1.5 ml. These solutions were equilibrated at 25 °C for at least 1 h, and then transferred to the cuvette for measurement. The final concentration of benzoylacetone in the cuvette was 0.07 mM, while the reference cuvette contained all reagents except for benzoylacetone.

Small angle X-ray scattering

Small-angle scattering experiments were performed with a Bruker Nanostar (Bruker AXS), equipped with a microfocus X-ray source (Incoatec IμS High Brilliance) with CuKα radiation with a wavelength of 0.1542 nm and a 2D-position sensitive detector (Vantec 2000). The scattering patterns were taken for 2 hours. The data were azimuthally integrated and corrected for background scattering to result in scattering intensities in dependence on the scattering vector $q = (4\pi/\lambda)\sin\theta$ in the range from $q = 0.1$ to 20 nm⁻¹, with 2θ being the scattering angle. In the figures, only the range up to $q = 5$ nm⁻¹ is shown for clarity.

Micellar catalysis

4-Nitrophenyl diphenyl phosphate (PNPDPP) was synthesized according to an already reported procedure⁶⁶ and crystallized from ethanol. Due to its low solubility in water a stock solution in acetonitrile was prepared (11.5 · 10⁻³ M). In parallel, a stock buffer solution of acetaldehydroxime (552 · 10⁻³ M) in water was prepared by using a NaOH solution as solvent in order to obtain a ratio NaOH:Oxime of 1:2.

Kinetic Measurements

Kinetics experiments were performed by UV-visible spectroscopy using a Shimadzu UV1800 spectrometer equipped with a thermostate at 25 °C. All kinetic runs were performed under pseudo first order conditions in which the concentration of the oxime was at least 100 times greater than the initial concentration [PNPDPP]₀. Reactions were always freshly prepared by first mixing the 2.4 mL ionic liquid solution with 10 μL oxime stock solutions. The kinetic run was then started by adding 2 μL of the PNPDPP stock solution to a 1 mL of ionic liquid-oxime solution in a quartz cuvette. The initial concentration of PNPDPP and oxime in the cuvette were 2.3 · 10⁻⁵ M and 2.3 · 10⁻³ M respectively. The appearance of the product *p*-nitrophenolate NP was monitored at a fixed wavelength 402.5 nm. Generally each reaction was followed until no change in the final absorbance was observed anymore. The rate constants for both the desired reaction, and the by-product formation, were calculated through least-squares exponential fitting, the system was assimilated to the case of parallel reactions. The following formula afforded the rate constants for the formation of product *p*-nitrophenolate NP:

$$[NP](t) = [PNPDPP]_0 \frac{k_{obs}}{k' + k_{obs}} (1 - e^{-(k' + k_{obs})t})$$

The reported values are an average of at least 3 runs. The values of the rate constant of the parallel reaction *k'* were always at least one order of magnitudes smaller than *k*_{obs}.

Theoretical final absorbance

The theoretical final absorbance values for the reaction between PNPDPP and the oxime were measured by injecting 2 μL of an acetonitrile solution of *p*-nitrophenol (11.5 · 10⁻³ M, equivalent to the PNPDPP in the kinetic study) in 1 mL of the corresponding IL-oxime solution. Each value was taken two times.

³¹P NMR measurements

Samples in DMSO-water [10:90 (w/w)] were measured containing either the starting material or the reaction mixture. The peaks were referred to a capillary with H₃PO₄ as external standard. The obtained chemical shifts are in good agreement with those reported in the literature, and no other products were identified.

Conclusions

We have reported a study on the counterion role in surface-active ionic liquids based on 1-dodecyl-3-methylimidazolium. The two investigated series based on halides and sulfonate anions show a correlation between the tendency of the anion to reside closer to the micellar interface and their surface activity. These properties were reflected also when the surface-active ionic liquids were implied in micellar catalysis, in fact, a different rate acceleration was obtained with the different ionic liquids in nucleophilic substitutions in micellar media.

Acknowledgements

Financial support by the Austrian Science Fund (FWF project P25504-N28) is gratefully acknowledged.

Notes and references

- 1 Y. Kohno and H. Ohno, *Chem. Commun.*, 2012, **48**, 7119.
- 2 S. Chen, S. Zhang, X. Liu, J. Wang, J. Wang, K. Dong, J. Sun and B. Xu, *Phys. Chem. Chem. Phys.*, 2014, **16**, 5893.
- 3 J. Łuczak, J. Hupka, J. Thöming and C. Jungnickel, *Colloids Surf. Physicochem. Eng. Asp.*, 2008, **329**, 125.
- 4 M. Blesic, M. H. Marques, N. V. Plechkova, K. R. Seddon, L. P. N. Rebelo and A. Lopes, *Green Chem.*, 2007, **9**, 481.
- 5 A. Cornellias, L. Perez, F. Comelles, I. Ribosa, A. Manresa and M. T. Garcia, *J. Colloid Interface Sci.*, 2011, **355**, 164.
- 6 O. A. El Seoud, P. A. R. Pires, T. Abdel-Moghny and E. L. Bastos, *J. Colloid Interface Sci.*, 2007, **313**, 296.
- 7 J. Sirieix-Plénet, *Talanta*, 2004, **63**, 979–986.
- 8 J. Bowers, C. P. Butts, P. J. Martin, M. C. Vergara-Gutierrez and R. K. Heenan, *Langmuir*, 2004, **20**, 2191.
- 9 I. Goodchild, L. Collier, S. L. Millar, I. Prokeš, J. C. D. Lord, C. P. Butts, J. Bowers, J. R. P. Webster and R. K. Heenan, *J. Colloid Interface Sci.*, 2007, **307**, 455.
- 10 Z. Miskolczy, K. Sebők-Nagy, L. Biczók and S. Göktürk, *Chem. Phys. Lett.*, 2004, **400**, 296.
- 11 Tariq, M.; Freire, M. G.; Saramago, B.; Coutinho, J. A. P.; Lopes, J. N. C.; Rebelo, L. P. N. *Chem. Soc. Rev.* 2012, **41**, 829.
- 12 K. R. Seddon, A. Stark and M.-J. Torres, *Pure Appl. Chem.*, 2000, **72**, 2275.
- 13 A. K. Ressmann, R. Zirbs, M. Pressler, P. Gaertner and K. Bica, *Z. Naturforsch. B*, 2013, **68b**, 1129.
- 14 K. Bica, P. Gärtner, P. J. Gritsch, A. K. Ressmann, C. Schröder and R. Zirbs, *Chem. Commun.*, 2012, **48**, 5013.
- 15 T. Inoue, H. Ebina, B. Dong and L. Zheng, *J. Colloid Interface Sci.*, 2007, **314**, 236.
- 16 J. Łuczak, C. Jungnickel, M. Joskowska, J. Thöming and J. Hupka, *J. Colloid Interface Sci.*, 2009, **336**, 111.
- 17 C. Jungnickel, J. Łuczak, J. Ranke, J. F. Fernández, A. Müller and J. Thöming, *Colloids Surf. Physicochem. Eng. Asp.*, 2008, **316**, 278.
- 18 G. Bai, A. Lopes and M. Bastos, *J. Chem. Thermodyn.*, 2008, **40**, 1509.
- 19 R. Vanyúr, L. Biczók and Z. Miskolczy, *Colloids Surf. Physicochem. Eng. Asp.*, 2007, **299**, 256.
- 20 P. D. T. Huibers, *Langmuir*, 1999, **15**, 7546.
- 21 M. Blesic, A. Lopes, E. Melo, Z. Petrovski, N. V. Plechkova, J. N. Canongia Lopes, K. R. Seddon and L. P. N. Rebelo, *J. Phys. Chem. B*, 2008, **112**, 8645.
- 22 Garcia, M. T.; Ribosa, I.; Perez, L.; Manresa, A.; Comelles, F. *Colloids and Surfaces B: Biointerfaces*, 2014, **123**, 318.
- 23 Garcia, M. T.; Ribosa, I.; Perez, L.; Manresa, A.; Comelles, F. *Langmuir* 2013, **29**, 2536.
- 24 Malek, N. I.; Vaid, Z. S.; More, U. U.; El Seoud, O. A. *Colloid and Polymer Science* 2015, **293**, 3213.
- 25 Q. Q. Baltazar, J. Chandawalla, K. Sawyer and J. L. Anderson, *Colloids Surf. Physicochem. Eng. Asp.*, 2007, **302**, 150.
- 26 Nacham, O.; Martín-Pérez, A.; Steyer, D. J.; Trujillo-Rodríguez, M. J.; Anderson, J. L.; Pino, V.; Afonso, A. M. *Colloids and Surfaces A: Physicochemical and Engineering Aspects* 2015, **469**, 224.
- 27 M. Ao and D. Kim, *J. Chem. Eng. Data*, 2013, **58**, 1529.
- 28 Frizzo, C. P.; Bender, C. R.; Gindri, I. M.; Salbego, P. R. S.; Villetti, M. A.; Martins, M. A. P. *Colloid and Polymer Science* 2015, **293**, 2901.
- 29 Nowicki, J.; Łuczak, J.; Stańczyk, D. *RSC Adv.* 2016, **6**, 11591.
- 30 Tourné-Péteilh, C.; Devoisselle, J.-M.; Vioux, A.; Judeinstein, P.; In, M.; Viau, L. *Phys. Chem. Chem. Phys.* 2011, **13**, 15523.
- 31 A. Heintz, J. K. Lehmann, S. A. Kozlova, E. V. Balantseva, A. B. Bazyleva and D. Ondo, *Fluid Phase Equilibria*, 2010, **294**, 187.
- 32 M. Blesic, M. Swadźba-Kwaśny, J. D. Holbrey, J. N. Canongia Lopes, K. R. Seddon and L. P. N. Rebelo, *Phys. Chem. Chem. Phys.*, 2009, **11**, 4260.
- 33 M. Blesic, M. Swadźba-Kwaśny, T. Belhocine, H. Q. N. Gunaratne, J. N. C. Lopes, M. F. C. Gomes, A. A. H. Pádua, K. R. Seddon and L. P. N. Rebelo, *Phys. Chem. Chem. Phys.*, 2009, **11**, 8939.
- 34 Anouti, M.; Jones, J.; Boisset, A.; Jacquemin, J.; Caillon-Caravancier, M.; Lemordant, D. *J. Colloid Interface Sci.* 2009, **340**, 104.
- 35 Anouti, M.; Sizaret, P.-Y.; Ghimbeu, C.; Galiano, H.; Lemordant, D. *Colloids and Surfaces A: Physicochemical and Engineering Aspects* 2012, **395**, 190.
- 36 A. B. Pereiro, J. M. M. Araújo, F. S. Teixeira, I. M. Marrucho, M. M. Piñeiro and L. P. N. Rebelo, *Langmuir*, 2015, **31**, 1283.
- 37 F. S. Teixeira, N. S. M. Vieira, O. A. Cortes, J. M. M. Araújo, I. M. Marrucho, L. P. N. Rebelo and A. B. Pereiro, *J. Chem. Thermodyn.*, 2015, **82**, 99.
- 38 J. Łuczak, M. Markiewicz, J. Thöming, J. Hupka and C. Jungnickel, *J. Colloid Interface Sci.*, 2011, **362**, 415.
- 39 R. Zana, *Langmuir*, 1996, **12**, 1208.
- 40 Milton J. Rosen and Joy T. Kunjappu, *Surfactants and interfacial phenomena*, John Wiley & Sons, 4th edn., 2012.
- 41 J. N. Israelachvili, in *Intermolecular and Surface Forces*, Elsevier, 2011, pp. 535.
- 42 C. Tanford, *J. Phys. Chem.*, 1972, **76**, 3020.
- 43 Milton J. Rosen and Joy T. Kunjappu, *Surfactants and interfacial phenomena*, John Wiley & Sons, 4th edn., 2012.
- 44 J. N. Israelachvili, in *Intermolecular and Surface Forces*, Elsevier, 2011, pp. 535.
- 45 C. Tanford, *J. Phys. Chem.*, 1972, **76**, 3020.
- 46 E. Iglesias, *J. Phys. Chem.*, 1996, **100**, 12592.
- 47 A. Dominguez, A. Fernandez, N. Gonzalez, E. Iglesias and L. Montenegro, *J. Chem. Educ.*, 1997, **74**, 1227.
- 48 P. Bartlett and R.H. Ottewill, *J. Chem. Phys.*, 1992, **96**, 3306.
- 49 G. La Sorella, G. Strukul and A. Scarso, *Green Chem.*, 2015, **17**, 644.
- 50 B. H. Lipshutz and S. Ghorai, *Green Chem.*, 2014, **16**, 3660.
- 51 T. Dwars, E. Paetzold and G. Oehme, *Angew. Chem. Int. Ed.*, 2005, **44**, 7174.
- 52 J. Łuczak, C. Jungnickel, M. Markiewicz and J. Hupka, *J. Phys. Chem. B*, 2013, **117**, 5653.
- 53 C. A. Bunton and L. Robinson, *J. Org. Chem.*, 1969, **34**, 773.
- 54 S. Couderc and J. Toullec, *Langmuir*, 2001, **17**, 3819.

- ⁵⁵ F. Terrier, P. Rodriguez-Dafonte, E. Le Guével and G. Moutiers, *Org. Biomol. Chem.*, 2006, **4**, 4352.
- ⁵⁶ V. K. Balakrishnan, X. Han, G. W. VanLoon, J. M. Dust, J. Toullec and E. Buncel, *Langmuir*, 2004, **20**, 6586.
- ⁵⁷ X. Han, V. K. Balakrishnan, G. W. vanLoon and E. Buncel, *Langmuir*, 2006, **22**, 9009.
- ⁵⁸ J. E. Omakor, I. Onyido, G. W. vanLoon and E. Buncel, *J. Chem. Soc. Perkin Trans. 2*, 2001, 324.
- ⁵⁹ B. A. Burnside, L. L. Szafraniec, B. L. Knier, H. D. Durst, R. A. Mackay and F. R. Longo, *J. Org. Chem.*, 1988, **53**, 2009.
- ⁶⁰ C. A. Bunton, F. Nome, F. H. Quina and L. S. Romsted, *Acc. Chem. Res.*, 1991, **24**, 357.
- ⁶¹ Nadia Quarti, Antonio Marques, Iva Blagoeva and Marie-Françoise Ruasse, *Langmuir*, 2000, **16**, 2157.
- ⁶² B. S. Souza, E. C. Leopoldino, D. W. Tondo, J. Dupont and F. Nome, *Langmuir*, 2012, **28**, 833.
- ⁶³ B. F. Gilmore, G. P. Andrews, G. Borberly, M. J. Earle, M. A. Gilea, S. P. Gorman, A. F. Lowry, M. McLaughlin and K. R. Seddon, *New J. Chem.*, 2013, **37**, 873.
- ⁶⁴ A. E. Bradley, C. Hardacre, J. D. Holbrey, S. Johnston, S. E. J. McMath and M. Nieuwenhuyzen, *Chemistry of Materials*, 2002, **14**, 629.
- ⁶⁵ A. E. Bradley, C. Hardacre, J. D. Holbrey, S. Johnston, S. E. J. McMath and M. Nieuwenhuyzen, *Chem. Mater.*, 2002, **14**, 629.
- ⁶⁶ S. Jones and D. Selitsianos, *Org. Lett.*, 2002, **4**, 3671.