

PCCP

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

Reaction of CF₂Cl₂ with gas-phase hydrated electrons

Jozef Lengyel,^{a*} Christian van der Linde,^a Michal Fárník^b and Martin K. Beyer^{a*}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

The reaction of dichlorodifluoromethane (CF₂Cl₂) with hydrated electrons (H₂O)_n⁻ (n = 30–86) in the gas phase was studied using Fourier transform ion cyclotron resonance (FT-ICR) mass spectrometry. The hydrated electron reacts with CF₂Cl₂, forming (H₂O)_nCl⁻ with a rate constant of (8.6 ± 2.2) × 10⁻¹⁰ cm³ s⁻¹, corresponding to an efficiency of 57 ± 15%. The reaction enthalpy was determined with nanocalorimetry, revealing a strongly exothermic reaction with ΔH(CF₂Cl₂, 298K) = -208 ± 41 kJ mol⁻¹. The combination of the measured reaction enthalpy with thermochemical data from the condensed phase yields a C–Cl bond dissociation enthalpy (BDE) ΔH_{C-Cl}(CF₂Cl₂, 298K) = 355 ± 41 kJ mol⁻¹ that agrees within error limits with the predicted values from quantum chemical calculations and published BDEs.

Introduction


Since the discovery of the atmospheric ozone hole the chlorofluorocarbons (CFCs) were recognized as one of the important players in ozone depletion.^{1,2} The most common CFC is dichlorodifluoromethane (CF₂Cl₂) that was used in refrigerants due to its high latent heat, non-toxicity and inertness. The other players in these processes are ice particles in polar stratospheric clouds (PSCs).³ Thus the chemistry which involves not only the CFC in the gas phase, but also the environment of the ice particles needs to be investigated.

The processes involved in ozone depletion are mostly driven by sunlight. Besides photochemistry on ice particles,⁴ electron-induced reactions have also received significant interest. The most prominent example is dissociative electron transfer (DET) on ice surfaces^{5–7} where electron transfer to CFCs was greatly enhanced by several orders of magnitude upon the adsorption on ice. The mechanism was explained via the presence of a 'self-trapped', solvated excess electron in a polar medium such as water or ammonia. Based on the large enhancement of Cl⁻ generation from the DET to CFCs on ices, Lu and Sanche^{8,9} proposed the *cosmic-ray-driven electron reaction model for ozone depletion*, short CRE mechanism, as an additional potential source of Cl[•] radicals. This mechanism initiated a controversial debate.^{8,10–15}

The Cl⁻ enhancement was also observed in other experiments^{16,17} and it was interpreted as an attachment of low energy secondary electrons to CF₂Cl₂ solvated in a polar medium. CF₂Cl₂ is then decomposed and Cl⁻ is generated due

to tunnelling of the solvated electron. In contrast to this interpretation, dissociative electron attachment (DEA) cross sections of mixed CF₂Cl₂/NH₃ clusters did not exhibit any enhancement for electron energies in the range of 0–20 eV.¹⁸ In condensed phase electrochemistry, DET of carbon-halogen bonds has been extensively studied.¹⁹ A sticky DET mechanism with the Cl⁻ and CCl₃[•] fragments bound in the solvent cage was established by Pause et al.²⁰ for DET of CCl₄ in *N,N'*-dimethylformamide, but the strength of the interaction was found to decrease with increasing polarity of the solvent.²¹

DEA to gas-phase CF₂Cl₂ is an efficient process due to the high electron affinity of the halogen atoms.²² Illenberger et al.^{23,24} observed that DEA to CF₂Cl₂ at electron energies close to 0 eV occurs via reaction (1), with large cross sections.



However, there are very few free electrons in the atmosphere below 50 km altitude,²⁵ because they are quickly captured by abundant molecules, in particular O₂.²⁶ Therefore, in the CRE mechanism it is assumed that the PSC particles are able to stabilize solvated electrons generated by ionizing radiation within the condensed phase particle. If CF₂Cl₂ is also condensed in the particle, DET with formation of Cl⁻ may take place. For the liquid phase, it has been shown already in 1971 by pulse radiolysis studies in bulk aqueous solution that thermalized hydrated electrons react rapidly with CFCs to produce Cl⁻ ions.²⁷

Previous studies in our group have shown that reductive cleavage of carbon-halogen bonds, i.e. DET, competes with the first step of Birch reduction in reactions of chlorobenzene as well as di- and trifluorobenzenes with gas-phase hydrated electrons (H₂O)_n⁻.^{28,29} In order to test whether fully thermalized solvated electrons are capable of inducing DET in CFCs, we examine the reaction of CF₂Cl₂ and hydrated electrons (H₂O)_n⁻ in the cluster size range n = 30–86. We report a systematic study on the reaction kinetics. Applying nanocalorimetry, we extract the reaction enthalpy from the

^a Institut für Ionenphysik und Angewandte Physik, Leopold-Franzens-Universität Innsbruck, Technikerstraße 25, 6020 Innsbruck, Austria. E-mail: jozef.lengyel@uibk.ac.at; martin.beyer@uibk.ac.at

^b J. Heyrovský Institute of Physical Chemistry v.v.i., Czech Academy of Sciences, Dolejškova 3, 18223 Prague, Czech Republic.

Electronic Supplementary Information (ESI) available: Conversion of ΔE_{inc} to ΔH_{298K}. See DOI: 10.1039/x0xx00000x

experimental data. In combination with literature thermochemistry from the condensed phase, the C–Cl bond dissociation enthalpy (BDE) of gas-phase CF_2Cl_2 is derived and compared with literature values as well as high-level quantum chemical calculations. We discuss the observed results in comparison with the DET studies in bulk ice and photodissociation experiments in water clusters.

Experiment

The experiments are performed using a modified Bruker/Spectrospin CMS47X FT-ICR mass spectrometer, equipped with a 4.7 T superconducting magnet, a Bruker infinity cell, and an APEX III data station.^{30,31} Hydrated electrons $(\text{H}_2\text{O})_n^-$ are generated in a home built external source^{31,32} by laser vaporization of a solid zinc target and supersonic expansion of the hot plasma in a helium/water gas pulse.^{33,34} The skimmed $(\text{H}_2\text{O})_n^-$ cluster beam is transferred via an electrostatic lens system through differential pumping stages into the ultra-high vacuum (UHV) region of the mass spectrometer, with a background pressure below 5×10^{-10} mbar, and stored in the ICR cell. CF_2Cl_2 is introduced into the UHV region of the mass spectrometer as a gas through a leak valve at constant pressures in the range of 0.5– 1.1×10^{-8} mbar. The purity of the reactant is checked via electron ionization and high resolution mass spectrometry directly in the ICR cell.

To determine the rate constant, reactions are monitored by recording mass spectra as a function of time. The intensities of reactant and product clusters in the mass spectra are summed over all cluster sizes. While the experiments are conducted at room temperature, the internal temperature of $(\text{H}_2\text{O})_n^-$ clusters is a result of the interplay between radiative heating by black-body radiation and evaporative cooling.³⁵ In combination with the caloric curves measured by Hock et al.³⁶ this places the internal temperature of the clusters in the range of 90–120 K.

The thermochemistry is investigated via nanocalorimetry.^{30,37} The heat released during the reaction is extracted by quantitative modelling of the average size of reactant and product clusters as a function of time, taking into account blackbody radiation induced dissociation (BIRD).^{35,38–41} The method was introduced by Höckendorf et al.³⁰ on reactions of $(\text{H}_2\text{O})_n^-$ with O_2 and CO_2 .

To extract the reaction enthalpy from the mass spectra, the average cluster size of reactant and product species is calculated. The results are fitted with a genetic algorithm with the following differential equations:

$$dN_R = -k_f(N_R - N_{0,R})dt \quad (2)$$

$$dN_P = -k_f(N_P - N_{0,P})dt + (N_R - \Delta N_{\text{vap}} - N_P) \left(\frac{k_{\text{IR}}}{T_P} \right) dt \quad (3)$$


Eq. (2) and the first term in eq. (3) describe BIRD of water clusters, with k_f describing the linear dependence on cluster size. $N_{0,R}$, $N_{0,P}$ account for the contribution of the ionic core to the IR absorption cross sections. The second term in eq. (3)

describes the evaporation of water molecules due to the reaction enthalpy released in the water cluster. The average number of evaporated water molecules ΔN_{vap} is the key result of the fit.

The experiments are assisted by quantum chemical calculations at the Gaussian-4 (G4) level⁴² using the Gaussian09 program package⁴³ to support the experimentally observed BDE of the CF_2Cl_2 molecule. In general, G4 level calculations exhibit an average absolute deviation from experiment of 3.5 kJ mol^{-1} .

Results

Figure 1 shows mass spectra of the reaction of CF_2Cl_2 with $(\text{H}_2\text{O})_n^-$. The reaction leads to the formation of hydrated chloride ions and CF_2Cl^* radicals, which evaporate from the water cluster, reaction (4).



At initial 0 s, the mass spectrum is dominated by hydrated electrons. However, a small amount of the product ions is present. Some clusters have reacted during the accumulation in the ICR cell, which takes 2 s. The intensity of the product ions increases with the delay time. After 1 s, the product ions represent more than 50% of the hydrated electron intensity. The product ions start to dominate after 1.5 s. At longer times, the shift of the cluster size distribution to smaller values is clearly visible. Both species undergo BIRD and continuously lose water ligands. After 30 s, the clusters have lost almost all water molecules, and $(\text{H}_2\text{O})_m\text{Cl}^-$, $m = 3-6$, is present in the mass spectrum.

To elucidate the reaction rate, the total intensities are plotted as a function of time. The intensities of reactant and product clusters in the mass spectra are summed over all cluster sizes and normalized. The reaction kinetics is quantitatively analysed for the first 4–8 s, depending on the initial cluster size. For $n < 30$, blackbody radiation induced electron detachment occurs, which interferes with the quantitative analysis.^{33,44} Thus all quantitative fits are stopped when the lower end of the cluster size distribution reaches $n = 30$. Figure 2a shows the kinetic fit using a pseudo-first-order rate law. The resulting first order rate constant $k_{\text{rel}} [\text{s}^{-1}]$ is converted to a pressure corrected absolute rate constant. A relative error of $\pm 25\%$ is determined by the uncertainty of the pressure gauge. Since the CF_2Cl_2 pressure is a critical parameter for the absolute rate constant determination, the measurements were performed at different pressures repeatedly on different days to minimize any uncertainty. The results of each measurement are shown in Table 1. The measured experimental rate constants k_{abs} are compared with calculated collision rates to determine the reaction efficiency. We have shown previously that average dipole orientation (ADO) theory,^{45–48} which describes the ion as a point charge, underestimates the collision rate of clusters with more than 10 water molecules ($k_{\text{ADO}} = 6.1 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$ for reaction with CF_2Cl_2). We therefore use models that account for the geometric size of the water cluster, in particular the hard

sphere average dipole orientation, $k_{\text{HSA}} = 1.0 \times 10^{-9} \text{ cm}^3 \text{ s}^{-1}$, and the surface charge capture model, $k_{\text{SSC}} = 2.0 \times 10^{-9} \text{ cm}^3 \text{ s}^{-1}$.⁴⁹ Earlier studies in our group indicate that the actual collision rate of ionic water cluster lies between the two models.^{50–54} The reaction efficiency can thus be determined via equation (5).

$$\Phi = 2 k_{\text{abs}} / (k_{\text{HSA}} + k_{\text{SSC}}) \quad (5)$$

Averaging over all experiments, we arrive at an absolute rate constant of $k_{\text{abs}} = 8.6 \pm 2.2 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$, which corresponds to an efficiency $\Phi = 57 \pm 15\%$. This means that about one out of two collisions is reactive.

The mass spectra reveal that the $(\text{H}_2\text{O})_m\text{Cl}^-$ ions have a slightly lower mean cluster size than the hydrated electrons $(\text{H}_2\text{O})_n^-$. This difference indicates that water molecules are lost due to the exothermic reaction. Nanocalorimetry is employed, in which the average number of evaporated water molecules is determined.³⁰ The mean cluster sizes for reactants and product as well as their difference were plotted as a function of time (Figure 2b,c). A nanocalorimetric fit reveals that the reaction leads to the evaporation of 4.9 ± 0.9 water molecules. The energy required to evaporate a single water molecule from the cluster is $\Delta E_{\text{vap}} = 43.3 \pm 3.1 \text{ kJ mol}^{-1}$.^{36,55} The total energy release is almost identical to the absolute value of the room temperature reaction enthalpy, with minor corrections and a small contribution to the uncertainty.^{30,50} Then ΔE_{nc} can be calculated via equation (6), which is converted to room temperature enthalpy $\Delta H_r(\text{CF}_2\text{Cl}_2, 298\text{K})$ as detailed in the accompanying ESI.

$$\Delta E_{\text{nc}} = -\Delta N_{\text{vap}} \Delta E_{\text{vap}} = -214 \pm 41 \text{ kJ mol}^{-1} \quad (6)$$

$$\Delta H_r(\text{CF}_2\text{Cl}_2, 298\text{K}) = -208 \pm 41 \text{ kJ mol}^{-1} \quad (7)$$

The observed electron transfer reaction of CF_2Cl_2 with $(\text{H}_2\text{O})_n^-$ is strongly exothermic. To the best of our knowledge, no thermochemical data on the reaction of hydrated electrons with CF_2Cl_2 have been reported so far.

To compare the measured results with literature thermochemistry, we use the observed ΔH_r in combination with reaction enthalpies from the condensed phase to calculate the BDE of $\text{Cl}-\text{CF}_2\text{Cl}$ bond cleavage. The same approach was successfully used previously on SF_6 as a benchmark for nanocalorimetry.⁵⁰ BDE is calculated from established data, namely the hydration energy of the electron,⁵⁶ the dissociation enthalpy of HCl ,⁵⁷ the solution enthalpy of gaseous HCl ,⁵⁸ and the ionization energy of the hydrogen atom.⁵⁹ A thermochemical cycle including all reaction steps is summarized in Table 2. BDE as the enthalpy change of the overall reaction is calculated as the sum of the reaction enthalpies of the partial equations. This results in the $\text{C}-\text{Cl}$ BDE of CF_2Cl_2 , $\Delta H_{\text{C-Cl}}(\text{CF}_2\text{Cl}_2, 298\text{K}) = 355 \pm 41 \text{ kJ mol}^{-1}$.

In addition, the thermochemistry of the $\text{C}-\text{Cl}$ bond cleavage of CF_2Cl_2 is derived by G4 calculations, in which BDE is obtained from the total enthalpies at 298.15 K. The calculated BDE amounts to 337 kJ mol^{-1} . This lies within 18 kJ mol^{-1} of the experimental value. Both values agree within error limits with

the published BDE of $346.0 \pm 13.4 \text{ kJ mol}^{-1}$ that was calculated from the standard enthalpies of formation.^{60,61}

Discussion

On first sight, the high exothermicity of reaction (4) may be surprising, given that the $\text{C}-\text{Cl}$ bond dissociation energy of $355 \pm 41 \text{ kJ mol}^{-1}$ is close to the electron affinity of the Cl atom, 348 kJ mol^{-1} . However, the additional energy is supplied by the much stronger interaction of the Cl^- ion with the solvent environment compared to the hydrated electron. Essentially, hydration promotes DEA.

The mechanism of the reaction is straightforward. In the first step, the solvated electron moves to the σ^* orbital of one of the $\text{C}-\text{Cl}$ bonds, reducing the bond order from 1 to 0.5. Water molecules rearrange to solvate the incipient chloride ion, which further weakens the bond until it is broken and the $\text{CF}_2\text{Cl}^\bullet$ radical is released. Whether a local bound minimum between the Cl^- and $\text{CF}_2\text{Cl}^\bullet$ exists, i.e. whether the DET corresponds to a sticky DET in aqueous solution,¹⁹ cannot be determined on the basis of our experiments. We therefore depicted potential curves for both scenarios in Figure 3. This mechanism can be discussed in connection with two other experiments mentioned in the introduction: the photodissociation of CF_2Cl_2 ,^{4,62} and the DET mechanism to CFCs on ices.^{5–9}

First, we discuss the photodissociation of CF_2Cl_2 : it is a similar process to DET in that sense that the electron is promoted by a UV photon to the antibonding σ^* orbital on one of the $\text{C}-\text{Cl}$ bonds.⁶² In our recent study of this process on large water clusters⁴ we have not seen any evidence for free Cl fragments. Accompanying theoretical calculations revealed a halogen bond⁶³ between Cl and O atom of water molecules. Thus the CF_2Cl_2 molecule was bound on the ice nanoparticle with the Cl atoms oriented towards the cluster and the Cl fragment was caged by the cluster after the photodissociation.⁴ This is consistent with the present observation that the Cl^- fragment of DET on water clusters remains with water clusters generating the observed $(\text{H}_2\text{O})_m\text{Cl}^-$ products, thus further supporting our previous proposal of halogen bonding between the CF_2Cl_2 molecule and water clusters. It also matches the observed efficiency of 57%, if one assumes that CF_2Cl_2 has to collide with the water cluster with the chlorine atoms facing the water network.

Concerning the DET mechanism to CFCs (and other molecules) on ices, it was proposed to proceed via so-called presolvated electrons with binding energies of about 1.3 eV below the vacuum level.^{8,64–67} The vertical detachment energy of solvated electrons in water clusters strongly depends on the cluster size. In the size range studied here, VDE of $(\text{H}_2\text{O})_n^-$, $n = 30\text{--}86$, ranges from 1.46 eV to 2.00 eV.⁶⁸ Extrapolation of cluster values to the bulk yields VDEs ranging from 3.3 eV to 4.0 eV in the literature.^{68–71} VDE of bulk water is directly accessible from liquid jet measurements, where values of 3.3 eV have been reported.^{72,73} No adiabatic values are available for neat water clusters, but Donald et al. studied the hydration of free electrons in $\text{La}(\text{H}_2\text{O})_n^{3+}$, $n = 42\text{--}160$.⁷⁴ From this study,

they extrapolated a bulk hydration enthalpy of -1.3 eV for the electron, identical to the value suggested for the presolvated electron. Direct measurements of the adiabatic hydration enthalpy of the electron in bulk water usually refer to the absolute hydration enthalpy of the proton, which is not precisely known. Taking e.g. the value reported by Shiraishi et al.⁵⁶ referenced to $\Delta H_{\text{hyd}}(\text{H}^+) = -1090$ kJ mol⁻¹, which is the textbook standard, results in $\Delta H_{\text{hyd}}(\text{e}^-) = -172$ kJ mol⁻¹ or -1.8 eV. It should be noted that the thermochemical analysis presented here does not rely on the absolute hydration enthalpy of the proton or the electron, but only on their combined value, which should be very reliable. Since hydrated electrons in the excited state relax within 400 fs ($n=25$) to 1 ps (bulk) to the electronic ground state,⁷⁵ we can safely conclude that in our clusters, the electrons are in the electronic ground state. This implies that dissociative electron attachment to CF₂Cl₂ in the condensed phase does not require a presolvated state, in agreement with the earlier results from pulse radiolysis in bulk aqueous solution.²⁷

Conclusions

As previously observed in pulsed radiolysis studies in bulk aqueous solution, our results with gas-phase hydrated electrons confirm that fully thermalized hydrated electrons in their electronic ground state are able to induce bond cleavage in CF₂Cl₂, analogous to the DET mechanism proposed for PSCs. The reaction is efficient and very exothermic. Nanocalorimetry combined with condensed phase literature thermochemistry yields thermochemical data that are consistent with literature values, as well as our own quantum chemical calculations. All these arguments together are consistent with the interpretation that CF₂Cl₂ undergoes DET in condensed aqueous environments, if thermalized hydrated electrons are present. Whether or not this mechanism is actually relevant for stratospheric ozone destruction is a different question, which cannot be answered on the basis of the present laboratory studies.

Acknowledgements

J.L. acknowledges support through the Lise Meitner Programme of the Austrian Science Fund (FWF): project No. M1983-N34. M.F. acknowledges the Czech Science Foundation grant No. 14-08937S.

References

1. F. S. Rowland and M. J. Molina, *Rev. Geophys.*, 1975, **13**(1), 1.
2. M. J. Molina and F. S. Rowland, *Nature*, 1974, **249**(5460), 810.
3. S. Solomon, R. R. Garcia, F. S. Rowland and D. J. Wuebbles, *Nature*, 1986, **321**(6072), 755.
4. V. Poterya, J. Kočíšek, J. Lengyel, P. Svrčková, A. Pysanenko, D. Hollas, P. Slavíček and M. Fárnik, *J. Phys. Chem. A*, 2014, **118**(26), 4740.
5. Q.-B. Lu and L. Sanche, *Phys. Rev. B*, 2001, **63**(15), 153403.
6. Q.-B. Lu and T. E. Madey, *Surf. Sci.*, 2000, **451**(1-3), 238.
7. Q.-B. Lu and T. E. Madey, *J. Chem. Phys.*, 1999, **111**(7), 2861.
8. Q.-B. Lu, *Phys. Rep.*, 2010, **487**(5), 141.
9. Q.-B. Lu and L. Sanche, *Phys. Rev. Lett.*, 2001, **87**(7), 78501.
10. J.-U. Grooß and R. Müller, *Atmos. Environ.*, 2011, **45**(20), 3508.
11. R. Müller, *Phys. Rev. Lett.*, 2003, **91**(5), 58502.
12. Q.-B. Lu and L. Sanche, *Phys. Rev. Lett.*, 2002, **89**(21), 219804.
13. P. K. Patra and M. S. Santhanam, *Phys. Rev. Lett.*, 2002, **89**(21), 219803.
14. Q.-B. Lu and L. Sanche, *Phys. Rev. Lett.*, 2002, **89**(21), 219802.
15. N. R. P. Harris, J. C. Farman and D. W. Fahey, *Phys. Rev. Lett.*, 2002, **89**(21), 219801.
16. N. S. Faradzhev, C. C. Perry, D. O. Kusmierek, D. H. Fairbrother and T. E. Madey, *J. Chem. Phys.*, 2004, **121**(17), 8547.
17. S. Solovev, D. O. Kusmierek and T. E. Madey, *J. Chem. Phys.*, 2004, **120**(2), 968.
18. J. Langer, S. Matt, M. Meinke, P. Tegeder, A. Stamatovic and E. Illenberger, *J. Chem. Phys.*, 2000, **113**(24), 11063.
19. A. Houmam, *Chem. Rev.*, 2008, **108**(7), 2180.
20. L. Pause, M. Robert and J.-M. Savéant, *J. Am. Chem. Soc.*, 2000, **122**(40), 9829.
21. L. Pause, M. Robert and J.-M. Savéant, *J. Am. Chem. Soc.*, 2001, **123**(48), 11908.
22. K. Graupner, S. A. Haughey, T. A. Field, C. A. Mayhew, T. H. Hoffmann, O. May, J. Fedor, M. Allan, I. I. Fabrikant, E. Illenberger, M. Braun, M.-W. Ruf and H. Hotop, *J. Phys. Chem. A*, 2010, **114**(3), 1474.
23. E. Illenberger, *Ber. Bunsenges. Phys. Chem.*, 1982, **86**(3), 252.
24. E. Illenberger, H.-U. Scheunemann and H. Baumgärtel, *Chem. Phys.*, 1979, **37**(1), 21.
25. J. Kaplan and C. A. Barth, *Proc. Natl. Acad. Sci.*, 1958, **44**(2), 105.
26. D. Smith and N. G. Adams, *Top. Curr. Chem.*, 1980, **89**, 1.
27. R. H. Schuler, T. I. Balkas and J. H. Fendler, *J. Phys. Chem.*, 1971, **75**(4), 455.
28. Y. Cao, R. F. Höckendorf and M. K. Beyer, *Int. J. Mass Spectrom.*, 2008, **277**(1-3), 206.
29. R. F. Höckendorf, O. P. Balaj and M. K. Beyer, *Phys. Chem. Chem. Phys.*, 2011, **13**(19), 8924.
30. R. F. Höckendorf, O. P. Balaj, C. van der Linde and M. K. Beyer, *Phys. Chem. Chem. Phys.*, 2010, **12**(15), 3772.

31. C. Berg, T. Schindler, G. Niedner-Schatteburg and V. E. Bondybey, *J. Chem. Phys.*, 1995, **102**(12), 4870.
32. S. Maruyama, L. R. Anderson and R. E. Smalley, *Rev. Sci. Instrum.*, 1990, **61**(12), 3686.
33. M. K. Beyer, B. S. Fox, B. M. Reinhard and V. E. Bondybey, *J. Chem. Phys.*, 2001, **115**(20), 9288.
34. M. Beyer, C. Berg, H. W. Görlitzer, T. Schindler, U. Achatz, G. Albert, G. Niedner-Schatteburg and V. E. Bondybey, *J. Am. Chem. Soc.*, 1996, **118**(31), 7386.
35. R. C. Dunbar, *Mass Spectrom. Rev.*, 2004, **23**(2), 127.
36. C. Hock, M. Schmidt, R. Kuhn, C. Bartels, L. Ma, H. Haberland and B. von Issendorff, *Phys. Rev. Lett.*, 2009, **103**(7), 73401.
37. W. A. Donald, R. D. Leib, J. T. O'Brien, A. I. S. Holm and E. R. Williams, *Proc. Natl. Acad. Sci.*, 2008, **105**(47), 18102.
38. M. Sena, J. M. Riveros and N. M. M. Nibbering, *Rapid Commun. Mass Spectrom.*, 1994, **8**(12), 1031.
39. T. Schindler, C. Berg, G. Niedner-Schatteburg and V. E. Bondybey, *Chem. Phys. Lett.*, 1996, **250**(3-4), 301.
40. P. D. Schnier, W. D. Price, R. A. Jockusch and E. R. Williams, *J. Am. Chem. Soc.*, 1996, **118**(30), 7178.
41. O. Hampe, T. Karpuschkin, M. Vonderach, P. Weis, Y. Yu, L. Gan, W. Klopper and M. M. Kappes, *Phys. Chem. Chem. Phys.*, 2011, **13**(20), 9818.
42. L. A. Curtiss, P. C. Redfern and K. Raghavachari, *J. Chem. Phys.*, 2007, **126**(8), 84108.
43. M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, Montgomery, J. A., Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski and D. J. Fox, *Gaussian 09: Revision D.01*, Gaussian, Inc, Wallingford CT, 2013.
44. S. T. Arnold, R. A. Morris and A. A. Viggiano, *J. Chem. Phys.*, 1995, **103**(21), 9242.
45. T. Su and M. T. Bowers, *Int. J. Mass Spectrom. Ion Processes*, 1975, **17**(2), 211.
46. T. Su and M. T. Bowers, *J. Am. Chem. Soc.*, 1973, **95**(5), 1370.
47. T. Su and M. T. Bowers, *J. Chem. Phys.*, 1973, **58**(7), 3027.
48. P. Langevin, *Ann. Chim. Phys.*, 1905, **5**, 245.
49. G. Kummerlöwe and M. K. Beyer, *Int. J. Mass Spectrom.*, 2005, **244**(1), 84.
50. A. Akhgarnusch, R. F. Höckendorf and M. K. Beyer, *J. Phys. Chem. A*, 2015, **119**(39), 9978.
51. A. Akhgarnusch and M. K. Beyer, *Int. J. Mass Spectrom.*, 2014, **365**, 295.
52. R. F. Höckendorf, K. Fischmann, Q. Hao, C. van der Linde, O. P. Balaj, C.-K. Siu and M. K. Beyer, *Int. J. Mass Spectrom.*, 2013, **354-355**, 175.
53. R. F. Höckendorf, Q. Hao, Z. Sun, B. S. Fox-Beyer, Y. Cao, O. P. Balaj, V. E. Bondybey, C.-K. Siu and M. K. Beyer, *J. Phys. Chem. A*, 2012, **116**(15), 3824.
54. A. Akhgarnusch, W. K. Tang, H. Zhang, C.-K. Siu and M. K. Beyer, *Phys. Chem. Chem. Phys.*, 2016. DOI: 10.1039/C6CP03324E
55. W. A. Donald, R. D. Leib, M. Demireva, B. Negru, D. M. Neumark and E. R. Williams, *J. Phys. Chem. A*, 2011, **115**(1), 2.
56. H. Shiraishi, G. R. Sunaryo and K. Ishigure, *J. Phys. Chem.*, 1994, **98**(19), 5164.
57. S. J. Blanksby and G. B. Ellison, *Acc. Chem. Res.*, 2003, **36**(4), 255.
58. V. B. Parker, *Thermal Properties of Aqueous Uni-Univalent Electrolytes: National Standard Reference Data Series*, National Bureau of Standards 2 (Category 5 - Thermodynamic and Transport Properties), Washington, DC, 1965.
59. P. J. Linstrom and W. G. Mallard, *NIST Chemistry WebBook: NIST Standard Reference Database Number 69*, National Institute of Standards and Technology, Gaithersburg, MD, 2005.
60. D. R. Lide, ed., *Handbook of Chemistry and Physics*, CRC Press, Boca Raton, 1997.
61. E. Tschuikow-Roux and S. Paddison, *Int. J. Chem. Kinet.*, 1987, **19**(1), 15.
62. V. Poterya, J. Kočišek, A. Pysanenko and M. Fárník, *Phys. Chem. Chem. Phys.*, 2014, **16**(2), 421.
63. G. Cavallo, P. Metrangolo, R. Milani, T. Pilati, A. Priimagi, G. Resnati and G. Terraneo, *Chem. Rev.*, 2016, **116**(4), 2478.
64. C.-R. Wang, J. Nguyen and Q.-B. Lu, *J. Am. Chem. Soc.*, 2009, **131**(32), 11320.
65. C.-R. Wang, K. Drew, T. Luo, M.-J. Lu and Q.-B. Lu, *J. Chem. Phys.*, 2008, **128**(4), 41102.
66. S. Ryu, J. Chang, H. Kwon and S. K. Kim, *J. Am. Chem. Soc.*, 2006, **128**(11), 3500.
67. Q.-B. Lu and L. Sanche, *J. Chem. Phys.*, 2004, **120**(5), 2434.
68. J. V. Coe, G. H. Lee, J. G. Eaton, S. T. Arnold, H. W. Sarkas, K. H. Bowen, C. Ludewigt, H. Haberland and D. R. Worsnop, *J. Chem. Phys.*, 1990, **92**(6), 3980.
69. L. Ma, K. Majer, F. Chirof and B. von Issendorff, *J. Chem. Phys.*, 2009, **131**(14), 144303.
70. J. R. R. Verlet, A. E. Bragg, A. Kamrath, O. Cheshnovsky and D. M. Neumark, *Science*, 2005, **307**(5706), 93.
71. T. Zeuch and U. Buck, *Chem. Phys. Lett.*, 2013, **579**, 1.

ARTICLE

Journal Name

72. Y. Tang, H. Shen, K. Sekiguchi, N. Kurahashi, T. Mizuno, Y.-I. Suzuki and T. Suzuki, *Phys. Chem. Chem. Phys.*, 2010, **12**(15), 3653.
73. K. R. Sieferrmann, Y. Liu, E. Lugovoy, O. Link, M. Faubel, U. Buck, B. Winter and B. Abel, *Nature Chem*, 2010, **2**(4), 274.
74. W. A. Donald, M. Demireva, R. D. Leib, M. J. Aiken and E. R. Williams, *J. Am. Chem. Soc.*, 2010, **132**(13), 4633.
75. R. M. Young and D. M. Neumark, *Chem. Rev.*, 2012, **112**(11), 5553.

Table 1 Kinetic and nanocalorimetric analysis of each data set

$(\text{H}_2\text{O})_n^-$ $n=$	$p(\text{CF}_2\text{Cl}_2)$ $\times 10^{-9}$ mbar	k_{abs} $\times 10^{-10}$ $\text{cm}^3 \text{s}^{-1}$	k_{HSA} $\times 10^{-9}$ $\text{cm}^3 \text{s}^{-1}$	k_{SCC} $\times 10^{-9}$ $\text{cm}^3 \text{s}^{-1}$	k_{ADO} $\times 10^{-10}$ $\text{cm}^3 \text{s}^{-1}$	Φ %	ΔN_{vap}	ΔH_{r} kJ mol^{-1}
31-47	7.2	7.1	0.9	1.8	6.2	53	4.99	-216
30-47	11	6.5	0.9	1.8	6.2	48	5.39	-233
36-50	4.7	6.6	0.9	1.9	6.2	47	6.02	-261
34-54	9.0	7.5	0.9	1.9	6.2	54	4.18	-181
35-58	8.1	7.4	0.9	1.9	6.2	53	3.41	-148
50-82	9.0	9.4	1.0	2.1	6.0	61	5.21	-226
51-82	11	10	1.0	2.1	6.0	65	5.13	-222
51-84	11	12	1.1	2.1	6.0	75	6.92	-300
51-84	8.3	9.5	1.1	2.1	6.0	59	2.57	-111
46-86	9.3	10	1.0	2.1	6.0	65	5.57	-241
Average	-	8.6	1.0	2.0	6.1	57	4.94	-214

Table 2 Thermochemical cycle for the Cl-CF₂Cl BDE.

Reaction	ΔH_r (298K) kJ mol ⁻¹	Ref.
$H^+(g) + e^-(g) \rightarrow H^+(aq) + e^-(aq)$	-1261.9	56
$H^+(aq) + Cl^-(aq) \rightarrow HCl(g)$	74.48	58
$HCl(g) \rightarrow H(g) + Cl(g)$	431.58	57
$H(g) \rightarrow H^+(g) + e^-(g)$	1318.4	59
$CF_2Cl_2(g) + e^-(aq) \rightarrow CF_2Cl(g) + Cl^-(aq)$	-208	Our work
$CF_2Cl_2(g) \rightarrow CF_2Cl(g) + Cl(g)$	355	Sum




Fig. 1 Mass spectra of the reaction of CF_2Cl_2 with hydrated electrons (blue line) after (a) 0.0, (b) 1.0, and (c) 3.6 s. $(\text{H}_2\text{O})_m\text{Cl}^-$ (red line) as product is present already at nominal 0.0 s due to the 2 s filling cycle.




Fig. 2 (a) Kinetic and (b,c) nanocalorimetric analysis of the reaction of CF_2Cl_2 with hydrated electrons $(\text{H}_2\text{O})_n^-$ at room temperature. Panel (a) represents the pseudo-first-order kinetic fit of $(\text{H}_2\text{O})_n^-$ (blue squares) as reactant and $(\text{H}_2\text{O})_n\text{Cl}^-$ (red circles) as product species. Panel (b) shows the fit of the cluster mean sizes for both species, and panel (c) illustrates the fit of their size difference (black diamonds).




Fig. 3 Schematic potential energy curves for sticky (dashed) or non-sticky (dotted) DET of a hydrated electron to CF_2Cl_2 .