

NJC

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Hexahalometallate salts of trivalent scandium, yttrium and lanthanum: cation–anion association in the solid state and in solution

Martin J. D. Champion, William Levason, David Pugh and Gillian Reid*

School of Chemistry, University of Southampton, Southampton SO17 1BJ, UK. Email: G.Reid@soton.ac.uk

Abstract

The hexahalide salts, $[N^rBu_4]_3[LaCl_6]$, $[BMPYRR]_3[LaCl_6]$ (BMPYRR = 1-butyl-1-methylpyrrolidinium, $[EMIM]_3[MX_6]$ (EMIM = 1-ethyl-3-methylimidazolium; M = La, X = Cl, Br, I; M = Sc, Y, Ce, X = Cl) and $[EDMIM]_3[MX_6]$ (EDMIM = 1-ethyl-2,3-dimethylimidazolium; M = Y, X = Cl; M = La, X = Cl, I) have been prepared and X-ray crystal structures determined for several of them, with a view to probing the effect of varying the trivalent metal ion, the halide and the counter-cation on the structures adopted in the solid state. The crystal structures of the EMIM and EDMIM salts show extensive H-bonding between the halide ligands and organic cations; based upon the H-bonding distances, this appears to be strongest for the $[EMIM]_3[MCl_6]$ salts, becoming progressively weaker for heavier metal ion or halide. In terms of the cations, changing from EMIM to EDMIM also reduces the strength of the H-bonding.

The strength of the cation-anion pairing in solution has also been probed in solution *via* NMR spectroscopy where possible (^{45}Sc , ^{89}Y and ^{189}La) and, for the EMIM salts, via the shift of $\delta(H_2)$ relative to $[EMIM]Cl$ at a standard concentration. The trends observed in solution mirror those determined in the solid state.

Introduction

In the course of our work on electrodeposition of metals and semiconductors from unusual low polarity solvents, such as CH_2Cl_2 ¹ and supercritical CH_2F_2 ,² we have established that halometallates can often offer both the solubility and extent of dissociation into discrete cations and anions to be highly effective as reagents for delivering the elements of choice. Taking germanium as an example, we have shown that the cation, halide and anion charge all play an important role in determining the extent of cation-halogermanate anion association both in the solids and in solution.³ While we have often used quaternary ammonium salts for this purpose, recent work on the electrodeposition of elemental germanium from supercritical CH_2F_2 has established that using the 1-ethyl-3-methylimidazolium (EMIM) cation leads to improved quality of the electrodeposited germanium,⁴ and hence this and other substituted imidazolium cation salts, which are also used extensively for electrodeposition from ionic liquids,⁵ have become of greater importance in this area of work.

In order to gain a better understanding of the cation-anion association in halometallate salts, we describe here a study of the $[\text{MX}_6]^{3-}$ trianions incorporating a range of trivalent metal ions from Group 3 and the lanthanides, in the anticipation that the trianionic charge would give rise to significant cation-anion association both in the solid state and in low polarity solvents. We report the synthesis of a series of halometallate anion salts of scandium, yttrium and several lanthanides. All complexes have been characterised by NMR spectroscopy (^1H , ^{45}Sc , ^{89}Y and ^{139}La as appropriate), IR spectroscopy, elemental analysis and, in several cases, single-crystal X-ray diffraction. To gain further insight into the cation-anion pairing in these species, we have explored the effects that systematic changes in the counter-cation, metal ion and halide co-ligands have on the solid-state structures adopted and the spectroscopic behaviour of these halometallate salts in MeCN (and where possible, CH_2Cl_2) solution.

A small number of earlier studies report structural data on halometallate salts of the Ln(III) ions. The examples most pertinent to the present study are $[\text{EMIM}]_3[\text{LnCl}_6]$ (Ln = La, Pr, Nd, Sm, Eu) grown from hydrated LnCl_3 in $[\text{EMIM}]\text{Cl}$ at $110\text{ }^\circ\text{C}$,^{6,7} $[\text{MPPYRR}]_3[\text{NdI}_6]$ (MPPYRR = 1-methyl-1-propylpyrrolidinium),⁸ and $[\text{SEt}_3]_3[\text{LnI}_6]$ (Ln = Nd, Sm), obtained from reaction of LnI_2 with the ionic liquid $[\text{SEt}_3][\text{NTf}_2]$ (NTf_2 = bis-(trifluoromethanesulfonyl)imide).⁹ A very recent study has described the structures of a series of $[\text{BMIM}]_3[\text{CeX}_6]$ salts (BMIM = 1-butyl-3-methylimidazolium), also obtained directly from CeX_3 in the $[\text{BMIM}]\text{X}$ ionic liquids, which are reported to exhibit intense Ce^{3+} based 5d-4f-centred emission.¹⁰

Results and Discussion

Preparations:

The hexahalide salts, $[\text{N}^w\text{Bu}_4]_3[\text{LaCl}_6]$, $[\text{BMPYRR}]_3[\text{LaCl}_6]$ (BMPYRR = 1-butyl-1-methylpyrrolidinium, $[\text{EMIM}]_3[\text{MX}_6]$ (M = La, X = Cl, Br, I; M = Sc, Y, Ce, X = Cl) and $[\text{EDMIM}]_3[\text{MX}_6]$ (EDMIM = 1-ethyl-2,3-dimethylimidazolium; M = Y, X = Cl; M = La, X = Cl, I), were selected for study on the basis that they would allow us to probe the effects that the cation (extent of H-bonding or not), the metal ion size and the halide type have upon the solid state and solution properties. The salts were readily prepared as powdered solids in good yield by treatment of anhydrous MX_3 with three (or slightly greater) molar equivalents of the appropriate halide salt of the organic cation, $[\text{N}^w\text{Bu}_4]^+$, $[\text{BMPYRR}]^+$, $[\text{EMIM}]^+$ or $[\text{EDMIM}]^+$ in MeCN, acetone or CH_2Cl_2 solution, and their formulations were confirmed by microanalyses. The isolated salts are moderately soluble in MeCN and, in some cases, in CH_2Cl_2 solution. Crystals of several examples with EMIM and EDMIM cations were obtained by vapour diffusion of Et_2O into a concentrated MeCN solution of the relevant salt.

Varying the halide:

In order to assess the effect of varying the halide on the nature of the hydrogen bonding interactions, the [EMIM]₃[LaX₆] system (X = Cl, Br, I) was chosen. [EMIM]₃[LaCl₆] had previously been synthesised and crystallographically characterised.^{6,7} Crystals of the bromide (Figure 1) and iodide (Figure 2) analogues were obtained by slow diffusion of Et₂O into an MeCN solution. Surprisingly these are rare examples of discrete [LaX₆]³⁻ anions with organic cations: [2,4,6-trimethylpyridinium]₄[LaCl₆]Cl¹¹ and [BMPYRR]₄[LaI₆][NTf₂]¹² are the only other structurally authenticated complexes.





Figure 1. ORTEP representation of [EMIM]₃[LaBr₆] showing the hydrogen bonding (dotted lines) surrounding the La1-centred cation (left) and the La2-centred cation (right). Displacement ellipsoids are drawn at the 50% probability level. Selected bond lengths (Å): La1–Br1 2.9371(13), La1–Br2 2.9296(16), La1–Br3 2.9534(13), La2–Br4 2.9477(15), La2–Br5 2.9484(14), La2–Br6 2.9524(13). Symmetry codes: (i) 2 – x, 1 – y, 1 – z; (ii) 1 – x, 1 – y, –z.



(a)




Figure 2. (a) ORTEP representation of [EMIM]₃[LaI₆] showing the hydrogen bonding (dotted lines) surrounding the La1-centred cation (left) and the La2-centred cation (right). Displacement ellipsoids are drawn at the 50% probability level and H atoms not involved in hydrogen bonding are omitted for clarity. Selected bond lengths (Å): La1–I1 3.1557(5), La1–I2 3.1448(4), La1–I3 3.1930(8), La2–I4 3.1988(5), La2–I5 3.1663(5), La2–I6 3.1733(10). Symmetry codes: (i) 1 – x, 3 – y, 1 – z; (ii) –x, 2 – y, 1 – z; (iii) x, 2.5 – y, 1.5 – z; (iv) 1 – x, 0.5 – y, 1 – z; (v) 1 – x, 0.5 – y, 1.5 – z. (b) Representation of [EMIM]₃[LaI₆] showing the extended structure generated by the hydrogen bonding (dotted lines).

[EMIM]₃[LaCl₆] and [EMIM]₃[LaBr₆] are isomorphous and isostructural, whereas [EMIM]₃[LaI₆] crystallises with a different unit cell, presumably reflecting the increased steric requirements of I[–] over Br[–] and Cl[–]. Nonetheless, in each case the [LaX₆]^{3–} anion is octahedrally coordinated with six halides comprising the primary coordination sphere. Two crystallographically independent [LaX₆]^{3–} anions were observed in the solid state with different arrangements of hydrogen bonding cations, including bifurcated hydrogen bonding, accounting for the difference. Most of the hydrogen bonding occurs through the acidic NCN proton of the imidazolium ring but interactions through the NCCN backbone protons (which are less acidic) were also observed, resulting in an extended structure being observed in the solid state (Figure 2(b)).

Here it should be noted that in our experience, the criteria used for assigning the strength of hydrogen bonds¹³ are not always applicable where metal complexes with heavy halogens are acting as acceptors. Other important factors such as crystal packing effects and π -bonding interactions can have a significant effect on the donor-acceptor distances and donor-hydrogen-acceptor (DHA) angles which are used to classify ‘strong’, ‘moderate’ or ‘weak’ hydrogen bonds. Typically, these DHA interactions are quantified via the crystallographically determined D...A (C...X) distances. The radii of Br[–] and especially I[–] also have an effect on the C...X bond lengths.

Nonetheless, for a given halide, the shortest C...X distances for the [EMIM]₃[LaX₆] series of complexes (Table 1) occur where the NCN proton (H2) acts as the donor, consistent with it being the most acidic proton on the imidazolium ring. The C...X distances where the backbone NCCN protons (H4/H5) act as donors are typically 0.1–0.2 Å longer, implying weaker hydrogen bonding. In all cases the DHA angles are also generally greater than 130°, hence we assign these as ‘moderate’ hydrogen bonds.

Table 1: Hydrogen bond data for the [EMIM]₃[LaX₆] series of compounds. ^a data from Reference 7

	[EMIM] ₃ [LaCl ₆] ^a		[EMIM] ₃ [LaBr ₆]		[EMIM] ₃ [LaI ₆]	
	Length (Å)	Angle (°)	Length (Å)	Angle (°)	Length (Å)	Angle (°)
C...X (H2)	3.557(8)	152.2	3.579(12)	134.2	3.942(5)	139.8
	3.479(6)	131.6	3.659(10)	153.8	3.758(5)	134.0
	3.393(4)	120.2	3.765(17)	134.7	3.864(5)	134.9
	3.612(4)	150.2			3.825(5)	160.4
C...X (H4/5)	3.635(5)	150.3	3.723(12)	121.6	3.974(5)	136.7
	3.614(6)	148.4	3.683(11)	139.8	3.972(5)	113.5
	3.692(5)	146.7	3.744(12)	118.3	3.796(5)	143.2
	3.632(5)	145.9	3.755(11)	146.8	3.894(5)	119.0
	3.564(6)	137.5	3.708(12)	145.1	3.830(5)	124.0
			3.693(12)	153.7	4.210(5)	172.5
			3.708(16)	143.9		

Varying the metal:

Another variable to assess in this series of complexes is the nature of the metal ion. By choosing a range of trivalent metals for [EMIM]₃[MCl₆] (M = Sc, Y, La, Ce) it is possible to vary the ionic radius, and hence the charge:radius ratio, and observe the effect on the hydrogen bonding. Addition of [EMIM]Cl to MCl₃ led to the formation of crystalline samples of [EMIM]₃[ScCl₆] (Figure 3), [EMIM]₃[YCl₆] (Figure 4) and [EMIM]₃[CeCl₆] (Figure 5).




Figure 3. ORTEP representation of $[\text{EMIM}]_3[\text{ScCl}_6]$ showing the hydrogen bonding (dotted lines). Displacement ellipsoids are drawn at the 50% probability level and hydrogen atoms not involved in hydrogen bonding are omitted for clarity. Selected bond lengths (Å): Sc–Cl2 2.479(1), Sc–Cl3 2.513(1), Sc–Cl4 2.5066(7), Sc–Cl5 2.520(1), Sc–Cl6 2.4735(8), Sc–Cl7 72.522(1). Symmetry codes: (i) $1 - x, -y, z - 0.5$; (ii) $x - 0.5, 0.5 - y, z$; (iii) $x, y - 1, z$.




Figure 4. ORTEP representation of $[\text{EMIM}]_3[\text{YCl}_6]$ showing the hydrogen bonding (dotted lines). Displacement ellipsoids are drawn at the 50% probability level and hydrogen atoms not involved in hydrogen bonding are omitted for clarity. Selected bond lengths (Å): Y–Cl1 2.6541(12), Y–Cl2 2.614(1), Y–Cl3 2.6185(12), Y–Cl4 2.640(1), Y–Cl5 2.6515(14), Y–Cl6 2.6524(14). Symmetry codes: (i) $1 - x, 1 - y, z - 0.5$; (ii) $0.5 - x, y - 0.5, z - 0.5$; (iii) $x, y - 1, z$; (iv) $x, y, z - 1$.




Figure 5. ORTEP representation of $[\text{EMIM}]_3[\text{CeCl}_6]$ showing the hydrogen bonding (dotted lines) surrounding the Ce1-centred cation (left) and the Ce2-centred cation (right). Displacement ellipsoids are drawn at the 50% probability level and H atoms not involved in hydrogen bonding are omitted for clarity. Selected bond lengths (Å): Ce1–C11 2.7872(16), Ce1–C12 2.7514(17), Ce1–C13 2.7738(16), Ce2–C14 2.7755(17), Ce2–C15 2.7752(17), Ce2–C16 2.7722(17). Symmetry codes: (i) $-x, -y, -z$; (ii) $-1 - x, -y, 1 - z$; (iii) $-x, 0.5 + y, 0.5 - z$; (iv) $x, -0.5 - y, 0.5 + z$; (v) $-x, y - 0.5, 0.5 - z$.

The Sc and Y complexes are isostructural and isomorphous, with a slight increase in cell parameters owing to the increase in metal ionic radius (Sc = 0.745 Å, Y = 0.90 Å for 6-coordinate complexes).¹⁴ The known lanthanide examples are structurally different, but within the series (including La, Eu, Gd, Nd, Pr and Sm analogues) they are all isomorphous and isostructural⁷ and the cell parameters for $[\text{EMIM}]_3[\text{CeCl}_6]$ fit well with this trend.

In the solid state all the metal centres are six-coordinate regular octahedra. There is only one metal environment in the asymmetric unit of $[\text{EMIM}]_3[\text{ScCl}_6]$ and $[\text{EMIM}]_3[\text{YCl}_6]$ and the hydrogen bonding interactions are very similar (Table 2). Despite the decreased charge:radius ratio for Y over Sc, the hydrogen bonds through H4/5 are identical within experimental error. However, there is a small increase in the hydrogen bond lengths involving H2 for $[\text{EMIM}]_3[\text{YCl}_6]$ compared to $[\text{EMIM}]_3[\text{ScCl}_6]$. These are notably shorter than the equivalent bonds in $[\text{EMIM}]_3[\text{LaCl}_6]$, continuing the same trend. Trifurcated hydrogen bonding is present in $[\text{EMIM}]_3[\text{CeCl}_6]$ through C13 (Figure 5).

Table 2: Hydrogen bond data for the [EMIM]₃[MCl₆] series of compounds prepared in this study.

	[EMIM] ₃ [ScCl ₆]		[EMIM] ₃ [YCl ₆]		[EMIM] ₃ [CeCl ₆]	
	Length (Å)	Angle (°)	Length (Å)	Angle (°)	Length (Å)	Angle (°)
C···X (H2)	3.506(3)	149.5	3.547(4)	146.6	3.547(7)	150.8
	3.496(3)	131.3	3.531(5)	130.0	3.425(7)	132.4
	3.407(3)	138.1	3.422(5)	135.9	3.333(7)	117.1
	3.604(3)	135.2	3.650(4)	134.6	3.569(8)	141.9
	3.548(3)	146.8	3.588(5)	145.0		
C···X (H4/5)	3.586(3)	166.7	3.584(5)	165.0	3.568(7)	150.4
	3.692(3)	152.5	3.695(4)	153.1	3.583(7)	143.2
	3.604(3)	135.6	3.604(5)	135.8	3.622(8)	142.7
	3.434(3)	131.7	3.459(4)	128.6	3.542(8)	147.2
	3.383(3)	135.1	3.365(5)	136.2	3.512(7)	139.7

Varying the cation:

Altering the nature of the cation can have a marked effect on the structural properties of the complexes. In order to probe this effect the [CATION]₃[LaCl₆] system was initially chosen (CATION = EMIM, EDMIM, NⁿBu₄ and BMPYRR). Complexes were prepared by adding a slight excess of [CATION]Cl to LaCl₃ in MeCN, resulting in a series of compounds where the hydrogen bonding interactions should be moderate (EMIM), weak (EDMIM) and non-existent (NⁿBu₄, BMPYRR). The difference between the [EDMIM]⁺ cation and the [EMIM]⁺ cation is the presence of a methyl group at the C2 position, thus the most acidic protons are now on the backbone of the imidazolium ring. This was expected to significantly influence the hydrogen bonding. The structure of [EDMIM]₃[LaCl₆] is shown in Figure 6.




Figure 6. ORTEP representation of $[\text{EDMIM}]_3[\text{LaCl}_6]$ showing the hydrogen bonding (dotted lines). Displacement ellipsoids are drawn at the 50% probability level. Selected bond lengths (Å): La–Cl1 2.7963(7), La–Cl2 2.7582(6), La–Cl3 2.8029(6), La–Cl4 2.7950(7), La–Cl5 2.8262(6), La–Cl6 2.7742(5). Symmetry codes: (i) $1 - x, 1 - y, 1 - z$; (ii) $x - 1, y, z$.

Structural characterisation of $[\text{EDMIM}]_3[\text{LaCl}_6]$ revealed that the $[\text{LaCl}_6]^{3-}$ trianion was stabilised by hydrogen bonding through five associated $[\text{EDMIM}]^+$ cations. The $\text{C}\cdots\text{Cl}$ distances were slightly shorter than the corresponding $\text{C}\cdots\text{Cl}$ distances for $[\text{EMIM}]_3[\text{LaCl}_6]$, suggesting that hydrogen bonding through the backbone protons may be slightly stronger when no H2 proton is present. However, ^1H NMR data (below) show that the chemical shifts for H4/5 are consistent with complete dissociation of the molecule in CD_3CN solution. The La–Cl bond lengths are comparable to those observed for $[\text{EMIM}]_3[\text{LaCl}_6]$.

A comparison between the $[\text{EMIM}]^+$ and $[\text{EDMIM}]^+$ cations can also be drawn for the $[\text{YCl}_6]^{3-}$ and $[\text{LaI}_6]^{3-}$ salts. The structures of $[\text{EDMIM}]_3[\text{YCl}_6]$ and $[\text{EDMIM}]_3[\text{LaI}_6]$ are shown in Figures 7 and 8, respectively.




Figure 7. ORTEP representation of $[\text{EDMIM}]_3[\text{YCl}_6]$ showing the hydrogen bonding (dotted lines). Displacement ellipsoids are drawn at the 50% probability level and hydrogen atoms not involved in hydrogen bonding are omitted for clarity. Selected bond lengths (Å): Y–Cl1 2.6580(5), Y–Cl2 2.6398(5), Y–Cl3 2.6496(5), Y–Cl4 2.6800(5), Y–Cl5 2.6464(5), Y–Cl6 2.6167(5). Symmetry codes: (i) $-x, 2 - y, -z$; (ii) $x + 0.5, 1.5 - y, z - 0.5$; (iii) $-x, 1 - y, -z$; (iv) $1 - x, 1 - y, -z$.




Figure 8. ORTEP representation of $[\text{EDMIM}]_3[\text{LaI}_6]$ showing the hydrogen bonding (dotted lines) surrounding the La1-centred cation (left) and the La2-centred cation (right). Displacement ellipsoids are drawn at the 50% probability level and H atoms not involved in hydrogen bonding are omitted for clarity. Selected bond lengths (Å): La1–I1 3.1800(7), La1–I2 3.1545(6), La1–I3 3.1903(7), La2–I4 3.1779(5), La2–I5 3.1566(5), La2–I6 3.1870(5). Symmetry codes: (i) $1 - x, -y, 1 - z$; (ii) $-x, -1 - y, -z$; (iii) $x, y - 1, z$; (iv) $-x, -1 - y, 1 - z$; (v) $1 - x, -1 - y, 1 - z$; (vi) $x, y, 1 + z$; (vii) $x, 1 + y, z$.

$[\text{EDMIM}]_3[\text{YCl}_6]$ and $[\text{EDMIM}]_3[\text{LaCl}_6]$ are isomorphous and isostructural, whereas $[\text{EMIM}]_3[\text{YCl}_6]$ and $[\text{EMIM}]_3[\text{LaCl}_6]$ crystallise in different space groups. This suggests that the hydrogen bonding interactions play a greater role in the crystal packing arrangements for the $[\text{EMIM}]^+$ salts. The $\text{C}\cdots\text{Cl}$ (H4/H5) distances (Table 3) are not notably different between $[\text{EMIM}]_3[\text{YCl}_6]$ and $[\text{EDMIM}]_3[\text{YCl}_6]$.

For $[\text{EDMIM}]_3[\text{LaI}_6]$ two different $[\text{LaX}_6]^{3-}$ environments are apparent, with one iodine acting as an acceptor to five different donor hydrogens. However, the $\text{C}\cdots\text{I}$ distances (Table 3) are not notably different between the EMIM and EDMIM compounds.

Table 3: Hydrogen bond data for $[\text{EDMIM}]_3[\text{YCl}_6]$, $[\text{EDMIM}]_3[\text{LaCl}_6]$ and $[\text{EDMIM}]_3[\text{LaI}_6]$.

	$[\text{EDMIM}]_3[\text{YCl}_6]$		$[\text{EDMIM}]_3[\text{LaCl}_6]$		$[\text{EDMIM}]_3[\text{LaI}_6]$	
	Length (Å)	Angle (°)	Length (Å)	Angle (°)	Length (Å)	Angle (°)
$\text{C}\cdots\text{X}$ (H4/5)	3.542(2)	142.4	3.580(2)	149.6	3.948(3)	118.8
	3.693(2)	123.8	3.515(2)	136.6	3.956(3)	118.5
	3.388(2)	129.8	3.521(2)	148.9	3.933(3)	139.2
	3.482(2)	145.1	3.686(2)	122.5	3.775(3)	125.2
	3.520(2)	149.9	3.455(2)	125.8	3.883(3)	124.3
	3.544(2)	150.2	3.601(2)	112.0	3.892(3)	124.8
	3.577(2)	122.6	3.524(2)	143.0	3.883(3)	115.3

Spectroscopic characterisation and comparisons:

In keeping with our previous observations for $[\text{GeCl}_6]^{2-}$ anions,³ the chemical shift of the H2 proton in the ^1H NMR spectrum was used to indicate the strength of the hydrogen bonding in solution (Table 4).

From the NMR data it is apparent that the chemical shift of the signal associated with the H2 proton shifts to low frequency for the heavier halides. This correlates with decreasing strength of the solution phase hydrogen bonding interactions upon changing from Cl to I. The chemical shifts of the less acidic H4/5 protons are not affected by changing the halide, suggesting that the salts are completely dissociated in solution. This is consistent with the trend observed for $[\text{EMIM}][\text{GeX}_3]$ ($X = \text{Cl}, \text{Br}, \text{I}$).³

For the $[\text{MX}_6]^{3-}$ salts, ^{45}Sc , ^{89}Y and ^{139}La NMR spectra were also obtained (Table 5). All complexes were soluble in MeCN at approximate 10–15 mM concentration. The $[\text{N}^n\text{Bu}_4]^+$ salt was soluble at a similar concentration in CH_2Cl_2 , whereas the $[\text{EMIM}]^+$, $[\text{BMPYRR}]^+$ and $[\text{EDMIM}]^+$ salts were poorly soluble in CH_2Cl_2 . For the $[\text{LaCl}_6]^{3-}$ salts a small high frequency shift in $\delta^{139}\text{La}$ occurs as the cation changes along the series: $[\text{N}^n\text{Bu}_4]^+ \rightarrow [\text{BMPYRR}]^+ \rightarrow [\text{EDMIM}]^+ \rightarrow [\text{EMIM}]^+$ in both MeCN and CH_2Cl_2 solution, but given the large ^{139}La NMR chemical shift range, these differences may not be significant. The ^{139}La NMR resonance shifts to high frequency as the halide becomes heavier; the $[\text{LaI}_6]^{3-}$ system showing a very broad resonance at ~ 1400 ppm ($w/2 \sim 12000$ Hz). The yttrium salts show no ^{89}Y NMR resonance at room temperature in MeCN, presumably due to chloride exchange. However, cooling to 233 K reveals a sharp singlet for each compound.

Table 4 Effect of $[\text{MX}_6]^{3-}$ on the N-CH-N ^1H NMR shift in CD_3CN solution

Compound	$\delta(^1\text{H})/\text{ppm}^a$	
	N-CH-N	Δ^b
$[\text{EMIM}]_3[\text{LaCl}_6]$	9.27	+0.45
$[\text{EMIM}]_3[\text{LaBr}_6]$	8.93	+0.11
$[\text{EMIM}]_3[\text{LaI}_6]$	8.61	-0.21
$[\text{EMIM}]_3[\text{CeCl}_6]$	9.77	+0.95
$[\text{EMIM}]_3[\text{YCl}_6]$	9.26	+0.44
$[\text{EMIM}]_3[\text{ScCl}_6]$	9.41	+0.59
$[\text{EMIM}]\text{Cl}$	8.82	-
$[\text{EMIM}]\text{Br}$	8.84	-

^a 6.8 mM solutions in CD_3CN ; ^b Change in shift compared with $[\text{EMIM}]\text{Cl}$ at the same concentration in CD_3CN

Table 5 Selected NMR spectroscopic data for [CATION]₃[MX₆].

Salt	$\delta(^{139}\text{La}, ^{89}\text{Y}, ^a\text{^{45}Sc})/\text{ppm}$	$W_{1/2}$ (Hz)	δ (ppm)	$W_{1/2}$ (Hz)
	MeCN		CH ₂ Cl ₂	
[N ⁿ Bu ₄] ₃ [LaCl ₆]	846	800	872	1200
[BMPYRR] ₃ [LaCl ₆]	855	500	880	300
[EDMIM] ₃ [LaCl ₆]	864	600	n.o. ^b	-
[EMIM] ₃ [LaCl ₆]	867	400	892	300
[EMIM] ₃ [LaBr ₆]	1095	1200	n.o. ^b	-
[EMIM] ₃ [LaI ₆]	~1400	12000	n.o.	-
[EMIM] ₃ [YCl ₆]	407	-	-	-
[EDMIM] ₃ [YCl ₆]	401	-	-	-
[EMIM] ₃ [ScCl ₆]	254	170	-	-

^a spectrum recorded at 233 K;

^b n.o. = not observed due to low solubility

Conclusions

All of the salts are based upon six-coordinate [MX₆]³⁻ units, and for those containing EMIM and EDMIM cations, significant C–H...X hydrogen-bonding interactions to the substituted imidazolium cations are evident from crystallographic studies. The shortest C...X distances for the [EMIM]₃[MX₆] series involve the NCN (H2) proton, with much weaker interactions to the H4/H5 protons. Despite the decreased charge:radius ratio along the series from Sc to Y to La, the hydrogen bonds through H4/5 are unaffected. However, there is a small increase in the hydrogen bond lengths involving H2 along the same series.

Substitution of the H2 proton with a Me group in the EDMIM cation has a significant influence on the cation-anion interactions in the solid state. The structure of [EDMIM]₃[LaCl₆] revealed five associated [EDMIM]⁺ cations and the C...Cl distances involving H4/H5 were shorter than those for [EMIM]₃[LaCl₆].

The solution ¹H NMR trends indicate that the hydrogen bonding to the EMIM (H2) cations becomes weaker as the halide becomes heavier, while the EDMIM salts appear to be completely dissociated in solution.

Experimental

All preparations were carried out using standard Schlenk techniques under a N₂ atmosphere. Anhydrous MCl₃ (M = La, Ce) were obtained by reacting hydrated MCl₃ (Sigma) with excess SOCl₂, then drying *in vacuo*. Anhydrous ScCl₃, YCl₃, LaBr₃ and LaI₃ were obtained from Alfa and used as received. [EMIM]X (X = Cl, Br, I), [EDMIM]Cl, [NⁿBu₄]Cl and [BMPYRR]Cl were obtained from Sigma, dried *in vacuo* at 100 °C for 4 hours and stored in a glove box. [EDMIM]I was prepared by stirring a solution of [EDMIM]Cl and NaI in water for 4 hours, then extracting into

CH₂Cl₂ followed by drying *in vacuo* at 100 °C for 4 hours. MeCN and CH₂Cl₂ were dried prior to use from CaH₂. Infrared spectra were recorded as Nujol mulls between CsI plates using a Perkin-Elmer Spectrum 100 spectrometer over the range 4000–200 cm⁻¹. Raman spectra were recorded from powdered solid samples using a Perkin-Elmer FT-Raman 2000R with a Nd:YAG laser. ¹H NMR spectra were recorded in CD₂Cl₂ and CD₃CN using a Bruker DPX400 spectrometer and are referenced to the residual protio solvent. ⁴⁵Sc, ⁸⁹Y and ¹³⁹La NMR spectra were also recorded in CD₂Cl₂ or CD₃CN on a Bruker DPX400 spectrometer and are referenced to 0.1 mol dm⁻³ solutions of the chlorides in D₂O solutions at pH = 1. For yttrium-89 NMR spectra a 3 s pulse delay was used and TEMPO (2,2,6,6-tetramethylpiperidine-oxyl) was added as relaxation agent. Microanalyses were undertaken by Medac Ltd or London Metropolitan University.

Preparations

[NⁿBu₄]₃[LaCl₆]: LaCl₃ (100 mg, 0.408 mmol) was suspended in dichloromethane (10 mL) in a Schlenk flask and a solution of [NⁿBu₄]Cl (396 mg, 1.43 mmol) in acetone (10 mL) added. The mixture was stirred until all the solids had dissolved. Diethyl ether (20 mL) was added slowly and a white solid precipitated which was filtered and dried under vacuum. Yield: 345 mg, 79%. Raman (solid): 262 cm⁻¹ (A_{1g} [LaCl₆]³⁻). ¹H NMR (CD₃CN, 300 K): δ = 3.08-3.14 (m, [2H]), 1.55-1.66 (m, [2H]), 1.30-1.42 (m, [2H]), 0.97 (t, *J* = 7.35 Hz, [3H]). ¹³⁹La NMR (CH₃CN/CD₃CN, 300 K): δ = 846 (s, *W*_{1/2} = 800 Hz). Anal. required for C₄₈H₁₀₈Cl₆LaN₃: C, 53.43; H, 10.09; N, 3.89%. Found: C, 53.29; H, 9.92; N, 4.04%.

[EDMIM]₃[LaCl₆]: Synthesised in an analogous way, using LaCl₃ (100 mg, 0.408 mmol) and [EDMIM]Cl (197 mg, 1.22 mmol). White crystalline product. Yield: 213 mg, 72%. Raman (solid): 244 cm⁻¹ (A_{1g} [LaCl₆]³⁻). ¹H NMR (CD₃CN, 300 K): δ = 7.36 (s, [2H]), 4.12 (q, *J* = 7.16 Hz, [2H]), 3.75 (s, [3H]), 2.54 (s, [3H]), 1.39 (t, *J* = 7.34 Hz, [3H]), (CD₂Cl₂, 300 K): δ = 7.64 (d, *J* = 2.20 Hz, [H]), 7.46 (d, *J* = 2.20 Hz, [H]), 4.23 (q, *J* = 7.5 Hz, [2H]), 3.97 (s, [3H]), 2.75 (s, [3H]), 1.50 (t, *J* = 7.34 Hz, [3H]). ¹³⁹La NMR (CH₃CN/CD₃CN): δ = 864 (s, *W*_{1/2} = 600 Hz). Anal. required for C₂₁H₃₉Cl₆LaN₆: C, 34.68; H, 5.41; N, 11.56%. Found: C, 34.55; H, 5.34; N, 11.45%.

[BMPYRR]₃[LaCl₆]: LaCl₃ (100 mg, 0.407 mmol) was suspended in MeCN (10 mL) and a solution of [BMPYRR]Cl, (255 mg, 1.43 mmol) in MeCN (10 mL) was added. The mixture was heated briefly and then stirred for one hour. The resulting suspension was filtered and diethyl ether (10 mL) was added to the filtrate, leading to the precipitation of a fine white crystalline product. The solvent was removed *via* cannula and the solid was dried under vacuum. Yield: 245 mg, 77%. Raman (solid): 240 cm⁻¹ (A_{1g} [LaCl₆]³⁻). ¹H NMR (CD₃CN, 300 K): δ = 3.44-3.59 (m, [4H]), 3.30-3.35 (m, [2H]), 3.03 (s, [3H]), 2.18 (br. s, [4H]), 1.68-1.78 (m, [2H]), 1.32-1.45 (m, [2H]), 0.96 (t, *J* = 7.35 Hz, [3H]). ¹³⁹La NMR (CH₃CN/CD₃CN): δ = 855 (s, *W*_{1/2} = 500 Hz). Anal. required for C₂₇H₆₀Cl₆LaN₃: C, 41.66; H, 7.77; N, 5.40%. Found: C, 41.81; H, 7.63; N, 5.50%.

[EMIM]₃[LaCl₆]: LaCl₃ (100 mg, 0.408 mmol) was suspended in MeCN (5 mL) and a solution of [EMIM]Cl, (210 mg, 1.43 mmol) in MeCN (5 mL) was added. The mixture was heated briefly and then stirred for 1 hour. The resulting suspension was filtered and diethyl ether (10 mL) was added to the filtrate, leading to the

precipitation of a white crystalline product. This was allowed to settle, the solvent removed *via* cannula and the solid was then dried under vacuum. Yield: 185 mg, 66%. Raman: 240 cm^{-1} (A_{1g} [LaCl_6] $^{3-}$). ^1H NMR (CD_3CN , 300 K): δ = 9.27 (s, [H]), 7.39 (s, [H]), 7.34 (s, [H]), 4.33 (q, J = 7.16 Hz, [2H]), 3.96 (s, [3H]), 1.46 (t, J = 7.16 Hz, [3H]). ^{139}La NMR ($\text{CH}_3\text{CN}/\text{CD}_3\text{CN}$): δ = 867 (s, $W_{1/2}$ = 400 Hz). Anal. required for $\text{C}_{18}\text{H}_{33}\text{Cl}_6\text{LaN}_6$: C, 31.56; H, 4.85; N, 12.27%. Found: C, 31.31; H, 4.67; N, 12.07%.

[EMIM] $_3$ [LaBr $_6$]: Synthesised in an analogous way, using LaBr_3 (100 mg, 0.264 mmol) and [EMIM]Br (151 mg, 0.792 mmol). White solid. Yield: 143 mg, 57%. ^1H NMR (CD_3CN , 300 K): 8.93 (s, [H]), 7.42 (s, [H]), 7.37 (s, [H]), 4.27 (q, J = 7.34 Hz, [2H]), 3.91 (s, [3H]), 1.47 (t, J = 7.34 Hz, [3H]). ^{139}La NMR ($\text{CH}_3\text{CN}/\text{CD}_3\text{CN}$, 300 K): δ = 1095 (s, $W_{1/2}$ = 1200 Hz). Anal. required for $\text{C}_{18}\text{H}_{33}\text{Br}_6\text{LaN}_6$: C, 22.71; H, 3.49; N, 8.83%. Found: C, 22.57; H, 3.39; N, 8.74%.

[EMIM] $_3$ [LaI $_6$]: Synthesised in an analogous way, using LaI_3 (150 mg, 0.289 mmol) and [EMIM]I (240 mg, 1.01 mmol). After filtering, the solvent volume was reduced to approximately 25% and diethyl ether was added. Large colourless crystals suitable for X-ray diffraction were grown overnight. Yield: 180 mg, 46%. ^1H NMR (CD_3CN , 300 K): δ = 8.61 (s, [H]), 7.41 (s, [H]), 7.36 (s, [H]), 4.19 (q, J = 7.16 Hz, [2H]), 3.84 (s, [H]), 1.46 (t, J = 7.35 Hz, [3H]). Raman: 110 cm^{-1} (A_{1g} [LaI_6] $^{3-}$). Anal. required for $\text{C}_{18}\text{H}_{33}\text{I}_6\text{LaN}_6$: C, 17.52; H, 2.70; N, 6.81%. Found: C, 17.38; H, 2.65; N, 6.94%.

[EDMIM] $_3$ [LaI $_6$]: Synthesised in an analogous way. Yield: 55%. White crystalline solid. ^1H NMR (CD_3CN , 300 K): δ = 7.36 (d, [2H], J = 1.96 Hz), 4.08 (q, J = 7.34 Hz, [2H]), 3.70 (s, [3H]), 2.51 (s, [3H]), 1.38 (t, J = 7.34 Hz, [3H]). Anal. required for $\text{C}_{21}\text{H}_{39}\text{I}_6\text{LaN}_6$: C, 19.77; H, 3.08; N, 6.59%. Found: C, 19.60; H, 3.15; N, 6.39%.

[EMIM] $_3$ [CeCl $_6$]: Synthesised in an analogous way, using CeCl_3 (100 mg, 0.406 mmol) and [EMIM]Cl (208 mg, 1.42 mmol), forming a white crystalline product. Yield: 175 mg, 63%. Raman (solid): 244 cm^{-1} (A_{1g} [CeCl_6] $^{3-}$). ^1H NMR (CD_3CN , 300 K): δ = 9.77 (s, [H]), 7.45 (s, [H]), 7.40 (s, [H]), 4.44 (q, J = 7.2 Hz, [2H]), 4.01 (s, [3H]), 1.51 (t, J = 6.6 Hz, [3H]). Anal. required for $\text{C}_{18}\text{H}_{33}\text{CeCl}_6\text{N}_6$: C, 31.50; H, 4.85; N, 12.24%. Found: C, 31.21; H, 4.58; N, 12.54%.

[EMIM] $_3$ [YCl $_6$]: Synthesised in an analogous way, using YCl_3 (100 mg, 0.512 mmol) and [EMIM]Cl (225 mg, 1.54 mmol). White crystalline product. Yield: 277 mg, 85%. Infrared (Nujol): 245 cm^{-1} (T_{1u} [YCl_6] $^{3-}$). Raman (solid): 264 cm^{-1} (A_{1g} [YCl_6] $^{3-}$). ^1H NMR (CD_3CN , 300 K): δ = 9.26 (s, [H]), 7.39 (s, [H]), 7.34 (s, [H]), 4.31 (q, J = 7.34 Hz, [2H]), 3.93 (s, [3H]), 1.46 (t, J = 7.34 Hz, [3H]). Anal. required for $\text{C}_{18}\text{H}_{33}\text{Cl}_6\text{N}_6\text{Y}$: C, 34.04; H, 5.24; N, 13.23%. Found: C, 33.90; H, 5.15; N, 13.18%.

[EDMIM] $_3$ [YCl $_6$]: Synthesised in an analogous way, 72%. White crystalline solid. Infrared (Nujol): 238 cm^{-1} (T_{1u} [YCl_6] $^{3-}$). ^1H NMR (CD_3CN , 300 K): δ = 7.40 (s, [2H]), 4.12 (q, J = 7.16 Hz, [2H]), 2.54 (s, [3H]), 3.75 (s, [3H]), 1.39 (t, J = 7.34 Hz, [3H]). (CD_2Cl_2 , 300 K): δ = 7.64 (d, J = 2.20 Hz, [H]), 7.46 (d, J = 2.20 Hz, [H]),

4.23 (q, $J = 7.5$ Hz, [2H]), 3.97 (s, [3H]), 2.75 (s, [3H]), 1.50 (t, $J = 7.34$ Hz, [3H]). ^{139}La NMR ($\text{CH}_3\text{CN}/\text{CD}_3\text{CN}$): $\delta = 864$ (s, $W_{1/2} = 600$ Hz). Anal. required for $\text{C}_{21}\text{H}_{39}\text{Cl}_6\text{N}_6\text{Y}$: C, 37.25; H, 5.80; N, 12.41%. Found: C, 37.12; H, 5.93; N, 12.20%.

[EMIM]₃[ScCl₆]: Synthesised in an analogous way, using ScCl_3 . White crystalline product, 85%. Infrared (Nujol): 290 cm^{-1} (T_{1u} [ScCl_6] $^{3-}$). ^1H NMR (CD_3CN , 300 K): $\delta = 9.41$ (s, [H]), 7.41 (s, [H]), 7.38, (s, [H]), 4.26 (q, $J = 7.34$ Hz, [2H]), 3.90 (s, [3H]), 1.46 (t, $J = 7.21$ Hz, [3H]). ^{45}Sc NMR ($\text{CH}_3\text{CN}/\text{CD}_3\text{CN}$): $\delta = 254.5$. Anal. required for $\text{C}_{18}\text{H}_{33}\text{Cl}_6\text{N}_6\text{Sc}$: C, 36.57; H, 5.63; N, 14.22%. Found: C, 36.39; H, 5.71; N, 14.09%.

X-ray Experimental: Crystals suitable for single crystal X-ray analysis were obtained as described above. Details of the crystallographic data collection and refinement parameters are given in Table 6. Data collections were carried out on a Rigaku AFC12 goniometer equipped with an enhanced sensitivity (HG) Saturn724+ detector mounted at the window of an FRE+ SuperBright molybdenum rotating anode generator ($\lambda_1 = 0.71073$ Å) with VHF Varimax optics (70 or 100 μm focus), apart from [EMIM]₃[LaI₆] which was collected on a Rigaku R-axis Spider including curved Fuji-film image plate and a graphite monochromated sealed tube Mo generator ($\lambda_1 = 0.71073$ Å). For all datasets: cell determination, data collection, data reduction, cell refinement and absorption correction were carried out using CrystalClear-SM Expert 2.0 r7.¹⁵

Structure solution and refinement were straightforward using WinGX and software packages within,¹⁶ except as detailed below. Although Q-peaks corresponding to the location of protons were observed in the Fourier difference map, hydrogen atoms were placed in geometrically assigned positions with C–H distances of 0.95 Å (CH), 0.98 Å (CH₃) or 0.99 Å (CH₂) and refined using a riding model, with $U_{iso}(\text{H}) = 1.2 U_{eq}(\text{C})$ (CH, CH₂) or $1.5 U_{eq}(\text{C})$ (CH₃). Mercury¹⁷ and enCIFer¹⁸ were used to prepare material for publication.

All the crystals of [EMIM]₃[CeCl₆] looked at in this study were multiply twinned and all attempts at multi-component twin integration failed. A satisfactory dataset was obtained by integrating on just the major component and ignoring the minor overlaps from the other domains. This caused a couple of large Q-peaks to appear close to the cerium atoms ($3.2\text{ e}\text{\AA}^{-3}$ at ~ 1.04 Å distance from Ce) as well as two large holes ($-5.9\text{ e}\text{\AA}^{-3}$ at ~ 0.84 Å from Ce). This explains the level A and B CheckCIF errors PLAT971_ALERT_2_A (holes) and PLAT971_ALERT_2_B (Q-peaks). The crystals of [EMIM]₃[LaBr₆] were all also multi-component twins and treated in a similar fashion.

Supplementary Information

Crystallographic data in cif format have been deposited with the Cambridge Crystallographic Data Centre (CCDC) and given reference numbers 1472564 ([EMIM]₃[ScCl₆]), 1472565 ([EMIM]₃[YCl₆]), 1472566 ([EMIM]₃[CeCl₆]), 1472567 ([EMIM]₃[LaBr₆]), 1472568 ([EMIM]₃[LaI₆]), 1472569 ([EDMIM]₃[YCl₆]), 1472570 ([EDMIM]₃[LaCl₆]) and 1472571 ([EDMIM]₃[LaI₆]). Copies of the data can be obtained free of charge

from The Director, CCDC, 12 Union Road, Cambridge, CB2 1EZ, UK, fax: +44 1223 366033, e-mail: deposit@ccdc.cam.ac.uk or on the web at <http://www.ccdc.cam.ac.uk>.

Acknowledgements

This work was funded by EPSRC through a Programme Grant (EP/I033394/1) and also through EP/K039466/1. The SCFED Project (www.scfed.net) is a multidisciplinary collaboration of British universities investigating the fundamental and applied aspects of supercritical fluids.

Table 6: X-ray crystallographic data^a

Compound	[EMIM] ₃ [ScCl ₆]	[EMIM] ₃ [YCl ₆]	[EMIM] ₃ [CeCl ₆]	[EMIM] ₃ [LaBr ₆]
Formula	C ₁₈ H ₃₃ Cl ₆ N ₆ Sc	C ₁₈ H ₃₃ Cl ₆ N ₆ Y	C ₁₈ H ₃₃ CeCl ₆ N ₆	C ₁₈ H ₃₃ Br ₆ LaN ₆
<i>M</i> /g mol ⁻¹	591.16	635.11	686.32	951.87
Temp./K	150(2)	100(2)	100(2)	100(2)
Crystal system	orthorhombic	orthorhombic	monoclinic	monoclinic
Space group (No.)	<i>Pna</i> 2 ₁ (33)	<i>Pna</i> 2 ₁ (33)	<i>P</i> 2 ₁ / <i>c</i> (14)	<i>P</i> 2 ₁ / <i>c</i> (14)
<i>a</i> /Å	19.411(2)	19.580(2)	15.492(3)	15.925(5)
<i>b</i> /Å	13.059(1)	13.036(2)	12.549(2)	12.787(4)
<i>c</i> /Å	11.046(1)	11.166(1)	14.708(3)	15.252(5)
α /°	90	90	90	90
β /°	90	90	90.482(3)	90.588(5)
γ /°	90	90	90	90
<i>U</i> /Å ³	2800.0(4)	2849.9(5)	2859.2(9)	3106(2)
<i>Z</i>	4	4	4	4
μ (Mo-K α)/mm ⁻¹	0.853	2.626	2.170	9.111
<i>F</i> (000)	1224	1296	1372	1800
Total reflections	15098	14268	12086	14407
Unique reflections	5189	6374	5811	6318
<i>R</i> _{int}	0.027	0.037	0.122	0.092
Goodness-of-fit on <i>F</i> ²	1.023	0.796	0.962	1.145
<i>R</i> ₁ ^b [<i>I</i> _o > 2 σ (<i>I</i> _o)]	0.026	0.028	0.063	0.082
<i>R</i> ₁ (all data)	0.029	0.043	0.090	0.116
<i>wR</i> ₂ ^b [<i>I</i> _o > 2 σ (<i>I</i> _o)]	0.063	0.045	0.167	0.134
<i>wR</i> ₂ (all data)	0.064	0.048	0.187	0.148

^a Common items: wavelength (Mo-K α) = 0.71073 Å; θ (max) = 27.5°.

^b $R_1 = \sum ||F_o| - |F_c|| / \sum |F_o|$; $wR_2 = [\sum w(F_o^2 - F_c^2)^2 / \sum wF_o^4]^{1/2}$

Table 6 cont.


Compound	[EMIM] ₃ [LaI ₆]	[EDMIM] ₃ [YCl ₆]	[EDMIM] ₃ [LaCl ₆]	[EDMIM] ₃ [LaI ₆]
Formula	C ₁₈ H ₃₃ I ₆ LaN ₆	C ₂₁ H ₃₉ Cl ₆ N ₆ Y	C ₂₁ H ₃₉ Cl ₆ LaN ₆	C ₂₁ H ₃₉ I ₆ LaN ₆
<i>M</i> /g mol ⁻¹	1233.81	677.19	727.19	1275.89
Temp./K	120(2)	100(2)	100(2)	100(2)
Crystal system	monoclinic	monoclinic	monoclinic	triclinic
Space group (No.)	<i>P</i> 2 ₁ / <i>c</i> (14)	<i>P</i> 2 ₁ / <i>n</i> (14)	<i>P</i> 2 ₁ / <i>n</i> (14)	<i>P</i> -1 (2)
<i>a</i> /Å	18.803(7)	9.9170(8)	10.027(2)	10.347(2)
<i>b</i> /Å	11.133(2)	16.274(1)	16.306(2)	11.075(3)
<i>c</i> /Å	17.017(2)	19.028(2)	19.133(3)	16.577(3)
α /°	90	90	90	92.594(3)
β /°	102.162(7)	99.973(1)	99.265(3)	90.255(3)
γ /°	90	90	90	106.930(4)
<i>U</i> /Å ³	3482(1)	3024.6(4)	3087.4(9)	1815.1(7)
<i>Z</i>	4	4	4	2
μ (Mo-K α)/mm ⁻¹	6.562	2.480	1.924	6.299
<i>F</i> (000)	2232	1392	1464	1164
Total reflections	34033	16980	26415	16557
Unique reflections	7929	6878	7055	8311
<i>R</i> _{int}	0.059	0.027	0.020	0.028
Goodness-of-fit on <i>F</i> ²	1.073	1.028	1.078	0.984
<i>R</i> ₁ ^b [<i>I</i> _o > 2 σ (<i>I</i> _o)]	0.031	0.028	0.023	0.019
<i>R</i> ₁ (all data)	0.036	0.039	0.026	0.021
<i>wR</i> ₂ ^b [<i>I</i> _o > 2 σ (<i>I</i> _o)]	0.075	0.064	0.055	0.041
<i>wR</i> ₂ (all data)	0.078	0.067	0.057	0.042

CONTENTS ENTRY

**Hexahalometallate salts of trivalent scandium, yttrium and lanthanum:
cation–anion association in the solid state and in solution**

Martin J. D. Champion, William Levason, David Pugh and Gillian Reid

School of Chemistry, University of Southampton, Southampton SO17 1BJ, UK



The hydrogen bonding of hexahalometallate anions with various organic cations is explored via crystallographic and solution NMR studies.

References

1. P. N. Bartlett, S. L. Benjamin, C. H. de Groot, A. L. Hector, R. Huang, A. Jolleys, G. P. Kissling, W. Levason, S. J. Pearce, G. Reid and Y. Wang, *Mater. Horiz.*, 2015, **2**, 420; P. N. Bartlett, D. A. Cook, C. H. de Groot, A. L. Hector, R. Huang, A. Jolleys, G. P. Kissling, W. Levason, S. J. Pearce and G. Reid, *RSC Advances*, 2013, **3**, 15645.
2. J. Ke, W. Su, S. M. Howdle, M. W. George, D. Cook, M. Perdjon-Abel, P. N. Bartlett, W. Zhang, F. Cheng, W. Levason, G. Reid, J. Hyde, J. Wilson, D. C. Smith, K. Mallik and P. Sazio, *Proc. Natl. Acad. Sci., USA* 2009, **106**, 14768; J. Ke, P. N. Bartlett, D. Cook, T. L. Easun, M. W. George, W. Levason, G. Reid, D. Smith, W. Su and W. Zhang, *Phys. Chem. Chem. Phys.* 2012, **14**, 1517; P. N. Bartlett, J. Burt, D. A. Cook, C. Y. Cummings, M. W. George, A. L. Hector, M. M. Hasan, J. Ke, W. Levason, D. Pugh, G. Reid, P. W. Richardson, D. C. Smith, J. Spencer, N. Suleiman and W. Zhang, *Chem. Eur. J.*, 2016, **22**, 302; C. Cummings, P. N. Bartlett, D. Pugh, G. Reid, W. Levason, M. M. Hasan, A. L. Hector, J. Spencer and D. C. Smith, *J. Electrochem. Soc.*, 2014, **162**, D619.
3. P. N. Bartlett, C. Y. Cummings, W. Levason, D. Pugh and G. Reid, *Chem. Eur. J.*, 2014, **20**, 5019.
4. C. Y. Cummings, P. N. Bartlett, D. Pugh, G. Reid, W. Levason, M. M. Hasan, A. L. Hector, J. Spencer, D. C. Smith, S. Marks and R. Beanland, DOI: 10.1002/celc.201500539, in press.
5. H. Ohno, *Electrochemical Aspects of Ionic Liquids*, Wiley, New York, 2005; *Electrodeposition from Ionic Liquids*, F. Endres, A. P. Abbot and D. R. Macfarlane (Eds.), Wiley, New York, 2008.
6. K. Matsumota, T. Tsuda, T. Nohira, R. Hagiwara, Y. Ito and O. Tamada, *Acta Cryst.*, 2002, **C58**, m186.
7. C. C. Hines, D. B. Cordes, S. T. Griffin, S. I. Watts, V. A. Cocalia and R. D. Rogers, *New. J. Chem.*, 2008, **32**, 872.
8. A. Babai and A.-V. Mudring, *Inorg. Chem.*, 2006, **45**, 4874.
9. A. Babai and A.-V. Mudring, *Inorg. Chem.*, 2005, **44**, 8168.
10. K. Pohako-Esko, T. Wehner, P. S. Schulz, F. W. Heinemann, K. Muller-Buschbaum and P. Wasserscheid, *Eur. J. Inorg. Chem.*, 2016, 1333.
11. J. Hallfeldt, PhD. Thesis, 2003, University of Hannover.
12. A. Babai, A.-V. Mudring, *J. Alloys Compd.*, 2006, **418**, 122.
13. G. A. Jeffrey, *An Introduction to Hydrogen Bonding*, Oxford University Press, Oxford, 1997; (b) T. Steiner, *Angew. Chem., Int. Ed.*, 2002, **41**, 48.

-
14. R. D. Shannon, *Acta Cryst. Sect. A*, 1976, **32**, 751.
 15. Rigaku Corporation, Tokyo, Japan, 2011
 16. L. J. Farrugia, *J. Appl. Crystallogr.*, 2012, **45**, 849.
 17. C. F. Macrae, P. R. Edgington, P. McCabe, E. Pidcock, G. P. Shields, R. Taylor, M. Towler and J. van de Streek, *J. Appl. Cryst.* 2006, **39**, 453.
 18. F. H. Allen, O. Johnson, G. P. Shields, B. R. Smith and M. Towler, *J. Appl. Cryst.* 2004, **37**, 335.