

RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Toward anticancer gold-based compounds targeting PARP-1: a new case study

Received 00th January 20xx,
Accepted 00th January 20xx

A. Citta,^{a,°} V. Scalcon,^{a,°} P. Göbel,^b B. Bertrand,^c M. Wenzel,^c A. Folda,^a M.P. Rigobello,^{a,*} E. Meggers^b and A. Casini^{c,d,*}

DOI: 10.1039/x0xx00000x

www.rsc.org/

A new gold(III) complex bearing a 2-((2,2'-bipyridin)-5-yl)-1H-benzimidazol-4-carboxamide ligand has been synthesized and characterized for its biological properties *in vitro*. In addition to showing promising antiproliferative effects against human cancer cells, the compound potently and selectively inhibits the zinc finger protein PARP-1, with respect to the seleno-enzyme thioredoxin reductase. The results hold promise for the design of novel gold-based anticancer agents disrupting PARP-1 function and to be used in combination therapies.

Gold compounds have recently gained increasing attention in the design of new metal-based anticancer therapeutics,^{1,4} including gold(III) complexes with multidentate N-donor or cyclometalating ligands, gold(III) dithiocarbamates, gold(I) *N*-heterocyclic (NHC) carbenes, as well as gold(I) alkynyl complexes.^{1, 5-7}

Concerning the possible mechanisms of action, early work suggested DNA as the anticancer target for gold complexes. However, later studies showed that actually the inhibition properties of different proteins and enzymes by gold compounds play major roles,^{8, 9} whereas interactions with nucleic acids appear to be markedly less relevant, with a few exceptions.^{10, 11} For example, thiol-containing enzymes such as glutathione reductase (GR), glutathione-S-transferase,¹² cysteine proteases,¹³ protein tyrosine phosphatases (PTP), and deubiquitinases (DUBs)¹⁴ were shown to be potently inhibited by gold complexes. Interestingly, recently the water and

glycerol membrane channels termed aquaporins (AQPs) have also been reported to be selectively targeted by certain families of gold(III) complexes,¹⁵⁻¹⁷ which could also be used to unravel the roles of AQPs in cancer cell proliferation.¹⁸ In this context, among the most studied and recognized targets for gold compounds, the seleno-enzyme thioredoxin reductase (TrxR) has been widely investigated.¹⁹ Human TrxR contains a cysteine-selenocysteine redox pair at the C-terminal active site, and the solvent-accessible selenolate group, arising from enzymatic reduction, constitutes a likely target for “soft” metal ions such as gold. Thus, a number of mono and dinuclear, as well as heteronuclear, gold(I) and gold(III) complexes have shown good correlation between cytotoxic activity and TrxR inhibition properties.²⁰⁻²⁹ In addition, mitochondria and the endoplasmic reticulum³⁰ have been proposed as potential targets for anticancer gold complexes.

Pursuing the search of novel protein targets for anticancer gold compounds, some of us reported on the inhibitory effects of different cytotoxic gold-based complexes with phosphine or bipyridyl ligands, towards the zinc finger (ZF) enzyme poly (adenosine diphosphate (ADP)-ribose) polymerase 1 (PARP-1).^{31, 32} Interestingly, Au(III) coordination complexes were among the most efficient in inhibiting PARP-1, at a nM level, followed by Au(I) compounds.

It is worth mentioning that PARPs are considered “the guardian angels” of DNA playing a key role in its repair by detecting DNA strand breaks and catalyzing poly (ADP-ribosylation).³³ Therefore, PARP inhibitors can be used in combination with conventional anticancer agents that act by damaging DNA, such as cytotoxic chemotherapy and radiotherapy, as the PARP inhibitors block the DNA-repair mechanisms that cancer cells use to resist destruction.³⁴

Concerning the molecular mechanisms of PARP-1 inhibition by metal complexes, gold ions in either oxidation state 3+ or 1+ are able to induce zinc substitution in ZF models, leading to the formation of the so-called *gold fingers*.^{31, 35, 36} Damage of the ZF domain responsible for DNA recognition leads to PARP-1 inhibition.

^a Department of Biomedical Sciences, University of Padova, via Ugo Bassi 58/b, 35131 Padova, Italy.

^b Fachbereich Chemie, Philipps-Universität Marburg, Hans-Meerwein-Strasse 4, 35043 Marburg, Germany.

^c Dept. of Pharmacokinetics, Toxicology and Targeting, Research Institute of Pharmacy, University of Groningen, Antonius Deusinglaan 1, 9713 AV Groningen, The Netherlands.

^d Cardiff School of Chemistry, Cardiff University, Main Building, Park place, Cardiff CF10 3A, United Kingdom.

[°] Valeria Scalcon and Anna Citta: these authors contributed equally

* Angela Casini and Maria Pia Rigobello: co-corresponding authors

† Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

Here, we report on the synthesis of a new gold(III) complex (**2**) bearing the bidentate N-donor ligand (**1**). Notably, the 1H-benzimidazole-4-carboxamide fragment of **1** has been designed as PARP-1 inhibitor acting on the catalytic site of the protein, and not on its ZF DNA binding domain by forming hydrogen bonds between the carboxamide and Ser904 as well as Gly863 within the catalytic site.³⁷ Ideally, the resulting gold complex should show enhanced properties as PARP-1 targeted agent profiting of the synergic inhibitory effects of both the Au(III) ions and the organic ligand. Thus, **1** and **2** were tested for their PARP-1 inhibition properties *in vitro* directly against the purified enzyme as well as in protein extracts from human cancer cells, against which the compounds also showed antiproliferative properties. Gold finger formation was observed by high-resolution ESI MS upon treatment of the PARP-1 zinc finger model with **2**.


In this context, where multiple protein targets have been identified for cytotoxic gold compounds, it is absolutely necessary to promptly assess selectivity of new families of complexes in order to avoid side-effects, and to construct solid and reliable structure-activity relationships which should orient the design of targeted chemotherapeutic agents. Therefore, the activity of the compounds as TrxR inhibitors was also tested on both purified enzyme and cell extracts, in comparison to auranofin, the gold(I) anti-arthritic drug with cytotoxic properties *in vitro*,³⁸ used here as the benchmark inhibitor of TrxR.³⁹

To further characterize the mechanisms of anticancer action of the gold(III) complex **2**, the study of its effects on the intracellular redox state was conducted measuring the total and oxidized glutathione content in cancer cells. Moreover, its effects on the mitochondrial membrane potential were also assessed in cancer cells, in comparison to ligand **1**. The obtained results allowed evaluating the selectivity of **2** for PARP-1 vs TrxR, with implications for the design of improved gold-based targeted agents.

Results and Discussion

A practical synthesis of ligand **1** was developed starting from 4-bromo-2,2'-bipyridine through reductive carbonylation⁴⁰ providing 2,2'-bipyridine-4-carbaldehyde⁴¹ in 96% yield, followed by condensation with 2,3-diaminobenzamide⁴² in 93% yield. Compound **2** was then synthesized adapting procedures used for previously reported Au(III) complexes with bidentate N-donor ligands⁴³ (Scheme 1) and characterized via different methods as described in the Experimental section (see Supplementary Information Available). Thus, 2-((2,2'-bipyridin-5-yl)-1H-benzimidazol-4-carboxamide (50 mg, 0.16 mmol) in suspension in ethanol (0.5 mL) was reacted with hydrogen tetrachloroaurate (1 eq, 54 mg, 0.16 mmol), also dissolved in ethanol (0.5 mL), in a round-bottom flask equipped with a condenser. The reaction mixture was refluxed overnight during which time a brown precipitate was formed. After cooling down, the precipitate was collected by filtration and washed twice with diethylether (68% yield). The product was characterized by various techniques including ¹H and ¹³C

NMR spectroscopy, mass spectrometry and elemental analysis (see Supplementary Information for details).



Scheme 1 – Synthesis of ligand **1** and of the related Au(III) complex **2**.

Initially, the stability of the gold(III) complex **2** was evaluated in PBS buffer (pH 7.4) using UV-visible spectrophotometry. The compound exhibits an intense transition in the 300-400 nm range, characteristic of the gold(III) chromophore, that may be straightforwardly assigned as LMCT bands (Fig. S1, supplementary information). Spectral changes are slowly observed with time that might be related to the occurrence of partial hydrolysis processes. In any case, the gold(III) complex is the dominant species in buffered aqueous solutions after several hours incubation.

The stability of **2** toward biologically occurring reducing agent glutathione (GSH) was also evaluated. Results show that GSH, present at a 2:1 molar ratio with respect to **2**, does not markedly affect the evolution of the main LMCT band of the complex with respect to its normal hydrolysis (Fig. S2). However, formation of soluble gold(I) thiolate species as a major product of gold(III) reduction, cannot be excluded.

Afterwards, the antiproliferative properties of the new gold complex **2** and ligand **1** were studied by monitoring their ability to inhibit cell growth using the MTT assay (see Experimental section). Cytotoxic activity of the compounds was determined after exposing for 72 h the human ovarian cancer A2780 cell line, and its cisplatin resistant variant (A2780cisR), the human ovarian cancer SKOV3 cell line, as well as the human non-small cell lung carcinoma A549 line, in comparison to cisplatin and auranofin (AF). The results are summarized in Table 1. The IC₅₀ values of **2** towards all tested cell lines are lower in comparison to the free ligand **1**. This may implicate that the gold(III) center plays an important role in the still unknown mechanism(s) of cytotoxic action. The IC₅₀ values towards the cisplatin resistant A2780cisR cell line is for **1** comparable to cisplatin, but **2** is markedly more effective. This observation support the idea that in general gold(III) complexes do not have the same mechanism of action as cisplatin, as discussed in the introduction. Both **1** and **2** are poorly toxic against the A549 cell. The greatest difference in IC₅₀ value between **1** and **2** is found against the SKOV3 cell line (**2** is ca. 4-fold more potent than **1**). Such discrepancy in the cytotoxic effects may be due to several factors, including different transport mechanisms (uptake and efflux) of complex **2** with respect to ligand **1** in the selected cancer cells, which may lead to decreased intracellular accumulation of **1**. Finally,

AF is certainly the most potent among the tested drugs; however, it is also the most unselective, again demonstrating differences of mechanisms of activity among different families of gold compounds.

Table 1 - IC₅₀ values of the Au complexes described in this study against human ovarian carcinoma cell lines SKOV3, cisplatin sensitive (A2780) and resistant (A2780cisR) and lung cancer cells (A549) compared to cisplatin and auranofin (AF).

Compound	IC ₅₀ (μM) ^a			
	SKOV3	A2780	A2780cisR	A549
1	84.4 ± 7.6	9.70 ± 3.06	33.1 ± 5.9	46.7 ± 17.5
2	22.7 ± 2.9	4.80 ± 2.35	13.0 ± 2.7	35.0 ± 6.5
AF	1.8 ± 0.4	1.25 ± 0.5	1.5 ± 0.3	2.5 ± 0.7
cisplatin	13.2 ± 3.5	5.2 ± 1.9	35.0 ± 5.9	10.8 ± 2.8

^a Data are the mean ± SD of at least four experiments.

Compounds **1-2** were then tested against purified PARP-1 using an established protocol.³¹ As expected, potent PARP-1 inhibition was observed with both compounds: **1** has an IC₅₀ = 5.0 ± 2.1 nM, and **2** has IC₅₀ = 6.0 ± 1.3 nM, in the same range as previously reported cytotoxic gold(III) complexes.³¹

Afterwards, PARP-1 activity was evaluated on cell extracts from A2780, A2780cisR and SKOV3 cells. Thus, incubation of protein cell extracts with the compounds for 24 h at room temperature was followed by PARP-1 activity determination. Fig. 1 shows the residual PARP-1 activity in protein extracts treated with the complexes at a fixed concentration (10 μM).

Attractively, both compounds can induce PARP-1 inhibition to a similar extent in A2780 and A2780cisR cell lines, while a marked difference could be detected in the case of SKOV3 cells, where **2** is able to inhibit PARP-1 until ca. 10% of its residual activity. Instead, **1** is practically ineffective on these cells, in line with the scarce anticancer effects observed above. Furthermore, PARP-1 activity was evaluated on protein extracts obtained from SKOV3 cells pre-treated with non-cytotoxic doses of each compound for 48 hours. Afterwards, the protein extracts were collected and analyzed for PARP-1 activity. Preliminary results indicate that only the gold complex **2** (20 μM) was able to induce ca. 70% reduction of PARP-1 activity, while ligand **1** was poorly effective.




Figure 1 - PARP-1 activity levels in human ovarian cancer cellular extracts. PARP-1 activity was measured in homogenates (50 μg of protein) treated with the compounds

(10 μM) over 24 h at room temperature. Data are the mean ± SD of at least three experiments each performed in triplicate.

In order to assess formation of adducts between the gold complex and the zinc finger domain of PARP-1, a peptide model corresponding to the N-terminal ZF domain sequence of PARP-1 was reacted with **2** and the sample was monitored by high-resolution ESI MS as described in the Experimental section. Figure S3 in the supplementary material shows the broadband mass spectrum of the **2**-ZF adduct. In agreement with previously reported studies on other Au(III) complexes, when **2** was incubated with the ZF domain in a 3 : 1 ratio for 10 min, partial displacement of Zn²⁺ from the ZF by gold ions leads already to formation of the so-called “gold-finger” adduct.

Afterwards, to evaluate if the different cytotoxic effects of **2** were related to differences in intracellular Au accumulation, ICP-MS analysis of cell extracts out of A2780 and SKOV3 cells, pre-treated with the gold compound for 24 h, demonstrated that the cytotoxicity is somehow proportional to the gold uptake, and the strongest antiproliferative effects correspond to higher values of intracellular gold concentration. In fact, the concentration of Au [pmol Au/10⁶ cells] measured in A2780 and SKOV3 cells is 1802 ± 209 and 1087 ± 322, respectively. Nevertheless, in spite the reduced accumulation of **2** in SKOV3 cells, the inhibition of PARP-1 activity is more pronounced than in the case of A2780 cells (Fig.1).

Since TrxR is also a potential target for gold complexes, *in vitro* inhibition of purified rat TrxR by the two compounds was studied using established protocols as described in the Experimental section. The results are summarized in Table 2 and Figure S4. Complex **2** inhibits cytosolic thioredoxin reductases (TrxR1) in the same range as auranofin (IC₅₀ = 14.32 ± 1.62 nM vs IC₅₀ = 6.88 ± 1.25 nM, respectively). Conversely, ligand **1** is completely ineffective, as expected since it is deprived of the Au(III) centre able to bind the selenol groups (Figure S1, supplementary information). Further studies demonstrated that **2** is also able to inhibit the TrxR closely related, but selenium-free, enzyme glutathione reductase (GR) with IC₅₀ = 0.40 ± 0.06 μM, about 28-fold less efficiently than in the case of TrxR (Table 2).

Afterwards, the effect of compounds on TrxR and GR activities was evaluated in cell lysates. For this purpose, SKOV3 cells where the two compounds showed markedly different cytotoxic effects, were pre-treated for 48 h with 20 and 40 μM of **1** and **2**, respectively. The obtained results show that **1** does not affect enzymes activities, while **2**, causes ca. 50% TrxR inhibition and a slight decrease of GR activity at 40 μM (Fig. 2). In addition, similar experiments were conducted in the A2780 cells, and the obtained results showed no statistically significant inhibition of TrxR at the tested compounds' concentrations (Figure S5). Notably, these latter results further corroborate the hypothesis of alternative pharmacological targets for the reported compounds.

Overall, the obtained results on both PARP-1 and TrxR activities, indicate that the cytotoxic gold complex **2** may operate via inhibition of PARP-1, whereas TrxR is only moderately affected. Concerning the observed differences in PARP-1 inhibition by **1** and **2**, it may be suggested that the ligand is not so selective for binding to PARP-1 as the gold complex **2**, once in the presence of other intracellular components. Nevertheless, in terms of the overall cytotoxic potency, differences in the uptake mechanisms and cellular accumulation between the two compounds should also be taken into account.

Table 2 IC₅₀ values of the inhibition of TrxR1 and GR on the isolated enzymes.

Complexes	IC ₅₀ (nM)	
	TrxR1	GR
1	>100	>10000
2	14.32 ± 1.62	400 ± 60
AF	6.88 ± 1.25	>10000

The glutathione redox pair (GSH/GSSG) is another fundamental component of the cell redox regulation in cisplatin resistant cells.⁴⁴ Therefore, our study continued with the analysis of total glutathione content (reduced + oxidized) and of the GSH/GSSG ratio in SKOV3, after treatment with the two compounds for 48 h in comparison to AF. The obtained results are shown in Figure S6 in the Supplementary material available. It can be observed that for all tested compounds no statistically significant variation of the total GSH content, as well as of the GSH/GSSG ratio occur, again made exception for **2**, which causes a slight increase of GSSG content at 40 μM, in accordance with the compound's above-mentioned inhibition effect of glutathione reductase. This behavior suggests that GSH does not particularly influence the cytotoxic potency of the gold complex, as for cisplatin in the case of certain resistant cancer cells.




Figure 2 - Effect of the compounds **1** and **2** on thioredoxin reductase and glutathione reductase activities in cell lysates. SKOV3 cells were treated for 48 h with 20 and 40 μM of **1** and **2**, respectively. * = p < 0.01

Mitochondrial membrane potential (MMP), a consequence of the electrochemical proton gradient maintained for the purpose of ATP synthesis, is an important indicator of functional mitochondria. Previously reported studies showed that gold(III) complexes are able to determine the decrease of MMP depending on the ligands. As an example, gold(III) Porphyrin **1a** induced apoptosis by mitochondrial death pathways related to reactive oxygen species.⁴⁵ Similarly, gold(III)-dithiocarbamate derivatives were shown to alter mitochondrial parameters, such as causing a drop of the mitochondrial membrane potential (MMP).⁴⁶ MMP evaluation was conducted monitoring the fluorescence of tetramethylrhodamine methyl esters (TMRM) according to established protocols (see Experimental for details). Thus, it was possible to determine if the complexes are able to induce a quick drop in mitochondrial membrane potential ($\Delta\psi_m$), detectable as a decline in the fluorescence intensity of TMRM.⁴⁷

Therefore, MMP of SKOV3 cells treated for 18 h with compounds **1** and **2** was measured by cytofluorometric analysis in comparison to auranofin (AF) and CCCP (Fig. 3). Cells were incubated with 25 nM TMRM for 20 min and then analyzed by flow cytometry utilizing an argon laser at 585 nm, as described in the Experimental section.




Figure 3 - Percentage of cells with a low and high mitochondrial membrane potential. Mitochondrial membrane potential of SKOV3 cells treated for 18 h with compounds **1** and **2** was measured by a cytofluorometric analysis. Cells were incubated with 25 nM TMRM for 20 min and then analyzed by flow cytometry utilizing an argon laser at 585 nm. Bars represent mean percentages \pm S.D. (n=3) of SKOV3 cells whose mitochondria maintained a high (gray) or low (dark gray) TMRM fluorescence, that correspond to their mitochondrial membrane potential.

From the obtained results it has been possible to determine that, at variance with AF and the classical uncoupling agent CCCP (Carbonyl cyanide m-chlorophenyl hydrazone), both complexes **1** and **2** do not affect the MMP values with respect to the controls, as it has instead been reported for gold(III) porphyrins and dithiocarbamate complexes.

Finally, since overexpression of PARP in cancer cells has been linked to drug resistance and PARP-1 inhibition has been shown to sensitize tumor cells to chemotherapeutic agents including platinum compounds, we decided to evaluate the cytotoxic effect of cisplatin administered in combination with different concentrations of **2**. Initial data were obtained for 72 h co-administration of cisplatin (7.5 μ M) and **2** at different concentrations (10-20-30 μ M) in SKOV3 cells (see experimental for details). In Table S1 (supplementary material) a comparison of the predicted survival rates (defined as the expected cell viability if the combined activities of the compounds are additive) and the experimentally determined values (the observed viabilities) is reported. Unfortunately, the observed survival rates for the combinations of **2** with cisplatin are similar to those predicted on the basis of an additive effect, ruling out the synergism. Further studies will be necessary to investigate possible synergic effects in different cancer cell types and to validate the possibility of using **2** in combination therapy.

Conclusions

We have reported here on the potent PARP-1 inhibition properties of a new cytotoxic gold(III) complex with a bidentate N-donor ligand. A series of biological and biochemical assays has shown that the compound targets preferentially PARP-1 with respect to the seleno-enzyme thioredoxin reductase, and in doing so it is more effective than the free ligand. The absence of effects of the gold(III)

compound on both MMP and intracellular glutathione redox state demonstrate that different mechanisms of action are in place for different families of gold-based cytotoxic agents, which holds promise for the design of targeted anticancer metallodrugs.

Notably, inhibition of PARP potentiates the activity of DNA-damaging agents, such as alkylators, platinum compounds, topoisomerase inhibitors, and radiation in *in vitro* and *in vivo* models. Thus, clinical development to date has focused on PARP inhibitors potential role in combination with DNA-damaging chemotherapy, where efficacy has been limited by enhanced normal tissue toxicity.

Olaparib, a highly potent PARP inhibitor, has recently been approved for ovarian cancer therapy by the FDA and European commission, in patients with platinum-sensitive, recurrent, high-grade serous ovarian cancer with BRCA1 or BRCA2 mutations.⁴⁸

Within this frame, gold(III) complexes such as **2** may constitute an alternative strategy to PARP-1 inhibition, acting on both the zinc finger DNA binding domain of the protein via gold binding, and on its catalytic domain; therefore, having enhanced efficacy. Further studies are necessary to fully validate this hypothesis and to design compounds with selectivity for PARP-1 with respect to other zinc finger proteins.

Acknowledgements

Authors thank EU COST Actions CM1105 for funding and for providing opportunities for discussion; M.P.R. acknowledges CPDA130272 granted by the University of Padova (Italy) and Consorzio Interuniversitario di Ricerca in Chimica dei Metalli nei Sistemi Biologici (CIRCSMB). E.M. thanks the German Research Foundation for financial support of this work (ME 1805/9-1).

Notes and references

‡ Footnotes relating to the main text should appear here. These might include comments relevant to but not central to the matter under discussion, limited experimental and spectral data, and crystallographic data.

§

§§

etc.

1. B. Bertrand and A. Casini, *Dalton Trans*, 2014, 43, 4209-4219.
2. L. Messori and A. Casini, *Curr Top Med Chem*, 2011, 11, 2647-2660.
3. T. Zou, C. T. Lum, C.-N. Lok, J.-J. Zhang and C.-M. Che, *Chem. Soc. Rev.*, 2015, 44, 8786-8801.
4. M. A. Cinellu, I. Ott and A. Casini, in *Bioorganometallic Chemistry*, Wiley-VCH Verlag GmbH & Co. KGaA, 2014, DOI: 10.1002/9783527673438.ch04, pp. 117-140.
5. E. M. Nagy, L. Ronconi, C. Nardon and D. Fregona, *Mini-Rev Med Chem*, 2012, 12, 1216-1229.
6. L. Oehninger, R. Rubbiani and I. Ott, *Dalton Trans.*, 2013, 42, 3269-3284.
7. A. Meyer, C. P. Bagowski, M. Kokoschka, M. Stefanopoulou, H. Alborzinia, S. Can, D. H. Vlecken, W. S. Sheldrick, S. Wolf and I. Ott, *Angew Chem Int Ed Engl*, 2012, 51, 8895-8899.

8. A. de Almeida, B. L. Oliveira, J. D. G. Correia, G. Soveral and A. Casini, *Coord. Chem. Rev.*, 2013, 257, 2689-2704.
9. A. Casini, G. Kelter, C. Gabbiani, M. A. Cinellu, G. Minghetti, D. Fregona, H. H. Fiebig and L. Messori, *J. Biol. Inorg. Chem.*, 2009, 14, 1139-1149.
10. B. Bertrand, L. Stefan, M. Pirrotta, D. Monchaud, E. Bodio, P. Richard, P. Le Gendre, E. Warmerdam, M. H. de Jager, G. M. Groothuis, M. Picquet and A. Casini, *Inorg Chem*, 2014, 53, 2296-2303.
11. C. Bazzicalupi, M. Ferraroni, F. Papi, L. Massai, B. Bertrand, L. Messori, P. Gratteri and A. Casini, *Angew Chem Int Ed Engl*, 2016, 55, 4256-4259.
12. A. De Luca, C. G. Hartinger, P. J. Dyson, M. Lo Bello and A. Casini, *J. Inorg. Biochem.*, 2013, 119, 38-42.
13. S. P. Fricker, *Metallomics*, 2010, 2, 366-377.
14. J.-J. Zhang, K.-M. Ng, C.-N. Lok, R. W.-Y. Sun and C.-M. Che, *Chem. Comm.*, 2013, 49, 5153-5155.
15. A. P. Martins, A. Marrone, A. Ciancetta, A. Galan Cobo, M. Echevarria, T. F. Moura, N. Re, A. Casini and G. Soveral, *Plos One*, 2012, 7, e37435.
16. A. P. Martins, A. Ciancetta, A. de Almeida, A. Marrone, N. Re, G. Soveral and A. Casini, *ChemMedChem*, 2013, 8, 1086-1092.
17. A. Casini and A. de Almeida, in *Aquaporins in Health and Disease*, CRC Press, 2016, doi:10.1201/b19017-19 pp. 297-318.
18. A. Serna, A. Galan-Cobo, C. Rodrigues, I. Sanchez-Gomar, J. J. Toledo-Aral, T. F. Moura, A. Casini, G. Soveral and M. Echevarria, *Journal of Cellular Physiology*, 2014, 229, 1787-1801.
19. A. Bindoli, M. P. Rigobello, G. Scutari, C. Gabbiani, A. Casini and L. Messori, *Coord. Chem. Rev.*, 2009, 253, 1692-1707.
20. E. Vergara, A. Casini, F. Sorrentino, O. Zava, E. Cerrada, M. P. Rigobello, A. Bindoli, M. Laguna and P. J. Dyson, *ChemMedChem*, 2010, 5, 96-102.
21. J. L. Hickey, R. A. Ruhayel, P. J. Barnard, M. V. Baker, S. J. Berners-Price and A. Filipovska, *J. Am. Chem. Soc.*, 2008, 130, 12570-12571.
22. M. P. Rigobello, L. Messori, G. Marcon, M. A. Cinellu, M. Bragadin, A. Folda, G. Scutari and A. Bindoli, *J. Inorg. Biochem.*, 2004, 98, 1634-1641.
23. C. Gabbiani, G. Mastrobuoni, F. Sorrentino, B. Dani, M. P. Rigobello, A. Bindoli, M. A. Cinellu, G. Pieraccini, L. Messori and A. Casini, *MedChemComm*, 2011, 2, 50-54.
24. R. Rubbiani, L. Salassa, A. de Almeida, A. Casini and I. Ott, *ChemMedChem*, 2014, 9, 1205-1210.
25. E. Schuh, C. Pfluger, A. Citta, A. Folda, M. P. Rigobello, A. Bindoli, A. Casini and F. Mohr, *J. Med. Chem.*, 2012, 55, 5518-5528.
26. A. Citta, E. Schuh, F. Mohr, A. Folda, M. L. Massimino, A. Bindoli, A. Casini and M. P. Rigobello, *Metallomics*, 2013, DOI: 10.1039/c3mt20260g, 1006-1015.
27. B. Bertrand, A. Citta, I. L. Franken, M. Picquet, A. Folda, V. Scalcon, M. P. Rigobello, P. Le Gendre, A. Casini and E. Bodio, *JBIC*, 2015, 20, 1005-1020.
28. N. Lease, V. Vasilevski, M. Carreira, A. de Almeida, M. Sanaú, P. Hirva, A. Casini and M. Contel, *J. Med. Chem.*, 2013, 56, 5806-5818.
29. K. Yan, C. N. Lok, K. Bierla and C. M. Che, *Chem Commun.*, 2010, 46, 7691-7693.
30. J. J. Zhang, R. W. Y. Sun and C. M. Che, *Chem. Comm.*, 2012, 48, 3388-3390.
31. F. Mendes, M. Groessl, A. A. Nazarov, Y. O. Tsybin, G. Sava, I. Santos, P. J. Dyson and A. Casini, *J. Med. Chem.*, 2011, 54, 2196-2206.
32. M. Serratrice, F. Edafe, F. Mendes, R. Scopelliti, S. M. Zakeeruddin, M. Gratzel, I. Santos, M. A. Cinellu and A. Casini, *Dalton Trans.*, 2012, 41, 3287-3293.
33. V. Schreiber, F. Dantzer, J. C. Ame and G. de Murcia, *Nat Rev Mol Cell Bio*, 2006, 7, 517-528.
34. K. Ratnam and J. A. Low, *Clin Cancer Res*, 2007, 13, 1383-1388.
35. U. A. Laskay, C. Garino, Y. O. Tsybin, L. Salassa and A. Casini, *Chem Commun (Camb)*, 2015, 51, 1612-1615.
36. A. Jacques, C. Lebrun, A. Casini, I. Kieffer, O. Proux, J. M. Latour and O. Seneque, *Inorg Chem*, 2015, 54, 4104-4113.
37. T. D. Penning, G.-D. Zhu, J. Gong, S. Thomas, V. B. Gandhi, X. Liu, Y. Shi, V. Klinghofer, E. F. Johnson, C. H. Park, E. H. Fry, C. K. Donawho, D. J. Frost, F. G. Buchanan, G. T. Bukofzer, L. E. Rodriguez, V. Bontcheva-Diaz, J. J. Bouska, D. J. Osterling, A. M. Olson, K. C. Marsh, Y. Luo and V. L. Giranda, *J. Med. Chem.*, 2010, 53, 3142-3153.
38. T. M. Simon, D. H. Kunishima, G. J. Vibert and A. Lorber, *Cancer Res*, 1981, 41, 94-97.
39. M. P. Rigobello, G. Scutari, A. Folda and A. Bindoli, *Biochem. Pharmacol.*, 2004, 67, 689-696.
40. L. Ashfield and C. F. J. Barnard, *Org Process Res Dev*, 2007, 11, 39-43.
41. J. Polin, E. Schmohel and V. Balzani, *Synthesis*, 1998, 1998, 321-324.
42. K. L. Arienti, A. Brunmark, F. U. Axe, K. McClure, A. Lee, J. Blevitt, D. K. Neff, L. Huang, S. Crawford, C. R. Pandit, L. Karlsson and J. G. Breitenbucher, *J. Med. Chem.*, 2005, 48, 1873-1885.
43. A. Casini, M. C. Diawara, R. Scopelliti, S. M. Zakeeruddin, M. Gratzel and P. J. Dyson, *Dalton Trans.*, 2010, 39, 2239-2245.
44. S. Okuno, H. Sato, K. Kuriyama-Matsumura, M. Tamba, H. Wang, S. Sohda, H. Hamada, H. Yoshikawa, T. Kondo and S. Bannai, *British journal of cancer*, 2003, 88, 951-956.
45. Y. Wang, He, Q.Y., Sun, R.W., Che, C.M., Chiu, J.F., *Cancer Res*, 2005, 65, 11553-11564.
46. D. Saggioro, M. P. Rigobello, L. Paloschi, A. Folda, S. A. Moggach, S. Parsons, L. Ronconi, D. Fregona and A. Bindoli, *Chem. Biol.*, 2007, 14, 1128-1139.
47. A. Folda, V. Scalcon, M. Ghazzali, M. H. Jaafar, R. A. Khan, A. Casini, A. Citta, A. Bindoli, M. P. Rigobello, K. Al-Farhan, A. Alsalmeh and J. Reedijk, *J Inorg Biochem*, 2015, 153, 346-354.
48. C. J. Lord, A. N. Tutt and A. Ashworth, *Annu Rev Med*, 2015, 66, 455-470.