

Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/chemicalscience

β -Glutamine-mediated Self-Association of Transmembrane β -Peptides within Lipid Bilayers

 U. Rost^a, C. Steinem^{*a} and U. Diederichsen^{*a}

 Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Transmembrane β -peptide helices and their association in lipid membranes are still widely unexplored. We designed and synthesized transmembrane β -peptides harboring different numbers of D- β^3 -glutamine residues (^hGln) by solid phase peptide synthesis. By means of circular dichroism spectroscopic measurements, the secondary structure of the β -peptides reconstituted into unilamellar vesicles was determined to be similar to a right-handed 3_{14} -helix. Fluorescence spectroscopy using D- β^3 -tryptophan residues strongly suggested a transmembrane orientation. Two respectively three ^hGln served as recognition units between the helices to allow helix-helix assembly driven by hydrogen bond formation. The association state of the transmembrane β -peptides as a function of the number of ^hGln residues was investigated by fluorescence resonance energy transfer (FRET). Therefore, two fluorescence probes (NBD, TAMRA) were covalently attached to the side chains of the transmembrane β -peptide helices. The results clearly demonstrate that only β -peptides with ^hGln as recognition units assemble into oligomers, presumably trimers. Temperature dependent FRET experiments further show that the strength of the helix-helix association is a function of the number of ^hGln residues in the helix.

Introduction

In the last three decades, there has been great interest in the design and synthesis of unnatural β -peptide oligomers in solutions, which are capable of forming stable secondary structures and which are resistant to proteolytic degradation by proteases and peptidases.^{1,2} Folding into well-defined secondary structures is a fundamental part of protein architecture and takes part in the recognition process between peptides.³ Even though integral membrane proteins represent nearly 30% of the human proteome and are involved in various cellular processes, the complexity of these protein structures often limits the precise analysis of their biological activities, functions and mechanisms on the molecular level.^{4–6} To accomplish their function, membrane proteins can be monomeric or they need to be assembled into oligomeric structures. Studies on the fundamental molecular aspects of membrane protein oligomerization have been mainly performed using model systems of transmembrane α -peptide helices.^{5–9} The organization and assembly of these transmembrane α -peptide helices depend on the lipid environment and the interacting molecular species themselves. These interactions are still not fully understood and synthetically readily available transmembrane peptide models are an appreciated tool to investigate peptide-lipid and peptide-peptide interactions occurring in membrane asso-

ciated processes.⁶ However, in contrast to membrane embedded α -peptide helices the field of transmembrane β -peptide helices and their association within a membrane is still widely unexplored. In contrast to α -peptides, β -peptides have one additional CH₂-group inserted in every amino acid residue, which offers an increased number of possible conformers. β -peptides are known to form stable secondary structure elements even with short chain lengths (> 6 amino acid residues) and they allow to vary the width of the helix, as well as the helical dipole moment.¹⁰ APPELLA *et al.* demonstrated that small 10-residue amphiphilic β -peptides form water soluble aggregates.^{11,12} In addition, specific β -peptide helix interactions in solution, controlled by the attachment of nucleobase pairing recognition units, have been reported.^{3,4} In the present study, we describe the rational design and synthesis of novel transmembrane β -peptides forming stable secondary structures in a membrane environment as analysed by means of circular dichroism (CD) spectroscopy. Membrane insertion was verified by fluorescence spectroscopy detecting the intrinsic tryptophan fluorescence.^{13,14} The transmembrane β -peptide helices were designed to allow for peptide oligomerization based on hydrogen bond formation within a lipid bilayer. Hydrogen bond formation is established by placing D- β^3 -glutamine (Fig. 1) at defined positions within the β -peptide in




Fig. 1 Schematic illustration of hydrogen bond formation of D- β^3 -glutamine residues.

^a Institute of Organic and Biomolecular Chemistry, Georg-August-University Goettingen, Tammannstr. 2, 37077 Goettingen, Germany. E-mail: csteinem@gwdg.de; udieder@gwdg.de

† Electronic Supplementary Information (ESI) available: Material and methods, syntheses and characterizations of the β^3 -amino acids and the β -peptides, enantiomeric purity, CD and UV spectroscopic methods, fluorescence emission spectra. See DOI: 10.1039/x0xx00000x




Fig. 2 Basic structure of the designed β -peptides ($H^h\text{Lys}_2^h\text{Trp}_2^h\text{Val}_{19}^h\text{Trp}_2^h\text{Lys}_2^h\text{NH}_2$).

analogy to asparagine, which has been shown to form hydrogen bonds and thus allows for a self-association of α -peptide helices.^{9,15} The peptide-peptide association of the synthesized β -peptides reconstituted into lipid bilayers was investigated by fluorescence resonance energy transfer (FRET), which is a powerful and widely used technique to analyse association processes.^{16,17}

Results and Discussion

Design of the Transmembrane β -Peptides

We followed the approach to synthesize β -peptides that are expected to form a 3_{14} -helix and can be inserted into lipid bilayers in a transmembrane configuration. In a 3_{14} -helix, three amino acid residues form one helix turn in which every fourth amino acid side chain is oriented on the same face of the helix.^{4,18} To span the hydrophobic core of a lipid bilayer, the β -peptides were composed of 19 D - β^3 -valines.^{19–22} Considering the peptide-lipid interactions, the lipid-water interface plays a vital role. Therefore, the β -peptides were flanked with two D - β^3 -tryptophan residues at the N - and C -terminal end of the sequence (Fig. 2). The aromatic indole moieties of these D - β^3 -tryptophans are known to arrange in the polar/apolar interface of the lipid membrane.²³ While the aromatic ring is preferentially localized in the apolar part of the membrane, the indole NH groups are adjacent to the lipid carbonyl moiety, where it forms hydrogen bonds via the NH-group to anchor and stabilize the peptide in the lipid membrane.^{24–26} Furthermore, two D - β^3 -lysine residues were attached at each side of the β -peptides to increase their solubility. It is known that lysine residues with the long and flexible side chains facilitate the solubility in aqueous systems.²³ In total, we designed the β -peptide $H^h\text{Lys}_2^h\text{Trp}_2^h\text{Val}_{19}^h\text{Trp}_2^h\text{Lys}_2^h\text{NH}_2$ as a core unit for our studies (Fig. 2). D - β^3 -Glutamine was introduced as recognition unit to control transmembrane β -peptide helix aggregation via hydrogen bond formation. We have chosen D - β^3 -glutamine, as its side chain has one additional CH_2 -group, which increases the side chains' flexibility compared to asparagine. D - β^3 -Valine residues were replaced by two respectively three D - β^3 -glutamines. Nucleobase functionalized β -peptides are known to prefer an antiparallel strand orientation in aqueous solutions.^{3,4} Hence, we anticipate that the transmembrane β -peptides also associate in an antiparallel formation and placed the respective D - β^3 -glutamine residues accordingly (Tab. 1, Fig. 3). In order to perform FRET-analysis, two different

fluorophores (NBD, TAMRA), serving as a donor-acceptor pair, were attached to the β -peptides. As free peptide termini are required for membrane insertion, and in order to avoid any conformational restrictions as a result of fluorophore labelling, the fluorophores were attached to the side chain of an additional D - β^3 -lysine residue that was added either to the C -terminal or the N -terminal end depending on the β -peptide sequence.⁶




Fig. 3 Schematic view on the antiparallel 3_{14} -helical β -peptides with three amino acids forming one turn and zero ^hGln (9/10/11, left), two ^hGln (12/13/14, middle) and three ^hGln (15/16/17, right) recognition units. The β -peptides were acetylated or labelled with NBD and TAMRA for FRET-analysis and synthesized from N - to C -terminus.

Table 1 Synthesized transmembrane β -peptides with zero, two or three D - β^3 -glutamine recognition units.

No.	Synthesized Transmembrane β -Peptides
Zero recognition units	
9	$H^h\text{Lys}(\text{acetyl})^h\text{Lys}_2^h\text{Trp}_2^h\text{Val}_{19}^h\text{Trp}_2^h\text{Lys}_2^h\text{NH}_2$
10	$H^h\text{Lys}(\text{NBD})^h\text{Lys}_2^h\text{Trp}_2^h\text{Val}_{19}^h\text{Trp}_2^h\text{Lys}_2^h\text{NH}_2$
11	$H^h\text{Lys}_2^h\text{Trp}_2^h\text{Val}_{19}^h\text{Trp}_2^h\text{Lys}_2^h\text{Lys}(\text{TAMRA})\text{-NH}_2$
Two ^hGln recognition units	
12	$H^h\text{Lys}(\text{acetyl})^h\text{Lys}_2^h\text{Trp}_2^h\text{Val}_9^h\text{Gln}^h\text{Val}_5^h\text{Gln}^h\text{Val}_3^h\text{Trp}_2^h\text{Lys}_2^h\text{NH}_2$
13	$H^h\text{Lys}(\text{NBD})^h\text{Lys}_2^h\text{Trp}_2^h\text{Val}_9^h\text{Gln}^h\text{Val}_5^h\text{Gln}^h\text{Val}_3^h\text{Trp}_2^h\text{Lys}_2^h\text{NH}_2$
14	$H^h\text{Lys}_2^h\text{Trp}_2^h\text{Val}_9^h\text{Gln}^h\text{Val}_5^h\text{Gln}^h\text{Val}_3^h\text{Trp}_2^h\text{Lys}_2^h\text{Lys}(\text{TAMRA})\text{-NH}_2$
Three ^hGln recognition units	
15	$H^h\text{Lys}(\text{acetyl})^h\text{Lys}_2^h\text{Trp}_2^h\text{Val}_9^h\text{Gln}^h\text{Val}_2^h\text{Gln}^h\text{Val}_2^h\text{Gln}^h\text{Val}_3^h\text{Trp}_2^h\text{Lys}_2^h\text{NH}_2$
16	$H^h\text{Lys}(\text{acetyl})^h\text{Lys}_2^h\text{Trp}_2^h\text{Val}_9^h\text{Gln}^h\text{Val}_2^h\text{Gln}^h\text{Val}_2^h\text{Gln}^h\text{Val}_3^h\text{Trp}_2^h\text{Lys}_2^h\text{NH}_2$
17	$H^h\text{Lys}_2^h\text{Trp}_2^h\text{Val}_9^h\text{Gln}^h\text{Val}_2^h\text{Gln}^h\text{Val}_2^h\text{Gln}^h\text{Val}_3^h\text{Trp}_2^h\text{Lys}_2^h\text{Lys}(\text{TAMRA})\text{-NH}_2$

Consequently, three different types of transmembrane β -peptides were designed for analysing β -peptide helix association: i) **9/10/11** with zero ^hGln , ii) **12/13/14** with two ^hGln as recognition units and iii) **15/16/17** three ^hGln as recognition units (Tab. 1).

CD-Spectroscopic Measurements

Structural studies on β -peptides using NMR- and circular dichroism (CD)-spectroscopy, X-ray crystallography as well as molecular dynamic simulation studies have shown that they adopt stable secondary structures in solution.^{1,27–32} In particular, β -peptides rich in β^3 -valine are well-known to form the 3_{14} -helical structure.^{21,22} Here, we used CD-spectroscopy to assess the secondary structure of the three synthesized β -peptides.^{29,33} First, CD-spectra of the β -peptides **9**, **12** and **15** were examined in 2,2,2-trifluoroethanol (TFE), which is known to stabilize secondary structure formation (peptide concentration 38 μM , 25 $^\circ\text{C}$).^{34,35} All three β -peptides show similar and very characteristic CD-spectra with a minimum at 193 nm, a zero crossing at 201 nm and a maximum at 211 nm (Electronic Supplementary Information Fig. S2). These bands are indicative of a 3_{14} -helix as reported for peptides in solutions.^{22,28,36} SEEBACH and co-workers were able to unambiguously demonstrate by a combination of CD-spectroscopy, NMR-spectroscopy, X-ray crystallography and molecular dynamic simulation studies that the characteristic CD bands are a result of a left-handed 3_{14} -helical structure in solution.^{1,29–32} As the recorded CD-spectra of the β -peptides **9**, **12** and **15** are mirrored compared to the published ones by SEEBACH and co-workers, we conclude that the β -peptides **9**, **12** and **15** fold into right-handed 3_{14} -helices. The 3_{14} -helix is the most frequently documented secondary structure of folded β -peptides. It consists of 14-membered hydrogen-bonded rings between N–H (*i*) and C=O (*i*+2) with a three-residue repeating arrangement.³⁷ The β -peptides **9**, **12** and **15** were then reconstituted in a membrane environment, i.e. in large unilamellar vesicles (LUVs) composed of DOPC at a β -peptide concentration of 38 μM and a P/L-ratio of 1/20 at 25 $^\circ\text{C}$. Again CD-spectra were recorded to determine the secondary structures (Fig. 4).




Fig. 4 CD-spectra of **9**, **12** and **15** in DOPC LUVs (peptides concentration: 38 μM , P/L-ratio = 1/20, 25 $^\circ\text{C}$).

The CD-spectra of all three β -peptides show again very similar bands with a minimum at 196 nm, a zero crossing near 199 nm and a maximum at 206 nm indicative of a right-handed 3_{14} -helix. HAMURO *et al.* have shown that the presence of micelles strongly stabilizes the 3_{14} -helical conformation of amphiphilic β -peptides and the mean residue ellipticity increases in a length-dependent manner consistent with our results.^{36,38} Even though the overall signatures of the CD-spectra confirms the right-handed 3_{14} -helix, the maxima and minima are slightly shifted compared to those spectra obtained in TFE. The slight shifts of the minima and maxima are a result of the differences in the dielectric constants of the two different solvents, TFE and DOPC lipid (TRIS[®] buffer).³⁹ These findings strongly support our notion that the three β -peptides form a stable secondary structure in membrane environment. CD-spectroscopy does not only depend on the backbone but also on the side chain conformation.⁴⁰ In helical peptides, aromatic side chains like tryptophan, show characteristic CD values in the 225 nm-region, even if they are positioned near the end of the helix.^{41–43} Indeed, in addition to the similar characteristic pattern of the right-handed 3_{14} -helix, the presented CD-spectra of the three β -peptides show a weak maximum at 229 nm due to the four D- β^3 -tryptophanes, which were attached to the β -peptides (two at each side).

It is expected that the secondary structure of the β -peptides is very stable, in particular, if embedded in a lipid bilayer. To analyse the thermal stability of the secondary structure, we measured CD-spectra of the β -peptides **12** and **15** reconstituted in DOPC-LUVs (peptide concentration: 38 μM and P/L-ratio = 1/20) at 25 $^\circ\text{C}$ and at 60 $^\circ\text{C}$ (Electronic Supplementary Information Fig. S3). The CD-spectra clearly indicate that the secondary structure of the β -peptides is almost unaffected by the increase in temperature from 25 $^\circ\text{C}$ to 60 $^\circ\text{C}$, supporting the thermal stability of the β -peptide secondary structures. Beside the neat secondary structure determination and the thermal stability, the CD-spectra also demonstrate that neither the recognition units nor the acetylated side chain influence the secondary structure of the β -peptides. However, structural conclusions for membrane incorporated β -peptides have only qualitative character since they are mainly based on CD-spectroscopic measurements and comparison with solution structures of 3_{14} -helices.

Tryptophan Fluorescence Spectroscopy

The two tryptophan residues were placed at both, the N- and C-terminal part of the β -peptides to position them in a transmembrane orientation with the tryptophan residues residing in the polar/apolar interface of the lipid membrane. As Trp-fluorescence is known to be sensitive to the polarity of its local environment, we monitored its maximum λ_{max} to determine the β -peptides' positions in a lipid bilayer.^{13,44} The fluorescence emission spectra of equimolar mixtures of the β -peptides **10/11**, **13/14** and **16/17** in DOPC (LUVs, peptide concentration: 12 μM , P/L-ratio = 1/500, 25 $^\circ\text{C}$) are shown in




Fig. 5 Fluorescence spectra of equimolar mixtures of 10/11, 13/14 and 16/17 in DOPC LUVs (peptide concentration: 12 μM , P/L-ratio = 1/500, 25 $^{\circ}\text{C}$).

Figure 5. In all three cases, the fluorescence emission λ_{max} of the tryptophan residues was determined to be 342 nm. If the tryptophan residues were located in a hydrophobic area like the inner membrane part, λ_{max} would be expected to be < 330 nm due to interactions of the indole ring with the acyl chains of the lipids.^{13,45,46} The tryptophan emission is, however red-shifted ($\lambda_{\text{max}} > 330$ nm), if the tryptophan residues are located in a more polar environment.¹³ Water exposure of the indole ring results in a fluorescence maximum at 350 nm.⁴⁷ From the obtained $\lambda_{\text{max}} = 342$ nm, we conclude that the tryptophan residues are located in a polar environment, i.e. at the polar/apolar interface of the lipid bilayer. This suggests that the β -peptides are orientated in a transmembrane fashion within the DOPC bilayer.

To conclude, the synthesized β -peptides form a stable secondary structure in membrane environment with a CD-signature characteristic for a right-handed 3_{14} -helix.^{22,28,36} From X-ray crystallography data it is known that an ideal 3_{14} -helix has three amino acid residues per turn with a pitch of 5.0 \AA .⁴⁸ D- β^3 -Tryptophan is located at the polar/apolar interface as determined by the intrinsic tryptophan fluorescence, which is in agreement with our recently published X-ray diffraction data obtained from very similar transmembrane β -peptides.⁴⁹ The X-ray analysis clearly demonstrate that the peptides insert in a lipid bilayer in a transmembrane fashion, with the $^{\text{h}}\text{Trp}$ localized at the lipid-water interface.⁴⁹ Consequently, the D- β^3 -lysine residues are expected to be localized in the aqueous environment. Assuming an ideal 3_{14} -helical structure, the membrane incorporated part of the β -peptide helix would have a length of 38.3 \AA . The thickness of a DOPC bilayer, given by the distance between the phosphate groups (D_{HH}) resulting from X-ray studies, is $D_{\text{HH}} = 36.7$ \AA .⁵⁰ Hence, the transmembrane part of the β -peptides is expected to be slightly longer than the thickness of the DOPC bilayer, which would result in a helical tilt angle of $\sim 16^{\circ}$, which was confirmed by the recently performed X-ray studies.⁴⁹

Determination of Peptide Aggregation State using FRET

$^{\text{h}}\text{Gln}$ residues were investigated as potential recognition units within the membrane in order to control the association state of the β -peptides reconstituted into DOPC vesicles. To analyse

the aggregation state, we made use of fluorescence resonance energy transfer (FRET). Therefore, β -peptides were labelled with a donor- (NBD) acceptor- (TAMRA) pair (Fig. 3).^{51,52} Figures 6A and 6B show fluorescence emission spectra of the β -peptides 12/13/14 (two $^{\text{h}}\text{Gln}$, Fig. 6A) and 15/16/17 (three $^{\text{h}}\text{Gln}$, Fig. 6B) at a P/L-ratio of 1/500 at 25 $^{\circ}\text{C}$ (see Electronic Supplementary Information for all fluorescence emission spectra, Fig. S4-S8).




Fig. 6 (A+B) Fluorescence emission spectra of NBD-labelled β -peptides (donor, 6.0 μM) and varying amounts of TAMRA-labelled species from $\chi_A = 0$ –0.5 determined at 25 $^{\circ}\text{C}$. The non-labelled compound was added to keep the total peptide concentration constant (12 μM) and the P/L-ratio at 1/500 (DOPC). (A) 12/13/14 with two $^{\text{h}}\text{Gln}$ and (B) 15/16/17 with three $^{\text{h}}\text{Gln}$. (C) Relative changes in NBD-fluorescence emission (F/F_0) as a function of increasing acceptor concentration (χ_A) are plotted for all three cases at 25 $^{\circ}\text{C}$ (9/10/11 with zero $^{\text{h}}\text{Gln}$, 12/13/14 with two $^{\text{h}}\text{Gln}$ and 15/16/17 with three $^{\text{h}}\text{Gln}$). The grey solid line is the result of a model according to WOLBER *et al.*, by taking only statistical occurrence of FRET in vesicles without the formation of aggregates into account.⁵³ The dashed lines are results of a global fit analysis, which takes a monomer-dimer equilibrium into account with $K_D = 1.7 \pm 0.2$ MF (two $^{\text{h}}\text{Gln}$) and $K_D = 1.0 \pm 0.2$ MF (three $^{\text{h}}\text{Gln}$). The solid lines are results of the global fit analysis, which takes a monomer-trimer equilibrium into account with $K_D = (3.5 \pm 1.2) \cdot 10^{-7}$ MF² (two $^{\text{h}}\text{Gln}$) and $K_D = (0.8 \pm 0.7) \cdot 10^{-7}$ MF² (three $^{\text{h}}\text{Gln}$).

The degree of β -peptide aggregation becomes accessible by measuring the ratio of NBD-fluorescence intensity F at 530 nm in the presence of TAMRA-labelled β -peptides and F_0 , NBD fluorescence intensity at 530 nm in its absence as a function of the TAMRA-labelled β -peptides concentration, while keeping the total peptide/lipid-ratio constant by adding the acetylated non-labelled β -peptide (Fig. 6A, Fig. 6B).¹⁵ It is expected that the β -peptide lacking ^hGln residues remains monomeric in lipid vesicles, while the β -peptides with two or three ^hGln residues tend to aggregate as a result of hydrogen bond formation in the membrane. The ratio F/F_0 as a function of the mole fraction (χ_A) of the acceptor of the β -peptides at 25 °C for all described cases (zero ^hGln, two ^hGln and three ^hGln) are shown in Figure 6C. In all cases, F/F_0 decreases with increasing mole fraction (χ_A) of the acceptor of the β -peptides. However, for the β -peptides with two ^hGln and three ^hGln, the decrease is significantly enhanced compared to that without ^hGln indicating a difference in the association behaviour of the three β -peptides. Assuming that the β -peptides without ^hGln (9/10/11) are monomeric in vesicles, the statistically occurring FRET in the vesicles can be calculated according to WOLBER *et al.* taking into account the area of 0.7 nm² for a DOPC lipid, a P/L ratio of 1/500 and a vesicle diameter of 100 nm.^{6,53} The grey solid line (Fig. 6C) is the result of a fitting routine, from which a FÖRSTER radius of $R_0 = 5.1 \pm 0.1$ nm was determined. This value is in excellent agreement with previously obtained FÖRSTER radii confirming our notion that the β -peptides without ^hGln are monomeric.⁶ For the two other β -peptide cases, a more pronounced FRET was observed, which indicates that the β -peptides indeed oligomerize. We analysed the data sets obtained for different P/L-ratios (1/500, 1/750, and 1/1000, Electronic Supplementary Information Fig. S7, Fig. S8, and Fig. S9) assuming a monomer-dimer and a monomer-trimer equilibrium (Fig. 6C).^{6,15,16,53} For a monomer-dimer equilibrium a $K_D = 1.7 \pm 0.2$ MF (two ^hGln) and $K_D = 1.0 \pm 0.2$ MF (three ^hGln) was found, respectively. The results of the global fit analysis are depicted in Figure 6C (dashed lines). Taking a monomer-trimer equilibrium into account, the data are in slightly better agreement between the fit with dissociation constants $K_D = (3.5 \pm 1.2) \cdot 10^{-7}$ MF² for β -peptides with two ^hGln (12/13/14) and $K_D = (0.8 \pm 0.7) \cdot 10^{-7}$ MF² for β -peptides with three ^hGln (15/16/17) (Fig. 6C, solid lines). The results provide strong evidence that the β -peptides 12/13/14 (two ^hGln) and 15/16/17 (three ^hGln) form aggregates as a result of hydrogen bond formation owing to the ^hGln residues. To investigate the impact of hydrogen bond formation on the aggregation properties in more detail, the temperature was varied. Hydrogen bonds are known to become weaker at higher temperatures.^{54,55} CD-spectroscopic measurements of the β -peptides 12 and 15 at 25 °C and at 60 °C have shown that the synthesized β -peptides form very stable secondary structures, even at higher temperature. Thus, if hydrogen bonds are the major driving force for aggregate formation, an increase in temperature should influence the aggregation state of the β -peptide helices. Fluorescence emission spectra of the β -peptides 9/10/11, 12/13/14 and 15/16/17 within lipid bilayers (P/L-ratio = 1/500) were measured at 25 °C and 60 °C (Electronic




Fig. 7 Relative changes in NBD-fluorescence emission (F/F_0) as a function of increasing acceptor concentration (χ_A) are plotted for 12/13/14 with two ^hGln (A) and 15/16/17 with three ^hGln (B) as recognition units at 25 °C and 60 °C. The bright grey solid line is the result of a model according to WOLBER *et al.*, taking only statistical occurrence of FRET in vesicles without the formation of aggregates into account.^{6,53} The dark grey solid lines are the results of a global fit analysis which takes a monomer-trimer equilibrium into account with $K_D = (3.5 \pm 1.2) \cdot 10^{-7}$ MF² (two ^hGln) and $K_D = (0.8 \pm 0.7) \cdot 10^{-7}$ MF² (three ^hGln).

Supplementary Information 9/10/11 in Fig. S4 at 25 °C, 12/13/14 in Fig. S5 at 25 °C and 60 °C, and 15/16/17 in Fig. S6 at 25 °C and 60 °C). Since the ratio F/F_0 is measured, any temperature effects such as the temperature dependency of the extinction coefficients of the fluorophores NBD and TAMRA are ruled out.⁶ Figure 7 shows the results obtained at 25 °C and 60 °C together with the expected F/F_0 (χ_A) for monomeric β -peptides. It is obvious that for both β -peptide species, the observed FRET is reduced at 60 °C compared to that at 25 °C. In case of 12/13/14 (two ^hGln), the β -peptides appear to be in the monomeric state at 60 °C (Fig. 7A), while for 15/16/17 (three ^hGln), the β -peptides do not fully dissociate into monomers, but still show an altered monomer-trimer equilibrium (Fig. 7B). As these β -peptides have one additional recognition unit compared to 12/13/14, the full dissociation does not occur, even at 60 °C owing to a larger number of hydrogen bonds in the aggregates. These results strongly support our notion that hydrogen bond formation via the α - β^3 -glutamine residues allows to control the aggregation state of the β -peptides in a membrane environment.

Conclusion

The organization and assembly of transmembrane peptide helices in lipid membranes is pivotal for proper function and is influenced by a number of different factors such as helix-helix and lipid-helix interactions. For transmembrane β -peptides with a defined secondary structure, these factors are barely understood and the control of helix-helix interaction in lipid bilayers has not yet been reported. We successfully designed, synthesized and reconstituted transmembrane β -peptides, modified with D- β^3 -glutamine as specific recognition unit to elucidate the impact of hydrogen bond formation within the lipid bilayer on helix-helix recognition and assembly. By varying the number of h Gln in the helix, we were able to tune the aggregation state (monomeric or oligomeric) and the strength of the helix-helix interaction. Owing to its similarity with the well-defined 3_{14} -helix where every third amino acid residue is oriented on the same face of the helix, a specific labelling of the transmembrane β -peptides makes it readily possible to precisely position a number of recognition units and hence tune the aggregation state and the strength of the interaction. This will open the avenue to rationally design different β -peptide assemblies with different functionalities.

Experimental

Syntheses of the D- β^3 -Amino Acid Residues and the β -Peptides

The D- β^3 -amino acids derivatives **1-7** were synthesized in excellent yields by an optimized ARNDT-EISTERT homologation method and were used without further purification.^{48,56,57} The synthesis of aromatic β^3 -amino acids is prone to racemization.⁵⁸ Thus, the enantiomeric purity of Fmoc-D- β^3 -Trp(Boc)-OH (**6**) was proven using MARFEYS reagent (Fig. S1).⁵⁹ The β -peptides **9-17** were synthesized using manual microwave-assisted Fmoc-solid phase peptide synthesis (SPPS).^{60,61} h Lys, h Trp and h Gln were used with orthogonally protected side chains to avoid side reactions. To attach the fluorophores 5(6)-carboxytetramethylrhodamine (5(6)-TAMRA) and 4-chloro-7-nitrobenzo-2-oxa-1,3-diazol (NBD-Cl) to the respective β -peptide, orthogonally protected h Lys were used. After Fmoc-deprotection, NBD-Cl was linked to the side chain of Boc-D- β^3 -Lys(Fmoc)-OH (**2**), which was coupled as final D- β^3 -amino acid to the N-terminus of the β -peptide. Fmoc-D- β^3 -Lys(Alloc)-OH (**3**) was introduced at the C-terminus of the β -peptide sequence providing orthogonality to the Boc-protecting group. However, during Alloc-deprotection the terminal Fmoc-group was also deprotected. Thus, Boc-D- β^3 -Lys(Boc)-OH (**8**) was coupled to the N-terminal end of the β -peptide and after Alloc-deprotection, 5(6)-TAMRA was attached to the β -peptide sequence. All experimental procedures are described in detail in the Supporting Information.

Peptide Reconstitution in Large Unilamellar Vesicles

Large unilamellar vesicles (LUVs) of 1,2-dioleoyl-*sn*-glycero-3-phosphatidylcholine (DOPC) were prepared in TRIS[®] buffer (100 mM NaCl, 25 mM TRIS, pH = 7.1).⁶² Lipids were dissolved in

chloroform and β -peptides in methanol. The β -peptide concentrations of the stock solutions were determined by UV absorption spectroscopy. Lipids and β -peptides were added in a test tube in the required concentrations to adjust the peptide/lipid-ratio (P/L-ratio). Removing the solvent in a nitrogen stream at room temperature produced almost clear peptide/lipid films at the test tube walls. To induce secondary structure formation the peptide/lipid films were treated with trifluoroethanol.^{50,35} Solvent was again removed in a nitrogen stream and the films were dried under reduced pressure at 40 °C overnight. Multilamellar vesicles were produced by hydration of the peptide/lipid film in TRIS buffer. After 2 h of incubation at 25 °C the hydrated peptide/lipid films were vortexed (30 s) and incubated (5 min) in five cycles. The milky suspensions were extruded 31 times through a polycarbonate membrane with 100 nm nominal pore size using the AVESTIN LIPOFAST mini extruder (Ottawa, Canada) to obtain clear vesicle suspensions. In case of fluorescence spectroscopy, DOPC dissolved in chloroform and the β -peptides **9-17** dissolved in methanol were mixed in the order: (i) DOPC, (ii) NBD labelled β -peptide (**10**, **13**, **16**), (iii) TAMRA labelled β -peptide (**11**, **14**, **17**), (iv) non-labelled β -peptide (**9**, **12**, **15**). For CD-spectroscopic analysis, the β -peptides **9**, **12** and **15** dissolved in methanol were mixed with DOPC dissolved in chloroform.

Circular Dichroism (CD) Spectroscopy

All CD-spectra were measured on a JASCO-1500 spectropolarimeter (Goß-Umstadt, Germany) equipped with a JULABO F250 (Seelbach, Germany) temperature control unit and recorded at 25 °C and at 60 °C in a wavelength range of 260-180 nm. The CD-spectra were background-corrected against pure vesicle suspension without incorporated β -peptides or TFE and expressed as molar ellipticity θ (deg·cm²·dmol⁻¹), according to GREENFIELD.⁶³

Tryptophan Fluorescence and Fluorescence Resonance Energy Transfer (FRET)

All fluorescence spectra were measured on a JASCO FP 6200 (Groß-Umstadt, Germany) under temperature control using a JASCO thermostat (model ETC-272T, Groß-Umstadt, Germany). Tryptophan fluorescence of the β -peptides was excited at 280 nm and the emission detected in the range of 300-400 nm. The association of the synthesized β -peptides in lipid bilayers was investigated by FRET. The β -peptides **10**, **13** and **16** were labelled with NBD as donor fluorophore and TAMRA served as acceptor in the β -peptides **11**, **14** and **17**. The concentration of the respective NBD-equipped β -peptide (6.0 μ M) as well as the total β -peptide concentration (12 μ M) were kept constant using the corresponding non-labelled β -peptide **9**, **12** or **15**. The mole fraction (χ_A) of the respective TAMRA-labelled β -peptide was varied from 0-0.5. After extrusion of the lipid vesicles (DOPC, LUVs, P/L-ratio = 1/500), the fluorescence emission of the samples was measured at 25 °C and 60 °C. The fluorescence was excited at 464 nm and the fluorescence emission was monitored between 500-650 nm. The theoretical

treatment of the obtained FRET data was performed as described previously.^{15,16,52,53,64,65}

Acknowledgements

Generous financial support of the Deutsche Forschungsgemeinschaft (SFB 803, project A01) is gratefully acknowledged.

References

- M. Rueping, Y. R. Mahajan, B. Jaun and D. Seebach, *Chem. - Eur. J.*, 2004, **10**, 1607–1615.
- U. Koert, *Angew. Chem. Int. Ed. Engl.*, 1997, **36**, 1836–1837.
- A. M. Brückner, P. Chakraborty, S. H. Gellman and U. Diederichsen, *Angew. Chem. Int. Ed.*, 2003, **42**, 4395–4399.
- P. Chakraborty and U. Diederichsen, *Chem. – Eur. J.*, 2005, **11**, 3207–3216.
- N. Naarmann, B. Bilgiçer, H. Meng, K. Kumar and C. Steinem, *Angew. Chem. Int. Ed.*, 2006, **45**, 2588–2591.
- P. E. Schneggenburger, S. Müller, B. Worbs, C. Steinem and U. Diederichsen, *J. Am. Chem. Soc.*, 2010, **132**, 8020–8028.
- B. Bilgiçer and K. Kumar, *Proc. Natl. Acad. Sci. U. S. A.*, 2004, **101**, 15324–15329.
- M. A. Lemmon and D. M. Engelman, *FEBS Lett.*, 1994, **346**, 17–20.
- C. Choma, H. Gratkowski, J. D. Lear and W. F. DeGrado, *Nat. Struct. Mol. Biol.*, 2000, **7**, 161–166.
- D. Seebach and J. Gardiner, *Acc. Chem. Res.*, 2008, **41**, 1366–1375.
- D. H. Appella, B. J. Joseph, S. R. Durell and S. H. Gellman, *J. Am. Chem. Soc.*, 1999, **121**, 2309–2310.
- T. L. Raguse, J. R. Lai, P. R. LePlae and S. H. Gellman, *Org. Lett.*, 2001, **3**, 3963–3966.
- J. T. Vivian and P. R. Callis, *Biophys. J.*, 2001, **80**, 2093–2109.
- D. Liu and W. F. DeGrado, *J. Am. Chem. Soc.*, 2001, **123**, 7553–7559.
- N. Naarmann, B. Bilgiçer, K. Kumar and C. Steinem, *Biochemistry (Mosc.)*, 2005, **44**, 5188–5195.
- B. D. Adair and D. M. Engelman, *Biochemistry (Mosc.)*, 1994, **33**, 5539–5544.
- J. M. G. Rogers, L. G. Lippert and F. Gai, *Anal. Biochem.*, 2010, **399**, 182–189.
- D. Seebach, A. K. Beck and D. J. Bierbaum, *Chem. Biodivers.*, 2004, **1**, 1111–1239.
- J.-C. Meillon and N. Voyer, *Angew. Chem.*, 1997, **109**, 1004–1006.
- J. A. Kritzer, J. Tirado-Rives, S. A. Hart, J. D. Lear, W. L. Jorgensen and A. Schepartz, *J. Am. Chem. Soc.*, 2005, **127**, 167–178.
- B. W. Gung, D. Zou, A. M. Stalcup and C. E. Cottrell, *J. Org. Chem.*, 1999, **64**, 2176–2177.
- T. L. Raguse, J. R. Lai and S. H. Gellman, *Helv. Chim. Acta*, 2002, **85**, 4154–4164.
- J. A. Killian, G. von Heijne and others, *Trends Biochem. Sci.*, 2000, **25**, 429–433.
- D. A. Kelkar and A. Chattopadhyay, *Biochim. Biophys. Acta BBA-Biomembr.*, 2007, **1768**, 2011–2025.
- M. R. R. de Planque, B. B. Bonev, J. A. A. Demmers, D. V. Greathouse, R. E. Koeppe, F. Separovic, A. Watts and J. A. Killian, *Biochemistry (Mosc.)*, 2003, **42**, 5341–5348.
- M. R. R. de Planque, J. A. W. Kruijtzter, R. M. J. Liskamp, D. Marsh, D. V. Greathouse, R. E. Koeppe, B. de Kruijff and J. A. Killian, *J. Biol. Chem.*, 1999, **274**, 20839–20846.
- X. Daura, D. Bakowies, D. Seebach, J. Fleischhauer, W. F. van Gunsteren and P. Krüger, *Eur. Biophys. J.*, 2003, **32**, 661–670.
- D. Seebach, J. V. Schreiber, S. Abele, X. Daura and W. F. van Gunsteren, *Helv. Chim. Acta*, 2000, **83**, 34–57.
- D. Seebach, M. Overhand, F. N. Kühnle, B. Martinoni, L. Oberer, U. Hommel and H. Widmer, *Helv. Chim. Acta*, 1996, **79**, 913–941.
- G. Guichard, S. Abele and D. Seebach, *Helv. Chim. Acta*, 1998, **81**, 187–206.
- X. Daura, W. F. van Gunsteren, D. Rigo, B. Jaun and D. Seebach, *Chem. Eur. J.*, 1997, **3**, 1410–1417.
- J. R. Allison, M. Müller and W. F. van Gunsteren, *Protein Sci. Publ. Protein Soc.*, 2010, **19**, 2186–2195.
- M. A. Andrade, P. Chacon, J. J. Merelo and F. Moran, *Protein Eng.*, 1993, **6**, 383–390.
- K. Kaur, T. Sprules, W. Soliman, R. Beleid and S. Ahmed, *Biochim. Biophys. Acta BBA - Proteins Proteomics*, 2008, **1784**, 658–665.
- F. D. Sonnichsen, J. E. Van Eyk, R. S. Hodges and B. D. Sykes, *Biochemistry (Mosc.)*, 1992, **31**, 8790–8798.
- R. P. Cheng, S. H. Gellman and W. F. DeGrado, *Chem. Rev.*, 2001, **101**, 3219–3232.
- D. Seebach, D. F. Hook and A. Glättli, *Biopolymers*, 2006, **84**, 23–37.
- Y. Hamuro, J. P. Schneider and W. F. DeGrado, *J. Am. Chem. Soc.*, 1999, **121**, 12200–12201.
- B. A. Wallace, *Protein Sci.*, 2003, **12**, 875–884.
- R. W. Woody, *Eur. Biophys. J. EBJ*, 1994, **23**, 253–262.
- R. W. Woody, *Biopolymers*, 1978, **17**, 1451–1467.
- A. Chakraborty, T. Kortemme, S. Padmanabhan and R. L. Baldwin, *Biochemistry (Mosc.)*, 1993, **32**, 5560–5565.
- H. E. Auer, *J. Am. Chem. Soc.*, 1973, **95**, 3003–3011.
- J. Ren, S. Lew, Z. Wang and E. London, *Biochemistry (Mosc.)*, 1997, **36**, 10213–10220.
- S. Lew, J. Ren and E. London, *Biochemistry (Mosc.)*, 2000, **39**, 9632–9640.
- J. Ren, S. Lew, J. Wang and E. London, *Biochemistry (Mosc.)*, 1999, **38**, 5905–5912.
- C. D. Geddes, *Reviews in Fluorescence 2007*, Springer Science & Business Media, 2009.
- D. Seebach and J. L. Matthews, *Chem. Commun.*, 1997, 2015–2022.
- U. Rost, Y. Xu, T. Salditt and U. Diederichsen, *ChemPhysChem*, 2016, DOI: 10.1002/cphc.201600289.
- N. Kučerka, S. Tristram-Nagle and J. F. Nagle, *J. Membr. Biol.*, 2006, **208**, 193–202.
- S. Sindbert, S. Kalinin, H. Nguyen, A. Kienzler, L. Clima, W. Bannwarth, B. Appel, S. Müller and C. A. M. Seidel, *J. Am. Chem. Soc.*, 2011, **133**, 2463–2480.
- D. M. Lilley and T. J. Wilson, *Curr. Opin. Chem. Biol.*, 2000, **4**, 507–517.
- P. K. Wolber and B. S. Hudson, *Biophys. J.*, 1979, **28**, 197–210.
- R. C. Dougherty, *J. Chem. Phys.*, 1998, **109**, 7372–7378.
- H. A. Scheraga, G. Nemethy and I. Z. Steinberg, *J. Biol. Chem.*, 1962, **237**, 2506–2508.
- P. I. Arvidsson, J. Frackenhohl and D. Seebach, *Helv. Chim. Acta*, 2003, **86**, 1522–1553.
- B. S. Patil, G.-R. Vasanthakumar and V. V. S. Babu, *Letts. Pept. Sci.*, 2002, **9**, 231–233.
- J. Podlech and D. Seebach, *Liebigs Ann.*, 1995, **1995**, 1217–1228.
- R. Bhushan and H. Brückner, *Amino Acids*, 2004, **27**, 231–247.
- J. K. Murray and S. H. Gellman, *Org. Lett.*, 2005, **7**, 1517–1520.
- O. Flögel, J. D. C. Codée, D. Seebach and P. H. Seeberger, *Angew. Chem. Int. Ed.*, 2006, **45**, 7000–7003.

ARTICLE

Chemical Science

- 62 R. C. MacDonald, R. I. MacDonald, B. P. M. Menco, K. Takeshita, N. K. Subbarao and L. Hu, *Biochim. Biophys. Acta BBA - Biomembr.*, 1991, **1061**, 297–303.
- 63 N. J. Greenfield, *Nat. Protoc.*, 2007, **1**, 2876–2890.
- 64 M. Li, L. G. Reddy, R. Bennett, N. D. Silva, L. R. Jones and D. D. Thomas, *Biophys. J.*, 1999, **76**, 2587–2599.
- 65 L. A. Chung, J. D. Lear and W. F. DeGrado, *Biochemistry (Mosc.)*, 1992, **31**, 6608–6616.