

Soft Matter

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Cite this: DOI: 10.1039/xxxxxxxxxx

Using Brownian motion to measure shape asymmetry in mesoscopic matter using optical tweezers[†]

Basudev Roy, Argha Mondal, Sudipta K. Bera, and Ayan Banerjee*

Received Date
Accepted Date

DOI: 10.1039/xxxxxxxxxx

www.rsc.org/journalname

We propose a new method for quantifying shape asymmetry in the mesoscopic scale. The method takes advantage of the intrinsic coupling between rotational and translational Brownian motion (RBM and TBM, respectively) which happens in the case of asymmetric particles. We determine the coupling by measuring different correlation functions of the RBM and TBM for single, morphologically different, weakly-trapped red blood cells in optical tweezers. The cells have different degrees of asymmetry that are controllably produced by varying the hypertonicity of their aqueous environment. We demonstrate a clear difference in the nature of the correlation functions both qualitatively and quantitatively for three types of cells having a varying degree of asymmetry. This method can have a variety of applications ranging from early stage disease diagnosis to quality control in microfabrication.

Measuring shape asymmetry can be almost trivial as long as the object is visualized easily and can be measured using standard length scales. However, as the size of the system reduces, visualization is rendered more and more difficult, and becomes extremely challenging once the size of the object is less than the diffraction limit for the wavelength of light used for imaging. Therefore, non-visual yet simpler techniques to study asymmetry need to be explored in this regime. This is even more so since asymmetry in the microscopic scales has far-reaching consequences, especially in the context of biological entities such as motor proteins or cells, where any biochemical processes often result in morphological changes to cell shape¹ or nuclear structure². Changes in shape of cells are also observed at the onset of diseases - as in the case of red blood cells (RBCs) due to malaria³, or sickle cell anaemia⁴. Thus, sensitive detection of small changes in asymmetry may lead to early stage disease detection, which is

a major technology challenge presently. Similarly, such methods may be useful for quality control in microfabrication processes, in order to check for small deviations from a preferred structure.

A mesoscopic particle suspended in a fluid undergoes both translational and rotational Brownian motion due to random collisions with solvent particles surrounding it⁵. While translational Brownian motion dominates for symmetric particles, for asymmetric ones - there is a strong rotational component. For the latter, the two types of motion also get coupled⁶, and the coupling can be expected to be stronger with increasing asymmetry. Optical tweezers are a very useful tool to study Brownian motion of micro-particles since they confine the particle spatially, and the trapped particle then executes Brownian motion with amplitude and frequency determined by the trap stiffness. In fact, the Brownian motion can be used to characterize the trap itself, as is very often done by measuring its power spectral density⁷. However, an important aspect of Brownian dynamics that remains largely ignored is the coupling between the translational and rotational Brownian motion[†] for asymmetric particles inside an optical trap, except in occasional studies⁶. It is this coupling that we exploit in this paper to study asymmetry, and we go on to show both theoretically and experimentally that the amount of coupling increases with the degree of asymmetry. As test particles, we use RBCs' due to the fact that shape of such cells can be controllably modified by varying the hypertonicity of a hypertonic aqueous solution - with healthy RBCs' in normal aqueous solutions being quite symmetric in shape³. Note that asymmetric particles also exhibit rotational motion (spin) inside an optical trap due to torque generated by asymmetric scattering^{8,9}. This rotation rate depends linearly on the light intensity. We avoid such motion in our work by maintaining a weak trap so that the light intensity is not enough to observe such motion. Rotation is also observed for birefringent particles with incident circularly polarized light, but this can be avoided by using linearly polarized light.

We first develop the theoretical premise for observing effects of the coupling between rotational and translational Brownian motion. It is well known that Brownian motion - both rotational

Indian Institute of Science Education and Research, Kolkata, Mohanpur 741246, India
*E-mail: ayan@iiserkol.ac.in

[†] Electronic Supplementary Information (ESI) available: Supplementary Information.pdf. See DOI: 10.1039/b000000x/

and translational - can be described by the Langevin equation⁷. In our theoretical approach, we add two coupling terms to cross-couple the rotation and the translation equations. Therefore, the final equation becomes:

$$\gamma_1 \frac{dx(t)}{dt} + \kappa_x x(t) + \gamma_2 \omega_1 \theta(t) = (2k_B T \gamma_1)^{1/2} \zeta_x(t), \quad (1)$$

$$\gamma_2 \frac{d\theta(t)}{dt} + \kappa_\theta \theta(t) - \gamma_1 \omega_2 x(t) = (2k_B T \gamma_2)^{1/2} \zeta_\theta(t), \quad (2)$$

where κ_x denotes the force constant of the optical trap for translational motion, while κ_θ is that for rotational motion that is polarization-dependent (i.e. it tries to orient an object with optics axis along the polarization of the input laser beam for both birefringence-driven rotation and that due to asymmetric scattering). $(2k_B T \gamma_1)^{1/2} \zeta_x(t)$ and $(2k_B T \gamma_2)^{1/2} \zeta_\theta(t)$ are two independent Gaussian random noise (k_B being the Boltzmann constant), which represent the Brownian forces at absolute temperature T for both x and θ coordinate systems. In the above equations $\gamma_2 \omega_1 \theta(t)$, and $\gamma_1 \omega_2 x(t)$ denote the coupling between x and θ .

After solving[†] for x and θ ¹⁰, we compute the auto-correlation functions[†] for the translational and rotational Brownian motions as¹¹:

$$\begin{aligned} \langle x(t)x(t+\delta t) \rangle &= \frac{Ak_B T}{\kappa_x} e^{-(\kappa_x/\gamma_1 + \kappa_\theta/\gamma_2)/2|\delta t|} \cos(C\delta t), \\ \langle \theta(t)\theta(t+\delta t) \rangle &= \frac{Ak_B T}{\kappa_\theta} e^{-(\kappa_x/\gamma_1 + \kappa_\theta/\gamma_2)/2|\delta t|} \cos(C\delta t), \end{aligned} \quad (3)$$

where A is a coupling amplitude whose exact form is not known to us, but which is strongly manifested in experimental data, as we show later. $B = (\kappa_x/\gamma_1 + \kappa_\theta/\gamma_2)/2$, $C = \sqrt{\omega_1 \omega_2 - (\kappa_x/\gamma_1 - \kappa_\theta/\gamma_2)^2/4}$, and characterizes the coupling strength. Here, κ_x/γ_1 is the translational corner frequency while the κ_θ/γ_2 is the rotational corner frequency. We then go on to define the well known cross-correlation function (CCF) between the translational and rotational Brownian motions¹¹ that is given by:

$$\begin{aligned} \langle x(t)\theta(t+\delta t) \rangle &= \frac{Ak_B T}{\sqrt{\kappa_x \kappa_\theta}} e^{-(\kappa_x/\gamma_1 + \kappa_\theta/\gamma_2)/2|\delta t|} \\ &\quad \cos(C\delta t + D), \\ \langle x(t)\theta(t-\delta t) \rangle &= \frac{Ak_B T}{\sqrt{\kappa_x \kappa_\theta}} e^{-B|\delta t|} \cos(C\delta t + D). \end{aligned} \quad (4)$$

Here, $\tan(D) = B/C$. The *CCF* differs from the *ACF* by the phase factor D , which has two consequences: (i) it renders the *CCF* asymmetric, (note that the *ACF* is necessarily symmetric due to causality), (ii) it introduces another parameter that may be used to quantify asymmetry, which is obviously advantageous to us. The form of the *CCF* is thus similar to a coupled harmonic oscillator system, and has been used in diverse areas of physics such as quantum optics¹⁰, cavity quantum electrodynamics¹², and torque calibration in Brownian motion¹¹. Thus, our task is to measure A, B, C , and D for different levels of asymmetry in microparticles executing Brownian motion in optical tweezers with high precision so that small changes in asymmetry may be de-

tected with enough signal to noise (S/N). For this reason, we proceed to introduce the differential cross-correlation, *DCCF*, the difference of two *CCFs*¹³, to yield better precision:

$$\begin{aligned} &\langle x(t)\theta(t+\delta t) - \theta(t+\delta t)x(t) \rangle = \\ &\frac{\alpha k_B T}{\sqrt{\kappa_x \kappa_\theta}} e^{-B|\delta t|} (\cos(C\delta t + D) - \cos(C\delta t - D)) \end{aligned} \quad (5)$$

The nature of the *DCCF* is anti-symmetric since the functional form is similar to the derivative of the *CCF*. Now, for dielectric particles with only form birefringence (or even for birefringent particles at low laser intensities), the translational corner frequency is much larger than the rotational one¹⁴. Thus, the product of coupling coefficients $\omega_1 \omega_2$ can be written as[†]

$$\omega_1 \omega_2 = B^2 + C^2. \quad (6)$$

The individual values of ω_1 and ω_2 cannot be ascertained from here since that would require exact information about the asymmetry causing the cross coupling. However, even the value of the product indicates how well the rotational Brownian motion and the translational Brownian motion are coupled. A very crucial point to note is that these equations work when the translational and rotational corner frequencies are lower than the coupling frequencies, so that one requires rather weak optical traps in order to probe the value of the coupling constants. Note that weak traps are also required to stop asymmetric particles from spinning, as has been described earlier. For our optical tweezers experiments[†],




Fig. 1 ((a). Normal RBC. (b) Moderately deformed RBC (hypertonic solution). (c) Maximally deformed RBC (hypertonic solution). (d) Typical measured time series of and translational (grey) and rotational (black) Brownian motion in RBC. The original data has been offset in voltage for visualization. (e) Histogram of time series data for rotational Brownian motion of normal RBC. (f) Power spectral density of translational (upper) and rotational (lower) motion of normal RBC.

we extracted RBCs from healthy donors, and typically started individual experiments after up to 1 hour from preparing the diluted solution in the hypertonic solvent[†], allowing the time for the RBCs to get deformed. The RBCs were also visually monitored continuously and experiments were started after the RBC reached the required degree of deformity. Fig. 1 shows RBCs at various degrees of deformation, starting with a normal RBC in aqueous solution (Fig. 1(a)), a moderately deformed RBC (Fig. 1(b)), and a maximally deformed RBC (Fig. 1(c)). With maximally deformed RBCs, we regularly obtained rotation (spin) at constant rates de-

pending on the trapping laser power, which demonstrated that the RBCs maintained a definite shape in the trap. However, at laser powers of around 15 mW after the microscope objective, the maximally deformed RBCs were stably trapped, but did not show any rotational motion, though they were typically aligned with the polarization direction of the linearly polarized laser. We used the same laser power for all RBCs during the experiments. The normal RBCs almost never showed rotation, while moderately deformed RBCs rarely showed rotation at high laser powers (less than 5% cases).




Fig. 2 Plots of the normalized CCF (NCCF) for (a) a spherical polystyrene bead, (b) Normal RBC, (c) Moderately deformed RBC, (d) Maximally deformed RBC. (b), (c), and (d) are fit (bold line) using Eqn. 4. The fit values of different parameters are given in boxes inside each plot, with the respective 1σ errors in parenthesis.

The measurements of Brownian motion are performed with a quadrant photodiode[†], and we use the technique we recently developed to detect rotational and translational motion simultaneously by measuring the sum and difference of the signals from the diagonal quadrants of the QPD¹⁴. The rotational and translational data are acquired at a sampling frequency of 40KHz taken for 5 seconds. Typical time series for translational and rotational Brownian motion for normal RBC are given in Fig. 1(d). The histograms of the time series data fit well to Gaussian distributions - a sample of which is shown in Figs. 1(e). Corresponding power spectral densities (PSD) for both translational and rotational motion are obtained by averaging over 25 individual spectra. We also check that our signal PSD for RBCs is substantially higher than that of the noise floor of the setup[†]. The measured PSDs are shown in Figs. 1(f), with the translational corner frequency (1.07 Hz) being 3 times higher than the rotational one (0.35 Hz). Before computing the CCF, we first find out the ACF for all types of RBCs and find that these are symmetric around $t = 0$, as is expected[†]. We now compute the CCF and DCCF for the different types of RBCs. Before proceeding with that, we normalize the functions in the following representative manner:

$$NCCF(x, \theta) = \frac{\langle x(t) \theta(t + \delta t) \rangle}{\sqrt{\langle x^2(t) \rangle} \sqrt{\langle \theta^2(t) \rangle}}, \quad (7)$$

where NCCF (NDCCF) is the normalized CCF (DCCF). We first proceed to measure the NCCF. As a control, we use a spheri-




Fig. 3 Plots of the normalized DCCF (NDCCF) for (a) a spherical polystyrene bead, (b) Normal RBC, (c) Moderately deformed RBC, (d) Maximally deformed RBC. (b), (c), and (d) are fit (bold line) using Eqn. 5. The fit values of different parameters (except y_0 , which is not of significance) are given in boxes inside each plot.

cal polystyrene bead of diameter $3 \mu\text{m}$ and determine the NCCF - as shown in Fig. 2(a). As expected, the signal has very small amplitude, since the rotational Brownian motion of the sphere is negligible. On the other hand, the NCCF from the least asymmetric NRBC has a clear structure. We can fit it well to Eqn. 4 with an additional offset term y_0 . It is also asymmetric in nature as seen from the non-zero values of D . The fit values of y_0 , A , B , C , and D with their respective error bars are given in the box inside Fig. 3(b). Similarly, the fit parameters for moderately and maximally deformed RBC are given in Figs. 3(c) and (d). Using these, we calculate the coupling coefficient products $\omega_1 \omega_2$ from Eq. 6, the values of which are given in Table 1. We observe that the amplitude A for the NCCF increases with increasing asymmetry, and is the most robust parameter in differentiating between the RBCs, even at the 3σ level of error bars. The errors in the fit parameters B and C (indicated in parenthesis) lead to the errors in $\omega_1 \omega_2$, and it is clear that we are not able to differentiate between normal RBC ($\omega_1 \omega_2 = 6.7 \pm 3.0$), and moderately deformed RBC ($\omega_1 \omega_2 = 9.3 \pm 3.5$) even at the 1σ level of error. Maximally deformed RBC ($\omega_1 \omega_2 = 17.1 \pm 5.1$), however, is different from the other types of RBCs at the 1σ level. Also, the value of D increases with increasing asymmetry, being the highest for a single maximally deformed RBC. Thus, even this could serve as a parameter that could be used to quantify asymmetry. Note that the value of D also acts as a consistency check for the values of B and C , since $\tan(D) = B/C$, and the fit values of D can thus be compared independently to that obtained from this relation. We obtain matches for all three types of RBCs considering the overlaps of error bars at the 3σ level[†]. Interestingly, the signs of C and D change for maximally deformed RBCs, and this needs further careful investigation at various levels of asymmetry. In addition, we check the value of B from the fit with that obtained by adding the translational and rotational corner frequencies, and as we show in the case of a normal RBC - the value of B from Fig. 1 comes out to be $1.15(15)$ Hz, matches well

Table 1 Table showing the values of different parameters obtained from fitting experimentally measured *NCCF* and *NDCCF* for Normal (NRBC), moderately deformed (MoRBC), and maximally deformed (MaRBC) RBC.

RBC type	<i>NCCF</i>					<i>NDCCF</i>				
	<i>A</i>	<i>B</i> (Hz)	<i>C</i> (Hz)	<i>D</i> (rad)	$\omega_1\omega_2$ (Hz ²)	<i>A</i>	<i>B</i> (Hz)	<i>C</i> (Hz)	<i>D</i> (rad)	$\omega_1\omega_2$ (Hz ²)
NRBC	0.013(3)	1.20(15)	-2.3(5)	-0.05(02)	6.7(3.0)	0.0075(1)	0.90(08)	-2.1(3)	-0.08(2)	5.2(1.0)
MoRBC	0.031(3)	1.4(2)	-2.7(5)	-0.30(08)	9.3(3.5)	0.011(1)	1.25(08)	-2.5(1)	-0.25(06)	7.8(7)
MaRBC	0.0503(4)	2.2(3)	3.5(5)	0.9(1)	17.1(5.1)	0.039(2)	2.05(10)	3.3(2)	0.75(06)	15.1(1.6)

with the value 1.20(15) in Table 1). We check this for all other cases and obtain similar results. The value of *B* increases for a maximally deformed RBC even for the same trapping powers as the other types of RBCs, and we believe that the reason may be the increase of both k_x and k_θ due to the asymmetry of the cell. Note that such increase of k_x has been reported earlier in Ref.¹⁵, where malaria-affected RBCs whose shape is distinctly different from normal RBCs (and similar to the maximally deformed RBCs in our case) showed a 25% increase in corner frequency over normal ones. The enhanced coupling of the two types of motion resultantly leads to an increase in k_θ , which leads to a higher value of *B*.

We now calculate the *NDCCF* to check whether error bars are reduced, and whether we can statistically differentiate the RBCs in terms of values of $\omega_1\omega_2$. Fig. 3(a)-(d) shows the plots of *NDCCF* for all four cases as in Fig. 2. While that for a sphere very small in amplitude, we obtain good signals (anti-symmetric as expected) for RBCs, and much better fits to the function for *NDCCF* defined in Eq. 5 compared to the *NCCF*. While *A* still remains the most robust parameter for differentiation between RBCs, we now have lower error bars for $\omega_1\omega_2$. Thus, as we observe from Table 1, we can differentiate normal RBC ($\omega_1\omega_2 = 5.2 \pm 1.0$) from moderately deformed RBC ($\omega_1\omega_2 = 7.8 \pm 0.7$) at the 1σ level and maximally deformed RBC ($\omega_1\omega_2 = 15.1 \pm 1.6$) from the other RBCs at the 3σ level. Thus, it is clear that the *NDCCF* is a much better tool to quantify shape asymmetry using our technique. Finally, we repeat these measurements for other RBCs having similar relative asymmetries, and obtain similar results. While it is not possible to control the shape asymmetry quantitatively for RBCs by varying the hypertonicity of the aqueous solution, we observe that single maximally deformed RBCs can be statistically differentiated from other RBCs using the *NDCCF*. Finally, we would like to point out that RBCs often exhibit membrane fluctuations due to thermal effects¹⁶, but such motional modes are mostly radially symmetric and would not be detected by our rotation measurement technique using the QPD[†], so that we may safely conclude that our technique for measuring the CCF is immune to membrane fluctuations.


In conclusion, we have developed a new technique for quantifying shape asymmetry by determining the cross-correlation between translational and rotational Brownian motion. The values of the coupling amplitude increase with increasing asymmetry, while other parameters also change appreciably. We have used RBCs as test particles, and our technique can differentiate

between RBCs that have different morphological states, which we control by changing the hypertonicity of the aqueous environment of the RBCs. This is a proof of concept, and we believe that the technique can be improved further by fabricating particles with controlled asymmetry using the various advanced lithography tools⁸ presently available, and could thus be useful in diverse applications in biology (early disease detection, etc.) and microfabrication (in terms of checking the symmetry of fabricated micro-particles).

The authors acknowledge Dr. Giovanni Volpe for useful discussions. This work was supported by the Indian Institute of Science Education and Research, Kolkata, an autonomous institute funded by the Ministry of Human Resource Development, Govt. of India.

References

- 1 S. J. Morrison and J. Kimble, *Nature*, 2006, **441**, 1068–1074.
- 2 A. Adjuk, S. B. Shihhare and M. Zernicka-Goetz, *Dev. Biol.*, 2014, **392**, 133–140.
- 3 A. K. Dharmadhikari, H. Basu, J. A. Dharmadhikari, S. Sharma and D. Mathur, *J. Biomed. Opt.*, 2013, **18**, 125001.
- 4 H. A. Pearson, *J. Infect. Dis.*, 1977, **136**, S25–S30.
- 5 R. Brown, *Phil. Mag.*, 1828, **4**, 161–173.
- 6 Y. Han, A. M. Alsayed, M. Nobili, J. Zhang, T. C. Lubensky and A. G. Yodh, *Science*, 2006, **314**, 626–630.
- 7 S. F. Norrelykke and H. Flyvberg, *Rev. Sci. Instrum.*, 2010, **81**, 075103.
- 8 P. Galajda and P. Ormos, *Appl. Phys. Lett.*, 2001, **78**, 249.
- 9 S. K. Mohanty, K. S. Mohanty and P. K. Gupta, *Opt. Exp.*, 2005, **13**, 4745.
- 10 H. J. Carmichael, R. J. Brecha and P. Rice, *Opt. Commun.*, 1991, **82**, 73–79.
- 11 G. Volpe and D. Petrov, *Phys. Rev. Lett.*, 2006, **97**, 210603.
- 12 G. T. Foster, S. L. Mielke and L. A. Orozco, *Phys. Rev. A*, 2000, **61**, 053821.
- 13 G. Volpe, G. Volpe and D. Petrov, *Phys. Rev. E*, 2008, **77**, 037301.
- 14 B. Roy, S. K. Bera and A. Banerjee, *Opt. Lett.*, 2014, **39**, 3316–3319.
- 15 V. Saraogi, P. Padmapriya, A. Paul, U. S. Tatu and V. Natarajan, *J. Biomed. Opt.*, 2010, **15**, 037003.
- 16 J. Evans, W. Gratzner, N. Mohandas, K. Parker and J. Sleep, *Biophys. J.*, 2008, **94**, 4134.



We quantify asymmetry of single optically trapped red blood cells from the crosscorrelation between their rotational and translational Brownian motion.