



ChemComm

**Nickel-Catalyzed Intermolecular Carboiodination of Alkynes
with Aryl Iodides**

Journal:	<i>ChemComm</i>
Manuscript ID	CC-COM-09-2018-007560.R2
Article Type:	Communication

SCHOLARONE™
Manuscripts

Nickel-Catalyzed Intermolecular Carboiodination of Alkynes with Aryl Iodides

Received 00th January 20xx,
Accepted 00th January 20xx

Toshifumi Takahashi,^{a,b} Daiki Kuroda,^{a,b} Toru Kuwano,^{a,b} Yuji Yoshida,^{a,b} Takuya Kurahashi,^{a*} and Seiji Matsubara^{a*}

DOI: 10.1039/x0xx00000x

www.rsc.org/

Nickel-catalyzed intermolecular carboiodination of alkynes with aryl iodides to form highly substituted and functionalized alkenyl iodides has been developed. The reaction involves radical-mediated formal alkyne insertion into the carbon–nickel bond and carbon–iodine reductive elimination facilitated by Ni(III) species.

Organohalogen compounds such as alkenyl iodides are versatile and essential building blocks in organic synthesis.^{1,2} Many established reactions (e.g. transition-metal-catalyzed cross-coupling reactions) require organohalogen compounds as coupling partners, and halogen atoms are easily replaced with other organic structures or heteroatoms to construct new chemical bonds.³ Although the carbometallation of an alkyne followed by reaction with a halogen source such as *N*-iodosuccinimide or I₂ is well known as a conventional synthetic method, the development of a direct, facile method to afford tetrasubstituted alkenyl halides remains a research topic of great interest. In the last decade, the transition-metal-catalyzed insertion of an alkyne into the carbon–halogen bond of an organohalogen compound, *i.e.* carbohalogenation, has been identified as a significant reaction which enables straightforward access to highly substituted alkenyl halides without by-product formation.⁴ However, based on Hartwig's study of the chemistry of ArPd(II)X complexes, the reductive elimination of a carbon–halogen bond from a transition metal is disfavored and largely promoted by sterically bulky ligands (Figure 1(a)).⁵ Thus, Lautens' group succeeded in developing intramolecular carbohalogenation reactions of alkynes by using bulky Q-Phos, tri-*tert*-butylphosphine, or aryl phosphadamantanes as ligands for the palladium catalysts, and sterically demanding mesityl or TIPS groups as substituents of the alkynes to facilitate the key reductive elimination step (Figure 1(b)).⁶ However, the steric bulkiness of both ligands and alkyne substituents retards alkyne insertion,

and therefore, the reaction was designed as an intramolecular process to overcome the steric hindrance. Thus, under this circumstance, there is a trade-off between the promotion of reductive elimination and alkyne insertion. Herein, we report the intermolecular carboiodination of alkynes with aryl iodides enabled by nickel catalysis and radical-involving reaction mechanism (Figure 1(c)).




Figure 1 Transition-metal-catalyzed carboiodination.





Figure 2 Working hypothesis for intermolecular carboiodination.

^a Department of Material Chemistry, Graduate School of Engineering, Kyoto University, Kyoto 615-8510, Japan

^b These authors contributed equally.

† Electronic Supplementary Information (ESI) available: See DOI: 10.1039/x0xx00000x

We initially hypothesized that the aryl radical species generated by homolysis of the carbon–nickel bond could react with alkynes, and recombination of the obtained alkenyl radical and Ni(II)I₂ forms an alkenylNi(III)I₂ complex, followed by facile reductive elimination to form the carbon–iodine bond from this high-valent complex would generate the desired alkenyl iodide. Accordingly, intermolecular formal alkyne insertion into carbon–halogen bonds would be achieved. To prove our hypothesis, we examined the nickel-catalyzed carboiodination of 4-octyne (**2a**) with iodobenzene (**1a**) in toluene at 50 °C (Table 1), and we found that alkenyl iodide **3aa** was obtained in 92% yield when dtbpy was employed as the ligand (Scheme 1).



To determine the scope of the reaction, we next performed the nickel-catalyzed carboiodination of alkynes with substituted iodobenzenes to obtain the corresponding tetrasubstituted alkenyl iodides (Scheme 2). The reaction of 4-tolyl iodide with 4-octyne **2a** gave **3ba** in 81% isolated yield. Even though a longer reaction time was required, *m*- or *o*-methyl-substituted phenyl iodides also afforded alkenyl iodides in high yields (**3ca**: 48 h, 71%; **3da**: 96 h, 91%). Naphthyl iodides also participated in the reaction to give the corresponding alkenyl iodides (**3ea**, **3fa**). Furthermore, various functional groups were tolerated under the reaction conditions; aryl iodides possessing alkoxy (**3ga**), trifluoromethyl (**3ha**), nitrile (**3ia**), ketone (**3ja**), or ester (**3ka**) substituents participated in the reaction to provide the corresponding alkenyl iodides in moderate to high yields. Of note, aryl iodides possessing bromine or chlorine substituents also participated in the reaction to give alkenyl iodides **3la** and **3ma** in 83% and 60% yields, respectively; clearly, bromine and chlorine atoms were tolerated under the reaction conditions, and the carboiodination proceeded in a chemoselective manner. Furthermore, boryl-substituted aryl iodides reacted with alkyne **2a** to afford the desired products in moderate-to-high yields (**3na**, **3oa**, and **3pa**) and the structure of **3pa** was confirmed with X-ray single crystal analysis (Figure 3). Alkynes possessing alkoxy or ester groups successfully participated to provide alkenyl iodides **3jb**, **3jc**, and **3jd** in 53%, 99%, and 90% yield, respectively. The reaction of phenyl bromide with **2a** resulted in low yield (<5%). The use of alkyl iodides in place of aryl iodide did not afford the products. The reaction using non-symmetrical alkynes afforded mixture of isomers and the reaction with terminal alkynes resulted in low yield.





Figure 3 ORTEP drawing of **3pa**



Scheme 2 Substrate scope in the carboiodination of alkynes with aryl iodides. All reactions were carried out using Ni(cod)₂ (5 mol%), ligand (5 mol%), **1** (0.5 mmol), and **2** (0.75 mmol) in 1 mL of toluene at 50 °C for 12 h, unless otherwise noted. Isolated yields are given. *Z/E* ratio were determined by ¹H NMR analysis and given in parentheses. ^a24 h. ^b48 h. ^c72 h. ^d96 h.




Figure 4 X-Band EPR merged-spectrum of the reaction mixture (red solid line) and simulated spectrum based on DFT calculations of Ni(I)I/dtbpy complex **4** (black dashed line). Spin density plot of computed Ni(I)I/dtbpy complex **4**. Mulliken spin density: Ni, 0.93; N, 0.05; N, 0.03; I, 0.02.

We next analyzed the catalytic reaction spectroscopically to gain insight into the reaction mechanism. The reaction mixture was subjected to electron paramagnetic resonance (EPR) measurements (Figure 4). The mixture of catalytic amount of Ni(0)/dtbpy, iodobenzene (**1a**) and 4-octyne (**2a**) in toluene (25mM) displays a signal at $g = 2.124$, which is significantly deviated from that of a free electron ($g = 2.002$). This result indicates the presence of paramagnetic nickel complex in the reaction mixture and unpaired electron is mostly nickel based.^{7,8,9,10} Further computational studies revealed that Ni(I)/dtbpy **4** simulates the experimentally observed EPR spectrum in good agreement.





Figure 5 Plausible catalytic cycle.

Based on these experiments, we propose a catalytic cycle involving odd-valent nickel complexes and radical species, e.g. Ni(I)/dtbpy **4** which is generated via comproportionation in situ (Figure 5).^{12,13} The proposed mechanism is as follows: 1) Oxidative addition of iodobenzene **1a** to Ni(I) **4** gives unstable PhNi(III)I₂ **5**, which 2) forms phenyl radical **6** via carbon–nickel bond homolysis. 3) The alkenyl radical **8** formed from phenyl radical **6** and alkyne **2a**, so that 4) recombination with the Ni(II) **7** complex gives alkenylNi(III)I₂ **9**. Finally, 5) alkenyl iodides **3aa** are obtained as final products via facile reductive elimination from the unstable high-valent Ni(III) complexes **9**. This is consistent with preliminary computational studies (Figure 6, see also SI); carbon–iodine reductive elimination from Ni(III) is exergonic with a sufficiently low activation barrier, in contrast to the endergonic reductive elimination from a Ni(II) complex with a high energy barrier. The observed reactivities of Ni(III) and Ni(II) in the reductive elimination are in good agreement with studies of the carbon–oxygen reductive elimination of nickel complexes reported by Hillhouse, MacMillan, Doyle, and their co-workers.^{8a,11,14}

In summary, we have developed the nickel-catalyzed intermolecular carboiodination of alkynes with simple aryl iodides to afford tetrasubstituted alkenyl iodides. The nickel-catalyzed radical-involving reaction mechanism affords a new way to access direct carbohalogenation in an intermolecular fashion with atom- and redox-economy. The mechanism is based on the unique properties of nickel catalysts with redox-active ligands such as bipyridine that allow Ni(I) and Ni(III) oxidation states.¹⁵ Further investigations on the reaction mechanism are underway, both experimentally and theoretically, and will be reported in due course.

– Favored path –

(a) Odd-valent mechanism of reductive elimination



– Disfavored path –

(b) Even-valent mechanism of reductive elimination




Figure 6 Plausible reductive elimination mechanisms. These calculations were performed with M06/SDD(2f)(Ni),SDD(I),6-311+G(2d,p)(C,H,N) // M06/LanI2dz(f)(Ni),LanI2dz(I),6-31G(d)(C,H,N) with CPCM(toluene) solvation. Further details are given in the SI.

We are grateful to Prof. Tatsuhiro Kato for valuable support in the EPR analysis. This work was supported by the Advanced Catalytic Transformation Program for Carbon Utilization (ACT-C, JPMJCR12YE) from the Japan Science and Technology Agency (JST), and Grants-in-Aid for Scientific Research (Nos. 18H04253, 17KT0006, and 15H03809) from the Ministry of Education, Culture, Sports, Science and Technology (Japan). T.K. also acknowledges the Uehara Memorial Foundation, and the Asahi Glass Foundation. T.T. acknowledges the Council of Human Resources Fostering Program in Chemistry and Japan Chemical Industry Association for fellowship support.


Conflicts of interest

There are no conflicts to declare.

References

- The Chemistry of Halides, Pseudo-Halides and Azides, Supplement D2, Parts 1 and 2.* (Eds.: P. Patai, Z. Rappoport), Wiley-VCH, Weinheim, **1995**.
- Transition Metals for Organic Synthesis.* (Eds.: M. Beller, C. Bolm), Wiley-VCH, Weinheim, **2004**.

- 3 *Metal-Catalyzed Cross-Coupling Reactions*. (Eds.: A. de Meijere, F. Diederich), Wiley-VCH, Weinheim, **2004**.
- 4 D. A. Petrone, J. Ye, M. Lautens, *Chem. Rev.*, **2016**, *116*, 8003–8104.
- 5 a) A. H. Roy, J. F. Hartwig, *J. Am. Chem. Soc.*, **2001**, *123*, 1232–1233; b) A. H. Roy, J. F. Hartwig, *J. Am. Chem. Soc.*, **2003**, *125*, 13944–13945; c) J. F. Hartwig, *Inorg. Chem.*, **2007**, *46*, 1936–1947.
- 6 For examples of Pd-catalyzed carbohalogenations of alkynes, see: C. M. Le, P. J. C. Menzies, D. A. Petrone, M. Lautens, *Angew. Chem. Int. Ed.*, **2015**, *54*, 254–257; For a mechanistic report, see: T. Sperger, C. M. Le, M. Lautens, F. Schoenebeck, *Chem. Sci.*, **2017**, *8*, 2914–2922.
- 7 Bipyridine/phenanthroline in Ni-catalyzed radical containing reactions; for examples, see: a) D. R. Heitz, J. C. Tellis, G. A. Molander, *J. Am. Chem. Soc.*, **2016**, *138*, 12715–12718; b) X. Zhang, D. W. C. MacMillan, *J. Am. Chem. Soc.*, **2016**, *138*, 13862–13865; c) V. B. Phapale, E. Buñuel, M. García-Iglesias, D. J. Cárdenas, *Angew. Chem. Int. Ed.*, **2007**, *46*, 8790–8795; d) D. A. Everson, R. Shrestha, D. J. Weix, *J. Am. Chem. Soc.*, **2010**, *132*, 920–921; e) A. García-Domínguez, Z. Li, C. Nevado, *J. Am. Chem. Soc.*, **2017**, *139*, 6835–6838; f) M. Börjesson, T. Moragas, R. Martin, *J. Am. Chem. Soc.*, **2016**, *138*, 7504–7507; g) T. Qin, J. Cornella, C. Li, L. R. Malins, J. T. Edwards, S. Kawamura, B. D. Maxwell, M. D. Eastgate, P. S. Baran, *Science*, **2016**, *352*, 801–805; h) A. Klein, A. Kaiser, W. Wielandt, F. Belaj, E. Wendel, H. Bertagnolli, S. Zális, *Inorg. Chem.*, **2008**, *47*, 11324–11333; i) S. Biswas, D. J. Weix, *J. Am. Chem. Soc.*, **2013**, *135*, 16192–16197; j) K. Nakajima, S. Nojima, Y. Nishibayashi, *Angew. Chem. Int. Ed.*, **2016**, *55*, 14106–14110.
- 8 For mechanistic reports on radical-like reactivity of Ni/bpy, see a) B. J. Shields, B. Kudisch, G. D. Scholes, A. G. Doyle, *J. Am. Chem. Soc.*, **2018**, *140*, 3035–3039; b) B. J. Shields, A. G. Doyle, *J. Am. Chem. Soc.*, **2016**, *138*, 12719–12722.
- 9 M. V. Joannou, M. J. Bezdek, K. Albahily, I. Korobkov, P. J. Chirik, *Organometallics*, **2018**, ASAP. (DOI: 10.1021/acs.organomet.8b00350).
- 10 For a report on characterization by ESR of NiI(PEt₃)₃ at *g* = 2.180, see: T. T. Tsou, J. K. Kochi, *J. Am. Chem. Soc.*, **1979**, *101*, 6319–6332.
- 11 a) P. T. Matsunaga, G. L. Hillhouse, A. L. Rheingold, *J. Am. Chem. Soc.*, **1993**, *115*, 2075–2077; b) P. T. Matsunaga, J. C. Mavropoulos, G. L. Hillhouse, *Polyhedron* **1995**, *14*, 175–185; c) R. Han, G. L. Hillhouse, *J. Am. Chem. Soc.*, **1997**, *119*, 8135–8136; d) J. A. Terrett, J. D. Cuthbertson, V. Shurtleff, D. W. C. MacMillan, *Nature*, **2015**, *524*, 330–334.
- 12 The stoichiometric experiment using Ni(0)/dtbpy, iodobenzene (2 equiv) and 4-octyne (3 equiv) resulted in formation of product in 33% yield. These results also supported the formation of Ni(I) via comproportionation.
- 13 For representative examples of comproportionation of Ni(0)/Ni(II) in catalytic reaction, see: a) I. Kalvet, Q. Guo, G. J. Tizzard, F. Schoenebeck, *ACS Catal.* **2017**, *7*, 2126–2132. a) A. B. Dürr, H. C. Fisher, I. Kalvet, K.-N. Truong, F. Schoenebeck, *Angew. Chem. Int. Ed.* **2017**, *56*, 13431–13445. b) K. Matsubara, Y. Fukahori, T. Inatomi, S. Takazaki, Y. Koga, S. Kanegawa, T. Nakamura, *Organometallics* **2016**, *35*, 3281–3287. c) A. Velian, S. Lin, A. J. M. Miller, M. W. Day, T. Agapie, *J. Am. Chem. Soc.* **2010**, *132*, 6296–6297. d) R. Beck, S. A. Johnson, *Organometallics*, **2013**, *32*, 2944–2951. e) Beck, R.; Shoshani, M.; Krasinkiewicz, J.; Hatnean, J. A.; Johnson, S. A. *Dalton Trans.* **2013**, *42*, 1461–1475. f) A. Velian, S. Lin, A. J. M. Miller, M. W. Day, T. Agapie, *J. Am. Chem. Soc.* **2010**, *132*, 6296–6297.
- 14 An addition of TEMPO radical scavenger retarded the reaction. Radical probe experiment with acetylene bearing two cyclopropyl group resulted in formation of product in trace yield.
- 15 For a selected review, see: S. Z. Tasker, E. A. Standley, T. F. Jamison, *Nature*, **2014**, *509*, 299–309.



Odd-valent nickel-catalyzed intermolecular carboiodination of alkynes with aryl iodides to form highly substituted and functionalized alkenyl iodides has been developed.