



**Synthesis of novel copper-lanthanide BINOLate frameworks
from a hydrogen bonding DBU-H⁺ lanthanide BINOLate
complex**

Journal:	<i>Dalton Transactions</i>
Manuscript ID	DT-COM-08-2018-003335
Article Type:	Communication
Date Submitted by the Author:	15-Aug-2018
Complete List of Authors:	Panetti, Grace; University of Pennsylvania, Department of Chemistry Robinson, Jerome; Brown University, Chemistry Carroll, Patrick; University of Pennsylvania, Chemistry Gau, Michael; University of Pennsylvania, Department of Chemistry Manor, Brian; University of Pennsylvania, Chemistry Walsh, Patrick; UPenn, chem Schelter, Eric J; University of Pennsylvania, Department of Chemistry

Synthesis of novel copper-lanthanide BINOLate frameworks from a hydrogen bonding DBU-H⁺ lanthanide BINOLate complex

Received 00th January 20xx,
Accepted 00th January 20xx


Grace B. Panetti, Jerome R. Robinson, Patrick J. Carroll, Michael R. Gau, Brain C. Manor, Patrick J. Walsh,* and Eric J. Schelter*

DOI: 10.1039/x0xx00000x

www.rsc.org/

The preparation of a novel H-bonding DBU-H⁺ BINOLate-lanthanide complex enabled the synthesis of the first copper-Rare Earth Metal BINOLate complex (CuDBU-REMB). CuDBU-REMB was compared to the analogous Li complex using X-ray crystallography and Exchange NMR spectroscopy (EXSY). The results provide insight into the role of the secondary metal cation in the framework's stabilization.

Asymmetric catalysis is the most important method for the preparation of optically active compounds, which are crucial in pharmaceutical and organic materials applications.¹⁻³ Specifically, chiral Rare Earth metal (RE) complexes continue to be advantageous for a number of Lewis-acid catalysed reactions.⁴⁻⁶ The REMB framework [M⁺₃(THF)_n][(BINOLate)₃RE] (REMB; RE = Sc, Y, La-Lu; M = Li, Na, K; B = 1,1'-bi-2-naphtholate; RE/M/B/ = 1/3/3; **Scheme 1**), pioneered by Shibasaki, is among the most enantioselective heterobimetallic catalysts, where changing of M, RE, or substitution of BINOL enables diverse reactivity.^{5, 7} We have shown that this framework is not only robust, but variable beyond Shibasaki's original concept. While the central cation is typically a trivalent rare earth cation, this site can also be occupied by a tetravalent cerium or tetravalent uranium ion.⁸⁻¹¹ The cation in the secondary coordination sphere can also be replaced with alternatives to alkali metals to include ethyl zinc (ZnEt⁺), and 1,1,3,3-tetramethyl guanidinium (TMG-H⁺), where the latter functions as a hydrogen-bond donor.^{10, 12, 13} By substituting the secondary coordination metal with zinc, we were able to increase the reactions capable by the REMB framework to include enantioselective ethylation of aldehydes.¹² In light of this variability, we expected that copper(I) could also occupy the secondary coordination sphere, due to its chemical similarities to Zn(II). We were also interested in copper because copper complexes are known to




Scheme 1: Left: previously synthesized REMB frameworks, which all adopt the same basic structure; Right: This work introducing copper from the corresponding H-bonding framework.

catalyse a host of organic reactions.¹⁴⁻¹⁸ Thus, we envisioned that the introduction of copper into the secondary coordination sphere would increase the library of reactions available to the REMB framework.

The preparation of previously reported REMB complexes with Li, Na, K, or Cs can be accomplished using protonolysis or transmetalation reactions.^{10, 19} However, [(ZnEt)₃(THF)₃][(BINOLate)₃RE] can only be prepared cleanly from protonolysis of a [(BINOLateH)₃RE] species generated *in situ* (**Scheme 2A**).¹² Despite this observation, for the current work, application of protonolysis to copper *tert*-butoxide (CuOtBu)₄, or mesitylcopper (CuMes)_n resulted in complex mixtures of products. Inspired by our previous demonstration that [TMG-H⁺]₃[(BINOLate)₃RE] acts as an air-stable precursor to many alkali metal-containing REMB complexes, we hypothesized that the copper analogue could be synthesized from a similar hydrogen-bonded framework with 1,8-diazabicyclo(5.4.0)undec-7-ene (DBU). Herein we disclose the synthesis of [Cu₃(DBU)₃][(BINOLate)₃RE] (**2-RE**) from the analogous [DBU-H⁺]₃[(BINOLate)₃RE] (**1-RE**) hydrogen-bonded precursor.

P. Roy and Diana T. Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, 231 South 34th Street, Philadelphia, Pennsylvania 19104, USA.
Email: pwalsh@sas.upenn.edu, schelter@sas.upenn.edu

Electronic Supplementary Information (ESI) available: X-ray structures of **1-La**, **1-Pr**, **2-La**, **2-Pr**; NMR data for all compounds; and additional EXSY-NMR data See DOI: 10.1039/x0xx00000x



Scheme 2 A. Previous methods of synthesis of REMB frameworks B. Synthesis of complexes **1-RE** from the corresponding RE amide bases. C Synthesis of complexes **2-RE** from **1-RE**.

Complexes **1-RE** were prepared by treatment of THF solutions of (*S*)-BINOL and (DBU) with a THF solution of the corresponding $\text{RE}[\text{N}(\text{SiMe}_3)_2]_3$ (Scheme 2B). The complexes were isolated by crystallization in good yields (79–88%). X-ray diffraction quality crystals of **1-RE** were grown from THF solutions layered with pentane, revealing a 6-coordinate lanthanide cation. In the cases of **1-Pr** and **1-Eu**, three HDBU^+ cations engage in hydrogen bonding, while in **1-La** only one HDBU^+ cation engages in this interaction (Figure 1; Figure S1). The ^1H NMR spectra of **1-RE** exhibited 6 signals attributed to BINOLate resonances and 16 broad resonances attributed to diastereotopic DBU resonances indicating a pseudo- D_3 symmetric complex on the NMR timescale.

Treatment of complexes **1-RE** with three equivalents of $(\text{CuMes})_n$ in toluene resulted in a new set of BINOLate resonances that were assigned as complexes **2-RE**. Crystallization of this species from hot toluene layered with pentane led to the isolation of **2-RE** in 74–88% yields (Scheme 2C). The NMR spectra indicated a pseudo- D_3 symmetric complex, consistent with previously described REMB complexes (Scheme 2C). However, the X-ray structure revealed a complex with pseudo- C_3 symmetry, in which the copper atoms coordinate to one of the two BINOLate oxygen atoms (Figure 2). This bridging coordination mode resulted in a lengthening of the Ln–O bond by an average of 0.16 Å, compared to oxygen atoms bound only to the lanthanide ion.




Figure 1: Thermal ellipsoids of $[\text{DBU-H}^+]_3[(\text{BINOLate})_3\text{Eu}]$ **1-Eu** at the 30% probability level, representative of **1-RE**. C-H hydrogen atoms and disorder removed for clarity. Selected bond lengths (Å): Eu(1)–O(1) 2.275(5), Eu(1)–O(2) 2.331(5), Eu(1)–O(3) 2.247(5), Eu(1)–O(4) 2.261(6), Eu(1)–O(5) 2.317(5), Eu(1)–O(6) 2.334(5), O(2)–N(1) 2.753(8), O(5)–N(3) 2.792(9), O(6)–N(5) 2.701(10). Selected angles (deg) N(1)–H(1)–O(1) 175.33, N(3)–H(3a)–O(5) 166.32, N(5)–H(5a)–O(6) 170.72.




Figure 2: Thermal ellipsoids of $[\text{Cu}(\text{DBU})]_3[(\text{BINOLate})_3\text{Eu}]$ **2-Eu** at the 30% probability level, representative of **2-RE**. Disorder, hydrogen atoms, and interstitial solvent removed for clarity. Selected bond lengths (Å): Eu(1)–O(1) 2.256(8), Eu(1)–O(2) 2.410(8), Eu(1)–O(3) 2.202(9), Eu(1)–O(4) 2.400(8), Eu(1)–O(5) 2.362(8), Eu(1)–O(6) 2.284(7), Cu(1)–O(2) 1.820(8), Cu(1)–N(5) 1.868(10), Cu(2)–O(4) 1.885(9), Cu(2)–N(1) 1.859(14), Cu(3)–O(5) 1.825(9), Cu(3)–N(3) 1.820(12). Selected angles (deg) N(1)–Cu(2)–O(4) 177.6(6), N(3)–Cu(3)–O(5) 169.2(4), N(5)–Cu(1)–O(2) 164.6(5).

In previous REMB frameworks, the secondary-coordination sphere metal was at least 3-coordinate, bound to two BINOLate oxygen atoms. However, in complexes **2-RE**, the copper atom was only 2-coordinate ($\angle\text{N-Cu-O} > 164^\circ$, closest non-bonding atom-Cu $> 2.8 \text{ \AA}$). Despite the wide use of DBU as a base in catalysis, there is only one other example of a Cu-DBU coordination characterized by X-ray diffraction.²⁰ The previously described Cu-DBU complex is a 3-coordinate Cu atom with two equivalents of DBU coordinated and one iodide atom.

To investigate the lability of the ligands in solution, an important feature of RE catalysis, we turned to EXSY-NMR to determine the ligand exchange rates, as previously demonstrated by our group.^{11, 21} The technique has found use for determining the exchange rates of ligands in a multitude of different catalytic systems.²² For EXSY-NMR to be applicable the two species in exchange must have sufficient spectral resolution.^{22, 23} Therefore, we chose to use a mixture of **2-Pr** and **2-Eu**, due to their solid state structures being isostructural, relatively short relaxation times, and the rare earth cations' anisotropic g-tensors having opposite signs.^{3, 24} In THF-*d*₈, the rate of transfer of BINOL ligands between **2-Pr** and **2-Eu** was 2.63 s^{-1} (300 K) (**Table 1**). Using variable temperature EXSY experiments, we also determined the activation parameters (**Table 1**).

Compared to the previously measured Li-REMB (0.759 s^{-1} for Li-REMB), we found that the BINOLate ligands of **2-RE** were more labile in solution.²¹ Likewise, the free energy, enthalpic barriers, and entropic barriers we measured for the exchange between **2-Pr** and **2-Eu** were lower than the analogous Li-REMB (**Table 1**). We postulate that the increased exchange rate of the BINOLate ligands was due to fewer metal-ligand interactions, in the secondary coordination sphere, increasing the lability of BINOLate ligands. In the case of Li-REMB, the rate of ligand exchange is already faster than the rate of catalysis in the Michael and Henry reactions.²¹

In summary, we have demonstrated that the isolation of a hydrogen-bonding REMB scaffolding enabled the synthesis of a Cu-REMB compound using simple ligands. The copper-containing complex was found to adopt an uncommon 2-coordinate geometry in the secondary coordination sphere. EXSY NMR studies show that the BINOLate ligands exchanged faster than the previously studied Li-REMB complex. We expect that these results will inspire use of hydrogen-bonding networks to build multi-metallic complexes. Future work will explore the utility of these complexes as catalysts.

Table 1: Associated EXSY Rate Data and Activation Parameters for **2-Pr/2-Eu**

	Cu-REMB ^a		Li-REMB ^b	
	2-Pr → 2-Eu	2-Eu → 2-Pr	Pr → Eu	Eu → Pr
$k'(\text{s}^{-1})^c$	2.63 ± 0.10^d	2.63 ± 1.02^d	0.759	0.806
$\Delta H^\ddagger (\text{kcal}\cdot\text{mol}^{-1})^e$	11.1 ± 2.3	9.0 ± 1.1	15.1	14.6
$\Delta S^\ddagger (\text{eu})^e$	-19.8 ± 7.8	-26.4 ± 2.7	-35.6	-41.3
$\Delta G^\ddagger (\text{kcal}\cdot\text{mol}^{-1})^e$	11.7 ± 2.3	9.7 ± 1.1	17.7	17.6

^a Values are reported as an average of 3 independently prepared samples ^b Previously reported values²¹ ^c At 300 K, $[\text{2-Pr}]/[\text{2-Eu}] = 15.0/14.6 \text{ mM}$. k' values determined using EXSYCALC. ^d Values obtained from H₂ at $t_{\text{mix}} = 170 \text{ ms}$. ^e At 298 K. Additional data can be found in the Supplemental Information.

Conflicts of interest

The authors declare no conflict of interests.


Notes and references

- E. N. Jacobson, A. Pfaltz, and H. Yamamoto *Comprehensive Asymmetric Catalysis*, Springer, New York, 1999 (supplement 2004).
- P. J. K. Walsh and M. C. Kozlowski, *Fundamentals of Asymmetric Catalysis*, University Science Books, Sausalito, CA, 2008.
- W. D. Horrocks and J. P. Sipe, *J. Am. Chem. Soc.*, 1971, **93**, 6800-6804.
- S. Matsunaga and M. Shibasaki, *Chem. Commun.*, 2014, **50**, 1044-1057.
- M. Shibasaki, M. Kanai, S. Matsunaga, and N. Kumagai, *Accounts Chem. Res.*, 2009, **42**, 1117-1127.
- J. Park and S. Hong, *Chem. Soc. Rev.*, 2012, **41**, 6931-6943.
- N. Kumagai, M. Kanai and H. Sasai, *ACS Catal.*, 2016, **6**, 4699-4709.
- J. R. Robinson, P. J. Carroll, P. J. Walsh and E. J. Schelter, *Organometallics*, 2013, **32**, 1493-1499.
- J. R. Robinson, P. J. Carroll, P. J. Walsh, and E. J. Schelter, *Angew. Chem. Int. Edit.*, 2012, **51**, 10159-10163.
- J. R. Robinson, Z. Gordon, C. H. Booth, P. J. Carroll, P. J. Walsh and E. J. Schelter, *J. Am. Chem. Soc.*, 2013, **135**, 19016-19024.
- J. R. Robinson, Y. Qiao, J. Gu, P. J. Carroll, P. J. Walsh and E. J. Schelter, *Chem. Sci.*, 2016, **7**, 4537-4547.
- I. Nieto, A. J. Wooten, J. R. Robinson, P. J. Carroll, E. J. Schelter and P. J. Walsh, *Organometallics*, 2013, **32**, 7431-7439.
- J. R. Robinson, X. Fan, J. Yadav, P. J. Carroll, A. J. Wooten, M. A. Pericàs, E. J. Schelter and P. J. Walsh, *J. Am. Chem. Soc.*, 2014, **136**, 8034-8041.
- X.-X. Guo, D.-W. Gu, Z. Wu and W. Zhang, *Chem. Rev.*, 2015, **115**, 1622-1651.
- C. Deutsch, N. Krause and B. H. Lipshutz, *Chem. Rev.*, 2008, **108**, 2916-2927.
- G. Evano, N. Blanchard and M. Toumi, *Chem. Rev.*, 2008, **108**, 3054-3131.
- S. Bhunia, G. Pawar Govind, S. V. Kumar, Y. Jiang and D. Ma, *Angew. Chem. Int. Edit.*, 2017, **56**, 16136-16179.
- A. P. Pulis, K. Yeung and D. J. Procter, *Chem. Sci.*, 2017, **8**, 5240-5247.
- A. J. Wooten, P. J. Carroll and P. J. Walsh, *J. Am. Chem. Soc.*, 2008, **130**, 7407-7419.

COMMUNICATION

Journal Name

20. R. Watari, Y. Kayaki, S. i. Hirano, N. Matsumoto and T. Ikariya, *Adv. Synth. Cat.*, 2015, **357**, 1369-1373.
21. J. R. Robinson, J. Gu, P. J. Carroll, E. J. Schelter and P. J. Walsh, *J. Am. Chem. Soc.*, 2015, **137**, 7135-7144.
22. C. L. Perrin and T. J. Dwyer, *Chem. Rev.*, 1990, **90**, 935-967.
23. I. Bertani and C. Luchinat, *Coordination Chemistry Reviews*, 1996, **150**, 185-220.
24. A. F. Cockerill, G. L. O. Davies, R. C. Harden and D. M. Rackham, *Chem. Rev.*, 1973, **73**, 553-588.



38x19mm (300 x 300 DPI)