



The Serendipitous Discovery of a Readily Available Redox-Bistable Molecule Derived from Cyclic(alkyl)(amino)carbenes

Journal:	<i>Organic Chemistry Frontiers</i>
Manuscript ID	QO-RES-05-2018-000447.R1
Article Type:	Research Article
Date Submitted by the Author:	18-May-2018
Complete List of Authors:	<p>Mahoney, Janell; UC San Diego, Chemistry Regnier, Vianney; Universite Grenoble Alpes, Department of Molecular Chemistry, CNRS, DCM (UMR 5250) Romero, Erik; UC San Diego, Chemistry Molton, Florian; Université Grenoble Alpes, Département de Chimie Moléculaire Royal, Guy; Universite Grenoble Alpes, Département de Chimie Moléculaire (UMR CNRS 5250) jazzar, rodolphe ; UCSD, chemistry Martin, David; Universite Grenoble Alpes, Department of Molecular Chemistry, CNRS, DCM (UMR 5250) Bertrand, Guy; UC San Diego, Chemistry</p>

The Serendipitous Discovery of a Readily Available Redox-Bistable Molecule Derived from Cyclic(alkyl)(amino)carbenes


Janell K. Mahoney,^{a,b} Vianney Regnier,^b Erik A. Romero,^a Florian Molton,^b Guy Royal,^b
Rodolphe Jazzar,^a David Martin*^b and Guy Bertrand*^a

^aUCSD-CNRS Joint Research Laboratory (UMI 3555), Department of Chemistry and Biochemistry, University of California, San Diego, La Jolla, California 92093-0358, United States.

^bUMR CNRS 5250, Département de Chimie Moléculaire, Université Grenoble-Alpes, CS 40700, 38058 GRENoble cedex 9, France

Abstract: Bis(azoliums) are readily available in one step from cyclic (alkyl)(amino)carbenes and bis(acyl chlorides). A two-electron reduction of the bis(azolium), featuring a *gem*-(dimethyl)malonoyl spacer, leads to the corresponding transient diradical, which undergoes an intramolecular cyclization. The latter can be re-oxidized at a higher potential to yield back the bis(azolium). The redox bistability of this simple organic molecular system is linked to the formation of a weak C–O bond (27 kcal mol⁻¹). Both redox forms can be isolated and stored for months without evidence of decay.


Graphical Abstract:



Redox bistable molecules (RBMs) have two oxidation states that can co-exist within a certain range of electrochemical potentials. They are designed based on the so-called square scheme,¹ in which the reduction (or oxidation) of the molecule triggers a reversible chemical transformation such as a conformational change, isomerism, or intramolecular bond formation. RBMs have attracted considerable attention due to their potential applications in molecular electronics.² For example, bistability prevents entropy-driven mutual electron exchanges between neighbors and therefore is critical for redox-based data storage at a molecular scale.³ Redox bistable molecules have also been used in the context of bottom-up approaches for the design of nano-sized devices.⁴

The vast majority of reported RBMs feature coordinated metals⁵ and/or complex supramolecular structures, such as rotaxanes and catenanes.⁶ In contrast, only a handful of simple organic patterns are known.⁷ They generally^{8,9} undergo a reversible intramolecular σ -bond formation upon electron transfer. The corresponding square scheme involves a diradical, which is formed upon a 2-electron oxidation of an activated diene (Scheme 1).


The coupling of the two radical units has to be exothermic enough to induce significant stability of the σ -bond, but not too much in order to preserve the chemical reversibility of the process. A rough qualitative analysis of square schemes indicates that the corresponding bond dissociation energies should ideally be in the 10 – 40 kcal mol⁻¹ range. Such low values are rarely encountered in standard organic species, which explains the scarcity of simple organic RBM representatives. Note that accessing such a non-standard σ -bond often requires a complex synthesis, which represents a serious drawback for further tuning and applications. Herein we report that the reversible formation of a carbon-oxygen σ -bond between two radical units led to the serendipitous discovery of readily available redox-bistable organic systems.




Scheme 1. Some organic patterns featuring redox bistability and the associated square scheme.

The combination of an electron-donating and an electron withdrawing substituent stabilizes C-centered radicals through capto-dative substitution,¹⁰ but usually not to such an extent that allows for the isolation of monomeric species. For example, only the corresponding C-C homodimer could be isolated in the case of the 2-oxomorpholin-3-yl radical **A** (Scheme 2).^{11,12} However, we recently

demonstrated that (amino)(carboxy)carbon radicals **B**, derived from cyclic (alkyl)(amino)carbenes (CAACs),^{13,14} can be isolated.^{15,16}



Scheme 2. Dimerization of capto-dative radical **A** and isolated CAAC-based radicals **B**.



Scheme 3. Synthesis of **1a²⁺** and **2a**.

Since paramagnetic species of type **B** are readily available in two steps from the reaction of a stable CAAC with acyl chlorides, followed by a one-electron reduction, we extrapolated this synthetic methodology to the preparation of diradicals¹⁷ such as **2a** (Scheme 3). The latter is stable at room temperature and has been characterized by an X-ray diffraction study.^{15a} The two radical units are not conjugated, as they correspond to π -systems that are orthogonal to the phenyl spacer. In addition, they are separated by about 700 pm, which limits through-space interactions. Therefore, it can be predicted that the first triplet and singlet states are nearly degenerate. In order to obtain a precise picture, low temperature EPR experiments were carried out. A frozen acetonitrile solution at 4 K displayed a weak half-field signal, which decreased when increasing the temperature, thus indicating the triplet nature of the ground state (Figure 1). The half-field band could hardly be detected above 10 K. Fitting our data with the Bleaney-Bowers equation¹⁸ showed that the triplet-singlet gap is less than 0.05 kcal mol⁻¹ (see supporting information).





Figure 1. X-band EPR spectra of diradical **2a** in frozen acetonitrile with zoom on the half-field absorption signals ($\Delta m = \pm 2$) at 5 K (left) and main transition ($\Delta m = \pm 1$) at 20 K (right).

To study the influence of the spacer on the singlet-triplet gap, we prepared the dication $\mathbf{1b}^{2+}$ from 1,1'-ferrocene dicarbonyl chloride in 93 % yield (Scheme 4). Cyclic voltammetry of $\mathbf{1b}^{2+}$ indicated two reversible reductions at $E_{1/2} = -0.73$ and -0.93 V (Figure 2), which were attributed to the successive formation of the two radical centers. Further reversible reductions below -1.5 V were attributed to the formation of the corresponding bis(enolate). We performed the quantitative electrolysis of a solution of $\mathbf{1b}^{2+}$ in acetonitrile at $E_{app} = -1.35$ V. The stoichiometry (two coulombs per mole of reagent) was consistent with two successive one-electron processes. Cyclic voltammograms of the resulting solution were similar to those of $\mathbf{1b}^{2+}$ indicating that, indeed, diradical $\mathbf{2b}$ was formed. The presence of a half-field band in the EPR spectra of frozen samples indicated the triplet nature of the ground state. However, in marked contrast with $\mathbf{2a}$, this signal could be easily detected from 4 K up to 80 K. On this broad range of temperature (T), the integrated signal (I), which is proportionate to the magnetization of the sample, followed the Curie law ($IT = \text{constant}$), thus indicating a more significant ferromagnetic coupling of the two spins than in $\mathbf{2a}$, with a triplet-singlet gap of at least $0.5\text{--}1$ kcal mol $^{-1}$. In addition, the main band features a zero-field splitting parameter $D \sim 42$ G. This value corresponds to an average distance of more than 8.5 Å between the two interacting $\frac{1}{2}$ spins,^[17] suggesting that the two radical centers are rather on opposite sides of the ferrocene linker, as depicted on Scheme 4. At higher temperatures, we observed the irreversible formation of a $\frac{1}{2}$ spin radical, likely due to the reaction of $\mathbf{2b}$ with oxygen, which was inevitably condensed when cooling the EPR tube.



Scheme 4: Synthesis of $\mathbf{1b}^{2+}$ and $\mathbf{2b}$.

Next, we synthesized dication $\mathbf{1c}^{2+}$ featuring an even shorter spacer (Scheme 5). This compound is perfectly stable as a solid, although slightly moisture sensitive in solution. An X-ray diffraction study of the tetrafluoroborate salt of $\mathbf{1c}^{2+}$ confirmed its structure (Figure 3).





Figure 2. Cyclic voltammograms of **1b**²⁺ in acetonitrile + 0.1M [Bu₄N]PF₆, scan rate: 100 mV s⁻¹ (left); X-band EPR spectra of **2b** in frozen acetonitrile (right) with zoom on the half-field absorption signals ($\Delta m = \pm 2$) at 5 K (top) and main transition ($\Delta m = \pm 1$) at 20 K (bottom).

The cyclic voltammogram of **1c**²⁺ is characterized by an irreversible reduction wave at $E_{pc} = -0.77$ V, with the corresponding re-oxidation wave at $E_{pa} = -0.16$ V (Figure 3). A quantitative electrolysis at $E_{app} = -1.1$ V showed that this reduction corresponds to a two-electron process. The resulting compound **3c** featured a similar cyclic voltammetry and, in turn, yielded back **1c**²⁺ upon quantitative oxidation at $E_{app} = +0.2$ V. We then performed the chemical reduction of **1c**²⁺ with zinc powder, which allowed for the isolation of **3c** in 92% yield. Both **1c**²⁺ and **3c** are EPR silent. An X-ray diffraction study revealed the spirocyclic structure of **3c**. Note that we also reacted **3c** with two equivalents of silver triflate and isolated the triflate salt of **1c**²⁺ in 87 % yield.



Scheme 5. Synthesis of **1c**²⁺ and **3c**.





Figure 3. Cyclic voltammogram of dication $1\mathbf{c}^{2+}$ in acetonitrile + 0.1M $[\text{Bu}_4\text{N}]\text{PF}_6$, scan rate: 100 mV s^{-1} (top, left) and of its reduced form $3\mathbf{c}$ (bottom, left), and solid-state structures of $1\mathbf{c}^{2+}$ (top, right) and $3\mathbf{c}$ (bottom, right) with 50 % ellipsoid probability. Hydrogen atoms, anions and solvent molecules were omitted for clarity.

It is important to note that the C-O cyclization is an extremely fast process. Indeed, even at wide sweep rate range (up to 12.8 V s^{-1}), no reversibility of either electrochemical wave could be evidenced (see supporting information). Similarly, even at -40°C , the cyclic voltammograms still showed irreversible waves at $E_{pa} = 0.03\text{ V}$ (oxidation) and $E_{pc} = -0.65\text{ V}$ (reduction).

Interestingly, capto-dative carbon radicals are traditionally depicted as C-centered radicals, which afford C-C homodimers as exemplified by the 2-oxomorpholin-3-yl radical **A** (Scheme 1). Although C-O homodimers have never been reported so far, their formation should not be surprising as (amino)(carboxy)carbon radicals also bear significant spin density on the oxygen atom. We considered the three possible dimerization products for model radical **I** at the B3LYP/6-311g** level of theory (Scheme 6).^{19,20} As expected, we found that the formation of the O-O dimer \mathbf{I}_{OO} is highly exothermic in enthalpy and endergonic in Gibbs energy, and that the C-C dimerization affording \mathbf{I}_{CC} is the most favorable process. Importantly, the formation of the C-O dimer \mathbf{I}_{CO} was found to be exothermic (-11 kcal mol^{-1}) but slightly endergonic ($+3.5\text{ kcal mol}^{-1}$). In the case of a diradical such as $2\mathbf{c}$, The C-O bond formation corresponds to an intramolecular process, in which most of the entropic contribution favoring dissociation is suppressed. Indeed, the ring opening of $3\mathbf{c}$ to afford diradical $2\mathbf{c}$ was found to be endergonic by $+27\text{ kcal mol}^{-1}$. This preliminary value corresponds to the homolytic cleavage of a weak C-O bond, perfectly in line with the observed redox-bistability of the system. Note that cyclization through the formation of a C-C bond would require the interpenetration of bulky substituents and no minimum on the hypersurface of energy could be found for the corresponding product.



Scheme 6: Energetics for intermolecular dimerization of model radical **I**.

In conclusion, **1c²⁺** was synthesized in one simple step from a cyclic (alkyl)(amino)carbene, which is itself readily available on large scales.¹⁴ The two-electron reduction triggers a cyclization, which accounts for the redox bistability of the system. Both redox forms could be isolated and stored for months without evidence of decay. We are currently combining in-depth DFT studies, as well as experimental evaluation of a variety of spacers and stable carbenes to determine the scope of this new approach towards simple organic redox-bistable molecules.

Corresponding Authors

Dr. David Martin; email: david.martin@univ-grenoble-alpes.fr

Pr Guy Bertrand; email: guybertrand@ucsd.edu

Conflicts of interest: The authors declare no competing financial interests.

Electronic Supplementary Information (ESI) available: ¹H and ¹³C NMR spectra of all new compounds, X-ray crystallographic data, VT-EPR plots, electrochemical study of **1c²⁺** and computational details.

Acknowledgement:

This work was supported by the NSF (CHE-1661518), the French National Agency for Research (ANR-14-CE06-0013-01 and ANR-17-ERC2-0015) and the French-American cultural ex-change foundation (FACE). Thanks are due to the “Centre de Calcul Intensif en Chimie de Grenoble” and the ICMG Chemistry Nanobio Platform of Grenoble. J. K. M. and E. A. R. benefited from a Chateaubriand Fellowship of the Office for Science & Technology of the Embassy of France in the United States and a GAANN fellowship from the US Department of Education, respectively.

Experimental Section

General considerations: All manipulations were performed under an inert atmosphere of dry argon, using standard Schlenk and drybox techniques. Dry and oxygen-free solvents were employed. Stable

carbenes were prepared as previously reported.¹⁴ 1,10-Ferrocene-di(carbonyl chloride) was prepared freshly in a two step sequence from 1,10-Diacetylferrocene as previously reported.²¹ Commercially available dimethylmalonyl chloride was purified by distillation before use. ¹H, ¹³C NMR spectra were recorded on Bruker Advance 300, Varian VX 500 and Jeol ECA 500 spectrometers. Chemical shifts are given relative to SiMe₄ and referenced to the residual solvent signal (¹H, ¹³C). Melting points were measured with Büchi Melting Point B-545 apparatus. Electrochemical experiments were carried out in CH₃CN + 0.1M [Bu₄N]PF₆ using a Bio-logic SP-300 potentiostat. Potentials were first referred to an Ag/0.01M AgNO₃ reference electrode in CH₃CN + 0.1M [Bu₄N]PF₆, and all redox potentials in the text are reported with respect to the E_{1/2} of the ferrocene/ferrocenium (Fc⁺/Fc) redox couple (Ferrocene was used as a standard, the Fc⁺/Fc potential is 0.07 V under our experimental conditions). The working electrode was a vitreous carbon disk (3 mm in diameter; polished with 1 μm diamond paste before each record) and the auxiliary electrode was a platinum wire. Bulk electrolysis were performed using reticulated vitreous carbon electrode as working electrode (A = 2 cm²). EPR spectra were recorded on a Bruker EMX+ spectrometer.

Synthesis of bis(tetrafluoroborate) salt of di(iminium) 1b²⁺: (NB: The following manipulations were performed in the absence of light as the compound was found to be light sensitive.) To a mixture of 1,10-Ferrocene-di(carbonylchloride) (0.300 mg, 0.95 mmol) and cyclic (alkyl)(amino)carbene (0.750 mg, 2.41 mmol) was added 5 ml of THF. The solution was stirred for 30 minutes and then filtered. After evaporation of the solvent, the remaining solid was washed with THF (5 mL), Et₂O (5 mL) and pentane (5 mL) to yield the bis(chloride) salt of **1b²⁺** as a purple powder (841 mg; 93 % yield). The anion exchange from chlorine to BF₄ was achieved by stirring **1b²⁺Cl²⁻** with excess NaBF₄ (4 eq.) at room temperature in CH₃CN for 12h (NB: to avoid decomposition during the anion exchange the NaBF₄ salt was melted twice under vacuum to remove adventitious water). After removal of the solvent, the BF₄ salt was extracted with dichloromethane affording, after filtration and evaporation, the title compound as a purple solid in nearly quantitative yield. mp: 228-234 °C (dec.) HRMS (m/z): calc. for C₅₆H₇₈FeN₂O₂⁺: 866.5413; found [M⁺], 866.5412. ¹H NMR (CD₃CN, 500 MHz): δ 0.77 (d, J = 6.00 Hz, 12H), 1.24-1.28 (m, 24H), 1.52 (s, 12H), 2.16-2.23 (m, 4H), 2.30-2.37 (m, 4H), 2.49 (sept, J = 7.00 Hz, 12H), 2.54 (s, 4H), 4.98 (br s, 4H), 5.02 (br s, 4H), 7.29 (d, J = 8.00 Hz, 4H), 7.47 (t, J = 8.00 Hz, 2H). ¹³C{¹H} NMR (CD₃CN, 125 MHz): δ 9.8, 25.2, 26.9, 29.6, 30.2, 31.4, 41.1, 59.6, 76.0, 76.1, 78.1, 78.2, 79.1, 87.0, 127.5 (br), 128.3, 132.9 (br), 145.8, 192.1, 194.9.

Synthesis of bis(tetrafluoroborate) salt of di(iminium) 1c²⁺: Dimethylmalonyl dichloride (99 μL, 0.75 mmol) was added to the cyclic (alkyl)(amino)carbene (0.45 g, 1.58 mmol) in solution in THF (10 mL). After stirring for twenty minutes, the solvent was removed by filtration and the precipitate washed with diethyl ether (2 x 10mL). The solid was then dissolved in water, and an aqueous solution of sodium tetrafluoroborate (0.69 g, 6.31 mmol) was added to the flask. After stirring for one hour, the

product was extracted in dichloromethane. The organics were washed with brine and dried over MgSO₄ before removing the solvent under vacuum. (272 mg; 43% yield). Single crystals were grown by slowly diffusing diethyl ether into a concentrated acetonitrile solution of the product. mp: 146–148°C. HRMS (m/z): calc. for C₄₅H₆₈N₂O₂²⁺, 334.2635; found [M²⁺], 334.2637. ¹H NMR (CDCl₃, 500 MHz): δ 1.01-1.04 (br, mult, 16H), 1.12-1.27 (br, mult, 22H), 1.47-1.50 (br, mult, 5H), 1.63-1.72 (br, 8H), 1.86 (br, 11H), 2.26-2.35 (br, 4H), 3.67 (br, 4H), 7.26-7.29 (br, mult, 4H), 7.52 (t, J = 8.00 Hz, 2H). ¹³C{¹H} NMR (CDCl₃, 125 MHz): δ 19.7, 25.5, 25.6, 25.8, 26.3, 26.7, 26.8, 27.0, 28.9, 29.3, 30.6, 47.6, 62.1, 89.0, 127.4 (br), 128.0 (br), 128.8, 133.5, 144.6 (br), 146.5 (br), 194.8, 200.1.

Synthesis of compound 3c: Zinc (20 mg, 0.3 mmol) and **1c**²⁺ (126 mg, 0.15 mmol) were combined under argon, and acetonitrile was added to the flask. After 10 minutes, the solvent was removed *in vacuo*. Subsequent extraction with hexanes (5.0 mL) led to the isolation of a white powder (93 mg; 92% yield). Single crystals for X-ray crystallography were isolated through slow evaporation of a dichloromethane solution of the product. Only the E isomer could be evidenced. mp: 217-221°C. HRMS (m/z): calc. for C₅₄H₆₈N₂O₂, 669.0336; found, [M+Na]⁺ = 691.5172. ¹H NMR (CDCl₃, 300 MHz): δ 7.18-7.11 (m, 3H), 7.08-6.99 (m, 3H), 4.05 (sept, J = 6 Hz, 1H), 3.59 (sept, J = 6 Hz, 1H), 3.53 (m, 2H), 2.14 (d, J = 12 Hz, 1H), 1.79 (s, 3H), 1.73 (s, 3H), 1.55 (s, 3H), 1.48 (s, 3H), 1.46 (d, J = 7 Hz, 1H), 1.32-1.29 (m, 10H), 1.19-1.10 (m, 23H), 0.81 (s, 3H), 0.27 (s, 3H). ¹³C NMR (CDCl₃, 125 MHz): δ 213.6 (C), 154.1 (C), 151.6 (C), 148.4 (C), 147.8 (C), 144.3 (C), 142.2 (C), 136.3 (C), 133.6 (C), 127.5 (CH), 125.9 (CH), 125.1 (CH), 124.5 (CH), 107.3 (C), 64.2 (C), 63.6 (C), 58.8 (CH₂), 55.7 (CH₂), 49.7 (C), 45.7 (C), 43.0 (C), 32.9 (CH₃), 30.6 (CH₃), 29.6 (CH₃), 29.4 (CH₃), 28.6 (CH), 28.5 (CH₃), 28.2 (CH), 27.9 (CH), 27.1 (CH₃), 26.9 (CH₃), 26.7 (CH₃), 26.7 (CH₃), 26.6, 26.4, 26.1, 25.9, 25.7, 25.0, 24.0, 21.9.

Synthesis of bis(triflate salt of 1c²⁺ from compound 3c : Silver triflate (69 mg, 0.3 mmol) and **3c** (100 mg, 0.15 mmol) were combined under argon. Acetonitrile (5 mL) was added to the flask and the mixture was stirred for 20 minutes. The supernatant was then removed via filtration. The solvent was removed *in vacuo* and the solid washed with hexanes to yield **1c**²⁺ (121 mg; 83% yield).

References:

- (1) (a) Jack, J., *Electroanal. Chem. Interfacial Electrochem.*, 1971, **29**, 149. (b) J. H. R Tucker and S. R. Collisson, *Chem. Soc. Rev.*, 2002, **31**, 147.

(2) (a) Y. C. Liang, A. S. Dvornikov and P. M. Rentzepis, *Proc. Natl. Acad. Sci.*, 2003, **100**, 8109. (b) A. H. Flood, J. F. Stoddart, D. W. Steuerman and J. R. Heath, *Science*, 2004, **306**, 2055. (c) K. Galatsis, K. Wang, Y. Botros, Y. Yang, Y. H. Xie, J. F. Stoddart, R. B. Kaner, C. Ozkan, J. L. Liu, M. Ozkan, C. W. Zhou and K. W. Kim, *IEEE Circuits & Devices*, 2006, **22**, 12. (d) J. E. Green, J. W. Choi, A. Boukai, Y. Bunimovitch, E. Johnston-Halperin, E. Delonno, Y. Luo, B. A. Sheriff, K. Xu, Y. S. Shin, H.-R. Tseng, J. F. Stoddart and J. R. Heath, *Nature*, 2007, **445**, 414.

(3) (a) F. Franceschi, E. Solari, R. Scopelliti and C. Floriani, *Angew. Chem. Int. Ed.*, 2000, **39**, 1685. (b) C. Floriani, E. Solari, F. Franceschi, R. Scopelliti, P. Belanzoni and M. Rosi, *Chem. Eur. J.*, 2001, **7**, 3052. (c) K. Venkatesan, O. Blacque and H. Berke, *Organometallics*, 2006, **25**, 5190.

(4) For examples, see: (a) C. P. Collier, E. W. Wong, M. Belohradský, F. M. Raymo, J. F. Stoddart, P. J. Kuekes, R. S. Williams and J. R. Heath, *Science*, 1999, **285**, 391. (b) X. Duan, Y. Huang and C. M. Lieber, *Nano Lett.*, 2002, **2**, 487. (c) M. Feng, X. Guo, X. Lin, X. He, W. Ji, S. Du, D. Zhang, D. Zhu and H.-J. Gao, *J. Am. Chem. Soc.*, 2005, **127**, 15338. (d) D. X. Shi, Y. L. Song, H. X. Zhang, P. Jiang, S. T. He, S. S. Xie, S. J. Pang and H.-J. Gao, *Appl. Phys. Lett.*, 2000, **77**, 3203.

(5) For some representative examples see: (a) D. A. Cummings, J. McMaster, A. L. Rieger and P. H. Rieger, *Organometallics*, 1997, **16**, 4362. (b) A. J. L. Pombeiro, M. F. C. Guedes da Silva and M. Lemos, *Coord. Chem. Rev.*, 2001, **219**, 53. (c) D. Ruiz-Molina, K. Wurst, D. N. Hendrickson and C. Rovira, *Adv. Funct. Mater.*, 2002, **12**, 347. (d) C. Bucher, J. C. Moutet, J. Pecaut, G. Royal, E. Saint-Aman and F. Thomas, *Inorg. Chem.*, 2004, **43**, 3777. (e) E. Evangelio, D. N. Hendrickson and D. Ruiz-Molina, *Inorg. Chim. Acta*, 2008, **361**, 3403. (f) E. Evangelio, M.-L. Bonnet, M. Cabanas, M. Nakano, J.-P. Sutter, A. Dei, V. Robert and D. Ruiz-Molina, *Chem. Eur. J.*, 2010, **16**, 6666. (g) M. Kepenekian, V. Vetere, B. Le Guennic, P. Maldivi and V. Robert, *Chem. Eur. J.*, 2011, **17**, 12045. (h) M. Bubrin, D. Schweinfurth, F. Ehret, S. Zaalis, H. Kvapilova, J. Fiedler, Q. Zeng, F. Hartl and W. Kaim, *Organometallics*, 2014, **33**, 4973.

(6) For reviews see: (a) *Electrochemistry of Functional Supramolecular systems*, Ed. P. Ceroni, A. Credi and M. Venturi, John Wiley & Sons, Hoboken, 2010. (b) C. Tock, J. Frey and J. P. Sauvage, In *Molecular Switches*, Ed. B. L. Feringa and R. W. Browne, Wiley-VCH, Weinheim, 2011, Vol. 1, pp. 97–120. (c) S. Saha and L. F. Stoddart, *Functional Org. Mater.*, 2007, **17**, 295. (d) A. C. Fahrenbach, C. J. Bruns, D. Cao and J. F. Stoddart, *Acc. Chem. Res.*, 2012, **45**, 1581. (e) R. Antonio, U. Chiacchio, A. Corsaro and G. Romeo, *Curr. Org. Chem.*, 2012, **16**, 127. (f) A. C. Fahrenbach, C. J. Bruns, H. Li, A. Trabolsi, A. Coskun and F. J. Stoddart, *Acc. Chem. Res.*, 2014, **47**, 482.

(7) (a) M. Horner and S. Hunig, *J. Am. Chem. Soc.*, 1977, **99**, 6120. (b) M. Horner and S. Hunig, *J. Am. Chem. Soc.*, 1977, **99**, 6122. (c) T. Suzuki, K. Wada, Y. Ishigaki, Y. Yoshimoto, E. Ohta, H. Kawai and K. Fujiwara, *Chem. Commun.*, 2010, **46**, 4100. (d) E. Ohta, H. Uehara, Y. Han, K. Wada, H. Noguchi, R. Katoono, Y. Ishigaki, H. Ikeda, Uosaki and T. Suzuki, *ChemPlusChem*, 2017, **82**, 1043. (e) J. Nishida, T. Miyagawa and Y. Yamashita, *Org. Lett.* 2004, **6**, 2523.

(8) For a rare example of an organic molecule whose redox bistability stems from a conformation change, see: O. Ivashenko, H. Logtenberg, J. Areephong, A. C. Coleman, P. V. Wesenhagen, E. M. Geertsema, N. Heureux, B. L. Feringa, P. Rudolf and W. R. Browne, *J. Phys. Chem. C*, 2011, **115**, 22965.

(9) Note that weakly bound π -dimers, such as TTF or viologen radical cations, afford an alternative design concept: (a) D.-W. Zhang, J. Tian, L. Chen, L. Zhang and Z.-T. Li, *Chem. Asian J.*, 2015, **10**, 56. (b) J. M. Spruell, *Pure Appl. Chem.*, 2010, **82**, 2281. (c) A. Iordache, M. Retegan, F. Thomas, G. Royal, E. Saint-Aman and C. Bucher, *Chem. Eur. J.*, 2012, **18**, 7648. (d) C. Kahlfuss, S. Denis-Quanquin, N. Calin, E. Dumont, M. Garavelli, G. Royal, S. Cobo, E. Saint-Aman and C. Bucher, *J. Am. Chem. Soc.*, 2016, **138**, 15234.

(10) (a) R. W. Baldock, P. Hudson, A. R. Katritzky, F. Soti, *J. Chem. Soc., Perkin Trans. 1*, 1974, 1422. (b) A. T. Balaban, M. T., Caproiu, N. Negotia and R. Baican, *Tetrahedron*, 1977, **33**, 2249. (c) H. G. Viehe, R. Mértnyi, L. Stella and Z. Janousek, *Angew. Chem. Int. Ed. Engl.*, 1979, **18**, 917. (d) H. G. Viehe, Z. Janousek, R. Mértnyi and L. Stella, *Acc. Chem. Res.*, 1985, **18**, 148. (e) H. Zipse, *Top Curr. Chem.*, 2006, **263**, 163. (f) G. Leroy, D. Peeters, M. Sana and C. Wilante, In *Substituent Effects in Radical Chemistry*, ed. V. H. Viehe, Z. Janousek and R. Mértnyi, Riedell, Boston, 1986, pp. 1–48. (g) A. R. Katritzky, M. C. Zerner and M. M. Karelson, *J. Am. Chem. Soc.*, 1986, **108**, 7213. (h) D. J. Pasto, *J. Am. Chem. Soc.*, 1988, **110**, 8164. (i) F. G. Bordwell and T.-Y. Lynch, *J. Am. Chem. Soc.* 1989, **111**, 7558. (j) J. S. Wright, H. Shadnia and L. L. Chepelev, *J. Comput. Chem.*, 2009, **30**, 1016. (k) *Radicals in organic chemistry*, ed. P. Renaud and M. P. Sibi, Wiley-VCH, Weinheim, 2001. (l) C. J. Easton, *Chem. Rev.*, 1997, **97**, 53.

(11) (a) T. H. Koch, J. A. Olesen and J. DeNiro, *J. Am. Chem. Soc.*, 1975, **97**, 7285. (b) R. J. Himmelsbach, A. D. Barone, D. L. Kleyer and T. H. Koch, *J. Org. Chem.*, 1983, **48**, 2989. (c) G. Gaudiano, K. Sweeney, R. C. Haltiwanger and T. H. Koch, *J. Am. Chem. Soc.*, 1984, **106**, 7628. (d) O. Benson Jr., S. H. Demirdji, R. C. Haltiwanger and T. H. Koch, *J. Am. Chem. Soc.*, 1991, **113**, 8879. (e) O. Benson Jr., G. Gaudiano, R. C. Haltiwanger and T. H. Koch, *J. Org. Chem.*, 1988, **53**, 3036. (f) D. L. Kleyer, R. C. Haltiwanger and T. H. Koch, *J. Org. Chem.*, 1983, **48**, 147. (g) F. Welle, S. P. Verevkin, M. Keller, H.-D. Beckhaus and C. Ruchardt, *Chem. Ber.*, 1994, **127**, 697.

(12) Such radicals have been described as “dynamically stable”; see discussion in: M. Frenette, C. Aliaga, E. Font-Sanchis and J. C. Scaiano, *Org. Lett.*, 2004, **6**, 2579; and references cited herein.

(13) For reviews on CAACs, see: (a) M. Melaimi, R. Jazzar, M. Soleilhavoup and G. Bertrand, *Angew. Chem. Int. Ed.*, 2017, **56**, 10046. (b) U. S. D. Paul and U. Radius, *Eur. J. Inorg. Chem.*, 2017, 3362. (c) S. Roy, K. C. Mondal and R. W. Roesky, *Acc. Chem. Res.*, 2016, **49**, 357. (d) M. Soleilhavoup and G. Bertrand, *Acc. Chem. Res.*, 2015, **48**, 256. (e) M. Melaimi, M. Soleilhavoup and G. Bertrand, *Angew. Chem. Int. Ed.*, 2010, **49**, 8810. (f) T. Droege and F. Glorius, *Angew. Chem. Int. Ed.*, 2010, **49**, 6940. (g) F. E. Hahn and M. C. Jahnke, *Angew. Chem. Int. Ed.*, 2008, **47**, 3122.

(14) For the synthesis of CAACs, see: (a) V. Lavallo, Y. Canac, C. Präsang, B. Donnadiu and G. Bertrand, *Angew. Chem. Int. Ed.*, 2005, **44**, 5705. (b) R. Jazzar, R. D. Dewhurst, J. B. Bourg, B. Donnadiu, Y. Canac and G. Bertrand, *Angew. Chem. Int. Ed.*, 2007, **46**, 2899. (c) R. Jazzar, J. B. Bourg, R. D. Dewhurst, B. Donnadiu and G. Bertrand, *J. Org. Chem.*, 2007, **72**, 3492. (d) X. Zeng, G. D. Frey, R. Kinjo, B. Donnadiu and G. Bertrand, *J. Am. Chem. Soc.*, 2009, **131**, 8690. (e) E. Tomás-Mendivil, M. M. Hansmann, C. M. Weinstein, R. Jazzar, M. Melaimi, G. Bertrand, *J. Am. Chem. Soc.*, 2017, **139**, 7753.

(15) (a) J. K. Mahoney, D. Martin, C. Moore, A. Rheingold and G. Bertrand, *J. Am. Chem. Soc.*, 2013, **135**, 18766. (b) J. K. Mahoney, D. Martin, F. Thomas, C. Moore, A. L. Rheingold and G. Bertrand, *J. Am. Chem. Soc.*, 2015, **137**, 7519. (c) J. K. Mahoney, R. Jazzar, G. Royal, D. Martin and G. Bertrand, *Chem. Eur. J.*, 2017, **23**, 6206. (d) C. L. Deardorff, E. R. Sikma, C. P. Rhodes and T. W. Hudnall, *Chem. Commun.*, 2016, **52**, 9024.

(16) For other organic radicals based on stable carbenes, see: (a) D. Martin, C. E. Moore, A. L. Rheingold and G. Bertrand, *Angew. Chem. Int. Ed.*, 2013, **52**, 7014. (b) Y. Li, K. C. Mondal, P. P. Samuel, H. Zhu, C. M. Orben, S. Panneerselvam, B. Dittrich, B. Schwederski, W. Kaim, T. Mondal, D. Koley and H. W. Roesky, *Angew. Chem. Int. Ed.*, 2014, **53**, 4168. (c) L. Jin, M. Melaimi, L. Liu and G. Bertrand, *Org. Chem. Front.*, 2014, **1**, 351. (d) S. Styra, M. Melaimi, C. E. Moore, A. L. Rheingold, T. Augenstein, F. Breher and G. Bertrand, *Chem. Eur. J.*, 2015, **21**, 8441. (e) D. Munz, J. Chu, M. Melaimi, and G. Bertrand, *Angew. Chem. Int. Ed.*, 2016, **55**, 12886. (f) M. M. Hansmann, M. Melaimi and G. Bertrand, *J. Am. Chem. Soc.*, 2017, **139**, 15620. (g) M. M. Hansmann, M. Melaimi and G. Bertrand, *J. Am. Chem. Soc.*, 2018, **140**, 2206. (h) M. M. Hansmann, M. Melaimi and G. Bertrand, *J. Am. Chem. Soc.*, 2018, **140**, 2546. (i) D. Rottschäfer, B. Neumann, H.-G. Stämmler, M. van Gastel, D. M. Andrada, and R. S. Ghadwal, *Angew. Chem. Int. Ed.*, 2018, **57**, 4765. (j) J. Böhnke, T. Dellermann, M. A. Celik, I. Krummenacher, R. D. Dewhurst, S. Demeshko, W. C. Ewing, K. Hammond, M. Heß, E. Bill, E. Welz, M. I. S. Röhr, R. Mitrić, B. Engels, F. Meyer and H. Braunschweig, *Nature Communications*, 2018, **9**, 1197.

(17) M. Abe, *Chem. Rev.*, 2013, **113**, 7011.

(18) B. Bleaney, K. D. Bowers, *Proc. Roy. Soc. London Ser. A*, 1952, **214**, 451.

(19) (a) A. D. Becke, *J. Chem. Phys.*, 1993, **98**, 5648. (b) C. Lee, W. Yang and R. G. Parr, *Phys. Rev. B*, 1988, **37**, 785. (c) S. H. Vosko, L. Wilk and M. Nusair, *Can. J. Phys.*, 1980, **58**, 1200. (d) P. J. Stephens, F. J. Devlin, C. F. Chabalowski and M. J. Frisch, *J. Phys. Chem.*, 1994, **98**, 11623.

(20) Calculations were performed on the Gaussian suite of programs: M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, *et al.* Gaussian 09, revision 4.2.0; Gaussian, Inc.: Wallingford, CT, **2009**. See supporting information for complete citation.

(21) B. Gao, B. Yang, T. Li and B. Zhang, *Synth. Comm.*, 2009, **39**, 2973.