Soft Matter



Nonequilibrium phase diagrams for actomyosin networks

Journal:	Soft Matter
Manuscript ID	SM-ART-04-2018-000741.R1
Article Type:	Paper
Date Submitted by the Author:	01-Aug-2018
Complete List of Authors:	Freedman, Simon; University of Chicago, Physics Hocky, Glen; University of Chicago, Chemistry; James Franck Institute Banerjee, Shiladitya; University College London, Physics and Astronomy Dinner, Aaron; James Franck Institute; University of Chicago, Chemistry

SCHOLARONE[™] Manuscripts

Soft Matter

ARTICLE TYPE

Cite this: DOI: 10.1039/xxxxxxxxx

Received Date Accepted Date

DOI: 10.1039/xxxxxxxxx

www.rsc.org/journalname

Nonequilibrium phase diagrams for actomyosin networks †

Simon L. Freedman,^a Glen M. Hocky,^b Shiladitya Banerjee,^c and Aaron R. Dinner^b

Living cells dynamically modulate the local morphologies of their actin cytoskeletons to perform biological functions, including force transduction, intracellular transport, and cell division. A major challenge is to understand how diverse structures of the actin cytoskeleton are assembled from a limited set of molecular building blocks. Here we study the spontaneous self-assembly of a minimal model of cytoskeletal materials, consisting of semiflexible actin filaments, crosslinkers, and molecular motors. Using coarse-grained simulations, we demonstrate that by changing concentrations and kinetics of crosslinkers and motors, as well as filament lengths, we can generate three distinct structural phases of actomyosin assemblies: bundled, polarity-sorted, and contracted. We introduce new metrics to distinguish these structural phases and demonstrate their functional roles. We find that the binding kinetics of motors and crosslinkers can be tuned to optimize contractile force generation, motor transport, and mechanical response. By quantitatively characterizing the relationships between the modes of cytoskeletal self-assembly, the resulting structures, and their functional consequences, our work suggests new principles for the design of active materials.

1 Introduction

Mechanical functions of living cells are determined by dynamic restructuring of the actin cytoskeleton, a highly conserved cellular machinery composed of filamentous actin (F-actin), myosin molecular motors, and crosslinking proteins¹. An enormous variety of F-actin binding proteins with diverse physicochemical properties² can combine with F-actin to assemble function-specific cellular structures. Spatiotemporal control over these structures is essential for coordinated force generation during cell migration^{3,4}, cell adhesion⁵, cytokinesis⁶, and intracellular transport^{7,8}. A quantitative understanding of how diverse cytoskeletal structures are assembled from a limited set of molecular building blocks presents an outstanding challenge at the interface of soft matter physics and cell biology.

Much effort has been devoted to understanding how contracted states arise in actomyosin networks that lack sarcomeric organization. One mechanism that has emerged is that actin filaments rectify randomly directed forces to select for contractile ones because they can buckle readily but cannot stretch significantly^{9–14}. Alternatively, crosslinking can arrest relative sliding of filaments as they increase their overlap^{15,16} (see also¹⁷ in the context of microtubules). The prevalence of these mechanisms and, in turn, contractile dynamics, depends on motor density and activity^{12,15,18–20}, network connectivity^{18,21–23}, actin filament turnover²⁴, and filament stiffness¹⁶.

Sufficient variation of system parameters leads to other states. Networks with low densities of motors exhibit bundled morphologies ^{18,25–27}; lattice-like (lamellar) and clustered morphologies have been seen in simulations as well²⁶. Consistent with experiments and simulations of microtubules^{28,29} as well as theories of active gels³⁰, polarity-sorted asters of crosslinked actin bundles have also been observed¹⁶. Understanding how these diverse structures relate and can be elicited requires systematically mapping phase diagrams for networks of filaments, motors, and crosslinkers in terms of both molecular parameters (e.g., association and dissociation kinetics, molecular sizes, binding geometries, and microscopic mechanical properties) and system parameters (e.g., molecular composition, turnover, system geometries, external forces).

In this paper, we take a step towards this goal. We investigate the structures accessible to a coarse-grained model that has been parameterized to represent actomyosin networks involving only a single crosslinking protein that binds without geometric restrictions. We consider the interplay of motor and crosslinker density, their binding kinetics, and filament length. We observe homo-



^a Department of Physics, The University of Chicago, 929 East 57th Street, Chicago, IL 60637, USA

^b James Franck Institute & Department of Chemistry, The University of Chicago, 929 East 57th Street, Chicago, IL 60637, USA, Chicago, IL, USA; Email: dinner@uchicago.edu

^c Department of Physics and Astronomy, University College London, Gower Street, London, WC1E-6BT E-mail: shiladitya.banerjee@ucl.ac.uk

geneous, contracted, bundled, and polarity sorted structures and introduce order parameters that quantify their prevalence. We then explore the materials properties of these structures. Specifically, we examine the viscoelastic response to shear strain and the motor transport properties. Our study thus unifies previously observed structures in a single model, reveals the dependence of these structures on molecular parameters, and links the structures to functional consequences.

2 Results and Discussion

2.1 Model

To study the spontaneous self-assembly of cytoskeletal structures at experimentally relevant length and time scales (microns and minutes), we use AFINES, a simulation framework we recently developed³¹. In brief, actin filaments are modeled as polar wormlike chains (represented by beads connected by springs) with defined barbed and pointed ends (Fig. 1); crosslinkers are modeled as linear springs with ends (heads) that can stochastically bind and unbind from F-actin via a kinetic Monte Carlo procedure that preserves detailed balance; molecular motors are modeled as active crosslinkers such that once bound, they walk toward the barbed ends of filaments at load-dependent speeds. While competing simulation frameworks are available and have their advantages^{32,33}, a unique feature of AFINES is its preservation of detailed balance in the absence of motors³¹. Treating the physics of passive systems correctly facilitates their comparison with active systems and, in turn, interpretation of the simulations.

We use Brownian dynamics to evolve the positions of constituents in 2D. The choice of dimension is consistent with the fact that actomyosin networks reconstituted in vitro are typically restricted to a region close to a supporting substrate^{12,16,25}. At the same time, to reflect the finite thickness of the experimental systems and enable rearrangement in 2D, we neglect excluded volume. For this reason, we restrict our attention to actin concentrations at which we expect the network connectivity to dominate the dynamics. This precludes modeling active nematic systems^{34,35}, previously studied by others^{36–38}; it may also lead to quantitative artifacts in the rates of structure formation (owing to a lack of entanglement³⁹) and in density-dependent statistics of states with features such as contracted clusters or bundles. Overall, however, the model yields results in good agreement with observations for many cases of experimental interest^{16,31}. The model is described in detail in Section S1, and Table S1 lists all simulation parameters.

2.2 Network structures

We observe three distinct network architectures formed from initially disordered mixtures of F-actin, motors, and crosslinkers: bundled, polarity-sorted, and contracted. Examples are shown for simulations of 500 10- μ m-long filaments in Fig. 1. When Factin is mixed with crosslinkers, thick bundles form and intersect to yield a well-connected mesh. When F-actin is mixed with motors, barbed ends aggregate to form a polarity-sorted network. Combining F-actin with both motors and crosslinkers results in macroscopic contraction of the filaments into dense and discon-



Fig. 1 Network structures. (lower left) Schematic of the model³¹ with F-actin (red), crosslinkers (green), and motors (black). Bound motors walk towards F-actin barbed ends (blue). (remaining panels) Network structures at 400 s for indicated motor and crosslinker densities. These include a bundled network (upper left) formed by filaments and crosslinkers, a polarity-sorted network (lower right) formed by filaments and motors, and a contracted network (upper right) formed from filaments, crosslinkers, and motors. For clarity, only the actin filaments are shown; the motors and crosslinkers are shown in Fig. S1. Cyan scale bar represents 10 μ m.

nected aggregates. These structures are qualitatively consistent with recent experimental observations^{16,40}.

2.3 Order parameters for characterization of network structures

To systematically explore how varying the properties of the network constituents affects structure formation, we introduce order parameters that characterize each of the observed structures. We compute the spatial extent of F-actin aggregation using the radial distribution function,

$$g(r) = P(r)/(2\pi r \delta r \rho_a) \tag{1}$$

where P(r) is the probability that two actin beads are separated by a distance in the range $[r, r + \delta r]$ (here, $\delta r = 0.05 \,\mu$ m), and ρ_a is the number density of actin beads. For a homogeneous network, $g(r) \approx 1$ at all distances (Fig. 2A; the small peaks at integer *r* arise from the spacing of beads within actin filaments). In contrast, for contracted networks, $g(r) \gg 1$ for $r < 10 \,\mu$ m, indicating F-actin exceeds the bulk density, as observed experimentally¹².

While Fig. 2A shows that actin filaments are nearly uniformly distributed in polarity-sorted networks, Fig. 1 indicates that their barbed ends are concentrated. To quantify their aggregation specifically, we compute the ratio $g(r_{barb})/g(r)$, where r_{barb} is the distance between barbed ends. Fig. 2B shows that in polarity-



Fig. 2 Order parameters. Bundled, polarity-sorted, and contracted cases correspond to the conditions highlighted in Fig. 1; the homogeneous case has no motors or crosslinkers. (A) Radial distribution function of actin, g(r). (B) Radial distribution function of F-actin barbed ends, normalized by g(r); the dashed line marks the motor rest length, 0.5 μ m. (C) Mesh size distribution. Inset: Evolution of the average mesh size, normalized by $\langle g(r) \rangle$. (D) Spatially averaged divergence (Section S2). All averages are over the final 50 s of 5 simulations of 400 s.

sorted networks F-actin barbed ends aggregate (and have a secondary peak at 0.5 μ m, the rest length of motors). In contracted networks, barbed ends also aggregate to a higher degree than in bundled networks, indicating a degree of polarity sorting. This is consistent with recent experiments that indeed show that contracted F-actin forms polarity sorted asters¹⁶.

The inset to Fig. 2A indicates that bundled networks aggregate at length scales comparable to the crosslinker rest length, $l_{xl} = 0.15 \ \mu$ m. To quantify the degree of bundling, and to distinguish it from contractility, we measure the distribution of network pore sizes by a procedure that is similar in spirit but simpler than that in Ref. 41. Namely, we grid the simulation box into $(0.25 \ \mu$ m)² bins and compute how many filaments pass through each. For each empty bin, we determine the lengths of the contiguous vertical and horizontal stretches of empty bins that intersect it (Fig. 3). We average these lengths over all empty bins to obtain an average mesh size for each structure. This procedure can be used for analysis of experimental images, in addition to the simulation structures in the present study.

In Fig. 2C, the distributions of mesh sizes for polarity-sorted and homogeneous networks are similar, indicating that the former does not coarsen significantly. The bundled and contracted networks exhibit larger pore sizes; indeed, contracted networks exhibit pore sizes spanning the simulation region, indicating that the network has ripped apart. We can distinguish these cases by normalizing the mesh size by $\langle g(r) \rangle = (1/R) \int_0^R g(r) dr$, (here, $R = 10 \ \mu m$ is the approximate size of a contracted aggregate under our maximally contractile conditions, as shown in Fig. 2A; see Fig. S2 for further details) which quantifies the extent of aggregation. The inset shows that, while the contracted networks initially



Fig. 3 Example of mesh size calculation described in the text. Actin is shown in white (compare with bundled structure in Fig. 1) and the size of the local mesh corresponds to the depth of color, as indicated by the scale.

bundle, at long times this effect is small compared to aggregation. In contrast, bundled networks have a steadily increasing normalized mesh size (Fig. 2C, inset), consistent with continuous coarsening of filamin crosslinked networks of long actin filament bundles⁴⁰. We thus have a metric for the degree of bundling.

To examine the relationship between actin network structure and contractility, we use the divergence of actin's velocity field, $\langle \nabla \cdot \mathbf{v} \rangle$, where $\langle ... \rangle$ indicates spatial averaging (Section S2)³¹. As shown in Fig. 2D, $\langle \nabla \cdot \mathbf{v} \rangle$ becomes significantly more negative for aggregating networks than for bundling or polarity-sorted networks. Comparison with Fig. 2A shows that extensive contraction is associated with large g(r).

2.4 Phase diagram for molecular composition

Using these order parameters, we map the structural phase diagram of actomyosin networks as functions of motor and crosslinker densities (Fig. 4). Consistent with Fig. 1, networks are contracted when motor and crosslinker densities are high (Fig. 4A), polarity-sorted when only motor density is high (Fig. 4B), and bundled when crosslinker density is high (Fig. 4C). These results echo observations in experiments and simulations that at low motor density, crosslinkers can bundle actin into meshes^{18,27}, and at high motor density, crosslinkers promote contractility by increasing network connectivity^{9,18,27,42}. While polarity sorting is not usually observed independent of contractility in actomyosin networks, our result that increasing motor density promotes this effect is consistent with previous experimental observations and simulations of microtubules interacting with plus-end oriented motors²⁹. Furthermore recent experiments on reconsituted actomyosin networks show that rigid F-actin bundles can form polarity-sorted structures without undergoing bulk contraction¹⁶.

Interestingly, we also observe one non-monotonic trend in this phase diagram: while high motor densities inhibit bundling, a small population of motors ($\rho_m \approx 0.02 \ \mu m^{-2}$) enhances bundling. Other non-monotonic trends have been reported, including that high crosslinker densities^{18,22}, and more surprisingly, high motor force²¹ can inhibit contractility and yield bundled or criti-



Fig. 4 Maps of network properties. Colors indicate the values of order parameters characterizing network contractility (A,D), polarity sorting (B,E), and bundling (C,F), at constant filament length, $L = 10 \ \mu m$, binding affinity $k_m^{\rm off} = k_{xl}^{\rm off} = 0.1 \ {\rm s}^{-1}$ (left), and densities $\rho_m = 0.2 \ \mu m^{-2}$, $\rho_{xl} = 1 \ \mu m^{-2}$ (right, plotted as a function of lifetime, $1/k_{m(x)}^{\rm off}$; structures shown in Fig. S5). Averages are over the last 50 s of five simulations of 400 s; order parameters that are functions of distance, e.g., g(r), are integrated over $0 < r \le 10 \ \mu m$. Error bars for representative curves are shown in Fig. S3.

cally connected networks. We did not observe these effects in the ranges of densities that we considered. This may be because our crosslinker-to-filament ratios (up to 7.5:1) are lower than observed thresholds for crosslinker inhibition (ranging from $15:1^{22}$ to >90:1¹⁸); we expect this threshold to depend on the kinetics of association, as we now discuss.

2.5 Phase diagram for binding kinetics

We also modulate molecular-level interaction parameters between F-actin and its binding partners (crosslinkers, motors), and dissect their relative roles in building different structures. These parameters are hard to control in experiment because they require modifying protein-protein interactions. At fixed motor and crosslinker densities, we find that cytoskeletal structures can be tuned by varying the dissociation constants, $k_{m(xl)}^{\text{off}}$ (Fig. 4D-F). The trends are non-monotonic, in contrast to those in Fig. 4A-B. In particular, contraction is highest for intermediate values of $k_{m(xl)}^{\text{off}}$ (Fig. 4D), similar to observations that increasing motor and crosslinker densities beyond a threshold inhibits contraction ^{18,21,22}. Bundling is highest for low values of k_{xl}^{off} with low or high k_m^{off} (Fig. 4F). Notably, this non-monotonic trend only arises for a fixed simulation time, t_F . For a fixed value of $t_F k_{m(xl)}^{\text{off}}$, chang-



Fig. 5 Effects of varying motor and crosslinker lifetimes, $1/k_{m(xl)}^{\text{off}}$, for filaments of different lengths, *L*, at $t_f = 400$ s (filled squares). (A) Contractility order parameter, radial distribution function, for mixtures of filaments, motors, and crosslinkers with $\rho_m = 0.2 \ \mu m^{-2}$ and $\rho_{xl} = 1 \ \mu m^{-2}$. (B) Polarity sorting order parameter, radial distribution function of barbed ends, for mixtures of filaments and motors, without crosslinkers. (C) Bundling order parameter, mesh size, for mixtures of filaments and crosslinkers, without motors. Normalizing quantities in (B) and (C) with respect to $\langle g(r) \rangle$ does not change the qualitative behaviors (Fig. S4). Error bars indicate standard error of the mean. The green dashed lines show how order parameters vary for one length, $L = 10 \ \mu m$, if the corresponding green squares are rescaled such that $t_f = 4/k_{stl(m)}^{\text{off}}$.

ing $1/k_{m(xl)}^{\text{off}}$ modulates structure formation in a monotonic manner (dashed lines in Fig. 5). The non-monotonic trends are important, however, for understanding how structures form in the presence of competing kinetic processes, such as turnover of different network elements^{7,43}.

2.6 Effect of filament length

As the length (*L*) of F-actin varies considerably within cells⁴⁴, we tested how modulating *L* and $k_{m(xl)}^{\text{off}}$ in tandem affects structure formation. In Fig. 5A, we show that increasing *L* favors aggregation by increasing network connectivity; short filaments ($L < 5 \,\mu$ m) do not organize. As in Fig. 4D, the dependence of aggregation on binding affinity is non-monotonic whenever there is significant aggregation (i.e., $L \ge 5 \,\mu$ m). In Fig. 5B, we show that for networks with only motors, at low k_m^{off} , increasing filament length promotes polarity sorting. By contrast, at high k_m^{off} , short lifetimes of motor attachment suppress polarity sorting. As evident from the representative network structures in Fig. S6, shortening the filaments also suppresses polarity sorting.

For networks with only crosslinkers, the mesh size (Fig. 5C) is non-monotonic with respect to both filament length and crosslinker affinity. Low crosslinker affinity and short filament lengths prevent forming stably crosslinked networks (Fig. S7). Conversely, assemblies with high crosslinker affinity or long filament lengths form crosslinked networks, but they rearrange slowly, so further coarsening is impeded, and the mesh size re-

mains small. As these non-monotonic trends only occur for L > 5 μ m, they are more likely to impact structures with longer actin filaments found in budding yeast ⁴⁵, stereocilia ⁴⁶, filopodia, or reconstituted *in vitro* ^{16,40} (though we note that long microtubules have been obseved to form different structures in the presence and absence of a membrane ⁴⁷). Structures with shorter filaments, as found in lamellipodia or the actin cortex ($< 2 \mu$ m)^{23,48}, are less likely to have a finite binding affinity that maximizes contractility. We discuss the effects of length further in conjunction with the mechanical response of networks, below.

2.7 Network structure tunes molecular transport

While the structures of contracted, polarity-sorted, and bundled networks clearly differ, their consequences for biophysical functions are not immediately apparent. To determine how these structures influence motor transport^{49,50}, we follow the dynamics of small numbers of motors introduced after the structures form (Fig. 6A). In Fig. 6B, we plot their mean-squared displacement,

$$MSD(t_f, \Delta) = \frac{1}{t_f - \Delta} \int_0^{t_f - \Delta} [\vec{v}_m(t, \Delta)]^2 dt$$
(2)

where t_f is the length of the trajectory and $\vec{v}_m(t,\Delta) = \vec{r}_m(t + \Delta) - \vec{r}_m(t)$ is the displacement of a motor with center of mass position $\vec{r}_m(t)$ at time *t* after a lag of Δ . While the scaling of the mean-squared displacement is consistent with simple diffusion (Fig. 6B), sample trajectories (Fig. 6A) indicate that motors in contracted and polarity-sorted structures spend significant amounts of time trapped in aggregates of barbed ends.

We quantify caging using a previously defined metric that can distinguish different kinds of motion⁵¹: we compute the distribution of angles, θ , between consecutive displacement vectors $\vec{v}_m(t,\Delta)$ and $\vec{v}_m(t+\Delta,\Delta)$, at different values of Δ . We find that for all structures, there is at least one time scale in which the distribution has a broad peak at $\theta = \pi$ (Fig. 6C), indicating that motors are reversing direction, consistent with confinement⁵¹. For polarity-sorted networks, barbed ends are tightly aggregated, and the motors exhibit caging at all time scales measured. Contracted networks are partially polarity-sorted, so filaments can direct motors both in and out of aggregates, making the caging more spatially extended. Because it takes longer to explore the extended length scale, the peak has is smaller than in the polarity sorted network. Bundled networks exhibit peaks at both $\theta = 0$ and $\theta = \pi$ depending on the time scale. The former corresponds to moving steadily along a filament, while the latter corresponds to switching between oppositely oriented filaments.

Consistent with our results, it was recently shown that actin networks with different structures result in different angle distributions⁵². As in our results for bundled networks, reconstituted networks of actin and filamin supported simultaneous peaks at $\theta = 0$ and $\theta = \pi$ at short times $(0.1 \text{ s})^{52}$. Interestingly, in our simulations, the peak at $\theta = \pi$ diminishes at intermediate times. A key feature of the experiments that is not represented in the present model is that the myosin minifilaments in the experiments have many heads^{51–53}, and this was previously shown to be important for observed glassy dynamics⁵³.



Fig. 6 Transport properties of motors on actin structures. (A) Example trajectories of three motors (each a different color) on actin networks (filaments shown in Fig. 1 and Fig. S1). Open circles show initial position of each motor, and open squares show their final position after 500 timesteps. (B) Mean squared displacement for motors on structures. Black dashed line shows diffusive behavior, $MSD(t_f, \Delta) \propto \Delta$. (C) Distributions of angles between subsequent motor displacements for different lag times, Δ^{51} . Averages in (B-C) are over 770 motors over the last 100 s of 5 independent trajectories of 200 s, at each condition.

2.8 Network structure tunes mechanical response

Next, we evaluated how structural rearrangements in actin networks affect their ability to propagate mechanical forces over long length scales. To this end, we subjected the final network configuration to a shear strain of magnitude $\gamma = 0.5$ (Fig. 7A, algorithm described in Section S3) and measured the resulting strain energy. In Fig. 7B, we show the strain dependence of the strain energy density, $w(\gamma) - w(0)$, where

$$w(\gamma) = (U_f(\gamma) + U_m(\gamma) + U_{xl}(\gamma))/V,$$
(3)

 U_f , U_m , and U_{xl} are the potential energies of the F-actin, motors, and crosslinkers, respectively, (Section S1) and $V = 250 \ \mu \text{m}^3$ is the simulation volume, assuming a thickness of 0.1 μ m. To determine the viscoelastic response of the material, we fit the computed strain energy density to the function $1/2G\gamma^2 + \eta \dot{\gamma}\gamma$ where *G* is the shear modulus, η is the dynamic viscosity, and $\dot{\gamma} = 1 \text{ s}^{-1}$ is the strain rate (see Section S3 for details)⁵⁴.

We found that bundled networks exhibited the most solidlike material response (G = 1.07 Pa, within range of experimentally measured values for actin-filamin networks²⁵, and $\eta =$ 0.007 Pa·s), while polarity-sorted and homogenous networks exhibited fluid-like responses ($G \approx 0, \eta = 0.080, 0.014$ Pa·s respectively). Contracted networks exhibited a viscoelastic response (G = 0.30 Pa, $\eta = 0.063$ Pa·s), consistent with theoretical predictions from active gel models^{55,56}.

We used this method to measure the material response of the broad spectrum of structures that resulted from varying motor and crosslinker densities, and the various bundled networks that resulted from varying filament length and crosslinker lifetime. We find that these structures can be elastic ($\eta \approx 0$), viscous ($G \approx 0$), or viscoelastic (nonzero *G* and η). The viscous systems exhibit no quadratic response to strain because they are not well-connected (indicating there exists a percolation threshold). For a given crosslinker lifetime $(1/k_{xl}^{\text{off}} = 10 \text{ s})$ and filament length $(10 \ \mu\text{m})$, we find that all networks below $ho_{xl} = 0.05 \ \mu m^{-2}$ have $G \approx 0$ and high viscosity (Fig. 7C,E). Therefore, $\rho_{xl} = 0.05 \ \mu \text{m}^{-2}$ defines the percolation threshold of the gel, which is independent of motor density. Above the percolation threshold, networks have a range of viscoleastic behavior, with low motor density yielding more elastic behavior, and high motor density yielding more viscous behavior. For networks without motors, increasing the crosslinker concentration yielded higher shear elastic moduli, consistent with theoretical predictions⁵⁷, experiments^{25,58}, and previous simulations⁵⁹.

In Fig. 7D,F we show that even at high crosslinker density (1.2 μ m⁻²), the network can respond as a fluid if the crosslinker lifetime or filament length is sufficiently low $(1/k_{xl}^{\text{off}} \le 2 \text{ s or } L \le 5 \mu$ m). At intermediate lifetimes, such as for the crosslinker filamin $(1/k_{xl}^{\text{off}} \approx 10 \text{ s})$, our results agree with experiments: below a critical length, networks respond as fluids⁴⁰, and increases in filament length lead to increases in the elastic modulus⁶⁰. At the longest filament lengths (15 μ m) and crosslinker lifetimes ($\ge 50 \text{ s}$), crosslinked networks are nearly solid with $\eta \approx 0$. A viscoelastic regime occurs at intermediate values of filament length and crosslinker lifetime, corresponding to the networks with the largest mesh sizes (Fig. 5C).

3 Conclusion and Outlook

We have shown how modulating the abundance and physicochemical properties of cytoskeletal constituents can tune emergent network structures and in turn their materials properties. We also introduced order parameters that should facilitate analysis of other models and images from experiments. We find that motor and crosslinker binding affinities, as well as filament lengths, have optimal values for maximizing contractility, bundling, and polarity sorting at finite times. From the perspective of materials design, our work demonstrates how a limited set of molecular building blocks can self-assemble diverse active materials.

In the present study, we consider only a single type of crosslinker (as defined by its stiffness and size), which binds without geometric restrictions. It would be straightforward to modify the energy function of the model to introduce geometric restrictions, and studies of passive systems indicate that such terms can introduce additional structural phases²⁶. The model can also be extended to include actin polymerization and turnover, which have been shown to modulate contractility^{42,61} and bundling³⁹. Finally, the model can be parameterized for other polymer assemblies, such as microtubules, kinesin, and dynein, which form vortices and polarity sorted asters^{29,62,63}.

Our framework can thus be built on to treat diverse types of crosslinkers, filaments, and motors. We expect that rich behavior can be achieved by simulating mixtures with four or more constituents. Because it is straightforward to vary the abundance and properties of the constituents computationally, they could be



Fig. 7 Elasticity of network structures. (A) Shearing a bundled network for 0.5 s at a strain rate of $\dot{\gamma} = 1$ /s. Depth of color indicates stretch of filament. (B) Viscoelastic response of the network to a simple shear, as measured by the dependence of the strain energy density on the strain. (C-D) Shear modulus of structures (shown in Fig. 1) formed with constant filament length ($L = 10 \ \mu$ m) and lifetimes ($1/k_{xl,m}^{\rm off} = 10$ s), but different motor and crosslinker densities (C), and for structures (shown in Fig. S7) formed without motors and constant crosslinker density ($\rho_{xl} = 1.2 \ \mu$ m⁻²), with different filament lengths and crosslinker lifetimes (D). (E-F) Viscosity of these networks. Strain calculations in (B-F) are averaged over 5 independent trajectories.

automatically optimized to achieve desired properties⁶⁴, and the results could inform the design and synthesis of new active materials.

4 Conflicts of Interest

There are no conflicts of interest to declare.

5 Acknowledgments

We thank Margaret Gardel, Jonathan Weare, Samantha Stam, and Kim Weirich for helpful conversations. This work was primarily supported by the University of Chicago Materials Research Science and Engineering Center, which is funded by National Science Foundation under award number DMR-1420709. Additional support was provided by the DoD through the NDSEG Program (to S.L.F.), a NIH Ruth L. Kirschstein NRSA award (1F32GM113415-01) (to G.M.H.), and a UCL Strategic Fellowship (to S.B.). Simulations resources were provided by the Research Computing Center at the University of Chicago and National Institutes of Health (NIH) Grant No. 5 R01 GM109455-02.

References

- 1 M. Murrell, P. W. Oakes, M. Lenz and M. L. Gardel, *Nat. Rev. Mol. Cell Biol.*, 2015, **16**, 486.
- 2 A. Michelot and D. G. Drubin, *Curr. Biol.*, 2011, **21**, R560–R569.
- 3 T. D. Pollard and G. G. Borisy, Cell, 2003, 112, 453-465.
- 4 A. J. Lomakin, K.-C. Lee, S. J. Han, D. A. Bui, M. Davidson, A. Mogilner and G. Danuser, *Nat. Cell Biol.*, 2015, **17**, 1435.
- 5 J. T. Parsons, A. R. Horwitz and M. A. Schwartz, *Nat. Rev. Mol. Cell Biol.*, 2010, **11**, 633–643.
- 6 J. Sedzinski, M. Biro, A. Oswald, J.-Y. Tinevez, G. Salbreux and E. Paluch, *Nature*, 2011, **476**, 462.
- 7 E. Munro, J. Nance and J. R. Priess, *Dev. Cell*, 2004, 7, 413–424.
- 8 S. M. A. Tabei, S. Burov, H. Y. Kim, A. Kuznetsov, T. Huynh, J. Jureller, L. H. Philipson, A. R. Dinner and N. F. Scherer, *Proc. Natl. Acad. Sci. USA*, 2013, **110**, 4911–4916.
- 9 M. S. e Silva, M. Depken, B. Stuhrmann, M. Korsten, F. C. MacKintosh and G. H. Koenderink, *Proc. Natl. Acad. Sci. USA*, 2011, **108**, 9408–9413.
- 10 M. Lenz, T. Thoresen, M. L. Gardel and A. R. Dinner, *Phys. Rev. Lett.*, 2012, **108**, 238107.
- 11 M. Lenz, M. L. Gardel and A. R. Dinner, New J. Phys., 2012, 14, 033037.
- 12 M. P. Murrell and M. L. Gardel, Proc. Natl. Acad. Sci. USA, 2012, 109, 20820–20825.
- 13 M. Lenz, Phys. Rev. X, 2014, 4, 041002.
- 14 P. Ronceray, C. P. Broedersz and M. Lenz, Proc. Natl. Acad. Sci. USA, 2016, 113, 2827–2832.
- 15 M. Murrell and M. L. Gardel, *Mol. Biol. Cell*, 2014, **25**, 1845– 1853.
- 16 S. Stam, S. L. Freedman, S. Banerjee, K. L. Weirich, A. R. Dinner and M. L. Gardel, *Proc. Natl. Acad. Sci. USA*, 2017, 114, E10037–10045.
- 17 K. T. Stanhope, V. Yadav, C. D. Santangelo and J. L. Ross, *Soft Matter*, 2017, **13**, 4268–4277.
- 18 P. M. Bendix, G. H. Koenderink, D. Cuvelier, Z. Dogic, B. N. Koeleman, W. M. Brieher, C. M. Field, L. Mahadevan and D. A. Weitz, *Biophys. J.*, 2008, **94**, 3126–3136.
- 19 I. Linsmeier, S. Banerjee, P. W. Oakes, W. Jung, T. Kim and M. Murrell, *Nat. Commun.*, 2016, 7, 12615.
- 20 J. M. Belmonte, M. Leptin and F. Nédélec, *Mol. Syst. Biol.*, 2017, **13**, 941.
- 21 J. Alvarado, M. Sheinman, A. Sharma, F. C. MacKintosh and G. H. Koenderink, *Nat. Phys.*, 2013, 9, 591–597.
- 22 H. Ennomani, G. Letort, C. Guérin, J.-L. Martiel, W. Cao, F. Nédélec, M. Enrique, M. Théry and L. Blanchoin, *Curr. Biol.*, 2016, **26**, 616–626.
- 23 P. Chugh, A. G. Clark, M. B. Smith, D. A. Cassani, K. Dierkes, A. Ragab, P. P. Roux, G. Charras, G. Salbreux and E. K. Paluch, *Nat. Cell Biol.*, 2017, **19**, 689–697.
- 24 T. Hiraiwa and G. Salbreux, Phys. Rev. Lett., 2016, 116, 188101.

- 25 K. Schmoller, O. Lieleg and A. Bausch, *Biophys. J.*, 2009, **97**, 83–89.
- 26 C. Cyron, K. Müller, K. Schmoller, A. Bausch, W. Wall and R. Bruinsma, *Europhys. Lett.*, 2013, **102**, 38003.
- 27 T. C. Bidone, W. Jung, D. Maruri, C. Borau, R. D. Kamm and T. Kim, *PLoS Comput. Biol.*, 2017, **13**, e1005277.
- 28 F. Nedelec, T. Surrey, A. C. Maggs and S. Leibler, *Nature*, 1997, **389**, 305.
- 29 T. Surrey, F. Nédélec, S. Leibler and E. Karsenti, *Science*, 2001, 292, 1167–1171.
- 30 K. Kruse, J.-F. Joanny, F. Jülicher, J. Prost and K. Sekimoto, *Phys. Rev. Lett.*, 2004, **92**, 078101.
- 31 S. L. Freedman, S. Banerjee, G. M. Hocky and A. R. Dinner, *Biophys. J*, 2017, **113**, 448–460.
- 32 F. Nedelec and D. Foethke, New J. Phys., 2007, 9, 427.
- 33 K. Popov, J. Komianos and G. A. Papoian, *PLoS Comput. Biol.*, 2016, **12**, e1004877.
- 34 T. Sanchez, D. T. Chen, S. J. DeCamp, M. Heymann and Z. Dogic, *Nature*, 2012, **491**, 431.
- 35 R. Zhang, N. Kumar, J. L. Ross, M. L. Gardel and J. J. de Pablo, Proc. Natl. Acad. Sci. USA, 2017, 201713832.
- 36 G. Letort, A. Z. Politi, H. Ennomani, M. Théry, F. Nedelec and L. Blanchoin, *PLoS Comput. Biol.*, 2015, **11**, e1004245.
- 37 R. Blackwell, O. Sweezy-Schindler, C. Baldwin, L. E. Hough, M. A. Glaser and M. Betterton, *Soft Matter*, 2016, **12**, 2676– 2687.
- 38 S. J. DeCamp, G. S. Redner, A. Baskaran, M. F. Hagan and Z. Dogic, *Nat. Mater.*, 2015, 14, 1110.
- 39 G. Foffano, N. Levernier and M. Lenz, *Nat. Commun.*, 2016, 7, 13827.
- 40 K. L. Weirich, S. Banerjee, K. Dasbiswas, T. A. Witten, S. Vaikuntanathan and M. L. Gardel, *Proc. Natl. Acad. Sci.* USA, 2017, **114**, 2131–2136.
- 41 W. Mickel, S. Münster, L. M. Jawerth, D. A. Vader, D. A. Weitz, A. P. Sheppard, K. Mecke, B. Fabry and G. E. Schröder-Turk, *Biophys. J.*, 2008, **95**, 6072–6080.
- 42 M. Mak, M. H. Zaman, R. D. Kamm and T. Kim, Nat. Commun., 2016, 7, 10323.
- 43 M. Fritzsche, C. Erlenkämper, E. Moeendarbary, G. Charras and K. Kruse, *Sci. Adv.*, 2016, **2**, e1501337.
- 44 L. Mohapatra, B. L. Goode, P. Jelenkovic, R. Phillips and J. Kondev, *Annu. Rev. Biophys.*, 2016, **45**, 85–116.
- 45 M. Chesarone-Cataldo, C. Guérin, H. Y. Jerry, R. Wedlich-Soldner, L. Blanchoin and B. L. Goode, *Dev. Cell*, 2011, 21, 217–230.
- 46 H. W. Lin, M. E. Schneider and B. Kachar, Curr. Opin. Cell Biol., 2005, 17, 55–61.
- 47 M. Pinot, F. Chesnel, J. Kubiak, I. Arnal, F. Nedelec and Z. Gueroui, *Curr. Biol.*, 2009, **19**, 954–960.
- 48 G. Salbreux, G. Charras and E. Paluch, *Trends Cell Biol.*, 2012, 22, 536–545.
- 49 C. M. Brawley and R. S. Rock, *Proc. Natl. Acad. Sci. USA*, 2009, 106, 9685–9690.
- 50 S. Kohler, V. Schaller, A. Bausch et al., Nat. Mater., 2011, 10,

462–468.

- 51 S. Burov, S. M. A. Tabei, T. Huynh, M. P. Murrell, L. H. Philipson, S. A. Rice, M. L. Gardel, N. F. Scherer and A. R. Dinner, *Proc. Natl. Acad. Sci. USA*, 2013, **110**, 19689–19694.
- 52 M. Scholz, K. L. Weirich, M. L. Gardel and A. R. Dinner, *bioRxiv*, 2018, 277947.
- 53 M. Scholz, S. Burov, K. L. Weirich, B. J. Scholz, S. M. A. Tabei, M. L. Gardel and A. R. Dinner, *Phys. Rev. X*, 2016, 6, 011037.
- 54 D. H. Boal, *Mechanics of the Cell*, Cambridge University Press, 2012.
- 55 T. B. Liverpool, M. C. Marchetti, J.-F. Joanny and J. Prost, *EPL* (*Europhys. Lett.*), 2009, **85**, 18007.
- 56 S. Banerjee and M. C. Marchetti, *Soft Matter*, 2011, 7, 463–473.
- 57 D. A. Head, A. J. Levine and F. C. MacKintosh, Phys. Rev. E,

2003, 68, 061907.

- 58 M. Gardel, J. Shin, F. MacKintosh, L. Mahadevan, P. Matsudaira and D. Weitz, *Science*, 2004, **304**, 1301–1305.
- 59 T. Kim, W. Hwang, H. Lee and R. D. Kamm, *PLoS Comput. Biol.*, 2009, **5**, e1000439.
- K. E. Kasza, G. H. Koenderink, Y. C. Lin, C. P. Broedersz,
 W. Messner, F. Nakamura, T. P. Stossel, F. C. MacKintosh and
 D. A. Weitz, *Phys. Rev. E*, 2009, **79**, 041928.
- 61 W. M. McFadden, P. M. McCall, M. L. Gardel and E. M. Munro, *PLoS Comput. Biol.*, 2017, **13**, e1005811.
- 62 F. Nédélec, T. Surrey and A. Maggs, *Phys. Rev. Lett.*, 2001, **86**, 3192.
- 63 F. Nédélec, J. Cell Biol., 2002, 158, 1005–1015.
- 64 M. Z. Miskin, G. Khaira, J. J. de Pablo and H. M. Jaeger, *Proc. Natl. Acad. Sci. USA*, 2016, **113**, 34–39.



Using simulation to explore the diverse structural and functional phases of the actin cytoskeleton.