



Applications of 1,3,5-trimethoxybenzene as a derivatizing agent for quantifying free chlorine, free bromine, bromamines, and bromide in aqueous systems

Journal:	<i>Analytical Methods</i>
Manuscript ID	AY-ART-07-2019-001443.R1
Article Type:	Paper
Date Submitted by the Author:	10-Sep-2019
Complete List of Authors:	Dias, Ryan; Towson University, Chemistry Schammel, Marella; Towson University, Chemistry Reber, Keith; Towson University, Chemistry Sivey, John; Towson University, Chemistry

1
2
3
4 **Applications of 1,3,5-trimethoxybenzene as a**
5 **derivatizing agent for quantifying free chlorine, free**
6 **bromine, bromamines, and bromide in aqueous**
7 **systems**
8
9
10

11
12
13 Ryan P. Dias,^{a,‡} Marella H. Schammel,^a Keith P. Reber,^a and John D. Sivey^{*ab}
14

15
16 ^a Department of Chemistry, Towson University, 8000 York Road, Towson, Maryland 21252,
17 United States
18

19 ^b Urban Environmental Biogeochemistry Laboratory, Towson University, 8000 York Road,
20 Towson, Maryland 21252, United States
21

22
23 * Corresponding author contact information: phone: 1-410-704-6087; e-mail: jsivey@towson.edu
24

25
26 ‡ Current affiliation : Department of Chemistry, University of Alberta, 11227 Saskatchewan Drive,
27 Edmonton, Alberta, Canada T6G 2G2
28

29
30
31 Submitted exclusively to *Analytical Methods*
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Free chlorine and free bromine (e.g., HOCl and HOBr) are employed as disinfectants in a variety of aqueous systems, including drinking water, wastewater, ballast water, recreational waters, and cleaning products. Yet, the most widely used methods for quantifying free halogens, including those employing *N,N*-diethyl-*p*-phenylenediamine (DPD), cannot distinguish between HOCl and HOBr. Herein, we report methods for selectively quantifying free halogens in a variety of aqueous systems using 1,3,5-trimethoxybenzene (TMB). At near-neutral pH, TMB reacted on the order of seconds with HOCl, HOBr, and inorganic bromamines to yield halogenated products that were readily quantified by liquid chromatography or, following liquid-liquid extraction, gas chromatography-mass spectrometry (GC-MS). The chlorinated and brominated products of TMB were stable, and their molar concentrations were used to calculate the original concentrations of HOCl (method quantitation limit (MQL) by GC-MS = $15 \text{ nmol L}^{-1} = 1.1 \text{ } \mu\text{g L}^{-1}$ as Cl_2) and HOBr (MQL by GC-MS = $30 \text{ nmol L}^{-1} = 2 \text{ } \mu\text{g L}^{-1}$ as Cl_2), respectively. Moreover, TMB derivatization was efficacious for quantifying active halogenating agents in drinking water, pool water, chlorinated surface waters, and simulated spa waters treated with 1-bromo-3-chloro-5,5-dimethylhydantoin. TMB was also used to quantify bromide as a trace impurity in 20 nominally bromide-free reagents (following oxidation of bromide by HOCl to HOBr). Several possible interferences were tested, and iodide was identified as impeding accurate quantitation of HOCl and HOBr. Overall, compared to the DPD method, TMB can provide lower MQLs, larger linear ranges, and selectivity between HOCl and HOBr.

1. Introduction


Free chlorine and free bromine serve as disinfectants (or are generated *in situ*) in drinking water, wastewater, spas, pools, ballast water in ships, as well as in household and industrial cleaning products.¹⁻³ Free chlorine (i.e., free available chlorine) describes all aqueous chlorine species in the +I or 0 oxidation state (except chloramines).⁴ Examples of free chlorine species include HOCl, ClO⁻, Cl₂, and Cl₂O.^{4,5} Solutions of free chlorine are commonly generated by dosing water with NaOCl(*aq*), Ca(OCl)₂(*s*), Cl₂(*g*), or stabilized chlorine, including various organic chloramines.^{1,6} Free bromine (i.e., free available bromine) refers to all aqueous bromine species in the +I or 0 oxidation state (except bromamines).⁷ Example of free bromine species include HOBr, BrO⁻, Br₂, BrCl, Br₂O, and BrOCl.⁷ Free bromine can be produced via hydrolysis of liquid bromine (Br₂) or organic bromamines, as well as by *in situ* oxidation of bromide (e.g., by HOCl, **eq 1**).^{1,7,8}



Ozone and hydrogen peroxide/peroxidase systems are also capable of oxidizing bromide into free bromine.^{9,10} At pH 7, hypochlorous acid (HOCl, $pK_a = 7.58$ at 20 °C)¹¹ and hypobromous acid (HOBr, $pK_a = 8.70$ at 20 °C)¹² are the most abundant forms of free chlorine and free bromine, respectively.

Accurate quantification of free halogen residuals in disinfected waters is important for protecting human and environmental health. At doses on the order of 1 – 4 mg L⁻¹, free halogens are capable of inactivating pathogens in drinking water, wastewater, and recreational waters.¹ Higher concentrations (e.g., 10 to >10,000 mg L⁻¹) of free chlorine are employed for disinfection processes associated with food service, medical/mortuary settings, and decontamination of biological warfare agents.¹³⁻¹⁶

1
2
3 Monitoring of free halogens in disinfected waters is also important due to the propensity
4 of free halogens to react with organic matter to form chlorinated and brominated disinfection
5 byproducts (DBPs).^{17,18} Brominated DBPs are generally more toxic than their chlorinated
6 analogues,¹⁹ which underscores the need for analytical methods capable of selectively quantifying
7 free chlorine and free bromine as precursors of DBPs. Quantification of free halogens is also
8 important in laboratory settings, including experiments investigating disinfection efficacy and
9 DBP (trans)formation.²⁰⁻²² Such studies frequently employ solutions containing mixtures of free
10 chlorine and free bromine generated via oxidation of bromide (e.g., from NaBr) by excess free
11 chlorine.²³ Free chlorine/bromine mixtures are also generated in waters (e.g., spas and toilet tanks)
12 disinfected with 1-bromo-3-chloro-5,5-dimethylhydantoin (BCDMH, **Scheme 1**),¹ as well as in
13 chlorinated solutions amended with salts (e.g., NaCl) in which bromide is present as an impurity.²⁴
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28




Scheme 1. Hydrolysis of 1-bromo-3-chloro-5,5-dimethylhydantoin (BCDMH) yielding HOBr and HOCl. Whether BCDMH exists principally as 1-bromo-3-chloro-5,5-dimethylhydantoin or as 3-bromo-1-chloro-5,5-dimethylhydantoin has been discussed in the literature.²⁵

41 Several methods exist for quantifying free halogen residuals in aqueous samples, including
42 iodometric titration, amperometric titration, and, perhaps most commonly, colorimetric methods
43 employing *N,N*-diethyl-*p*-phenylenediamine (DPD).^{26,27} Methods of quantifying free chlorine
44 involving chemiluminescent reactions²⁸ and quantum dots²⁹⁻³¹ have also been reported. None of
45 these methods, however, are able to selectively and concurrently quantify free chlorine and free
46 bromine in samples containing mixtures of these analytes.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 To distinguish between free chlorine and free bromine, electron-rich aromatic compounds
4 can serve as derivatizing agents (i.e., probes) for these free halogens. Such methods leverage the
5 propensity of aromatic compounds containing “activating” substituents (e.g., hydroxy and
6 methoxy) to form organohalides via electrophilic substitution.^{32,33} When selecting among possible
7 probes, several characteristics of the candidate probe and of its halogenated products need to be
8 considered, including water solubility, nucleophilicity, stability, ease of analysis, molecular
9 symmetry, acid/base behavior, and toxicity. Probes (and their halogenation products) must be
10 sufficiently water soluble so as to preclude precipitation. Probes should demonstrate adequate
11 nucleophilicity in reactions with free halogens so as to promote stoichiometric conversion of free
12 halogens to the corresponding organohalides in reasonable time frames (ideally on the order of
13 seconds). Candidate probes must generate stable halogenated products amenable to selective
14 quantitation (e.g., by liquid chromatography (LC) and/or gas chromatography (GC)) so that non-
15 halogenated, chlorinated, and brominated probe species can be individually quantified. Probes
16 with higher degrees of symmetry have fewer unique substitution positions, generate fewer
17 products, and thereby simplify analysis of product mixtures compared to probes of lower
18 symmetry. Acid-base behavior (if any) should be chosen so as to enhance (or, at a minimum, not
19 diminish) the probe’s performance in terms of water solubility, nucleophilicity, and ease of
20 analysis. Importantly, probes should also be selected so as to minimize toxicity hazards.


21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45 Previous studies have demonstrated the ability of electron-rich aromatic compounds
46 (including phenol, 2,6-dimethylphenol, 2,6-dichlorophenol, and 3,5-dimethyl-1*H*-pyrazole
47 (DMPyr), **Figure 1**) to serve as probes of free chlorine and/or free bromine via quantification of
48 their halogenated products by LC or GC.³⁴⁻³⁹ Notably, all of these compounds are ionizable, with
49 greater nucleophilicity under alkaline conditions⁴⁰⁻⁴⁴ whereas the electrophilicity of free chlorine
50
51
52
53
54
55
56
57
58
59
60

1
2
3 and free bromine increases under acidic conditions.^{45,46} Collectively, these factors can complicate
4
5 pH-optimization efforts for electrophilic aromatic substitutions involving ionizable nucleophiles.
6
7 Use of 2,6-dichlorophenol as a derivatizing agent for free bromine was also confounded by
8
9 precipitate formation in systems disinfected with peracetic acid; precipitation was not observed in
10
11 systems disinfected with free chlorine or ozone.³⁴ Recently, DMPyr was shown to provide an
12
13 exceptionally low method limit of quantification (MQL = 18 pmol L⁻¹) as a probe for free bromine
14
15 and other reactive bromine species in conjunction with LC-tandem mass spectrometry; however,
16
17 the ability of DMPyr to concurrently quantify free chlorine and free bromine was not
18
19 investigated.³⁹
20
21
22
23
24



32 **Figure 1.** Examples of aromatic compounds previously examined as probes of free chlorine and
33 free bromine.³⁴⁻³⁹
34
35

36 Herein, we propose a derivatization method wherein the aromatic ether 1,3,5-
37 trimethoxybenzene (TMB) serves as a free halogen trap (**Scheme 2**). The water solubility of TMB
38 (estimates ranging from 4.6 to 19 mmol L⁻¹ at 25 °C)⁴⁷ is sufficient to permit its use as a
39 derivatizing agent across a wide range of free halogen concentrations. The presence of three
40 methoxy groups activates TMB toward electrophilic substitution.³³ The *D*_{3h} symmetry of TMB
41 minimizes the number of possible regioisomeric products that can form via electrophilic aromatic
42 substitution. The absence of proton donor/acceptor (e.g., hydroxy or amino) groups renders the
43 nucleophilicity of TMB essentially invariant as a function of pH and permits quantification of
44 TMB and its products by both LC and GC.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Scheme 2. Derivatization of free chlorine (HOCl) and free bromine (HOBr) by TMB to give Cl-TMB and Br-TMB, respectively.

Previous studies demonstrated that TMB reacts with HOCl ($k_{\text{TMB,HOCl}} (20\text{ }^{\circ}\text{C}) = 697\text{ M}^{-1}\text{ s}^{-1}$) and monochloramine ($k_{\text{TMB,NH}_2\text{Cl}} (20\text{ }^{\circ}\text{C}) \approx 3 \times 10^{-3}\text{ M}^{-1}\text{ s}^{-1}$) to give 1-chloro-2,4,6-trimethoxybenzene (Cl-TMB).^{5,21} Reaction of TMB with HOBr is more rapid ($k_{\text{TMB,HOBr}} (20\text{ }^{\circ}\text{C}) = 3.35 \times 10^6\text{ M}^{-1}\text{ s}^{-1}$) and yields 1-bromo-2,4,6-trimethoxybenzene (Br-TMB).²¹ In waters containing mixtures of free bromine and ammonium ions (e.g., recreational waters disinfected with free bromine or chlorinated wastewater containing bromide), bromamines can form.⁴⁸ The reactivity of TMB toward bromamines has not been previously characterized and is examined herein. Previous work in our laboratory demonstrated that TMB can be used to selectively quench and quantify free chlorine and free bromine;²¹ however, these studies were limited to synthetic waters whose matrices were notably less complex than natural or recreational waters.

The purpose of this study is to develop methods for using TMB as a probe to (1) quantify residual free halogens in synthetic waters, chlorinated surface waters, municipal drinking water, and pool water; (2) monitor free halogens generated by BCDMH present in disinfecting tablets designed for spas; and (3) quantify bromide as an impurity in ostensibly bromide-free reagents

upon oxidation by HOCl to HOBr. Overall, this work affords selective and sensitive methods for concurrent quantification of HOCl and HOBr that are amenable to either LC or GC analysis.

2. Experimental

All aqueous solutions were prepared using deionized water further purified with a NANOpure system (Thermo Scientific) to a resistivity $\geq 18 \text{ M}\Omega \text{ cm}$. Laboratory-grade sodium hypochlorite (NaOCl, ~6% w/w, Fisher Scientific) served as the source of free chlorine and was standardized via iodometric titration.²⁶ Working solutions of free chlorine were prepared fresh daily by diluting 6% w/w NaOCl solution with 11 mM NaOH and were standardized via UV-vis spectrophotometry ($\epsilon_{\text{OCl}^-} = 365.8 \pm 0.7 \text{ M}^{-1} \text{ cm}^{-1}$, $T = 21 \pm 1 \text{ }^\circ\text{C}$, see Electronic Supplementary Information, ESI, for more details). Additional reagents are described in **Table S1** of the ESI.

Under most experimental conditions employed herein, TMB is added in sufficient excess of the total free halogen concentration so as to render di- and tri-halogenation of TMB negligible ($\leq 0.3\%$ of the initial concentration of TMB). Under such conditions, the TMB mass balance can be calculated as:

$$[\text{TMB}]_{\text{tot}} = [\text{TMB}] + [\text{Cl - TMB}] + [\text{Br - TMB}]$$

(2)

Measured concentrations of Cl-TMB and Br-TMB can be used to determine the residual free chlorine ($[\text{HOCl}]_{\text{res}}$) and free bromine ($[\text{HOBr}]_{\text{res}}$) concentrations, respectively, in the original samples.

$$[\text{HOCl}]_{\text{res}} = [\text{Cl - TMB}] \quad (3)$$

$$[\text{HOBr}]_{\text{res}} = [\text{Br - TMB}] \quad (4)$$

We note, however, that when free chlorine is dosed to solutions containing bromide, the mass balance of total free chlorine must account for the conversion of free chlorine into free bromine (eq 1). In such instances, the total (added) free chlorine ($[\text{HOCl}]_{\text{tot}}$) mass balance is given by:

$$[\text{HOCl}]_{\text{tot}} = [\text{Cl - TMB}] + [\text{Br - TMB}] \quad (5)$$

When free chlorine is added in excess of the initial bromide concentration, bromide can be stoichiometrically converted into free bromine and, upon addition of TMB, into Br-TMB. Thus, Br-TMB can serve as a useful surrogate for the bromide originally present in a sample treated with free chlorine (assuming that free bromine does not appreciably react with components of the system other than TMB). Additional considerations pertaining to mass balances are provided in the ESI.

As described below, concentrations of TMB and its halogenated products can be determined by high-performance liquid chromatography (HPLC) and, following extraction into toluene, by gas chromatography-mass spectrometry (GC-MS). Analyte concentrations determined by these methods were corrected for any sample preparation steps resulting in dilution or preconcentration.

2.1 Comparison of free halogen quantification by TMB and DPD

The utility of quantifying HOCl and HOBr via derivatization with TMB was compared to the DPD method²⁶ using laboratory solutions of free chlorine/bromine as well as chlorinated swimming pool water. Laboratory solutions of free chlorine/bromine (25 mL) were prepared in 40-mL amber glass vials containing borate buffer (20 mM, adjusted to pH 6.0 or 7.0 using HNO_3), NaNO_3 (90 mM), NaCl (10 mM), and NaBr (0 or 40 μM). NaNO_3 served as an ionic strength adjuster. NaCl was added to maintain a uniform $[\text{Cl}^-]$, noting that the speciation of free chlorine and free bromine can vary with $[\text{Cl}^-]$.^{49,50} After dosing with NaOCl (20 μM), reactors were

1
2
3 incubated in a circulating water bath at 20.00 ± 0.01 °C for ~4 min to permit thermal equilibration
4
5 and (for solutions amended with NaBr) bromide oxidation.
6
7

8 Pool water samples were obtained in 250-mL clear plastic bottles from an indoor
9
10 swimming pool (volume ≈ 900 m³) in Baltimore County, Maryland, whose treatment includes
11
12 filtration and UV irradiation followed by chlorination. Water samples were collected 0.3 m below
13
14 the surface when no swimmers were present; the pool had not been occupied for 21 h prior to
15
16 sampling. Samples were stored in a cooler chilled with ice during transit back to the laboratory
17
18 and were analyzed for free chlorine/bromine within 1 h of sampling.
19
20


21 For TMB workup, triplicate aliquots of free halogen solutions (1.00 mL) were transferred
22
23 to 4-mL amber glass vials pre-amended with methanolic TMB solution (0.15 mL at 2.62 mM to
24
25 give a diluted TMB concentration of 342 μ M). The 4-mL vials were capped and shaken manually
26
27 for 10 s. After all samples were quenched with TMB, toluene (0.50 mL, containing 2-
28
29 chlorobenzonitrile at 10.2 μ M as the internal standard) was added to each 4-mL vial as the
30
31 extraction solvent. Vials were capped and shaken manually for 30 s. After phase separation was
32
33 re-established, an aliquot of the toluene phase (0.20 mL) was transferred to a 0.3-mL glass insert
34
35 seated inside an amber glass 2-mL autosampler vial. The 2-mL vials were capped with a screw-
36
37 top plastic cap fitted with a PTFE-lined septum. Concentrations of TMB and its halogenated
38
39 products in toluene extracts were determined using GC-MS (see ESI, **Table S2**, for additional GC-
40
41 MS method details). Example GC-MS chromatograms are shown in **Figure S5**.
42
43
44
45

46 For DPD workup, triplicate aliquots of free halogen solutions (3.9 mL) were added to
47
48 amber glass vials (capacity = 4.4 mL) pre-amended with 0.2 mL of DPD indicator solution and
49
50 0.2 mL of phosphate buffer (0.5 M, pH 7.0). Vials were capped, shaken manually for 10 s, and
51
52 allowed to stand for 30 s to permit development of a magenta color. Absorbance measurements at
53
54
55
56
57
58
59
60

1
2
3 515 nm were obtained using a Cary 60 UV-Vis spectrophotometer following established
4 methods.²⁶
5
6

7 8 9 **2.2 Assessing the reactivity of TMB with aqueous combined bromine**

10
11 The reactivity of TMB towards inorganic combined bromine (i.e., bromamines) was
12 assessed using batch reactors. Reactor solutions (total volume = 25 mL) were prepared in 40-mL
13 amber glass vials containing borate buffer (20 mM, pH 8.0), NaNO₃ (90 mM), NaCl (10 mM), and
14 NaBr (100 μM). Reactors were incubated at 20.00 ± 0.01 °C in a circulating water bath for 4 min
15 prior to dosing with NaOCl, which was added to the reactor vial to achieve a final concentration
16 of 150 μM. The vial was capped, shaken manually for 10 s, and returned to the water bath for 5
17 min to allow bromide oxidation to HOBr by the 1.5-molar excess of free chlorine. Following this
18 equilibration period, NH₄Cl was added (as a source of NH₄⁺ + NH₃) to the reactor vial to achieve
19 a final concentration of 300 μM. The vial was capped, shaken manually for 10 s, and returned to
20 the water bath for 2 min to permit bromamine formation via:
21
22
23
24
25
26
27
28
29
30
31
32



33
34
35
36
37
38
39 At t = 0, a spike of methanolic TMB solution (0.150 mL at 100 mM) was added to the reaction
40 solution such that [TMB]₀ in the reaction solution was 630 μM. Aliquots (0.8 mL) were obtained
41 periodically and transferred to 4-mL amber glass vials pre-amended with aqueous Na₂S₂O₃ (0.075
42 mL at 10 mM) to quench residual halamines.
43
44
45
46

47
48 After all samples were quenched, liquid-liquid extractions as described in Section 2.1 were
49 performed, using 0.80 mL of toluene containing 2-chlorobenzonitrile (10.2 μM) as the internal
50 standard. TMB and halogenated products were analyzed via GC-MS (**Table S2**).
51
52
53
54

55 56 **2.3 Quantification of free halogens in chlorinated surface waters and drinking water**

57
58
59
60

1
2
3 The ability of TMB to quench free halogens was investigated in surface waters amended
4 with free chlorine and in chlorinated municipal tap waters. Surface water samples were collected
5 from four different water bodies: the Loch Raven Reservoir (freshwater, near Timonium,
6 Maryland, on 6 Jun 2017), the Susquehanna River (freshwater, near Havre de Grace, Maryland,
7 on 6 Jun 2017), the Chesapeake Bay (brackish, at Cove Point, Maryland, on 5 Jun 2017), and the
8 Atlantic Ocean (at Bethany Beach, Delaware, on 6 Jun 2017). A summary of water quality metrics
9 for each water source is provided in **Table S3**. Surface water samples were obtained approximately
10 0.3 m below the water surface and within 3 m of the shore in 2-L clear glass bottles with plastic
11 caps. The 2-L bottles were pre-rinsed with free chlorine followed by 18 M Ω cm water prior to
12 sampling. At each sampling location, bottles were rinsed with surface water several times before
13 samples were obtained (volume = 2 L, collected in triplicate). Samples were stored in a cooler
14 chilled with ice-packs during transit. Upon arrival at Towson University, samples were stored at 4
15 °C prior to use.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33 Aliquots of surface water (25.0 mL) were added to 40-mL amber glass vials and placed in
34 a recirculating water bath at 20.00 ± 0.01 °C for 4 min to permit thermal equilibration. Vials
35 containing Loch Raven Reservoir water and Susquehanna River water, respectively, were
36 amended with NaBr (150 μ M) to normalize bromide content between the two waters. Vials were
37 dosed with NaOCl (100 μ M), capped, and shaken manually for 10 s before being returned to the
38 water bath. Triplicate aliquots (0.90 mL) were obtained at each sampling time over 1 h and added
39 to vials pre-amended with TMB (0.332 mL at 2.60 mM in methanol). Vials were capped and
40 shaken manually for 10 s to facilitate quenching. After all samples were quenched, similar liquid-
41 liquid extractions as described in Section 2.1 were performed, using 1.00 mL of toluene containing
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 2-chlorobenzonitrile (10.2 μM) as the internal standard. TMB and its halogenated products were
4 analyzed via GC–MS (**Table S2**).
5
6

7
8 Municipal drinking water samples were obtained from three locations on Towson
9 University’s main campus in Towson, Maryland (**Table S4**). Samples were stored in a cooler
10 chilled with ice-packs during transit back to the laboratory and were treated with TMB within 1 h
11 of sampling. Water samples were obtained in 125-mL clear plastic bottles covered in aluminum
12 foil. Aliquots (1.00 mL) were added to clear glass 2-mL HPLC autosampler vials pre-amended
13 with TMB solution (0.238 mL at 2.60 mM in methanol) and quenched at room temperature ($21 \pm$
14 1°C). Vials were capped and shaken manually for 10 s. Concentrations of unreacted TMB and Cl-
15 TMB were determined using HPLC with a diode array detector (DAD). Additional method details
16 are available in the ESI (**Figure S6**).
17
18
19
20
21
22
23
24
25
26
27
28

29 **2.4 Determination of trace bromide in laboratory salts and acids**

30
31 The ability of TMB to react with free bromine to generate Br-TMB was utilized to detect
32 trace bromide impurities in 20 reagents, including salts, acids, and bases (**Table S5**). Solutions
33 (25.0 mL) of each reagent were prepared in 40-mL amber glass vials with initial concentrations
34 ranging from 0.31 to 1.4 M (**Table S5**). An aliquot of a NaOCl primary dilution (0.45 mL at 7.5
35 mM) was added to each 40-mL vial, which were capped, shaken manually for 10 s, and allowed
36 to stand for 5 min at room temperature ($21 \pm 1^\circ\text{C}$) to permit bromide oxidation. Aliquots from
37 these vials (0.700 mL) were transferred to 4-mL amber glass vials containing methanolic TMB
38 solution (0.150 mL at 2.60 mM). The 4-mL vials were capped, shaken manually for 30 s, and
39 allowed to stand for 2 min to facilitate derivatization.
40
41
42
43
44
45
46
47
48
49
50
51

52 To assess whether nominally bromide-free salts can interfere with quantification of
53 bromide, experiments were performed in which 25-mL solutions were prepared in 40-mL amber
54
55
56
57
58
59
60

1
2
3 glass vials containing NaBr (30.0 μM , obtained from a 1003 mg $\text{Br}^- \text{L}^{-1}$ certified standard solution)
4
5 and a second (background) salt, typically at 1.0 M (**Table 1**). Examined background salts included
6
7 NaCl (99.999%), NaHCO_3 (99.7+%), Na_2SO_4 (99.9%), NaNO_3 (99.0+%), and $\text{NaI}\cdot 2\text{H}_2\text{O}$ (99+%).
8
9 To each 25-mL solution containing a background salt was added NaOCl (0.20 mL at 12.5 mM).
10
11 Vials were capped, shaken manually for 10 s, and allowed to stand at room temperature (21 ± 1
12
13 $^\circ\text{C}$) to permit bromide oxidation. Control experiments were performed with each background salt
14
15 in the absence of added NaBr. Three aliquots (0.800 mL each) were obtained from each salt +
16
17 NaOCl solution and transferred to individual 4-mL amber glass vials, which were pre-amended
18
19 with TMB (0.175 mL at 2.24 mM in methanol). The 4-mL vials were capped, shaken manually for
20
21 30 s, and allowed to stand for 2 min to facilitate reaction of TMB with free bromine and residual
22
23 free chlorine.
24
25
26
27
28
29

30
31 **Table 1.** Experimental Systems Investigating the Effects of Background Salts
32 on Quantification of Bromide as Br-TMB Following Oxidation by NaOCl

System #	Background Salt	[Background Salt] ₀ (mM)	[NaBr] ₀ (mM)	[NaOCl] ₀ (mM)
1a	none	not applicable	0.0300	0.100
1b			none added	
2a	NaCl	1.00×10^3	0.0300	0.100
2b			none added	
3a	NaHCO_3	1.00×10^3	0.0300	0.100
3b			none added	
4a	Na_2SO_4	1.00×10^3	0.0300	0.100
4b			none added	
5a	NaNO_3	1.00×10^3	0.0300	0.100
5b			none added	
6a	$\text{NaI}\cdot 2\text{H}_2\text{O}$	0.0300	0.0300	0.100
6b		0.200	0.0300	
6c		0.200	none added	

53
54
55
56
57
58
59
60

1
2
3 After all samples were quenched, similar liquid-liquid extractions as described in Section
4 2.1 were performed, using 0.80 mL of toluene containing 2-chlorobenzonitrile (10.2 μM) as the
5 internal standard. TMB and halogenated products were analyzed via GC–MS (**Table S2**).
6
7
8
9
10
11

12 **2.5 Quantification of halogens generated by spa disinfecting tablets**

13

14 The ability of TMB to serve as a probe of halogens was assessed using simulated spa
15 solutions treated with disinfecting tablets containing BCDMH. Simulated spa solutions (30 L)
16 were prepared in a 39-L clear glass aquarium containing municipal (chlorinated) tap water.
17 Simulated spa solutions were left open to the air and incubated in a circulating water bath at 39.0
18 \pm 0.1 $^{\circ}\text{C}$ for 1 h prior to dosing with one spa disinfecting tablet (Clorox spa brominating tablets,
19 28 g, 96.0 wt% BCDMH). After tablet addition (at $t = 0$), solutions were gently stirred manually
20 for 20 s. Aliquots (1.00 mL) were obtained prior to addition of a disinfecting tablet and hourly
21 thereafter over 8 h. Prior to each sampling event, solutions were stirred manually for 10 s. Four
22 aliquots were obtained at each time point; two aliquots were added to separate 2-mL vials pre-
23 amended with TMB (0.14 mL at 5.56 mM in methanol), and two aliquots were added to separate
24 2-mL vials pre-amended with 2,6-dichlorophenol (0.13 mL at 6.04 mM in methanol). Following
25 addition of an aliquot, the 2-mL vials were capped and shaken manually for 10 s. After all samples
26 were quenched, TMB, 2,6-dichlorophenol, and their halogenated products were quantified by
27 HPLC-DAD (see ESI for method details).
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 **3. Results and discussion**

49

50
51 Many traditional methods of quantifying free halogens are unable to distinguish between
52 free bromine and free chlorine.^{26,51} The principal goal of this work was to design methods for
53 selectively quantifying free chlorine and free bromine in a variety of aqueous systems via
54
55
56
57
58
59
60

1
2
3 derivatization with TMB. As is common in the disinfection literature, mass-based concentrations
4
5 of both free chlorine and free bromine are expressed herein as mg (or μg) of Cl_2 per L .
6
7
8
9

10 11 **3.1 Comparison of free halogen quantification by TMB and DPD** 12

13 In **Table 2**, analytical metrics associated with the TMB GC-MS and TMB HPLC methods
14 are compared with those of three previously reported methods, including the DPD method. For
15 HOCl and HOBr, the TMB GC-MS method provides substantially lower method quantitation
16 limits (MQLs) relative to the TMB HPLC method, due in part to the preconcentration step
17 associated with the GC-MS method. The MQL for GC-MS determination of free chlorine by TMB
18 is a factor of 7 lower than that of the DPD method. The MQL associated with free bromine
19 derivatization by TMB (60 nM by GC-MS) is greater than that previously reported for 2,6-
20 dimethylphenol (1.4 nM by GC-MS)³⁷ and for DMPyr (0.018 nM by LC-MS/MS).³⁹
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2. Comparison method detection limits (MDL), method quantitation limits (MQL), and instrument upper limits of linearity (LOL) of selected free halogen quantification methods^a

Analyte	Method	MDL (μM)	MQL (μM)	LOL (μM)	Ref
Free Chlorine	Cl-TMB by HPLC ^b	1.4	5	600	current work
	Cl-TMB by GC-MS ^b	0.010	0.03	120	current work
	2,4,6-trichlorophenol by HPLC ^c	5	18	600	current work
	Cl-DMP by GC-MS ^d	8×10^{-4}	0.003	59	37
	DPD ^e	0.06	0.19	70	51
Free Bromine	Br-TMB by HPLC ^b	1.0	3	600	current work
	Br-TMB by GC-MS ^b	0.017	0.06	100	current work
	4-bromo-2,6-dichlorophenol by HPLC ^c	6	19	600	current work
	Br-DMP by GC-MS ^d	4×10^{-4}	0.0014	11	37
	Br-DMPyr by LC-MS ^f	6×10^{-6}	1.8×10^{-5}	0.12	39
Bromide	Br-TMB by GC-MS ^b	0.017	0.06	100	current work

^a MDL (3s/m) and MQL (10s/m) values account for preconcentration (e.g., associated with liquid-liquid extraction prior to GC-MS analysis) and extraction efficiencies (Table S2). Concentration units of $\mu\text{g L}^{-1}$ for free chlorine and free bromine correspond to $\mu\text{g L}^{-1}$ as Cl_2 . Uncertainties denote 95% confidence intervals.

^b MDL, MQL, LOL, and slope values for TMB are provided in Table S6.

^c This method involves the use of 2,6-dichlorophenol to derivatize free chlorine (2,4,6-trichlorophenol) and free bromine (yielding 4-bromo-2,6-dichlorophenol).

^d This method pertains to the use of 2,6-dimethylphenol (DMP) to derivatize free chlorine (yielding 4-chloro-2,6-dimethylphenol, Cl-DMP) and free bromine (yielding 4-bromo-2,6-dimethylphenol, Br-DMP) in aquarium seawater treated with ozone.

^e The DPD method is not selective between free chlorine and free bromine. For a more extensive list of free chlorine detection methods, including comparisons of MDL values, see ref 51.

^f This method pertains to the use of 3,5-dimethyl-1H-pyrazole (DMPyr) to derivatize free bromine and other reactive bromine species (yielding 4-bromo-3,5-dimethyl-1H-pyrazole, Br-DMPyr); the use of DMPyr to derivatize free chlorine was not reported in ref 39.

The TMB GC-MS method and the DPD method were compared (with respect to measured total free halogen concentrations) using synthetic waters containing free chlorine, free bromine, an equimolar mixture of free chlorine + free bromine, as well as chlorinated pool water (**Table 3**).

Overall, treatment with TMB or DPD did not yield significantly different results at the 95% confidence level, which suggests that the TMB method performs comparably to the DPD method for quantifying free halogens. Furthermore, the use of both methods permitted recovery of approximately 100% of free bromine and/or free chlorine in synthetic waters.

Table 3. Summary of paired Student's t test results between DPD and TMB GC-MS methods^a

Sample	Average Measured Total Free Halogen Concentration (μM)			Significantly Different at 95% Confidence Level?
	DPD method	TMB method	Difference	
Synthetic water HOCl only ^b	14.72	16.31	-1.59	No (p-value = 0.11)
	15.70	16.15	-0.46	
	15.62	16.38	-0.75	
Synthetic water HOBr only ^c	21.85	20.61	1.24	No (p-value = 0.06)
	21.74	21.07	0.67	
	21.79	20.07	1.72	
Synthetic water HOCl + HOBr ^c	19.77	20.75	-0.98	No (p-value = 0.07)
	19.94	20.36	-0.42	
	19.50	19.99	-0.49	
Chlorinated Pool Water ^e	27.92	28.13	-0.21	No (p-value = 0.70)

^a Conditions (unless specified otherwise): pH = 7.0, [phosphate buffer] = 20 mM, $[\text{HOCl}]_{\text{tot},0} \approx 20 \mu\text{M}$, $[\text{NaCl}] = 10 \text{ mM}$, $[\text{NaNO}_3] = 90 \text{ mM}$, $T = 20.0 \text{ }^\circ\text{C}$. Total free halogen concentration by the TMB method calculated as $[\text{Cl-TMB}] + [\text{Br-TMB}]$; p-values are two-tailed.

^b No NaBr added.

^c $[\text{NaBr}]_0 = 40 \mu\text{M}$.


^d $[\text{NaBr}]_0 = 10 \mu\text{M}$.

^e Pool water samples obtained from an indoor swimming pool in Baltimore County, Maryland. Conditions: pH = 7.4, $T = 21 \text{ }^\circ\text{C}$. Values denote averages of four sampling locations in the pool that were measured in triplicate. The full data set is provided in Table S7.


3.2 Assessing the reactivity of TMB towards aqueous combined bromine

To determine the inherent reactivity of TMB toward combined bromine, bromamines were prepared by dosing free bromine solutions (prepared from $\text{NaOCl} + \text{Br}^-$) with ammonium chloride at pH 8.0 and a $\text{HOBr}_{\text{tot}}:\text{NH}_3_{\text{tot}}$ molar ratio of approximately 1:3. The principal bromamine species formed in such solutions is anticipated to be monobromamine (NH_2Br , eq 6).

At near-neutral pH, solutions of ammonia and free bromine can also rapidly generate dibromamine, which can form via disproportionation of monobromamine.^{9,48}



Similarly, disproportionation of dibromamine can generate tribromamine (NBr_3).



Consequently, bromination of TMB in solutions containing bromamines will reflect the net reactivity of all active bromine species in solution, including free bromine species and various bromamines. Under our solution conditions, the equilibrium position of **eq 6** favors the products and $[\text{NH}_2\text{Br}(aq)]/[\text{HOBr}_{\text{tot}}(aq)]$ is calculated to be 4×10^4 . The ratio of [Br-TMB] generated from bromamines to [Br-TMB] generated from free bromine ranges from 74 – 87% and does not appear to vary regularly as a function of sampling time (**Table 4**). These findings suggest that bromamines are reactive brominating agents towards TMB and that bromamination of TMB occurs under approximately the same timescale as bromination by free bromine (half-lives of seconds). Unlike bromamines, chloramines react sluggishly with TMB (half-lives of hours);²¹ this result is consistent with previous reports of bromamines serving as more labile halogenating agents compared to their chlorinated counterparts.⁴⁸

Table 4. Summary of average Br-TMB yields from reactions of TMB with bromamines and with free bromine^a

Sampling Time	Average [Br-TMB] (μM)		$\frac{[\text{Br-TMB}]_{\text{Bromamines}}}{[\text{Br-TMB}]_{\text{HOBr}}} \times 100 (\%)$
	TMB + Bromamines	TMB + HOBr	
10 s	77.4 ± 1.3	92.7 ± 1.3	83.5 ± 1.8
20 s	75 ± 2	93.3 ± 1.1	80 ± 2
30 s	65.6 ± 1.3	79.8 ± 1.6	74 ± 2
60 s	69.5 ± 1.6	83 ± 2	87 ± 3

^a Conditions: pH = 8.0, [borate buffer] = 20 mM, $[\text{NaBr}]_0 = 100 \mu\text{M}$, $[\text{NaOCl}]_0 = 155 \mu\text{M}$, $[\text{NH}_4\text{Cl}]_0 = 0$ (HOBr reactors) or $300 \mu\text{M}$ (bromamine reactors), $[\text{NaCl}] = 10 \text{ mM}$, $[\text{NaNO}_3] = 90 \text{ mM}$, $T = 20.0 \text{ }^\circ\text{C}$. Uncertainties denote 95% confidence intervals. The decreased recovery of bromine in the bromamine reactors compared to the free bromine reactors could result from irreversible decomposition of bromamines, which is subject to base catalysis.^{9,52}

3.3 Quantification of free halogens in chlorinated surface waters and drinking water

Free chlorine was added to surface water samples to model disinfection of drinking water and saline ballast water. Following a free chlorine contact time of 30 s, aliquots were taken from each water sample and quenched with TMB. Recoveries of total free halogens in Chesapeake Bay and Susquehanna River water were somewhat less than recoveries in Atlantic Ocean and Loch Raven Reservoir water (**Table 5**), owing to possible reaction of free chlorine and free bromine with natural organic matter (NOM) or other nucleophilic species in each water source.⁵³ Free chlorine residuals measured in municipal drinking water ranged from 0.2 – 1.2 mg L⁻¹ as Cl₂ (**Table S4**), consistent with the range (0.2 – 4 mg L⁻¹ as Cl₂) mandated by the U.S. Environmental Protection Agency.⁵⁴ Overall, these findings demonstrate the ability of TMB to facilitate selective quantification of HOCl and HOBr in chlorinated waters.

Table 5. Total free halogen recoveries measured in natural water during simulated disinfection.^a

Natural Water Source	[HOCl] _{tot} + [HOBr] _{tot} Recovery (%) ^b	Relative Standard Deviation (%)
Chesapeake Bay	91 ± 2	1.2
Atlantic Ocean	97 ± 4	2.6
Susquehanna River	92 ± 3	2.5
Loch Raven Reservoir	100 ± 4	2.7

^a Conditions: contact time = 30 s, [NaBr]_{added} = 150 μM (only in Susquehanna River and Loch Raven Reservoir water), [HOCl]_{tot,o} = 100 μM. Error estimates represent 95% confidence intervals.

^b [HOCl]_{tot} + [HOBr]_{tot} Recovery = $\frac{[\text{HOCl}]_{\text{TMB}} + [\text{HOBr}]_{\text{TMB}}}{[\text{HOCl}]_{\text{tot,o}}} \times 100\%$

3.4 Background salt interference studies

The potential for nominally bromide-free background salts to interfere with the detection of bromide was examined with NaCl, NaHCO₃, Na₂SO₄, NaNO₃, and NaI•2H₂O (**Figure 2**). Compared to a reactor receiving no background salt, measured Br⁻ concentrations appeared to increase by 8% in the presence of NaCl (one-tailed p-value = 0.08) and decreased by ~10% in the

presence of NaHCO_3 , Na_2SO_4 , and NaNO_3 (all present at 3.3×10^5 molar excess relative to NaBr ; all one-tailed p-values < 0.05). The modest increase in the presence of NaCl could result from the presence of Br^- as an impurity in the NaCl , as evidenced by the detection of Br^- in a no-added NaBr control (**Figure S7**). Bromide was not detected in NaBr -free controls for any of the other examined background salts.




Figure 2. Influence of background salts on quantification of bromide and total free chlorine via derivatization with TMB to give Br-TMB and Cl-TMB . Conditions: $[\text{NaBr}]_0 = 30 \mu\text{M}$, $[\text{background salt}]_0 = 1.00 \text{ M}$ (excepting NaI , which was added at 0.200 mM as $\text{NaI} \cdot 2\text{H}_2\text{O}$), $[\text{NaOCl}]_0 = 0.100 \text{ mM}$. Free chlorine concentration was calculated as $[\text{Cl-TMB}] + [\text{Br-TMB}]$ (eq 5). Error bars denote 95% confidence intervals. Asterisks (*) denote not detected. Data for control experiments performed in the absence of added NaBr are shown in Figure S7. Of the background salts examined in the absence of added NaBr , bromide was only detected in NaCl (Cl^-/Br^- molar ratio = 1.2×10^5 , corresponding to a Br^- concentration in the NaCl of 11 ppm by weight).

1
2
3 The modest decrease in bromide measured in the presence of large excesses of NaHCO_3 , Na_2SO_4 ,
4 and NaNO_3 could result from low-level impurities capable of scavenging free bromine. The
5
6 apparent incomplete recovery of free chlorine in **Figure 2** is discussed in the ESI.
7
8
9

10 NaI (0.200 mM) completely precluded detection of bromide (added at 0.030 mM) and free
11 chlorine (added at 0.100 mM). When the concentration of NaI was decreased to 0.030 mM, 0.7
12 μM of bromide was measured (data not shown in **Figure 2**), which corresponds to a spike recovery
13 of 2%. These results indicate that iodide can interfere with bromide quantification as Br-TMB.
14 Iodide is a known reductant of free chlorine/bromine, which can generate free iodine (e.g., HOI)
15 and other electrophilic iodine species.^{55,56} Free iodine can react with nucleophilic organic
16 compounds,⁵⁷⁻⁵⁹ including TMB,⁶⁰ to yield iodinated products. Thus, iodide could adversely
17 impact free chlorine/bromine quantitation by (1) serving as a reductant of free chlorine/bromine
18 and, subsequently, (2) generating free iodine, which could subsequently iodinate TMB (or other
19 derivatizing agents) and their chlorinated/brominated products.
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 **3.5 Determination of trace bromide in laboratory salts and acids**

35
36 Reagents from chemical suppliers commonly report bromide as an impurity in salts such
37 as NaCl and acids such as HCl. The presence of bromide as an impurity in these reagents can
38 complicate experiments involving free chlorine, chloramines, and ozone due to the ability of these
39 disinfectants to oxidize bromide into free bromine, which can be several orders of magnitude more
40 inherently reactive as a halogenating agent relative to free chlorine.^{22,53,61} Indeed, low levels of
41 bromide introduced by NaCl have been shown to enhance rates of aromatic compound
42 consumption in experiments involving free chlorine.^{24,62} However, high background
43 concentrations of anions (e.g., chloride) can compromise the accuracy of some methods when
44 measuring bromide.^{63,64}
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

When solutions of 20 reagents (**Table S5**) were treated with free chlorine followed by TMB, bromide impurity levels were greater than the MQL (60 nM = 0.005 ppm) in 7 of the 20 examined reagents (**Table 6**). The TMB method employed herein was capable of detecting ppm and sub-ppm levels of bromide present in a high chloride background (≥ 1 M) with relative standard deviations $< 9\%$. Quantified bromide levels in high-purity NaCl (3 ppm Br⁻) are approximately three times less than the upper limit reported by the manufacturer.

Table 6. Summary of trace bromide impurity levels in reagent solutions treated with NaOCl and TMB.^a

Reagent	Description	[Br ⁻] (ppm) Reported By Manufacturer	Average [Br ⁻] (ppm) Measured As Br-TMB	Relative Standard Deviation (%)
NaCl	Higher purity (99.999%)	< 10	3	8.5
	Lower purity (99.0%)	< 100	37	6.4
	Pool salt	not reported	146	2.6
	Table salt	not reported	167	2.6
HCl	Fisher Scientific	< 50	34	7.8
	Fisher Chemical	< 50	50	7.2
	J.T. Baker (ultra-pure)	not reported	0.72	6.0

^a Of the 20 examined reagents listed in Table S5, bromide levels exceeded the method quantitation limit (0.005 ppm) for the seven reagents shown above.

The ability of TMB to react with free bromine to yield a stable product in a high chloride background permits precise quantification of bromide as a trace impurity in a variety of laboratory reagents. Ion-selective electrodes (ISEs)⁶³ and ion chromatography (IC)⁶⁴ techniques can also be used to quantify bromide in solutions containing chloride. A bromide-ISE permits bromide quantification, but is limited to a maximum mole ratio of Cl⁻:Br⁻ of 400:1 and a relatively high MQL (1 ppm).⁶³ Conversely, IC can accurately determine bromide levels in aqueous solutions containing a Cl⁻:Br⁻ mole ratio of 2500:1 (MQL = 30 ppb).⁶⁴ The solution conditions employed herein included relatively high concentrations of NaCl and HCl (~1.0 M) compared to the concentrations of bromide (3–167 μ M); a lesser chloride concentration would be expected in

1
2
3 typical halogenation studies mirroring surface water conditions (median $[\text{Cl}^-] = 0.3 \text{ mM} = 11 \text{ mg}$
4 L^{-1}).^{5,22} Accordingly, the ratio of $\text{Cl}^-:\text{Br}^-$ in the examined salt solutions ($\sim 10^6:1$) far exceeds the
5
6 ratios observed in surface waters ($\sim 250:1$; see **Table S3**). Our results indicate that the TMB method
7
8 described herein can afford trace quantitation of bromide in the presence of high chloride
9
10 backgrounds without the need for more time-intensive methodologies (e.g., standard addition).
11
12
13
14

15 **3.6 Quantification of active halogens generated by spa disinfecting tablets**

16
17
18 The ability of TMB to serve as a halogen trap was applied to tap water treated with spa
19
20 disinfecting tablets. The active ingredient in the tablets was BCDMH, which hydrolyzes to yield
21
22 free bromine and free chlorine (**Scheme 1**). In waters treated with BCDMH, TMB can conceivably
23
24 be halogenated by free chlorine, free bromine, or by BCDMH itself. Accordingly, for such
25
26 systems, Cl-TMB and Br-TMB serve as surrogates of *active* chlorine and *active* bromine,
27
28 respectively, not just free chlorine and free bromine.
29
30

31
32 In the simulated spa water quenched with TMB, active bromine residuals increased
33
34 continually over 8 h to a maximum of $79 \mu\text{M}$ ($5.6 \text{ mg Cl}_2 \text{ L}^{-1}$, **Figure 3A**). In the same system,
35
36 active chlorine residuals increased up to $40 \mu\text{M}$ ($2.9 \text{ mg Cl}_2 \text{ L}^{-1}$) over 8 h. When 2,6-dichlorophenol
37
38 was used as the derivatizing agent for aliquots obtained from the same simulated spa water, the
39
40 active bromine profile (**Figure 3B**) was in strong agreement with that obtained from samples
41
42 quenched with TMB. Indeed, a paired t test (two-tailed) indicated that the active bromine data
43
44 obtained with TMB and with 2,6-dichlorophenol were not significantly different at the 95%
45
46 confidence level ($p = 0.38$).
47
48
49

50
51 Notably, active chlorine was not detected in any of the samples treated with 2,6-
52
53 dichlorophenol. The somewhat greater water solubility of 2,4,6-trichlorophenol (4.3 mM)⁶⁵
54
55 relative to 4-bromo-2,6-dichlorophenol (3.4 mM)⁶⁶ suggests that solubility limitations cannot
56
57
58
59
60

account for the failure to detect active chlorine when 2,6-dichlorophenol served as the derivatizing agent. These results suggest that 2,6-dichlorophenol is less nucleophilic than TMB under the conditions of the simulated spa solutions (e.g., pH 7.3). Under these conditions, 2,6-dichlorophenol ($pK_a = 6.78$, presumably corresponding to 25 °C)⁶⁷ exists predominantly in the more nucleophilic (anionic) form and HOCl ($pK_a = 7.46$ at 35 °C)¹¹ exists predominantly in the more electrophilic (neutral) form. Accordingly, the simulated spa conditions should ostensibly favor reactions between free chlorine and 2,6-dichlorophenolate. That 2,6-dichlorophenol was reactive toward active brominating (but not chlorinating) agents is consistent with the greater inherent electrophilicity of active bromine species compared to their chlorinated counterparts.⁶¹ Overall, these findings illustrate the ability of the TMB method to selectively quantify active bromine and active chlorine in waters disinfected with BCDMH.




Figure 3. Active halogen residuals generated by spa disinfecting tablets (28 g, 96.0 wt% BCDMH) quantified using (A) TMB and (B) 2,6-dichlorophenol as the derivatizing agent. Tap water served as the reaction matrix. Error bars denote standard deviations ($n = 2$) and are smaller than symbols if not shown. Solution conditions: pH = 7.3, $T = 39.0 \pm 0.1$ °C, total volume = 30 L. Neither active chlorine nor active bromine were detected in the water prior to addition of the spa tablet. In frame A, active bromine and active chlorine were not detected until 2 h and 3 h, respectively, after tablet addition at $t = 0$. In frame B, active chlorine was not detected at any of the sampling times.

4. Summary of Methodological Advantages and Limitations

TMB can be used to derivatize free chlorine and free bromine into stable chlorinated and brominated products that can be quantified using GC-MS and HPLC-DAD. TMB can be used to quantify free chlorine and free bromine in solutions of varying matrix complexity, including synthetic waters treated with NaOCl, drinking water, pool water, and a variety of chlorinated surface waters.

Performance of TMB as a free chlorine probe gave results that were on par with those of the traditional DPD method. Unlike the DPD method, however, the halogenation products of TMB are stable and therefore do not require immediate analysis. Moreover, while the DPD method cannot distinguish between free chlorine and free bromine, derivatization with TMB permits simultaneous and selective quantification in solutions containing both free halogens. The large linear range of the TMB method for free chlorine and for free bromine can be tailored to meet the requirements of individual samples by varying the extent of preconcentration (i.e., during solvent extraction prior to GC-MS analysis) or dilution. For the GC-MS method reported herein, MDLs for free chlorine and free bromine are on the order of $1 \mu\text{g L}^{-1}$ as Cl_2 .

The formation of Br-TMB in solutions treated with free chlorine can be used to quantify trace bromide present as an impurity in laboratory reagents with substantially lower MQLs than traditional assays. Unlike alternative methods (e.g., ion-selective potentiometry and IC), the presence of high chloride backgrounds does not appear to limit the performance of the TMB method described herein when quantifying bromide.

While our results demonstrate that TMB is sufficiently nucleophilic to serve as a derivatizing agent for inorganic bromamines, TMB is unlikely to be able to distinguish between free bromine and inorganic bromamines. As with other derivatization methods relying on LC or

1
2
3 GC,^{37,68} the TMB method reported herein is not suitable for applications requiring remote
4 monitoring, continuous analysis, or in laboratories lacking access to both HPLC-DAD and GC-
5 MS. The TMB method is also susceptible to interference by iodide, which is likely to affect other
6 derivatizing agents that are reactive toward free iodine.
7
8
9
10

11
12 Together with our recent work,²¹ this study establishes a range of applications by which
13 TMB can be used to quantify free chlorine, free bromine, inorganic bromamines, and bromide.
14 The potential utility of TMB as a derivatizing agent of aqueous free iodine and iodide (upon
15 oxidation to free iodine) merits future investigation.
16
17
18
19
20
21
22

23 **Conflicts of interest**

24
25
26 There are no conflicts to declare.
27
28
29
30
31

32 **Acknowledgements**

33
34
35 The authors gratefully acknowledge the helpful comments received from the anonymous peer
36 reviewers. The authors also thank Prof. Jay Nelson (Towson University, Department of Biology)
37 for his technical assistance. Funding to J. D. S. is acknowledged from the American Chemical
38 Society Petroleum Research Fund (54560-UNI4) and the U.S. National Science Foundation
39 (CBET-1651536). Funding to J. D. S. and K. P. R. was received from Jess and Mildred Fisher
40 Endowed Professorships. Funding to R. P. D. and M. H. S. is acknowledged from the Fisher
41 College of Science and Mathematics and the Office of Undergraduate Research and Creative
42 Inquiry at Towson University. Any views reported herein are those of the authors and do not
43 necessarily reflect the views of the American Chemical Society Petroleum Research Fund or the
44 U.S. National Science Foundation.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


References

1. *Black & Veatch Corporation. White's Handbook of Chlorination and Alternative Disinfectants*, Wiley, Hoboken, NJ, 5th edn., 2010.
2. T. K. Das, Disinfection, in *Kirk-Othmer Encyclopedia of Chemical Technology*, Wiley, New York, 2002.
3. E. Tsolaki and E. Diamadopoulos, Technologies for ballast water treatment: a review, *J. Chem. Technol. Biotechnol.*, 2010, **85**, 19-32.
4. M. Deborde and U. von Gunten, Reactions of chlorine with inorganic and organic compounds during water treatment - Kinetics and mechanisms: A critical review, *Water Res.*, 2008, **42**, 13-51.
5. J. D. Sivey and A. L. Roberts, Assessing the reactivity of free chlorine constituents Cl₂, Cl₂O, and HOCl toward aromatic ethers, *Environ. Sci. Technol.*, 2012, **46**, 2141-2147.
6. A. Dong, Y.-J. Wang, Y. Gao, T. Gao and G. Gao, Chemical insights into antibacterial N-halamines, *Chem. Rev.*, 2017, **117**, 4806-4862.
7. J. D. Sivey, M. A. Bickley and D. A. Victor, Contributions of BrCl, Br₂, BrOCl, Br₂O, and HOBr to regiospecific bromination rates of anisole and bromoanisoles in aqueous solution, *Environ. Sci. Technol.*, 2015, **49**, 4937-4945.
8. R. Frim and S. D. Ukeles, Bromine, Inorganic Compounds, in *Kirk-Othmer Encyclopedia of Chemical Technology*, John Wiley & Sons, Inc., 2000.
9. M. B. Heeb, J. Criquet, S. G. Zimmermann-Steffens and U. von Gunten, Oxidative treatment of bromide-containing waters: Formation of bromine and its reactions with inorganic and organic compounds — A critical review, *Water Res.*, 2014, **48**, 15-42.
10. S. J. Weiss, S. T. Test, C. M. Eckmann, D. Roos and S. Regiani, Brominating oxidants generated by human eosinophils, *Science*, 1986, **234**, 200-203.
11. J. C. Morris, The Acid Ionization Constant of HOCl from 5 to 35 °C, *J. Phys. Chem.*, 1966, **70**, 3798-3805.
12. M. Eigen and K. Kustin, The kinetics of halogen hydrolysis, *J. Am. Chem. Soc.*, 1962, **84**, 1355-1361.
13. J. L. Guentzel, K. L. Lam, M. A. Callan, S. A. Emmons and V. L. Dunham, Reduction of bacteria on spinach, lettuce, and surfaces in food service areas using neutral electrolyzed oxidizing water, *Food Microbiol.*, 2008, **25**, 36-41.
14. W. A. Rutala, D. J. Weber and Healthcare Infection Control Practices Advisory Committee, *Guideline for Disinfection and Sterilization in Healthcare Facilities*, U.S. Centers for Disease Control and Prevention, 2008.
15. B. Kim, J. Anderson, S. Mueller, W. Gaines and A. Kendall, Literature review—efficacy of various disinfectants against *Legionella* in water systems, *Water Res.*, 2002, **36**, 4433-4444.
16. Funeral Service Academy, OSHA Compliance Guidance for Funeral Homes, <https://funeralcourse.com/wp-content/uploads/coursebooks/FuneralBook-OSHA-Compliance-for-Funeral-Homes.pdf>, (accessed 5 Mar 2019).
17. D. Sedlak and U. von Gunten, The chlorine dilemma, *Science*, 2011, **331**, 42-43.
18. X.-F. Li and W. A. Mitch, Drinking Water Disinfection Byproducts (DBPs) and Human Health Effects: Multidisciplinary Challenges and Opportunities, *Environ. Sci. Technol.*, 2018, **52**, 1681-1689.

19. Y. Yang, Y. Komaki, S. Y. Kimura, H.-Y. Hu, E. D. Wagner, B. J. Mariñas and M. J. Plewa, Toxic impact of bromide and iodide on drinking water disinfected with chlorine or chloramines, *Environ. Sci. Technol.*, 2014, **48**, 12362-12369.
20. U. Pinkernell, B. Nowack, H. Gallard and U. Von Gunten, Methods for the photometric determination of reactive bromine and chlorine species with ABTS, *Water Res.*, 2000, **34**, 4343-4350.
21. S. S. Lau, R. P. Dias, K. R. Martin-Culet, N. A. Race, M. H. Schammel, K. P. Reber, A. L. Roberts and J. D. Sivey, 1,3,5-Trimethoxybenzene (TMB) as a new quencher for preserving redox-labile disinfection byproducts and for quantifying free chlorine and free bromine, *Environ. Sci.: Water Res. Technol.*, 2018, **4**, 926-941.
22. J. D. Sivey, J. S. Arey, P. R. Tentscher and A. L. Roberts, Reactivity of BrCl, Br₂, BrOCl, Br₂O, and HOBr toward dimethenamid in solutions of bromide + aqueous free chlorine, *Environ. Sci. Technol.*, 2013, **47**, 1330-1338.
23. M. B. Heeb, J. Criquet, S. Zimmermann-Steffens and U. von Gunten, Bromine production during oxidative water treatment of bromide-containing waters and its reactions with inorganic and organic compounds – A critical review, *Water Res.*, 2014, **48**, 15-42.
24. J. D. Sivey, C. E. McCullough and A. L. Roberts, Chlorine monoxide (Cl₂O) and molecular chlorine (Cl₂) as active chlorinating agents in reaction of dimethenamid with aqueous free chlorine, *Environ. Sci. Technol.*, 2010, **44**, 3357-3362.
25. S. D. Worley, D. E. Williams and R. A. Crawford, Halamine water disinfectants, *Crit. Rev. Environ. Control*, 1988, **18**, 133-175.
26. E. W. Rice, R. B. Baird, L. S. Clesceri and A. D. Eaton, eds., *Standard Methods for the Examination of Water and Wastewater*, American Public Health Association, American Water Works Association, Water Environment Federation, Washington, D.C., 2012.
27. Y. Xiong, J. Tan, S. Fang, C. Wang, Q. Wang, J. Wu, J. Chen and M. Duan, A LED-based fiber-optic sensor integrated with lab-on-valve manifold for colorimetric determination of free chlorine in water, *Talanta*, 2017, **167**, 103-110.
28. M. M. Nakamura, N. Coichev, J.-M. Lin and M. Yamada, Flow-injection investigation of the chemiluminescent reaction of bis (2, 4, 6-(trichlorophenyl) oxalate) with free chlorine, *Anal. Chim. Acta*, 2003, **484**, 101-109.
29. Y. Dong, G. Li, N. Zhou, R. Wang, Y. Chi and G. Chen, Graphene quantum dot as a green and facile sensor for free chlorine in drinking water, *Anal. Chem.*, 2012, **84**, 8378-8382.
30. Y. Tang, Y. Su, N. Yang, L. Zhang and Y. Lv, Carbon nitride quantum dots: A novel chemiluminescence system for selective detection of free chlorine in water, *Anal. Chem.*, 2014, **86**, 4528-4535.
31. K. Singh and S. K. Mehta, Luminescent ZnO quantum dots as an efficient sensor for free chlorine detection in water, *Analyst*, 2016, **141**, 2487-2492.
32. P. B. D. De la Mare, *Electrophilic halogenation: Reaction pathways involving attack by electrophilic halogens on unsaturated compounds*, Cambridge University Press, Cambridge, 1976.
33. E. V. Anslyn and D. A. Dougherty, *Modern Physical Organic Chemistry*, University Science, Sausalito, CA, 2005.
34. A. D. Shah, Z.-Q. Liu, E. Salhi, T. Höfer, B. Werschkun and U. von Gunten, Formation of disinfection by-products during ballast water treatment with ozone, chlorine, and

- peracetic acid: influence of water quality parameters, *Environ. Sci. Water Res. Technol.*, 2015, **1**, 465-480.
35. A. D. Shah, Z.-Q. Liu, E. Salhi, T. Höfer and U. von Gunten, Peracetic acid oxidation of saline waters in the absence and presence of H₂O₂: Secondary oxidant and disinfection byproduct formation, *Environ. Sci. Technol.*, 2015, **49**, 1698-1705.
36. Z.-Q. Liu, A. D. Shah, E. Salhi, J. Bolotin and U. von Gunten, Formation of brominated trihalomethanes during chlorination or ozonation of natural organic matter extracts and model compounds in saline water, *Water Res.*, 2018, **143**, 492-502.
37. Z. Qiang, Y. Jiang, W. Ben, C. Adams and H. Dong, Monitoring free chlorine and free bromine in aquarium seawater treated by ozone, *Anal. Methods*, 2012, **4**, 3646-3652.
38. H.-S. Shin, Simple and simultaneous determination of free chlorine, free bromine and ozone in water by LC, *Chromatographia*, 2010, **71**, 647-651.
39. P. Lu, K. Lin, X. Huang and D. Yuan, A sensitive method for the determination of ultra trace levels of reactive bromine species in water using LC-MS/MS, *Talanta*, 2019, **199**, 567-572.
40. O. Tee, M. Paventi and J. Bennett, Kinetics and mechanism of the bromination of phenols and phenoxide ions in aqueous solution. Diffusion-controlled rates, *J. Am. Chem. Soc.*, 1989, **111**, 2233-2240.
41. O. S. Tee and N. R. Iyengar, Kinetics and mechanism of the bromination of phenols in aqueous solution. Evidence of general base catalysis of bromine attack, *Can. J. Chem.*, 1990, **68**, 1769-1773.
42. H. Gallard and U. von Gunten, Chlorination of phenols: Kinetics and formation of chloroform, *Environ. Sci. Technol.*, 2002, **36**, 884-890.
43. H. Gallard, F. Pellizzari, J. P. Croue and B. Legube, Rate constants of reactions of bromine with phenols in aqueous solution, *Water Res.*, 2003, **37**, 2883-2892.
44. J. Catalán and J. Elguero, Basicity of azoles. Part 2. Theoretical study of the basicity of methylpyrazoles and methylimidazoles, *J. Chem. Soc. Perkin Trans. 2*, 1983, 1869-1874.
45. C. G. Swain and D. R. Crist, Mechanisms of chlorination by hypochlorous acid. The last of chlorinium ion, Cl⁺, *J. Am. Chem. Soc.*, 1972, **94**, 3195-3200.
46. E. A. Voudrias and M. Reinhard, Reactivities of hypochlorous and hypobromous acid, chlorine monoxide, hypobromous acidium ion, chlorine, bromine, and bromine chloride in electrophilic aromatic substitution reactions with *p*-xylene in water, *Environ. Sci. Technol.*, 1988, **22**, 1049-1056.
47. U.S. Environmental Protection Agency, Chemistry Dashboard, <https://comptox.epa.gov/dashboard>, (accessed 7 Mar 2019).
48. M. B. Heeb, I. Kristiana, D. Trogolo, J. S. Arey and U. Von Gunten, Formation and reactivity of inorganic and organic chloramines and bromamines during oxidative water treatment, *Water Res.*, 2017, **110**, 91-101.
49. M. W. Beach and D. W. Margerum, Kinetics of oxidation of tetracyanonickelate(II) by chlorine monoxide, chlorine, and hypochlorous acid and kinetics of chlorine monoxide formation, *Inorg. Chem.*, 1990, **29**, 1225-1232.
50. Q. Liu and D. W. Margerum, Equilibrium and kinetics of bromine chloride hydrolysis, *Environ. Sci. Technol.*, 2001, **35**, 1127-1133.
51. R. E. Wilson, I. Stoianov and D. O'Hare, Continuous chlorine detection in drinking water and a review of new detection methods, *Johnson Matthey Technol. Rev.*, 2018, **63**, 103-118.

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
52. H. Lei, B. J. Mariñas and R. A. Minear, Bromamine decomposition kinetics in aqueous solutions, *Environ. Sci. Technol.*, 2004, **38**, 2111-2119.
53. P. Westerhoff, P. Chao and H. Mash, Reactivity of natural organic matter with aqueous chlorine and bromine, *Water Res.*, 2004, **38**, 1502-1513.
54. *Guidance Manual for Compliance with the Filtration and Disinfection Requirements for Public Water Systems Using Surface Water Sources*, U.S. Environmental Protection Agency, Washington, DC, 1991.
55. Y. Bichsel and U. Von Gunten, Oxidation of iodide and hypiodous acid in the disinfection of natural waters, *Environ. Sci. Technol.*, 1999, **33**, 4040-4045.
56. R. C. Troy and D. W. Margerum, Non-metal redox kinetics: hypobromite and hypobromous acid reactions with iodide and with sulfite and the hydrolysis of bromosulfate, *Inorg. Chem.*, 1991, **30**, 3538-3543.
57. E. Grovenstein, N. S. Aprahamian, C. J. Bryan, N. Gnanapragasam, D. C. Kilby, J. M. McKelvey and R. J. Sullivan, Aromatic halogenation. IV. Kinetics and mechanism of iodination of phenol and 2'6-dibromophenol, *J. Am. Chem. Soc.*, 1973, **95**, 4261-4270.
58. J. A. Warner, W. H. Casey and R. A. Dahlgren, Interaction kinetics of I₂ (aq) with substituted phenols and humic substances, *Environ. Sci. Technol.*, 2000, **34**, 3180-3185.
59. P. J. Vikesland, E. M. Fiss, K. R. Wigginton, K. McNeill and W. A. Arnold, Halogenation of bisphenol-A, triclosan, and phenols in chlorinated waters containing iodide, *Environ. Sci. Technol.*, 2013, **47**, 6764-6772.
60. J. Pavlinac, M. Zupan and S. Stavber, 'Green' Iodination of Dimethoxy- and Trimethoxy-Substituted Aromatic Compounds Using an Iodine-Hydrogen Peroxide Combination in Water, *Synthesis*, 2006, **2006**, 2603-2607.
61. M. A. Broadwater, T. L. Swanson and J. D. Sivey, Emerging investigators series: comparing the inherent reactivity of often-overlooked aqueous chlorinating and brominating agents toward salicylic acid, *Environ. Sci.: Water Res. Technol.*, 2018, **4**, 369-384.
62. K. Inaba, T. Doi, N. Isobe and T. Yamamoto, Formation of bromo-substituted triclosan during chlorination by chlorine in the presence of trace levels of bromide, *Water Res.*, 2006, **40**, 2931-2937.
63. Thermo Scientific, *Bromide Ion Selective Electrode User Guide*, 2008.
64. R. D. Rocklin and E. L. Johnson, Determination of cyanide, sulfide, iodide, and bromide by ion chromatography with electrochemical detection, *Anal. Chem.*, 1983, **55**, 4-7.
65. R. P. Schwarzenbach, P. M. Gschwend and D. M. Imboden, *Environmental Organic Chemistry*, John Wiley & Sons, 2016.
66. Data predicted using the U.S. Environmental Protection Agency's EPISuite program., <http://www.chemspider.com>, (accessed Sept. 6, 2019).
67. F. Muller and L. Caillard, Chlorophenols, in *Ullmann's Encyclopedia of Industrial Chemistry*, 2011.
68. A. Mayeno, A. Curran, R. Roberts and C. Foote, Eosinophils preferentially use bromide to generate halogenating agents, *J. Biol. Chem.*, 1989, **264**, 5660-5668.



19 Chlorine and bromine can be concurrently and selectively quantified via derivatization by 1,3,5-
20 trimethoxybenzene followed by GC-MS or HPLC-DAD analysis.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60