



ChemComm

Designing New Togni Reagents by Computation

Journal:	<i>ChemComm</i>
Manuscript ID	CC-COM-02-2019-001320.R2
Article Type:	Communication

SCHOLARONE™
Manuscripts

Designing New Togni Reagents by Computation

Heming Jiang,^{a,†,§} Tian-Yu Sun,^{b,a,c,†,§} Ying Chen,^{a,†,§} Xinhao Zhang,^{a,†,§} Yun-Dong Wu,^{*a,§} Yaoming Xie,^c and Henry F. Schaefer III^{*c}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/chemcomm

New trifluoromethylating reagents are designed via computational studies inspired by the Togni's experimental research. *Trans* influence and steric effects are important for this rational design. We have found that the Togni derivative with X = SO₂ is the best choice based on its high reactivity and its potential synthesis.


Fluorinated organic compounds play a conspicuous and increasingly important role in pharmaceuticals, agrochemicals and materials science.¹⁻⁶ However, few fluorinated organic compounds have been identified in nature.^{6,7} Therefore, it is very important to develop new methods for introducing fluorine into various organic molecules.

Trifluoromethylation is one of the most important reactions among such methods.⁸⁻¹¹ In recent years, the two reagents, Togni I and Togni II,¹² developed by Togni's group (see Fig. 1), have been used widely as trifluoromethylating reagents. Their effectiveness is in part due to their highly polarized hypervalent I-C (or I-O) bonds, and they are commercially available.^{10,13}

Electrophilic trifluoromethylation by Togni's reagents often needs activation by Lewis acids (see Fig. 2), Brønsted acids, or metals to weaken the I-O bond and promote the release of the CF₃⁺


unit.^{10,13,14} Lewis acids, Brønsted acids, and metals may disrupt the reaction system, and hence we desire to modify the structures of Togni's reagents to perhaps improve upon their intrinsic reactivity.

Trans influence plays an important role in the reactivity of hypervalent iodine reagents. The *trans* ligand has an influence on the stability of the ground state of the molecule, and the *trans* influence can be analyzed in terms of the bond lengths between the central atom and the ligand *trans* to a *trans*-influencing ligand.¹⁵⁻¹⁷ From Togni's research, shortening of the I-CF₃ bond, which is synonymous with lengthening of the I-O bond at the *trans* position, leads to increased electrophilic reactivity.^{18,19} A weaker *trans* influence of the I-O bond will cause a shorter I-CF₃ bond. Previous theoretical research²⁰⁻²² indicated changing the linker of the hypervalent iodine-based five-membered ring could change the *trans* influence (see Fig. 3). Accordingly, the reactivity of the Togni reagents will be changed. Therefore, we designed new Togni derivatives with different linkers to increase reactivity. Moreover, following previous work, the *steric* effect of the methyl group at the *ortho*-position of the aromatic ring may also enhance the reactivity (see Fig. 3).^{23,24} Thus, here we have designed additional new Togni derivatives using the *ortho*-methyl group.



Togni's reagent I Togni's reagent II

Fig. 1 Qualitative structures of Togni's reagents.



Electrophilic trifluoromethylation

Fig. 2 Togni's reagent II activated by Lewis acids.



Trans influence

Steric effect

Fig. 3 Designed Togni derivatives showing *trans* influence and *steric* effect.

^a Lab of Computational Chemistry and Drug Design, State Key Laboratory of Chemical Oncogenomics, Peking University Shenzhen Graduate School, Shenzhen 518055, China. E-mail: zhangxh@pkusz.edu.cn, wuyd@pkusz.edu.cn.

^b Key Laboratory of Bio-Inspired Smart Interfacial Science and Technology of Ministry of Education, School of Chemistry, Beihang University, Beijing 100191, China.

^c Center for Computational Quantum Chemistry, University of Georgia, Athens, GA 30602, USA. E-mail: ccq@uga.edu.

[†] Heming Jiang, Tian-Yu Sun and Ying Chen contributed equally.

[§] Institutions a and b contributed equally to this work.

Electronic Supplementary Information (ESI) available. See DOI: xx.xxxx/xxxxxxxxxx.

In the present research, computational studies were conducted to verify our hypotheses. The M06-2X²⁵ method was chosen as the functional, and cc-pVQZ(-PP)²⁶⁻²⁸ [the (-PP) suffix here represents the pseudopotential for iodine] used as the basis set. Structures were optimized in acetonitrile solvent with Truhlar's SMD method (Solvation Model based on the Quantum Mechanical Charge Density).²⁹ Our previous work showed that M06-2X/cc-pVQZ(-PP) describes the structures and energies of hypervalent iodine reagents reliably.^{20,21}

The highly electronegative linkers are expected to weaken the *trans* influence (lengthening the I-O bond and shortening the I-CF₃ bond), leading to the more active electrophilic trifluoromethylation reactions (Fig. 2).¹⁸⁻¹⁹ In the present study, eight Togni derivatives (1 - 8) with high electronegativity linkers are investigated (Fig. 4 and Table 1). we choose them based on a wide range of *trans* influence distribution. Among these, the newly designed Togni reagents, as well as the known Togni reagent I (8) and Togni reagent II (4) are included.

The *trans* influence of the linkers may be assessed in terms of the length of either the I-CF₃ bond or the I-O bond, and these bond distances for the eight Togni derivatives are reported in Table 1. From compound 1 to compound 8, the I-CF₃ bond distance (bond 5-6 in Table 1) sequentially increases from 2.194 Å to 2.257 Å, and the I-O bond (bond 5-4 in Table 1) decreases from 2.478 Å to 2.105 Å. This indicates that *trans* influence for the linkers of compound 1 is the weakest, while the compound 8 is the strongest. Table 1 also reports the dihedral angles (2-1-5-4) for the structures (1-8), which are measures of the torsion for the five-membered ring. In two of these (X = CO and BOH), the carbon atom in C=O and the boron atom in BOH display sp² hybridization to form a Π₅ conjugation system, leading to the planar five-membered rings. For other structures, since the linker central atoms take the sp³ hybridization without forming Π₅ cyclic conjugation systems, the five-membered rings are not necessarily planar. In fact, most of these structures have nonplanar five-membered rings with the dihedral angles (2-1-5-4) in the range from 7° to 21°. Table 1 shows that there is no correlation between the dihedral angle and the reactivity. For the known Togni I and Togni II, our computational geometries are consistent with their crystal data^{12,30} with the bond distance differences less than 0.02 Å and the bond angle differences less than 3° (Figure S4).

We studied the reactivity of a model trifluoromethylation reaction (shown at the top of Fig. 4), which includes the I-O bond breaking and the I-CF₃ bond hetero-cleavage. Since we have found that no energy barrier for such reactions exist, the reaction free energies were computed to measure the reactivity, which was supported by Xue, Cheng, and co-workers.³¹⁻³³ Table 1 shows that the reaction free energies steadily increase from 1 to 8, corresponding to the I-CF₃ and I-O distance changes, indicating the correlation between the electrophilic reactivity and the *trans* influence.

This strong correlation between the electrophilic reactivity and the *trans* influence may be seen more clearly in Fig. 4 (red line). That is, the weaker the *trans* influence, the more reactive the Togni derivative should be. This conclusion is supported by results for the known Togni reagent II (4) and Togni reagent I (8). Our predicted reaction free energy for Togni reagent II (4, 39.3 kcal mol⁻¹) is lower than that for Togni reagent I (8, 63.7 kcal mol⁻¹) by 24 kcal mol⁻¹, and this is consistent with the experimental result that Togni II is

more reactive than Togni I.¹⁸ To examine the eight Togni derivatives,



Reagents	1(X=SO ₂)	2(X=PO ₂ H)	3(X=CF ₂)	4(X=CO)	5(X=BOH)	6(X=NCH ₃)	7(X=CH ₂)	8(X=C(CH ₃) ₂)
bond (Å) 5-4	2.478	2.315	2.265	2.263	2.184	2.158	2.118	2.105
bond (Å) 5-6	2.194	2.207	2.207	2.212	2.232	2.239	2.254	2.257
angle (°) 2-1-5-4	16.8	6.9	0.0	0.0	0.0	20.8	15.1	14.3
reaction energy	25.1	32.7	37.0	39.3	48.1	55.7	57.4	63.7

Reagents	1'(Y=SO ₂)	2'(Y=PO ₂ H)	3'(Y=CF ₂)	4'(Y=CO)	5'(Y=BOH)	6'(Y=NCH ₃)	7'(Y=CH ₂)	8'(Y=C(CH ₃) ₂)
bond (Å) 5'-4'	2.658	2.401	2.297	2.284	2.212	2.156	2.125	2.108
bond (Å) 5'-6'	2.198	2.206	2.217	2.224	2.239	2.254	2.266	2.272
angle (°) 2'-1'-5'-4'	25.3	26.6	19.7	14.5	17.8	27.1	24.8	22.6
reaction energy	22.4	28.2	30.0	32.8	40.7	52.6	53.2	59.5

Table 1. The geometries and the trifluoromethylation reaction energies (see Fig. 4) of eight Togni derivatives.

we found that the newly designed derivative with the -SO₂- linker (1) has the lowest reaction free energy (only 25.1 kcal mol⁻¹), even lower than that of Togni's reagent II (4) by 14 kcal mol⁻¹. This suggests that our predicted structure 1 has the potential to be the most reactive reagent among 1 - 8.

Steric effects were further considered, and thus the ortho-methyl group was introduced into 1 - 8 to form 1' - 8'. The strong correlation between the electrophilic reactivity and the *trans* influence is maintained (blue line in Fig. 4). Furthermore, the steric effect of the ortho-methyl group in 1' - 8' makes the reaction free energies lower than those of structures 1 - 8 by 2.7 - 7.4 kcal mol⁻¹ (Table 1 and Fig. 4). This change is thought from the steric effect of




Fig. 4 Correlation between reaction free energies and *trans*

influence for eight Togni derivatives. The red line is for those structures without the steric effect, while the blue line is for those with the steric effect of the *ortho*-methyl group.




Fig. 5 Relative free energies (kcal/mol) of **TS1** (X = SO₂) and **TS2** (X = CO).³⁷

the *ortho*-methyl group, and it makes the electrophilic trifluoromethylation reaction easier. From our computational results, we suggest the Togni derivative **1'** to be the most reactive reagent among **1' - 8'**.

The dihedral angles (2'-1'-5'-4') for structures **1' - 8'** are reported in **Table 1**. These angles (from 15° to 27°) are generally somewhat larger than those of their analogues (**1 - 8**). This is expected, because the CF₃ group is driven out-of-the-plane by the added methyl group to reduce the steric hindrance, and this would cause more torsion of the five-membered ring.

Furthermore, the reactivity of Togni reagents can also be evaluated kinetically. In electrophilic trifluoromethylation, the Togni's reagents are generally activated by Lewis acids, such as Sc(OTf)₃,³⁴ Cu(OTf)₂,³⁵ Zn(NTf₂)₂,³⁶ etc. For example, in 2009, Togni's group reported the Zn(NTf₂)₂ mediated trifluoromethylation of aliphatic alcohols with Togni II (**4**), and their ESI-MS measurements detected a cationic species of [Zn(**4**)₂(NTf₂)]⁺.³⁶ Combining ¹⁹F NMR measurements, they suggested the reactive intermediate in this reaction to be [Zn(**4**)₂(NTf₂)₂]. Soon after, Li and co-authors conducted computational research of the mechanism for this Zn(NTf₂)₂ mediated trifluoromethylation reaction. With the Togni's reagent activated by the Lewis acid Zn²⁺, it readily undergoes an S_N2 type nucleophilic attack at the CF₃ group by pentanol with a reasonable energy barrier.¹³

In the present study, we predict such a mechanism of trifluoromethylation for our designed new Togni reagents (derivative **1**), and compared with that for the known Togni II (**4**). The energy barriers for the S_N2 zinc-mediated trifluoromethylation reactions are shown in **Fig. 5**. The free energy of the transition state relative to the intermediate [Zn(**1**)₂(NTf₂)₂] for derivative **1** is predicted to be 20.9 kcal mol⁻¹ (**TS1**, **Fig. 5**), which is about 8.1 kcal mol⁻¹ lower than that (29.0 kcal mol⁻¹) of **TS2** for the known Togni II (**4**). This kinetic result confirms that our newly suggested Togni reagent (derivative **1**) is more reactive for trifluoromethylation than the known Togni reagents.

In summary, based on our understanding of trans influence and steric effect in hypervalent iodine reagents, we have designed new Togni derivatives. Comparing the reaction free energy difference (14.2 kcal mol⁻¹) and the transition state energy barrier difference (8.1 kcal mol⁻¹) between **1** and **4**, we conclude that the Togni derivative **1** (**Fig. 4**) is the best choice, based on its highest reactivity

and its potential for synthesis.^{10,38-41} In addition to **1**, Togni derivatives **2**, **3**, **1' 2'** and **3'** (**Fig. 4**) are also rational choices for trifluoromethylation with high reactivity.

We are grateful for financial support from the Shenzhen STIC (JCYJ20170412150343516) and the Shenzhen San-Ming Project (SZSM201809085), and the U.S. National Science Foundation, Grant CHE-1661604. Tian-Yu Sun thanks the University of Georgia for hosting his 2015–2016 scientific visit.

Notes and references

- D. O'Hagan and D. B. Harper, *Nat. Prod. Rep.*, 1994, **11**, 123.
- T. Hiyama, *Organofluorine Compounds, Chemistry and Applications*. Springer-Verlag: Berlin Heidelberg, 2000.
- P. Kirsch, *Modern Fluoroorganic Chemistry*. Wiley-VCH: Weinheim, 2013.
- J. Wang, M. Sánchez-Roselló, J. L. Aceña, C. del Pozo, A. E. Sorochinsky, S. Fustero, V. A. Soloshonok and H. Liu, *Chem. Rev.*, 2013, **114**, 2432.
- X. Yang, T. Wu, R. J. Phipps and F. D. Toste, *Chem. Rev.*, 2014, **115**, 826.
- J.-P. Bégue and D. Bonnet-Delpon, *J. Fluorine Chem.*, 2006, **127**, 992.
- D. O'Hagan and D. B. Harper, *J. Fluorine Chem.*, 1999, **100**, 127.
- L. Chu and F. L. Qing, *Acc. Chem. Res.*, 2014, **47**, 1513.
- A. A. Studer, *Angew. Chem. Int. Ed.*, 2012, **51**, 8950.
- J. Charpentier, N. Früh and A. Togni, *Chem. Rev.*, 2015, **115**, 650.
- P. Chen and G. Liu, *Synthesis*, 2013, **45**, 2919.
- P. Eisenberger, S. Gischig and A. Togni, *Chem. Eur. J.*, 2006, **12**, 2579.
- L. Ling, K. Liu, X. Li and Y. Li, *ACS Catal.*, 2015, **5**, 2458.
- O. Sala, H. P. Lüthi, A. Togni, M. Iannuzzi and J. Hutter, *J. Comput. Chem.*, 2015, **36**, 785.
- T. G. Appleton, H. C. Clark and L. E. Manzer, *Coord. Chem. Rev.*, 1973, **10**, 335.
- (a) M. Ochiai, T. Sueda, K. Miyamoto, P. Kiprof and V. V. Zhdankin, *Angew. Chem. Int. Ed.*, 2006, **118**, 8383; (b) M. Ochiai, T. Sueda, K. Miyamoto, P. Kiprof and V. V. Zhdankin, *Angew. Chem., Int. Ed.*, 2006, **45**, 8203-8206.
- (a) P. K. Sajith and C. H. Suresh, *Inorg. Chem.*, 2012, **51**, 967; (b) P. K. Sajith and C. H. Suresh, *Inorg. Chem.*, 2013, **52**, 6046.
- K. Niedermann, J. M. Welch, R. Koller, J. Cvengroš, N. Santschi, P. Battaglia and A. Togni, *Tetrahedron*, 2010, **66**, 5753.
- N. Santschi, R. C. Sarott, E. Oth, R. Kissner and A. Togni, *Beilstein J. Org. Chem.*, 2014, **10**, 1.
- T.-Y. Sun, X. Wang, H. Geng, Y. Xie, Y.-D. Wu, X. Zhang and H. F. Schaefer, *Chem. Commun.*, 2016, **52**, 5371.
- H. Jiang, T.-Y. Sun, X. Wang, Y. Xie, X. Zhang, Y.-D. Wu and H. F. Schaefer, *Org. Lett.*, 2017, **19**, 6502.
- S. Koichi, B. Leuthold and H. P. Lüthi, *Phys. Chem. Chem. Phys.*, 2017, **19**, 32179.
- J. T. Su and W. A. Goddard, *J. Am. Chem. Soc.*, 2005, **127**, 14146.
- J. N. Moorthy, K. Senapati, K. N. Parida, S. Jhulki, K. Sooraj and N. N. Nair, *J. Org. Chem.*, 2011, **76**, 9593.
- Y. Zhao and D. G. Truhlar, *Theor. Chem. Acc.*, 2008, **120**, 215.
- T. H. Dunning, *J. Phys. Chem. A*, 2000, **104**, 9062.
- R. A. Kendall, T. H. Dunning and R. J. Harrison, *J. Chem. Phys.*, 1992, **96**, 6796.
- K. A. Peterson, R. A. Kendall and T. H. Dunning, *J. Chem. Phys.*, 1993, **99**, 1930.
- A. V. Marenich, C. J. Cramer and D. G. Truhlar, *J. Phys. Chem. B*, 2009, **113**, 637.

- 30 I. Kieltsch, P. Eisenberger and A. Togni, *Angew. Chem., Int. Ed.*, 2007, **46**, 754.
- 31 M. Li, X.-S. Xue, J. Guo, Y. Wang and J.-P. Cheng, *J. Org. Chem.*, 2016, **81**, 3119.
- 32 M. Li, H. Zheng, X.-S. Xue and J.-P. Cheng, *Tetrahedron Lett.*, 2018, **59**, 1278.
- 33 J. Zhang, J.-D. Yang, H. Zheng, X.-S. Xue, H. Mayer and J.-P. Cheng, *Angew. Chem. Int. Ed.*, 2018, **57**, 1.
- 34 A. E. Allen and D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2010, **132**, 4986.
- 35 Q.-H. Deng, H. Wadepohl and L. H. Gade, *J. Am. Chem. Soc.*, 2012, **134**, 10769.
- 36 R. Koller, K. Stanek, D. Stolz, R. Aardoom, K. Niedermann and A. Togni, *Angew. Chem. Int. Ed.*, 2009, **48**, 4332.
- 37 In **Table 1**, the single-point energies are evaluated with the cc-pVQZ(-PP) basis sets using the SMD model (in acetonitrile solvent) at the geometries optimized with the 6-31+G(d) basis sets except for iodine (cc-pVTZ-PP) and zinc (LANL2DZ).
- 38 M. Uyanik, M. Akakura and K. Ishihara, *J. Am. Chem. Soc.*, 2009, **131**, 251.
- 39 G. F. Koser, G. Sun, C. W. Porter and W. J. Youngs, *J. Org. Chem.*, 1993, **58**, 7310.
- 40 M. Justik, *Tetrahedron Lett.*, 2007, **48**, 3003.
- 41 Y. Li, D. P. Hari, M. V. Vita and J. Waser, *Angew. Chem. Int. Ed.*, 2016, **55**, 4436.




Table of contents (TOC)

New trifluoromethylating reagents are designed based on trans influence and steric effect.