



ChemComm

**Low-spin 1,1'-diphoshametalloenates of Chromium and  
Iron**

Journal:	<i>ChemComm</i>
Manuscript ID	CC-COM-09-2020-006518.R1
Article Type:	Communication

SCHOLARONE™  
Manuscripts

## COMMUNICATION

## Low-spin 1,1'-diphospha-metalloenates of Chromium and Iron

Samuel M. Greer,<sup>a,b,c</sup> Ökten Üngör,<sup>b</sup> Ross J. Beattie,<sup>a</sup> Jaqueline L. Kiplinger,<sup>\*a</sup> Brian L. Scott,<sup>a</sup> Benjamin W. Stein,<sup>\*a</sup> and Conrad A. P. Goodwin<sup>\*a,d</sup>

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

We report two anionic diphospha-metalloenates,  $[K(2.2.2\text{-crypt})][M(PC_4Me_4)_2]$  ( $M = Cr, 2\text{-Cr}; Fe, 2\text{-Fe}$ ). Both are low-spin ( $S = 1/2$ ) by EPR spectroscopy and SQUID magnetometry. This contrasts the high-spin ( $S = 3/2$ ) ferrocenate,  $[K(2.2.2\text{-crypt})][Fe(C_5H_2-1,2,4\text{-}^tBu)_2]$  ( $4\text{-Fe}$ ). Quantum chemical calculations suggest this is due to significant differences in ligand field splitting of the d-orbitals which also explain structural features in the 2-M complexes.

The chemistry and electronic structure of transition metal (TM) metallocenes<sup>1,2</sup> has fascinated chemists since the discovery of ferrocene,  $[Fe(Cp)_2]$  ( $FcH$ ,  $Cp = \text{cyclopentadienyl}, \{C_5H_5\}$ ).<sup>3</sup> Oxidation of  $FcH$  leads to the ferrocenium cation,  $[Fe(Cp)_2]^+$  ( $FcH^+$ ),<sup>4</sup> which finds use in organic synthesis,<sup>5</sup> while the  $FcH^{+/0}$  couple serves as an electrochemical standard.<sup>6</sup> "Heterometallics" are related complexes that feature heterocycles in lieu of Cp ligands (e.g.  $\eta^5\text{-pyrrolides}, \{NC_4R_4\}^-$ ;  $\eta^5\text{-stannolides}, \{SbC_4R_4\}^-$ ; poly-phospholides  $\{P_nC_{5-n}R_{5-n}\}^-$  etc).<sup>7</sup> The exchange of C atoms for heteroatoms can lead to drastic changes in reactivity.<sup>1,2</sup> For example, ferrocene undergoes deprotonation by alkyl-lithium reagents,<sup>8,9</sup> but 1,1'-diphosphaferrocenes undergo nucleophilic attack at the P atom.<sup>10,11</sup>

The oxidation of TM metallocenes to metallocenium cations is a defining feature of their chemistry, with examples across the first row transition metals (V–Ni).<sup>1,12</sup> The corresponding reductive chemistry is underdeveloped outside of electrochemical studies.<sup>13</sup> Exceptions include the synthesis of  $[Mn(Cp^*)_2]^-$  salts,<sup>14</sup> the reduction of  $[Co\{\eta^5\text{-}C_9H_5\text{-}1,3\text{-}(SiMe_3)_2\}_2]$  to a formally anionic "cobaltate",<sup>15</sup> and an anionic iron bis-stannolide.<sup>7c</sup> Some of us have recently described the chemical reduction of  $[M(Cp^{ttt})_2]$  (**3-M**;  $M = Mn, Fe, Co$ ;  $Cp^{ttt} = \{C_5H_2\text{-}1,2,4\text{-}^tBu\}$ ) to  $[K(2.2.2\text{-crypt})][M(Cp^{ttt})_2]$  (**4-M**;  $M = Mn, Fe, Co$ ) giving an 18, 19 and 20 valence electron (VE) series,<sup>16</sup> similar to the 17, 18, 19 VE series  $[Fe(Cp^R)(\text{arene})]^{2+/+0}$  complexes.<sup>17</sup>

The synthesis of 3d metallocene anions could open up the possibility of using them as nucleophiles like "A[Re(Cp)<sub>2</sub>]" ( $A = Li,^{18} K^{19}$ ). While the  $[Mn(Cp^*)_2]^-$  anion is temperature stable,<sup>14c,14a</sup> analogous  $[FcH]^-$  and  $[Fe(Cp^*)_2]^-$  anions are not easily accessed chemically,<sup>13b,16</sup> and **4-M complexes** are extremely temperature sensitive. We sought ligands that would impart greater stability to the anionic species. Reports of  $[M(PC_4R_nH_{4-n})_2]$  for a range of metals include the Fe/Ru/Os triad,<sup>7d,20</sup> and exhibit facile redox chemistry.<sup>21</sup> For example,  $[M(TMP)_2]$  ( $M = Cr, 1\text{-Cr};^{22} Fe, 1\text{-Fe};^{23} TMP = \{PC_4Me_4\}$ ) have  $M^{+0}$  couples (vs  $FcH^{+/0}$ :  $M = Cr, -0.77\text{ V};^{24} Fe, 0.06\text{ V}$ ), and  $M^{0/-}$  (vs  $FcH^{+/0}$ :  $M = Cr, -2.4\text{ V}; Fe, -3.02\text{ V}$ ) at less cathodic potentials than **3-M**.<sup>16</sup> Here we report the synthesis of 1,1'-diphospha-metalloenates,  $[K(2.2.2\text{-crypt})][M(TMP)_2]$  ( $M = Cr, 2\text{-Cr}; Fe, 2\text{-Fe}$ ), by low-temperature reduction of **1-M** by  $KC_8$  in the presence of 2.2.2-cryptand.


<sup>a</sup> Chemistry Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA. Email: kiplinger@lanl.gov, bstein@lanl.gov, cgoodwin@lanl.gov.

<sup>b</sup> National High Magnetic Field Laboratory, Florida State University, Tallahassee, FL 32310, USA.


<sup>c</sup> Department of Chemistry & Biochemistry, Florida State University, Tallahassee, FL 32306, USA.

<sup>d</sup> Department of Chemistry, School of Natural Sciences, The University of Manchester, Oxford Road, Manchester, M13 9PL, UK.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x



Scheme 1 Synthesis of **2-M** ( $M = Cr, Fe$ ) from **1-M**,  $KC_8$  and 2.2.2-cryptand.



**Fig. 1** Molecular structure of **2-Fe** with selective atom labelling (C = grey, P = purple, K = violet, Fe = orange, N = blue, O = red). Displacement ellipsoids set at 50 % probability level and H-atoms omitted for clarity.


Precursor **1-M** (M = Cr, Fe) were synthesized from  $MCl_2$  and base-free KTMP by modification of literature procedures<sup>21b,23b</sup>. During the course of this work we have also structurally characterized  $[Zr(Cp)_2(C_4Me_6)]$  (**5**) and Ph-TMP (**6**) for the first time, which are precursors in the synthesis of KTMP (See Supporting Information). Complexes **1-M** were then reduced under conditions established for the synthesis of low-valent metallocenes.<sup>16,25</sup> Addition of solid  $KC_8$  to cooled ( $-35^\circ C$ ) solutions of **1-M** and 2.2.2-crypt in THF caused rapid color changes (from red to red-brown, **2-Cr**; or red to green-black, **2-Fe**). Intensely colored crystals of **2-Cr** and **2-Fe** were obtained upon workup (Figure 1, and Supporting Information).

These specific Cr and Fe complexes (**1-Cr** and **2-Cr**) were chosen as they have established redox chemistry and thus provided an ideal testbed for the comparison of spin-states in reduced metallocenes.<sup>22-23</sup> We were unable to synthesize **1-Mn** and **1-Co** in order to furnish a redox series from  $Cr^{2+}$  ( $d^4$ ) to  $Co^{1+}$  ( $d^8$ ). Reactions between  $Mn^{2+}$  halides and KTMP gave intractable mixtures. For  $Co^{2+}$  we could isolate traces of impure  $[Co(\eta^5-TMP)(\mu:\eta^1:\eta^1-TMP)]_2$  (**7**) (Supporting Information).

Salient structural parameters for **1-M** are shown along with those of **2-M** in Table 1. Complexes **2-M** (M = Cr, Fe) are isomorphous, their ATR-FTIR spectra are superimposable, and elemental analysis results were in excellent agreement with prediction. The structures contain discrete  $[K(2.2.2-crypt)]$  cations, and  $[M(TMP)_2]$  anions. In **2-M** the TMP-P atoms are each situated opposite a  $\beta$ -carbon of the second ring, similar to neutral  $[Fe(PC_4H_2-2,5-Me_2)_2]$ .<sup>26</sup> Upon reduction there is a small decrease in M–P distances for Cr ( $\Delta = -0.022(1)$  Å) but an increase for Fe ( $\Delta = +0.103(1)$  Å); and changes in the  $TMP_{cent}\cdots M$  distances follow the same trend (Cr,  $\Delta = -0.057$  Å; Fe,  $\Delta = +0.053$  Å). The  $TMP_{cent}\cdots Cr$  change between  $d^4$  **1-Cr** (1.795(1) Å) and  $d^5$  **2-Cr** (avg. 1.738(1) Å) is similar to the  $Cp^*_{cent}\cdots Mn$  change between  $d^4$  and  $d^5$  manganocenes.<sup>27</sup> The  $TMP_{cent}\cdots Fe$  change between  $d^6$  **1-Fe** (1.660(1) Å) and  $d^7$  **2-Fe** (avg. 1.713(1) Å) is

**Table 1** Selected structural parameters for **1-M** and **2-M**.

	<b>1-Cr</b>	<b>2-Cr</b>	<b>1-Fe</b>	<b>2-Fe</b>
P–M / Å	2.3812(4)	2.3596(8) 2.3595(7)	2.2932(4)	2.3842(8) 2.4078(7)
$TMP_{cent}\cdots M$ / Å	1.795(1)	1.739(1) 1.737(1)	1.660(1)	1.714(1) 1.711(1)
$PC_{2plane}\cdots C_{4plane}$ / °	4.45(12)	3.57(2) 4.64(2)	1.27(13)	6.58(2) 8.73(2)
TMP–P <sub>2</sub> twist angle / °	180	139.82(5)	180	146.35(5)



**Fig. 2 a)** Temperature dependence of  $\chi T$  for **2-Fe**; **b)** Experimental (black) and simulated (red) EPR spectra of **2-Fe** recorded at 118 GHz and 5 K. The spectrum is simulated with  $g_1 = 2.033$ ,  $g_2 = 1.999$ , and  $g_3 = 1.943$ . Baseline features in the **2-Fe** spectrum result from polycrystallinity. A minor signal arising from an unidentified impurity at lower field is omitted (see Figure S28); **c)** Experimental (black) and simulated (red)  $^{57}Fe$  Mössbauer spectra of **1-Fe** and **2-Fe**. Simulations were generated for **1-Fe** using  $\delta = 0.48(3)$ ,  $\Delta E_Q = 2.02(2)$  mm  $sec^{-1}$ , and for **2-Fe** with  $\delta = 0.65(4)$ ,  $\Delta E_Q = 1.28(4)$  mm  $sec^{-1}$ .

larger than the increase from **3-Fe** (avg. 1.715(2) Å) to **4-Fe** (avg. 1.750(3) Å), likely as the distances in **3-Fe** are due to the bulky ligands rather than electronic effects.<sup>16</sup>

The  $^1H$ ,  $^{13}C\{^1H\}$ , and  $^{31}P\{^1H\}$  NMR spectra of **2-M** were largely uninformative (Supporting Information) due to paramagnetic broadening and diamagnetic impurities which dominate the spectra. However, in the  $^1H$  NMR spectrum of **2-Fe** we observed three broad peaks ( $\nu_{1/2} < 17$  Hz) in the range 3.37 – 3.30 ppm corresponding to 2.2.2-crypt; a doublet at 2.19 ppm ( $^3J_{HP} = 10.29$  Hz) and a singlet at 1.94 ppm correspond to TMP ligand resonances. Complex **2-Cr** could be heated in THF solution (up to  $50^\circ C$ ) and recrystallized with >80% recovery of material, whereas THF solutions of **2-Fe** show decomposition in the  $^1H$  NMR spectrum over the course of an hour at room temperature. Data for **1-M** have been reported previously.

We have employed SQUID magnetometry to measure the susceptibility of the open-shell complexes. **1-Cr** and **2-Cr** exhibit  $\chi T$  (298 K) values of  $\sim 1.24$  and  $\sim 0.38$  emu·K  $mol^{-1}$  (Figure S34) which suggest  $S = 1$  as previously reported,<sup>21b</sup> and  $S = 1/2$  respectively. The observed  $\chi T$  (298 K) value of **2-Fe** ( $\sim 0.26$  emu·K  $mol^{-1}$ ) is less than the spin-only value for an  $S = 1/2$  system (0.375 emu·K  $mol^{-1}$ ) (Figures 2, and S35), which we attribute to some thermal decomposition of paramagnetic **2-Fe** to diamagnetic **1-Fe**, as well as weighing errors and diamagnetic corrections affecting this weakly paramagnetic system. As magnetic measurements were inconclusive for **2-Fe**, we turned to high-field electron paramagnetic resonance (EPR) spectroscopy to determine the ground state spin of **2-Cr** and **2-Fe**. These measurements reveal spectra that are typical of complexes where  $S = 1/2$ . The spectrum of **2-Cr** is nearly axial with  $g_1 = 2.023(5)$ ,  $g_2 = 1.993(5)$ , and  $g_3 = 1.985(5)$  (Figure S32); and **2-Fe** exhibits a rhombic signal (Figure 2b) with  $g_1 = 2.033(5)$ ,  $g_2 = 1.999(5)$ , and  $g_3 = 1.943(5)$ , which is similar to other low-spin  $Fe^{1+}$  sandwich complexes.<sup>17</sup>

**Table 2** Mössbauer parameters for **1-Fe** to **4-Fe**.

		<b>1-Fe</b>	<b>2-Fe</b>	<b>3-Fe<sup>b</sup></b>	<b>4-Fe<sup>b</sup></b>
$\delta$ (mm sec <sup>-1</sup> )	Exp.:	0.48(3)	0.65(4)	0.66(2)	1.25(2)
	Theory:	0.50	0.66	0.64	1.09
$\Delta E_Q$ (mm sec <sup>-1</sup> )	Exp.:	2.02(2)	1.28(4)	2.60(2)	1.23(2)
	Theory:	1.91	1.46	2.52	0.65

<sup>a</sup> Calculated with BP86 and Fe (CP(PPP)), C (def2-tzvp), H (def2-svp) basis set combination. This gave the best agreement with experiment and multiple basis sets were previously examined for complexes **3-Fe** and **4-Fe**. <sup>b</sup> From reference 16.


<sup>57</sup>Fe Mössbauer spectroscopy was used to further investigate the differences between **1-Fe** and **2-Fe** which display a single quadrupole doublet in each spectrum (at 120 K, Figure 2c). The parameters for **1-Fe**, **2-Fe**, **3-Fe** and **4-Fe** are summarized in Table 2. The more positive isomer shift ( $\delta$ ) of **2-Fe** suggests that there is decreased s-electron density at the Fe nucleus compared to that in **1-Fe**. The parameters of **1-Fe** are similar to those reported for derivatized ferrocenes.<sup>28</sup>  $\delta$  for **2-Fe** is slightly more positive than many characterized Fe(I) complexes ( $\sim 0.2 - \sim 0.6$  mm sec<sup>-1</sup>),<sup>29</sup> but in good agreement with the low-spin 19 VE [Fe(Cp<sup>R</sup>)(arene)] complexes,<sup>17,30</sup> in contrast to high-spin **4-Fe** (1.25 mm sec<sup>-1</sup>).<sup>16</sup> We also found excellent agreement between the DFT calculated  $\delta$  and  $\Delta E_Q$  parameters and the experimental data (Table 2) for both **1-Fe** and **2-Fe**. Data for **3-Fe** and **4-Fe** is shown for completeness.

To rationalize the structural features of **2-M** with their ground-state electronic structures, we have performed CASSCF/NEVPT2 calculations (see Supporting Information for details). The *ab-initio* ligand field theory (AILFT) derived molecular orbital (MO) diagrams for **1-M**, **2-M**, **3-Fe**, and **4-Fe**, are shown in Figure 3.<sup>16</sup> The most striking observation is the significant difference in ligand field strength between **2-Fe** and **4-Fe**, that results in a drastic difference between the energetic separation of the  $d_{z^2}$  and  $d_{xz}/d_{yz}$  orbitals which is responsible for the change in spin-state of **2-Fe** compared to **4-Fe**. This can be rationalized by comparing the  $L_{\text{cent}} \cdots \text{Fe}$  distances ( $L = \text{TMP}$ , avg 1.713(1) Å;  $L = \text{Cp}^{\text{ttt}}$ , avg 1.750(3) Å); the closer TMP ligand provides a stronger ligand field than  $\text{Cp}^{\text{ttt}}$ , similar to the situation in high-spin/low-spin manganocenes.<sup>31</sup>

By considering the MO diagram of a 1,1'-diphosphametalocene,<sup>32</sup> which is similar to that of FcH, we can rationalize some of the bond changes.<sup>33</sup> In the MO scheme of  $D_{5d}$  symmetric FcH the LUMO is comprised of a doubly degenerate set of antibonding orbitals with ligand  $e_1''$  (for a  $D_{5h}$  Cp ring) contributions that mix with the  $d_{xz}/d_{yz}$  ( $e_{1g}$ ) orbitals. The LUMO of 1,1'-diphosphametalocenes is also comprised of metal  $d_{xz}/d_{yz}$  orbitals, along with a single ligand HOMO ( $\pi_p$  here, Figure S36) that approximates the Cp  $e_1''$  MOs by having a single nodal plane perpendicular to the ring plane, but has significant P-character. The SOMO of **2-Fe** is principally  $d_{xz}$  in character, and is anti-bonding with respect to  $\pi_p$  which explains the increased Fe–P and  $\text{TMP}_{\text{cent}} \cdots \text{Fe}$  distances. Conversely, as the SOMO for **2-Cr** is comprised of the non-bonding  $d_{z^2}$  orbital as it is in **1-Cr**, the structural changes on reduction are minimal. The slight shortening of  $\text{TMP}_{\text{cent}} \cdots \text{Cr}$  from **1-Cr** to **2-Cr** is due to the spin change from  $S = 1$  to  $S = 1/2$  which lowers the energy of the  $d_{x^2-y^2}$ ,  $d_{xy}$  orbitals, increasing covalency as in  $[\text{Cr}(\text{Cp})_2]$ .<sup>34</sup>

We suggest that the removal of degeneracy in the ligand HOMO orbitals for TMP vs the equivalent orbitals in Cp lifts the near-degeneracy between the  $d_{xz}/d_{yz}$  orbitals (as in **2-Fe** compared to **4-Fe**); and furthermore, the large change in ligand steric profile has drastically changed the separation between  $d_{z^2}$  and the  $d_{xz}/d_{yz}$  orbitals. These combined effects have stabilized the low-spin state in **2-Fe** and should prove instructive towards chemical control of the spin state in reduced metallocenes and heterometallocenes for the 3d series. For example, by significantly changing the gap between the  $d_{z^2}$  and  $d_{xz}/d_{yz}$  orbitals in **4-Fe**, a low-spin ferrocenate might be realized.

To summarize, we have demonstrated that octamethyl-1,1'-diphosphametalocenes for Cr and Fe can be reduced to afford crystalline anionic complexes. Comparison of **2-Fe** to authentic formal Fe(I) complexes suggests that it also contains a formal



**Fig. 3** AILFT orbital energies for the frontier orbitals with principally d-character for **1-M**, **2-M**, **3-Fe** and **4-Fe**.<sup>16</sup> The indicated electronic configuration corresponds to the dominant configuration of the CASSCF+NEVPT2 calculated ground state, and all six complexes feature the same orbital ordering (from low- to high-energy:  $d_{x^2-y^2}$ ,  $d_{xy}$ ,  $d_{z^2}$ ,  $d_{xz}$ ,  $d_{yz}$ ). The molecular z-axis was chosen to align approximately with the vector formed by the metal and ligand ring-centroids. The SOMO orbital(s) of the reduced anionic complexes has been indicated for **2-M** and **4-Fe**.

Fe(I). We targeted complexes that might be temperature stable; while **2-Fe** is somewhat thermally unstable, **2-Cr** appears stable indefinitely in the solid-state at room temperature. Both were more amenable to characterization than **4-Fe** which decomposes in the solid-state even at  $-35^{\circ}\text{C}$ .<sup>16</sup> We have found that both anionic species studied here exhibit low spin ( $S = 1/2$ ) ground states, in contrast to high-spin ( $S = 3/2$ ) **4-Fe**, and that these results are well-explained by changes in the ligand frontier orbitals. This suggests that synthetic control of the ground spin state of 3d metallocenes is possible and that with judicious planning, it should be possible to target desired high- or low-spin examples of this family.

SMG acknowledges support from a Director's Postdoctoral Fellowship (20180759PRD4), and CAPG was sponsored by a J. Robert. Oppenheimer Distinguished Postdoctoral Fellowship (20180703PRD1), both through the Los Alamos National Laboratory – Laboratory Directed Research and Development program (LANL-LDRD). CAPG also thanks the UK Engineering and Physical Sciences Research Council (EPSRC) for a Doctoral Prize Fellowship (UoM). An EPSRC grant (EP/K039547/1) provided a single-crystal X-ray diffractometer (UoM). JLK also thanks the LANL-LDRD program. We also acknowledge the University of California and LANL Postdoctoral Entrepreneurial Fellowship and the Feynman Center for Innovation for support of RJB. BWS acknowledges funding provided by the Director, Office of Science, Office of Basic Energy Sciences, Division of Chemical Sciences, Geosciences, and Biosciences, Heavy Element Chemistry Program of the U.S. Department of Energy (DOE). We thank Prof. Michael Shatruk for supervising collection of magnetometry data; and Dr David P. Mills, Dr Fabrizio Ortu, Dr Andrew J. Gaunt and Dr Michael T. Janicke for helpful discussions. We also thank Prof. James M. Boncella for the provision of synthetic work space. A portion of this work was supported by the NSF award CHE-1955754, and performed at the National High Magnetic Field Laboratory under National Science Foundation Cooperative Agreement No. DMR-1644779 with the State of Florida. LANL, an affirmative action/equal opportunity employer, is managed by Triad National Security, LLC, for the NNSA of the U.S. Department of Energy (89233218CNA000001). Raw research data files supporting this publication are available from Mendeley Data at DOI: 10.17632/tw78xbvc7f.1. The Supplementary Information contains ORCA input files used in this work.

### Author Contributions

CAPG lead the project and performed all synthesis, UV-vis-NIR and NMR characterization; and also X-ray diffraction measurements with supervision by BLS. SMG performed EPR and Mössbauer spectroscopies and interpretation and all theoretical work, with supervision by BWS. OU performed SQUID magnetometry measurements and interpretation. RJB sealed samples for analysis and was supervised by JLK. The manuscript was prepared by CAPG with input from all authors.

### Conflicts of interest

There are no conflicts to declare.


### Notes and references


- N. J. Long, *Metallocenes: An Introduction to Sandwich Complexes*, Wiley, 1998.
- A. Togni, R. L. Halterman, *Metallocenes: Synthesis Reactivity Applications*, 1998.
- a) T. J. Kealy, P. L. Pauson, *Nature*, 1951, **168**, 1039; b) S. A. Miller, J. A. Tebboth, J. F. Tremaine, *J. Chem. Soc.*, 1952, 632; c) J. D. Dunitz, L. E. Orgel, A. Rich, *Acta Crystallogr.*, 1956, **9**, 373.
- G. Wilkinson, M. Rosenblum, M. C. Whiting, R. B. Woodward, *J. Am. Chem. Soc.*, 1952, **74**, 2125.
- Š. Toma, R. Šebesta, *Synthesis*, 2015, **47**, 1683.
- R. R. Gagne, C. A. Koval, G. C. Lisensky, *Inorg. Chem.*, 1980, **19**, 2854.
- a) N. Kuhn, E. M. Horn, R. Boese, N. Augart, *Angew. Chem., Int. Ed.*, 1988, **27**, 1368; b) O. J. Scherer, T. Brück, *Angew. Chem., Int. Ed.*, 1987, **26**, 59; c) M. Saito, N. Matsunaga, J. Hamada, S. Furukawa, T. Tada, R. H. Herber, *Chem. Lett.*, 2019, **48**, 163; d) F. Mathey, *Coord. Chem. Rev.*, 1994, **137**, 1.
- M. Walczak, K. Walczak, R. Mink, M. D. Rausch, G. Stucky, *J. Am. Chem. Soc.*, 1978, **100**, 6382.
- W. Erb, F. Mongin, *Synthesis*, 2019, **51**, 146.
- B. Deschamps, J. Fischer, F. Mathey, A. Mitschler, *Inorg. Chem.*, 1981, **20**, 3252.
- B. Deschamps, J. Fischer, F. Mathey, A. Mitschler, L. Ricard, *Organometallics*, 1982, **1**, 312.
- M. Malischewski, M. Adelhardt, J. Sutter, K. Meyer, K. Seppelt, *Science*, 2016, **353**, 678.
- a) A. J. Bard, E. Garcia, S. Kukharensko, V. V. Strelets, *Inorg. Chem.*, 1993, **32**, 3528; b) Y. Mugnier, C. Moise, J. Tirouflet, E. Laviron, *J. Organomet. Chem.*, 1980, **186**, C49; c) W. E. Geiger, *J. Am. Chem. Soc.*, 1974, **96**, 2632; d) J. D. L. Holloway, W. L. Bowden, W. E. Geiger, *J. Am. Chem. Soc.*, 1977, **99**, 7089; e) N. Ito, T. Saji, S. Aoyagui, *J. Organomet. Chem.*, 1983, **247**, 301.
- a) M. Malischewski, K. Seppelt, *Dalton Trans.*, 2019, **48**, 17078; b) J. L. Robbins, N. M. Edelstein, S. R. Cooper, J. C. Smart, *J. Am. Chem. Soc.*, 1979, **101**, 3853; c) J. C. Smart, J. L. Robbins, *J. Am. Chem. Soc.*, 1978, **100**, 3936.
- F. Hung-Low, C. A. Bradley, *Inorg. Chem.*, 2013, **52**, 2446.
- C. A. P. Goodwin, M. Giansiracusa, S. M. Greer, H. M. Nicholas, P. Evans, M. Vonci, S. Hill, N. F. Chilton, D. P. Mills, *Nat. Chem.*, 2020.
- a) D. Astruc, *Tetrahedron*, 1983, **39**, 4027; b) M. Rajasekharan, S. Giezynski, J. Ammeter, N. Oswald, J. Hamon, D. Astruc, P. Michaud, *J. Am. Chem. Soc.*, 1982, **104**, 2400.
- D. Baudry, M. Ephritikhine, *J. Organomet. Chem.*, 1980, **195**, 213.
- B. M. Gardner, J. McMaster, W. Lewis, S. T. Liddle, *Chem. Commun.*, 2009, 2851.
- a) R. Loschen, C. Loschen, W. Frank, C. Ganter, *Eur. J. Inorg. Chem.*, 2007, **2007**, 553; b) F. Mathey, *J. Organomet. Chem.*, 2002, **646**, 15; c) M. Ogasawara, T. Nagano, K. Yoshida, T. Hayashi, *Organometallics*, 2002, **21**, 3062; d) M. Ogasawara, M. Shintani, S. Watanabe, T. Sakamoto, K. Nakajima, T. Takahashi, *Organometallics*, 2011, **30**, 1487.
- a) Y. Cabon, D. Carmichael, L. Ricard, *Chem. Commun.*, 2011, **47**, 11486; b) P. Lemoine, M. Gross, P. Braunstein, F. Mathey, B. Deschamps, J. H. Nelson, *Organometallics*, 1984, **3**, 1303; c) N. G. Connelly, W. E. Geiger, *Chem. Rev.*, 1996, **96**, 877.

22. R. Feher, F. H. Köhler, F. Nief, L. Ricard, S. Rossmayer, *Organometallics*, 1997, **16**, 4606.
23. a) A. J. Ashe III, S. Al-Ahmad, S. Pilotek, D. B. Puranik, C. Elschenbroich, A. Behrendt, *Organometallics*, 1995, **14**, 2689; b) F. Nief, F. Mathey, L. Ricard, F. Robert, *Organometallics*, 1988, **7**, 921.
24. **1-Cr** was reported vs the  $[\text{Co}(\text{Cp})_2]^{+0}$  couple, and corrected to values vs  $\text{FcH}^{+0}$ .
25. W. J. Evans, *Organometallics*, 2016, **35**, 3088.
26. G. De Lauzon, B. Deschamps, J. Fischer, F. Mathey, A. Mitschler, *J. Am. Chem. Soc.*, 1980, **102**, 994.
27. a) N. Augart, R. Boese, G. Schmid, *Z. Anorg. Allg. Chem.*, 1991, **595**, 27; b) W. E. Broderick, J. A. Thompson, E. P. Day, B. M. Hoffman, *Science*, 1990, **249**, 401; c) D. P. Freyberg, J. L. Robbins, K. N. Raymond, J. C. Smart, *J. Am. Chem. Soc.*, 1979, **101**, 892.
28. a) R. M. G. Roberts, J. Silver, B. Yamin, *J. Organomet. Chem.*, 1984, **270**, 221; b) R. A. Stukan, S. P. Gubin, A. N. Nesmeyanov, V. I. Gol'danskii, E. F. Makarov, *Theor. Exp. Chem.*, 1966, **2**, 581; c) G. K. Wertheim, R. H. Herber, *J. Chem. Phys.*, 1963, **38**, 2106.
29. J. P. Mariot, P. Michaud, S. Lauer, D. Astruc, A. X. Trautwein, F. Varret, *J. Physique*, 1983, **44**, 1377.
30. P. Gütlich, E. Bill, A. X. Trautwein, *Mössbauer Spectroscopy and Transition Metal Chemistry*, 2011.
31. M. D. Walter, C. D. Sofield, C. H. Booth, R. A. Andersen, *Organometallics*, 2009, **28**, 2005.
32. N. M. Kostic, R. F. Fenske, *Organometallics*, 1983, **2**, 1008.
33. A. Haaland, *Acc. Chem. Res.*, 1979, **12**, 415.
34. M. Swart, *Inorg. Chim. Acta*, 2007, **360**, 179.



**SUMMARY FOR TABLE OF CONTENTS**

We report two low-spin diphospha-metalloenes  $[K(2.2.2\text{-crypt})][M(\text{PC}_4\text{Me}_4)_2]$  ( $M = \text{Cr}, \text{Fe}$ ) and propose structural features that allow spin-control in 3d metalloenes.





$\xrightarrow{\text{KC}_8}$   
**222-crypt**



$S =$	$\text{Cr}^{2+}\text{L}_2$	$\text{Fe}^{2+}\text{L}_2$	$[\text{CrL}_2]^-$	$[\text{FeL}_2]^-$
	1	0	$1/2$	$1/2$

**Low-spin diphospha-metalloccenates**