



Natural Product  
Reports

**Terpene synthases and pathways in animals: enzymology  
and structural evolution in the biosynthesis of volatile  
infochemicals**

Journal:	<i>Natural Product Reports</i>
Manuscript ID	NP-REV-11-2022-000076.R1
Article Type:	Review Article
Date Submitted by the Author:	10-Feb-2023
Complete List of Authors:	Tholl, Dorothea; Virginia Polytechnic Institute and State University Rebholz, Zarley; Virginia Polytechnic Institute and State University Morozov, Alexandre; Rutgers University O'Maille, Paul; SRI International

SCHOLARONE™  
Manuscripts

1 Terpene synthases and pathways in animals: enzymology and structural evolution in the  
2 biosynthesis of volatile infochemicals

3

4 Dorothea Tholl<sup>\*a</sup>, Zarley Rebholz<sup>a</sup>, Alexandre V. Morozov<sup>b</sup>, and Paul E. O'Maille<sup>c</sup>

5

6 <sup>a</sup>Department of Biological Sciences, Virginia Tech, Blacksburg, VA 24060, USA

7 <sup>b</sup>Department of Physics and Astronomy and Center for Quantitative Biology, Rutgers University,  
8 Piscataway, NJ 08854, USA

9 <sup>c</sup>SRI International, San Diego, CA 92131, USA

10

11

12

13 \*Corresponding author: Dorothea Tholl, Department of Biological Sciences, Virginia Tech,  
14 [tholl@vt.edu](mailto:tholl@vt.edu), +1-540-231-4567

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

**Abstract**

35  
36  
37 Many animals release volatile or semi-volatile terpenes as semiochemicals in intra- and inter-  
38 specific interactions. Terpenes are important constituents of pheromones or serve as chemical  
39 defenses to ward off predators. Despite the occurrence of terpene specialized metabolites from  
40 soft corals to mammals, the biosynthetic origin of these compounds has largely remained  
41 obscure. An increasing number of animal genome and transcriptome resources is facilitating the  
42 identification of enzymes and pathways that allow animals to produce terpenes independent of  
43 their food sources or microbial endosymbionts. Substantial evidence has emerged for the  
44 presence of terpene biosynthetic pathways such as in the formation of the iridoid sex pheromone  
45 nepetalactone in aphids. In addition, terpene synthase (TPS) enzymes have been discovered that  
46 are evolutionary unrelated to canonical plant and microbial TPSs and instead resemble precursor  
47 enzymes called isoprenyl diphosphate synthases (IDSs) in central terpene metabolism. Structural  
48 modifications of substrate binding motifs in canonical IDS proteins presumably facilitated the  
49 transition to TPS function at an early state in insect evolution. Other arthropods such as mites  
50 appear to have adopted their TPS genes from microbial sources via horizontal gene transfer. A  
51 similar scenario likely occurred in soft corals, where TPS families with closer resemblance to  
52 microbial TPSs have been discovered recently. Together, these findings will spur the identification  
53 of similar or still unknown enzymes in terpene biosynthesis in other lineages of animals. They will  
54 also help develop biotechnological applications for animal derived terpenes of pharmaceutical  
55 value or advance sustainable agricultural practices in pest management.

56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68

69	<b>1 Introduction</b>
70	<b>2 Volatile terpenes in the animal kingdom</b>
71	<b>2.1 Invertebrates</b>
72	<b>2.1.1 Mollusks and Corals</b>
73	<b>2.1.2 Arthropods – Millipedes and arachnids</b>
74	<b>2.1.3 Arthropods – Insects</b>
75	<b>2.2 Vertebrates</b>
76	<b>2.2.1 Amphibians</b>
77	<b>2.2.2 Reptiles</b>
78	<b>2.2.3 Mammals</b>
79	<b>3 Terpene biosynthetic pathways and enzymes in insect pheromone/defense</b>
80	<b>biosynthesis</b>
81	<b>3.1 Biosynthesis of iridoids</b>
82	<b>3.2 Horizontal gene transfer of terpene synthases in mites</b>
83	<b>3.3 IDS-like terpene synthases in insects</b>
84	<b>3.3.1 FPP synthase type terpene synthases</b>
85	<b>3.3.1.1 <i>Phyllotreta striolata</i></b>
86	<b>3.3.1.2 Pentatomids</b>
87	<b>3.3.1.3 Genomic organization and evolution of stink bug and hemipteran IDS-like genes</b>
88	<b>3.3.2 GGPP synthase type terpene synthases</b>
89	<b>4 Structural and mechanistic evolution of insect IDS-like TPSs</b>
90	<b>5 Terpene synthases in corals</b>
91	<b>6 Conclusions and Outlook</b>
92	<b>7 Acknowledgements</b>
93	<b>8 Author Contributions</b>
94	<b>9 Conflict of Interest</b>
95	<b>10 References</b>
96	
97	
98	
99	
100	
101	
102	

103

104 **1 Introduction**

105

106 All organisms use small molecules for communication within their own species or in interactions  
107 with other species.<sup>1</sup> Depending on their physicochemical properties, these specialized  
108 metabolites facilitate efficient interactions at short and long distance both in water and on land.  
109 Natural products of the large class of terpenes or terpenoids (> 80,000 structures known;  
110 <https://dnp.chemnetbase.com>; <sup>2</sup>) represent the most structurally diverse group of molecules used  
111 in chemical interactions. Due to their high vapor pressure and volatility, low molecular weight  
112 terpenes are ideal “infochemicals” for mediating airborne messages. While volatile terpenes have  
113 been studied largely in plants and microbes<sup>3-5</sup>, they have received comparatively little attention in  
114 animals. This is surprising given the occurrence of volatile terpenes in invertebrates, especially  
115 arthropods, but also in different lineages of vertebrates. Even fewer information is available on  
116 the enzymatic formation of volatile terpenes in animals, which is partly due to the general notion  
117 that specialized metabolites in animals are predominantly derived from other organisms such as  
118 host plants or microbial endosymbionts.<sup>6,7</sup> Therefore, pathways and reactions involved in de novo  
119 biosynthesis of terpene infochemicals in animals are not well understood.

120

121 All organisms are capable of making the 5-carbon precursors of terpenes, isopentenyl  
122 diphosphate (IPP, **1**) and its allylic isomer dimethylallyl diphosphate (DMAPP, **2**) by conserved  
123 core enzymatic pathways. Plants produce IPP and DMAPP via the plastidial methylerythritol  
124 phosphate (MEP) and the non-plastidial mevalonic acid (MVA) pathway.<sup>8</sup> Cyanobacteria, parasitic  
125 protozoa, and most eubacteria share the MEP pathway with plants, while animals, fungi, and most  
126 other eukaryotes rely solely on the MVA pathway in providing the C5-diphosphate precursors for  
127 terpene primary and specialized metabolism (Fig. 1).<sup>9-11</sup> Isoprenyl diphosphate synthase (IDS)  
128 enzymes consecutively combine DMAPP with IPP units in a (1'–4) head-to-tail condensation  
129 reaction to synthesize 10-carbon geranyl diphosphate (GPP, **3**), 15-carbon farnesyl diphosphate  
130 (FPP, **4**), 20-carbon geranylgeranyl diphosphate (GGPP, **5**), as well as longer prenyl  
131 diphosphates required for the formation of non-volatile 30-carbon triterpenes, 40-carbon  
132 tetraterpenes, and longer polyprenyl terpenes (e.g. the 50-carbon tail of ubiquinone-10 derived  
133 from decaprenyl diphosphate, DPP, **6**).<sup>8,12,13</sup> Condensation reactions can occur in *cis*- or *trans*  
134 configuration and are catalyzed by structurally unrelated *cis*- and *trans*-IDS enzymes.<sup>14</sup> Enzymatic  
135 reactions catalyzed by terpene synthases (TPS) then convert DMAPP, GPP, FPP, and GGPP to  
136 acyclic or cyclic hemiterpenes (e.g. isoprene, **7**), monoterpenes, sesquiterpenes, and diterpenes,

137 respectively (Fig. 1). [Place Fig. 1 here] Depending on secondary modifications such as  
138 hydroxylation, acylation, methylation, glycosylation, and others, the produced terpenes remain  
139 volatile or semi-volatile or are converted into non-volatile derivatives. Canonical class I and II  
140 enzymes of the TPS superfamily have been studied extensively in plants and microbes.<sup>15-17</sup>  
141 However, there is growing evidence for the role of various non-canonical TPS enzymes in terpene  
142 biosynthesis as recently described by Rudolf and Chang<sup>14</sup>, raising questions about the evolution  
143 of “unconventional TPSs” in animals. In addition, microbial-type TPSs may have been integrated  
144 in animal genomes through horizontal gene transfer as it was found previously for TPS enzymes  
145 in lower land plants.<sup>18</sup> In this review, we will first provide an overview of the diversity of volatile  
146 and semi-volatile terpenes in the animal kingdom. We will then describe recent findings of  
147 pathways and TPSs involved in the de novo formation of terpenes in animals, with a particular  
148 focus on non-canonical IDS-type enzymes in insects that have adopted TPS activities. Moreover,  
149 we will discuss the structural modifications that are likely involved in the transition of these  
150 enzymes from IDS to TPS function and compare them with structural features of microbial-type  
151 TPSs discovered in octocorals and mites.

152

## 153 **2 Volatile terpenes in the animal kingdom**

154

### 155 **2.1 Invertebrates**

156

#### 157 **2.1.1 Mollusks and Corals**

158 The largest number of terpene specialized metabolites among invertebrate organisms have been  
159 identified in the species-rich phyla of arthropods and mollusks. Terpenes including monoterpenes,  
160 sesquiterpenes, and diterpenes, are particularly abundant in marine mollusks such as sponges  
161 and opisthobranch gastropods (e.g. brasudol, **8**, an eudesmane-type sesquiterpene from the sea  
162 hare *Aplysia brasiliiana*, Fig. 2).<sup>19,20</sup> While many of these compounds share core linear or cyclic  
163 scaffolds with terpenes made by terrestrial animals, they typically carry groups with other  
164 characteristic atoms such as halogens, nitrogen, and sulfur that affect water solubility and lead to  
165 considerable structural diversity.<sup>19</sup> Similar to mollusks, octocorals including soft corals are known  
166 as a rich source of bioactive sesquiterpenes and diterpenes (e.g. lophotoxin, **9**, a cembranoid  
167 diterpene from *Leptogorgia chilensis*, Fig. 2).<sup>21,22</sup> Overall, these specialized terpenes are believed  
168 to benefit the producer in the form of chemical feeding deterrents (e.g. Ben et al.<sup>23</sup>). It should be  
169 noted that this function does not necessarily require extensive water solubility. For instance,  
170 Giordano et al.<sup>24</sup> showed that volatile hydrophobic furanosesquiterpenes (e.g. isofuranodiene, **10**,

171 Fig. 2) released by a Mediterranean octocoral function as olfactory cues to ward off predators.

172 [Place Fig. 2 here]

173 Despite the profound structural diversity of terpenes found in marine invertebrates, the pathways  
174 of their formation remain obscure. In most cases, it is believed that mollusks sequester these  
175 metabolites from their algal diet prior to further biotransformation. For instance, species within the  
176 opisthobranch family Aplysiidae (sea hares) are known to derive sesquiterpenes from precursors  
177 sequestered from their algal prey.<sup>19</sup> Similar dietary sequestrations likely occur for sesquiterpene  
178 and diterpene skeletons in nudibranch-sponge predator-prey interactions (e.g. Shen et al.<sup>25</sup>) and  
179 even between specific mollusks and their soft coral prey.<sup>19</sup> On the other hand, marine  
180 invertebrates are generally believed to accumulate terpenes from microbial sources.<sup>26</sup> In  
181 particular, it has been suggested that symbiotic dinoflagellates may be the sources of diterpenes  
182 found in octocorals.<sup>27</sup> While many of the sequestered terpenes may indeed be derived from food  
183 sources, it remains unclear how the host further modifies individual skeletons. In other cases, the  
184 ability of marine gastropods to synthesize terpene skeletons de novo might have simply been  
185 overlooked. A striking example of such capability was recently brought to light with the  
186 identification of terpene synthases in coral genomes (see section 5). Despite the significant  
187 diversity of terpenes in marine mollusks, no accumulation of terpenes has, to the best of our  
188 knowledge, been reported in panpulmonate gastropods inhabiting terrestrial or freshwater  
189 environments. Moreover, examples of terpene compounds occurring within panpulmonate  
190 species are scarce compared to examples in other gastropods and appear restricted to marine  
191 genera.<sup>28</sup> This astonishing difference clearly indicates dissimilarities in the chemical ecology of  
192 food webs and chemically mediated defenses of mollusks in marine and terrestrial ecosystems.

193

### 194 **2.1.2 Arthropods – Millipedes and arachnids**

195 In contrast to terrestrial mollusks, terpenes are substantially more abundant in terrestrial  
196 arthropods in conjunction with intra-specific communication or chemical defense. Millipedes  
197 (arthropod class Diplopoda), which represent the oldest fully terrestrial group of animals, are  
198 known for their evolution of complex chemical defenses, some of which include terpene-derived  
199 compounds. A nitrogen-containing terpene called polyzonimine (**11**) and its related compound  
200 nitropolyzonamine (**12**) have been identified in polyzoniidan millipedes (Fig. 3).<sup>29-31</sup> More recently,  
201 another alkaloid named gosodesmine (**13**), a 7-substituted hexahydroindolizine carrying an  
202 isoprenyl moiety was detected in the millipede *Gosodesmus claremontus* of the related order  
203 Platydesmida.<sup>32</sup> Shear also found volatile monoterpenes such as limonene (**14**) in polyzoniidan  
204 and siphonophoridan millipedes.<sup>29</sup> How or whether any of these compounds are derived from

205 millipede-specific biosynthetic pathways or dietary sources is currently unclear. A microbial origin  
206 of terpenes is more likely in some species of polydesmid millipedes (*Niponia nodulosa*), where  
207 adult stages release 2-methyl-isoborneol (**15**) and the sesquiterpene geosmin (**16**) as volatile  
208 defense compounds or possible alarm pheromones.<sup>33</sup> These compounds are typically produced  
209 by cyanobacteria and Actinomycetes, which are possibly acquired by the millipedes via food  
210 intake.<sup>34</sup>

211 [Place Fig. 3 here]

212 Among the arachnids, the occurrence of volatile terpenes has been reported in the order Opiliones  
213 (harvestmen) and different groups of mites. Harvestmen of the species *Sclerobunus robustus*  
214 carry bornyl esters (e.g. bornyl acetate, **17**, stereoisomer not reported, Fig. 3) and small amounts  
215 of other monoterpenes in their defense secretions.<sup>35</sup> Oribatid mites (beetle or moss mites) are  
216 known to release the iridoid monoterpene chrysomelidial (**18**) and the diterpene  $\beta$ -springene (**19**)  
217 from exocrine oil glands.<sup>36</sup> Moreover, dust mites (Acariformes, Epidermoptidae) emit the  
218 monoterpene ester neryl formate (**20**) as an aggregation pheromone, and the monoterpene  $\beta$ -  
219 acaridial (**21**) has been found in other acarid mites as sex, aggregation, and alarm pheromone.<sup>37,38</sup>  
220 It can also be assumed that trombidid mites such as chiggers produce terpenes for chemical  
221 interactions based on the surprising finding of TPS genes in the mites' genomes, which  
222 presumably have been acquired from microbial sources via horizontal gene transfer (see 3.2).<sup>39</sup>

223

### 224 **2.1.3 Arthropods – Insects**

225 The by far largest group of terrestrial invertebrates known to release terpenes as infochemicals  
226 are insects. Insects employ volatile monoterpenes, sesquiterpenes, and semi-volatile diterpenes  
227 in various intra- and inter-specific interactions such as mate-finding (sex and aggregation  
228 pheromones), predator-avoidance (alarm pheromones), facilitation of eusocial living, food finding  
229 (trail pheromones) and chemical defense.<sup>40-44</sup> The large number of different compounds that have  
230 been reported makes it impossible to list them all in this review. Therefore, we will focus on the  
231 most characteristic and recent findings of compounds in the context of their biosynthesis and gene  
232 discovery. Comprehensive listings of terpenes and other insect semiochemicals are accessible  
233 through the Pherobase database (<https://www.pherobase.com/>).

234

235 One of the most basal insect orders where terpene defenses have been reported are the  
236 **Phasmatodea** or stick insects. Monoterpene iridoids such as actinidine (**22**) and nepetalactone  
237 (*cis,trans* and/or *trans,cis*) (**23**) are disseminated by these insects in defense secretions to deter  
238 predators (Fig. 4).<sup>45,46</sup> Nepetalactone and other monoterpene iridoids also occur as defenses in

239 rove beetles and as sex pheromones in aphids.<sup>42,47</sup> Positioned in the same larger taxonomic clade  
240 as stick insects, the **Blattodea** including cockroaches and termites are well known for releasing  
241 terpenes as pheromones and defense metabolites. Roaches of the genus *Periplaneta*, notably  
242 the American cockroach *Periplaneta americana*, use cyclic sesquiterpenoids called periplanones  
243 (e.g. periplanone B, **24**) as female-specific aggregation/sex pheromones.<sup>48-51</sup> In families of  
244 advanced termites (Termitidae and Rhinotermitidae) soldiers release blends of mono-, sesqui-,  
245 and/or diterpenes, including the cembrene A- (**25**) derived multi-cyclic secotrinervitane- (**26**),  
246 trivernitane- (**27**), and kempene-type (**28**) diterpenes, as part of their frontal gland defense  
247 secretions.<sup>52-54</sup> Other constituents of these secretions such as (+)- $\alpha$ -pinene (**29**) and (*E,E*)- $\alpha$ -  
248 farnesene (**30**) are thought to act as alarm pheromones<sup>55-57</sup> or function as primer pheromones  
249 involved in the developmental differentiation of members of the soldier caste.<sup>58</sup> Additional  
250 functions of terpenes in higher termites include roles as queen sex pheromones [(3*R*,6*E*)-  
251 nerolidol] (**31**) and trail pheromones (e.g. cembrene A, **25**) in *Prorethra simplex*.<sup>44,59</sup>  
252 [place Fig. 4 here]

253 Another large and diversified group of insects, which has been documented for its dissemination  
254 of terpene semiochemicals, comprises the **Hemiptera** (also referred to a true bugs) and their  
255 sister lineage, thrips (**Thysanoptera**). Many species of thrips use monoterpenes or their  
256 derivatives as pheromone or defensive components. For example, the gall-forming thrips,  
257 *Thlibothrips isunoki*, produces a defensive secretion/alarm pheromone containing  $\beta$ -myrcene (**32**)  
258 (Fig. 4) upon disturbance of the host gall, and the monoterpene  $\beta$ -acardial (**21**) has been  
259 identified in secretions from other gall-forming thrips.<sup>60,61</sup> Notably, male aggregation pheromones  
260 of flower thrips such as *Frankliniella occidentalis* are composed of the irregular monoterpene (*R*)-  
261 lavandulol (**33**) and its ester derivatives<sup>62,63</sup>, raising questions about whether these compounds  
262 are biosynthesized by irregular TPSs similarly to those identified in plants.<sup>64</sup> Interestingly,  
263 derivatives of lavandulol and irregular cyclopropane and cyclobutane monoterpenes and  
264 sesquiterpenes are also used by hemipteran scale insects and mealy bugs as sex pheromones  
265 (e.g. oleanderlure, **34**, from the oleander scale, *Aspidiotus nerii*).<sup>65-67</sup> In addition to these  
266 compounds, a large number of acyclic and cyclic regular monoterpenes and sesquiterpenes serve  
267 as sex or aggregation pheromones and defense compounds in several other hemipteran lineages  
268 with stink bugs and related species being the most prominent group (e.g. murgantiol, **35**, from the  
269 harlequin bug *Murgantia histrionica*).<sup>68</sup> These terpenes will be discussed in more detail below in  
270 the context of recent findings of their de novo biosynthesis (see 3.3).

271

272 Eusocial species of the **Hymenoptera** rely on complex communication and defense systems that  
273 are mediated by chemicals released from exocrine/secretory glands. The released compounds  
274 largely facilitate social organization and colony defense<sup>69</sup>, often through the emission of terpenes.  
275 For example, the red imported fire ant, *Solenopsis invicta*, uses isomers of the linear  
276 sesquiterpene (*Z,E*)- $\alpha$ -farnesene (**36**) in its worker trail pheromone.<sup>70</sup> Similar emissions of  
277 monoterpenes such as (*E*)- $\beta$ -ocimene (**37**) occur in species of army ants<sup>71</sup>, and blends of linear  
278 monoterpenes and sesquiterpenes including (*Z*)-citral (neral) (**38**), geraniol (**39**), and farnesol (**40**)  
279 compose the trail pheromone blend secreted from the Nasonov gland of the honey bee *Apis*  
280 *mellifera* (Fig. 5).<sup>72</sup> Other examples include (*S*)-citronellol (**41**), a pheromone of male *Bombus*  
281 bumblebees to attract virgin queens<sup>73</sup>, and the monoterpenes (+)-limonene [(+)-**14**] and  $\alpha$ -  
282 phellandrene (**42**) that serve as alarm pheromones and solvents for toxic alkaloids in the poison  
283 glands of *Myrmecaria* ant species.<sup>74</sup>

284 [place Fig. 5 here]

285 Monoterpenes and sesquiterpenes further occur as aggregation pheromone constituents and  
286 defense compounds in **Coleoptera**. Monoterpene alcohols, ketones and acetals from bark  
287 beetles of the genera *Ips* and *Dendroctonus* (Scolytidae) are among the best studied examples  
288 of aggregation pheromones. Investigation of their biosynthetic origins revealed that they are either  
289 derived from host tree-specific monoterpenes [*cis/trans*-verbenol (**43**) and verbenone (**44**) from  
290  $\alpha$ -pinene (**29**)] or formed de novo from endogenous terpene precursors made by the MVA  
291 pathway [e.g. ipsdienol (**45**), sulcatone (**46**), frontalin (**47**)] (Fig. 5).<sup>43</sup> Elucidation of the ipsdienol  
292 biosynthetic pathway, which now has been completed<sup>43</sup>, led to the identification of the first  
293 endogenous IDS-type TPS enzymes in insects (see 3.3). In other Coleoptera, non-oxygenated  
294 sesquiterpenes represent predominant components of their pheromone blends. The invasive,  
295 cosmopolitan Asian lady beetle *Harmonia axyridis* releases a female-specific sex/aggregation  
296 pheromone primarily consisting of (*E*)- $\beta$ -caryophyllene (**48**) and isomers.<sup>75</sup> Flea beetles of  
297 *Phyllotreta* and *Aphthona* genera also release sesquiterpenes in the form of himachalene and  
298 cadinene isomers as male-specific aggregation pheromones (e.g. **49**, **50**).<sup>76</sup> Characteristic  
299 terpene defense compounds found in beetles are the iridoid monoterpene chrysolmelidial (**18**)  
300 secreted by larvae of leaf beetles, the irregular monoterpene grandisol (**51**), an aggregation  
301 pheromone in the cotton boll weevil, *Anthonomus grandis*<sup>77</sup>, and the highly toxic sesquiterpene  
302 cantharidin (**52**), which is produced by male blister beetles (Meloidae) and transferred to females  
303 and offspring through nuptial gifts.<sup>78</sup>

304

305 Lastly, terpene semiochemicals also occur in butterflies (**Lepidoptera**) and flies (**Diptera**). In  
306 particular, larvae of swallowtail butterflies in the genus *Papilio* use blends of linear and cyclic  
307 mono- and sesquiterpenes such as germacrene A (**53**) as defense compounds (Fig. 5). Late-  
308 instar larvae emit these volatiles from osmeteria, which are fork like defense organs that are  
309 everted upon threat.<sup>79,80</sup> In another case, the monoterpene (*E*)- $\beta$ -ocimene (**37**) is used by the  
310 males of *Heliconus* butterflies as an anti-aphrodisiac pheromone transferred to females during  
311 copulation to repel conspecific males.<sup>81</sup> The compound was recently found to be made by an IDS-  
312 type TPS enzyme as discussed in 3.3.2. Examples of terpenes found in dipteran sex pheromones  
313 include monoterpene and sesquiterpene blends from fruit flies (*Ceratitis capitata*<sup>82</sup>, *Anastrepha*  
314 *suspensa*<sup>83</sup>) and homosesquiterpene or diterpene constituents (e.g. 3-methyl-himachalene, **54**)  
315 from the Brazilian sand fly *Lutzomyia longipalpis*, which is a vector of trypanosome parasites  
316 responsible for leishmaniasis disease in humans.<sup>84</sup>

317  
318 Taken together, volatile and semi-volatile terpenes of substantial structural diversity are released  
319 from species of at least nine insect orders (Fig. 6) ranging from simple acyclic monoterpenes in  
320 bumble and honey bees to multi-cyclic diterpenes in termites. The appearance of iridoids and  
321 other terpene pheromones and defenses in basal lineages of stick insects and the Blattodea  
322 raises questions about how early terpene specialized metabolism emerged in insect evolution.  
323 Biosynthetic studies of volatile terpenes in invertebrates (see section 3) will provide answers to  
324 these questions and may also help understand the formation of these infochemicals in  
325 vertebrates.

326 [place Fig. 6 here]

327

## 328 **2.2 Vertebrates**

329 In comparison to the diverse nature of terpenes found in invertebrate animals, the occurrence of  
330 terpene semiochemicals in vertebrates appears rather limited. While chemical communication  
331 and scent marking are integral components of vertebrate communication, the reduced abundance  
332 of terpenes is not entirely surprising given the expansion of other mechanisms of communication  
333 including acoustic, visual, and tactile signals. However, there may be other reasons for the limited  
334 use of terpenes in higher animals such as possible adverse physiological effects of specialized  
335 terpene compounds. Other causes may include competition and confusion with terpene signals  
336 that evolved earlier in plants and lower animals or simply to escape scent-guided predation by  
337 other animals. However, there is currently no clear evidence for any of these assumptions.

338

### 339 **2.2.1 Amphibians**

340 Some of the most interesting findings of terpene semiochemicals in recent years have been made  
341 in amphibians, more specifically in **anuran amphibians** comprising frogs and toads. African reed  
342 frogs of the family Hyperoliidae and species in the family Mantellidae, which is endemic to  
343 Madagascar, use pheromones for chemical communication.<sup>85-87</sup> Mantellid males release volatile  
344 compounds from scent glands on the ventral sides of their shanks. These compounds consist of  
345 alcohols and macrocyclic lactones (macrolides)<sup>86</sup>, which can be perceived by the olfactory system  
346 of the frogs.<sup>88</sup> Males of African red frogs attract females by inflating vocal sacs that carry colorful  
347 glands. Unexpectedly, these glands release sesquiterpenes besides aliphatic macrolides in  
348 complex species-specific mixtures. One of the sesquiterpenes was identified as the sesquiterpene  
349 macrolactone (S)-3,7,11-dodec-6,10-dien-12-olide named frogolide (**55**) because of its broader  
350 occurrence in both hyperoliid and mantellid families (Fig. 7).<sup>85</sup> Sesquiterpene macrolides are not  
351 unique to frogs but occur also as tris-norsesquiterpene lactone pheromones in insects such as  
352 the compound cucujolide I (**56**) in cucujid grain beetles and *Pieris* butterflies and its isomer  
353 suspensolide (**57**) in males of the Caribbean fruit fly *Anastrepha suspense*.<sup>89-91</sup> It is, therefore,  
354 possible that frogs obtain their sesquiterpene lactones from their insect diet. However, Mencke et  
355 al. assume that frogolide is synthesized de novo since the compound was present in frogs fed  
356 with frogolide-free insect diet.<sup>85</sup> Similarly, Shear notes that poison frogs carry the terpene alkaloid  
357 polyzonimine (**11**) even though their diet does not contain millipedes, which are known to  
358 accumulate this compound for defense.<sup>29</sup> It might also be possible that microbes associated with  
359 the frog's skin or gland represent the source of these metabolites, which warrants further  
360 investigation. Finally, the presence of the common sesquiterpene (*E*)- $\beta$ -caryophyllene (**48**) in skin  
361 secretions of the Australian green tree frog *Litoria caerulea* has been shown to result from the  
362 sequestration of this compound from insect diets.<sup>92</sup>

363 [place Fig. 7 here]

### 364 **2.2.2 Reptiles**

365 Volatile terpenes in reptiles have, to the best of our knowledge, only been studied in the context  
366 of crocodilian chemical ecology. In particular, secretions from the paracloacal glands of crocodiles  
367 have been investigated, which are believed to produce nesting and/or mating pheromones. Early  
368 studies on smooth-fronted caimans (*Paleosuchus trigonatus*) and the Chinese alligator (*Alligator*  
369 *sinensis*) detected the diterpenes  $\beta$ -springene (**19**) and cembrene A (**25**) as well as its ketone  
370 derivative 11,12-dihydrocembren-10-one (**58**) (Fig. 7).<sup>93,94</sup> Schulz et al. later identified additional  
371 novel acyclic monoterpene and sesquiterpene hydrocarbons in these and other alligatorid  
372 species<sup>95</sup>, and García-Rubio et al. found citronellyl esters (e.g. citronellyl acetate, **59**) among other

373 unidentified terpenes in the gland secretions of the American crocodile (*Crocodylus acutus*).<sup>96</sup> It  
374 is tempting to speculate that crocodiles have evolved their own enzymes for the formation of these  
375 terpene compounds; however, there is currently no immediate genomic evidence for the presence  
376 of such proteins.

377

### 378 **2.2.3 Mammals**

379 Volatile terpenes have been documented in glandular secretions of several mammalian species.  
380 Exudates of the dorsal gland of the springbok (*Antidorcas marsupialis*) are known to contain a  
381 series of C8 to C30 terpene hydrocarbons and ketones with the diterpene hydrocarbons  $\alpha$ - and  
382  $\beta$ -springene (**19**, **60**) being the most common compounds (Fig. 7).<sup>97</sup> The secretions are believed  
383 to serve as conspecific alarm signals. In African elephants (*Loxodonta africana*), the simple  
384 sesquiterpene alcohol (*E,E*)-farnesol (**40**), its hydrate derivatives (e.g. **61**) and the cyclic  
385 sesquiterpene alcohol drima-8 $\alpha$ ,11-diol (**62**) are constituents of secretions from temporal glands,  
386 which are modified facial sweat glands that are particularly active in stressed and aggressive  
387 animals.<sup>98,99</sup> More recent reports of volatile terpenes in mammals include the finding of the  
388 monoterpene alcohol linalool (**63**) and linalool oxides in pheromone secretions from shoulder  
389 glands of male Northern yellow-shouldered-bats (*Sturnira parvidens*).<sup>100</sup> Moreover, common  
390 monoterpene hydrocarbons and alcohols were detected in ano-genital odor secretions used for  
391 scent marking by crowned lemurs (*Eulemur coronatus*).<sup>101</sup> A number of monoterpenes and  
392 sesquiterpenes including compounds **64-68** (Fig. 7) have also been found in sternal gland  
393 secretions of male koalas (*Phascolarctos cinereus*).<sup>102</sup> Except for the volatiles released by the  
394 springbock and elephants, most of the compounds reported in the other cases likely originate  
395 from diet sources such as terpene-rich fruits and leaves. For instance, 1,8-cineole (**64**) found in  
396 scent secretions of male koalas is the predominant monoterpene constituent of the leaf essential  
397 oil of eucalyptus trees, the primary food source of koalas.<sup>103</sup> No monoterpene derivatives of 1,8-  
398 cineole were reported in the scent secretions making it unlikely that this compound is further  
399 converted by *P. cinereus* endogenous pathways.

400

401 In summary, this survey of specialized terpenes in animals shows that several lineages of  
402 animals, especially invertebrates, have integrated terpene compounds in their semiochemical  
403 repertoire for intra- and interspecific interactions (Fig. 8). Despite the diversity of terpenes in  
404 different animal species, there is also overlap in the constituents of chemical blends. For example,  
405 the acyclic diterpene  $\beta$ -springene occurs in mites as well as the springbock. Whether these  
406 metabolites are produced by similar enzymes and if terpene infochemicals are more broadly

407 synthesized de novo in the animal kingdom is in many cases not well understood, with discoveries  
408 of genes and enzymes just beginning to emerge.

409 [place Fig. 8 here]

410

### 411 **3 Terpene biosynthetic pathways and enzymes in insect pheromone/defense**

#### 412 **biosynthesis**

413 Since the majority of terpene specialized metabolites has been identified in plants and microbes,  
414 terpene biosynthetic enzymes have largely been elucidated in these organisms, and  
415 comparatively little attention has been given to the discovery of equivalent enzymatic steps in  
416 animals. The fact that animals can sequester specialized metabolites from their food sources or  
417 microbial symbionts has complicated the search for de novo biosynthetic pathways. However, an  
418 increasing number of high-quality transcriptomes and genomes, which allow detailed genomic  
419 and phylogenetic comparisons, facilitate the discovery of terpene biosynthetic genes in animals.  
420 Here we review recent findings of enzymes involved in terpene de novo biosynthesis in insects  
421 and arachnids.

422

#### 423 **3.1 Biosynthesis of iridoids**

424 Methylcyclopentanoid monoterpenes or iridoids act as defensive compounds and sex  
425 pheromones in a number of different insects. Several of the same compounds also occur in plants  
426 where they are involved in defensive activities.<sup>104</sup> For instance, nepetalactone (**23**) is best known  
427 as the characteristic iridoid compound of catnip (*Nepeta cataria*) and functions as an insect  
428 repellent, but it also serves as a sex pheromone in aphids.<sup>42,105</sup> Gene cluster analysis in *Nepeta*  
429 led to the identification of the nepetalactone biosynthetic pathway in this species.<sup>106</sup> An equivalent  
430 pathway has been elucidated for the formation of *cis-trans*-nepetalactol (**69**), which is a precursor  
431 in the biosynthesis of pharmacologically important monoterpene indole alkaloids in Madagascar  
432 periwinkle (*Catharanthus roseus*).<sup>107,108</sup> The identification of these pathways in plants raised the  
433 question of the existence of similarly evolved pathways in insects. Several enzymatic steps were  
434 initially characterized in the production of iridoid-related dialdehydes such as chrysomelidial (**18**),  
435 which are made by larvae of the chrysomelid leaf beetle *Phaedon cochleariae*.<sup>109-113</sup> Most of these  
436 steps have been verified by the just completed identification of the entire biosynthetic pathway of  
437 the nepetalactone sex pheromone in the pea aphid *Acyrtosiphon pisum* (Fig. 9).<sup>114</sup> Sexual  
438 females of *A. pisum* exclusively secrete (1*R*,4*aS*,7*S*,7*aR*)-*cis-trans*-nepetalactol (**69**) and  
439 (4*aS*,7*S*,7*aR*)-*cis-trans*-nepetalactone (**23**) from glands of their hind legs. The elegant study by  
440 Köllner et al. determined pathway specific gene candidates by differential gene expression

441 analysis of the hind legs and the non-pheromone producing front legs of sexual females as well  
442 as the hind legs of asexual females and males.<sup>114</sup> Functional characterization of target gene  
443 candidates (Table 1) established a 7-step pathway (Fig. 9), which starts with the formation of GPP  
444 (**3**) by a GPP synthase homolog of the *P. cochleariae* bifunctional GPP/FPP synthase enzyme  
445 that shares the same  $\text{Co}^{2+}/\text{Mg}^{2+}$  metal dependency. This initial step is followed by the conversion  
446 of GPP to geraniol (**39**) catalyzed by a dolichyldiphosphatase-type homologue (ApGES), and a  
447 subsequent hydroxylation to 8-hydroxygeraniol (**70**) by a cytochrome P450 enzyme of clan 3  
448 (ApG8H). The alcohol is then converted in a two-step oxidation by an NADP-dependent short-  
449 chain dehydrogenase (ApHGO) to form 8-oxo-geraniol (**71**). The resulting aldehyde is  
450 subsequently reduced to 8-oxo-citronellyl enol (**72**) and cyclized to *cis-trans*-nepetalactol by a  
451 membrane-bound reductase (ApISY) related to polyprenol type reductases. The final oxidation to  
452 *cis-trans*-nepetalactone is catalyzed by a GMC-type oxidase (ApNEPO).<sup>114</sup>

453 [Place Fig. 9 here]

454  
455 Interestingly, with the exception of the GPP synthase, which is likely localized in mitochondria,  
456 several enzymes of the pathway are presumably associated with the ER membrane (ApGES,  
457 ApG8H, ApISY) or targeted to the ER lumen (ApNEPO). This membrane-specific association  
458 suggests that the proteins could be organized in the form of a metabolon. Metabolons that  
459 modulate metabolic flux and facilitate efficient channeling of intermediates have been  
460 documented for a number of secondary metabolic pathways in plants,<sup>115</sup> whereas still little is  
461 known about such protein complexes in insects.<sup>116</sup> Comparison of the aphid-specific enzymes  
462 with those identified in leaf beetles indicates that several of the pathway genes evolved  
463 independently in these insect lineages. Moreover, a comparison with the plant-specific formation  
464 of nepetalactone in *Catharanthus* and *Nepeta* reveals that plants and insects employ the same  
465 enzymatic steps but use unrelated enzymes. For instance, in plants, TPS enzymes catalyze the  
466 formation of geraniol,<sup>106,107</sup> whereas this step is mediated by a phosphatase in aphids. In addition,  
467 aphids use a membrane bound polyprenol reductase-like protein for the formation of nepetalactol;  
468 by contrast, plants employ members of the short-chain dehydrogenase/reductase (SDR) family  
469 to catalyze this step.<sup>106,108</sup> The finding of these independent iridoid biosynthetic pathways  
470 represents a powerful example of convergent evolution in the metabolism of volatile terpene  
471 semiochemicals in plants and animals.

472 [Place Table 1 here]

473

474 **3.2 Horizontal gene transfer of terpene synthases in mites**

475 A surprising finding of TPS gene families has recently been made in the genomes of trombidid  
476 mites. Larvae in the superfamilies of the Trombiculoidea (chiggers) and Trombidioidea (velvet  
477 mites) feed as ectoparasites on vertebrates or other arthropods, respectively.<sup>39</sup> Genomes of the  
478 chigger *Leptotrombidium deliense* and the velvet mite *Dinotrombium tinctorium* were found to  
479 contain a family of 39 putative sesqui-TPS genes and a related family of 21 TPS genes,  
480 respectively. An additional group of 17 TPS genes was identified in *L. delicense*. These genes  
481 and their encoded proteins are most closely related to fungal and bacterial TPSs, albeit with  
482 sequence identity of less than 30%. The absence of homologs in other arthropods or metazoans  
483 suggests that the TPS genes are the result of ancient lateral gene transfers from soil-derived  
484 bacteria and fungi. This horizontal gene transfer is similar to that of carotenoid biosynthetic genes,  
485 which are responsible for the orange coloration in both types of mites.<sup>39</sup> The biochemical function  
486 of the detected TPS genes and the role of their putative terpene products as pheromones or  
487 defense compounds is currently unknown. It remains to be determined why such large gene  
488 families have been maintained in the mite genomes and to what extent functionally active genes  
489 may be correlated with the release of terpene blends.

490

### 491 **3.3 IDS-like terpene synthases in insects**

492 With the exception of the recently discovered TPS genes in mite genomes, insects have generally  
493 been thought to lack homologs of canonical microbial and plant TPSs and thus the ability to  
494 synthesize specialized terpenes via endogenous pathways. However, early isotope-labeling  
495 experiments questioned this notion. For instance, experiments with <sup>14</sup>C-labeled precursors in the  
496 bark beetle *Ips pini* provided evidence that the monoterpene aggregation pheromone ipsdienol  
497 (**45**) is synthesized de novo via the MVA pathway.<sup>117</sup> A combination of biochemical and  
498 transcriptome analyses further determined a coordinated regulation of terpene biosynthetic genes  
499 with tissue specificity in the midgut and elevated expression in *I. pini* males, and upon treatment  
500 with juvenile hormone III.<sup>43</sup> This approach led to the identification of a bifunctional IDS/TPS  
501 enzyme, which makes GPP (**3**) from IPP (**1**) and DMAPP (**2**) and subsequently converts GPP to  
502 the ipsdienol precursor myrcene (**32**) (Fig. 10A, Table 1).<sup>118,119</sup> It should be noted that all  
503 subsequent steps from myrcene to ipsdienol have also been elucidated.<sup>43</sup> The GPP/myrcene  
504 synthase was found to be structurally related to canonical IDS enzymes<sup>118</sup> and carries two  
505 aspartate-rich motifs (DDIMD, NDFKD). These motifs, typically called first and second aspartate  
506 rich motifs (FARM and SARM), are characteristic of IDS proteins. The *I. pini* synthase might be  
507 targeted to peroxisomes based on computational predictions of its transit peptide. Despite its  
508 similarity to canonical IDS proteins, the *I. pini* enzyme shares 20% or lower amino acid sequence

509 identity with the *I. pini* FPP synthase and insect GGPP synthases such as *Drosophila*  
510 *melanogaster* GGPP synthase, which indicates an early divergence of this protein from canonical  
511 insect IDS enzymes.

512 [place Fig. 10 here]

### 513 **3.3.1 FPP synthase type terpene synthases**

#### 514 **3.3.1.1 *Phyllotreta striolata***

515 Almost ten years after the discovery of an IDS-type TPS gene in *I. pini*, a family of similar genes  
516 was identified in a leaf beetle, the striped flea beetle *Phyllotreta striolata* (Chrysomelidae).<sup>120</sup> Four  
517 out of five IDS-type genes were functionally characterized as sesquiterpene synthases. Among  
518 those, the recombinant protein of the male-expressed *PsTPS1* gene was found to convert (*Z,E*)-  
519 FPP (**73**) to (*6R,7S*)-himachala-9,11-diene (**74**), a major constituent of the *P. striolata* aggregation  
520 pheromone, together with five other sesquiterpenes including **75** and  $\gamma$ -cadinene (**76**) (Fig. 10B,  
521 Table 1). Interestingly, *PsTPS1* requires a (*Z,E*)-FPP isomer as a substrate, which is made by an  
522 unusual, *cis*-double bond forming IDS (*PsIDS3*) from GPP and IPP (Fig. 10B).<sup>120</sup> The other  
523 functionally active TPS enzymes converted (*E,E*)-FPP to (*E*)-nerolidol (**31**) (*PsTPS4*) and  
524 mixtures of sesquiterpenes (*PsTPS2* and 3). These enzymatic products were not detected *in vivo*  
525 since the corresponding genes are expressed at low levels in males and females. *PsTPS1* and  
526 the *P. striolata* (*E,E*)-FPP synthase (*PsIDS1*) carry putative mitochondrial targeting sequences,  
527 indicating a subcellular compartmentalization in the formation of the volatile sesquiterpenes. The  
528 study also provided insight into the evolution of the *P. striolata* TPS genes. Gene structures of  
529 *PsTPS1*, *PsIDS1* and *PsIDS3* comprised four exons and three introns. The positions of these  
530 introns and the intron phases were conserved in canonical IDS genes from other Coleoptera,  
531 Lepidoptera and Diptera, which indicated an emergence of IDS-type TPS genes from an IDS  
532 progenitor, presumably a FPP synthase.<sup>120</sup> In addition, *PsIDS1*, *PsIDS3*, and homologs from other  
533 insects were found to be under strong purifying selection, indicating a selective removal of  
534 deleterious variants to preserve IDS function in core metabolism. In contrast, the IDS-type TPS  
535 genes are under relaxed constraints, which is in agreement with the neofunctionalization and  
536 diversification of these genes in the evolution of pheromones and chemical communication.

537

#### 538 **3.3.1.2 Pentatomids**

539 Tholl and collaborators have investigated the presence of IDS-type TPS genes in stink bugs  
540 (Pentatomidae), a diverse family in the order of pierce-sucking hemipteran insects, which  
541 comprises herbivorous and carnivorous species.<sup>121</sup> Due to their ability to easily adapt to different  
542 environmental conditions, several species of stink bugs have become important pests with

543 economic impact on agricultural crops in the Neotropics and worldwide.<sup>41</sup> Mature males of at least  
544 ten genera of pentatomids emit sesquiterpenes with a bisabolane-type skeleton as sex or  
545 aggregation pheromones such as (6*S*,7*R*)- $\beta$ -sesquiphellandrene (**77**) in the red banded stink bug,  
546 *Piezodorus guildinii*, *cis*-zingiberenol [(3*R*,6*R*,7*S*)-1,10-bisaboladien-3-ol] (**78**) in the rice stink  
547 bug, *Oebalus poecilus*, 10,11-epoxy-1-bisabolen-3-ol (**35**) in the harlequin bug, *M. histrionica* and  
548 the brown marmorated stink bug, *Halyomorpha halys*, and *trans/cis*-(*Z*)- $\alpha$ -bisabolene epoxide (**79**)  
549 in the Southern green stink bug, *Nezara viridula* (Fig. 11).<sup>41,122</sup> The compounds are released as  
550 mixtures of distinct stereoisomeric composition, sometimes in combination with fatty acid  
551 derivatives.<sup>41</sup> The structural relationships of the terpene constituents of stink bug pheromones  
552 suggests that they could be synthesized *de novo* by evolutionary related pathways instead of  
553 being made from sequestered precursors. This notion is supported by the fact that stink bug  
554 specialists and generalists feed on different host plants, many of which do not synthesize  
555 bisabolane type sesquiterpenes or make them only in limited amounts.

556 [Place Fig. 11 here]

557  
558 The first investigation of IDS-like TPS genes was performed in the harlequin bug, *Murgantia*  
559 *histrionica* (tribe Strachiini), which is a crucifer specialist native to Central America and invasive  
560 in the southeast of the United States. Mature males of *M. histrionica* emit an aggregation  
561 pheromone, which is composed of the (3*S*,6*S*,7*R*,10*S*)- and (3*S*,6*S*,7*R*,10*R*)-stereoisomers of  
562 10,11-epoxy-1-bisabolen-3-ol (**35**) named murgantiol (Fig. 12A).<sup>123,124</sup> The pheromone attracts  
563 both males and females as well as nymphs. Sex- and development-specific transcriptome  
564 analyses led to the identification of a canonical (*E,E*)-FPP synthase (*MhFPPS*) and an IDS-type  
565 TPS (*MhTPS*), which converts (*E,E*)-FPP (**4**) to (1*S*,6*S*,7*R*)-1,10-bisaboladien-1-ol, called  
566 sesquipiperitol (**80**), as an intermediate in the pathway leading to murgantiol (Fig. 12A, Table  
567 1).<sup>123,125</sup> Sesquipiperitol is also produced in plant species of the Asteraceae, Zingiberaceae, and  
568 Cupressaceae families (e.g. Sy and Brown<sup>126</sup>). *MhTPS* presumably catalyzes a carbocation  
569 mediated reaction typical of a type I TPS enzyme.<sup>8</sup> In this reaction, a nerolidyl carbocation is first  
570 formed by a metal ion-catalyzed cleavage of the carbon-oxygen bond to release the  
571 pyrophosphate moiety of FPP. Next, a bisabolyll cation is generated by a 1,6 ring closure followed  
572 by a hydride shift to form a sesquipiperityl cation and subsequent quenching of the carbocation  
573 with water (Fig. 12A). Analysis of *MhFPPS* and *MhTPS* transcript abundances showed an equal  
574 expression of *MhFPPS* at nymphal and adult stages. By contrast, *MhTPS* is most highly  
575 expressed in mature males and exhibits highest transcript levels in epithelial cells associated with  
576 the cuticle of the ventral abdominal sternites, from which the pheromone is likely released.<sup>123</sup>

577  
578 Interestingly, the pheromone constituents of the invasive brown marmorated stink bug  
579 *Halyomorpha halys* (Stål) (tribe Cappaeini), which is native to Asia, share the same skeleton as  
580 murgantiol but with a different stereoisomeric composition of (3*S*,6*S*,7*R*,10*S*)-10,11-epoxy-1-  
581 bisabolen-3-ol and (3*R*,6*S*,7*R*,10*S*)-10,11-epoxy-1-bisabolen-3-ol being (**35**) emitted in a 3.5:1  
582 mixture (Fig. 12A).<sup>127</sup> To investigate whether the pheromones of *M. histrionica* and *H. halys*, which  
583 are native to different geographical regions, are produced by closely related enzymes or may be  
584 the result of convergent evolution, IDS-like genes were mined in *H. halys* genome and  
585 transcriptome resources.<sup>128,129</sup> A family of seven IDS-like genes was discovered, of which two  
586 were characterized as functionally active sesqui-TPSs (*HhTPS1*, *HhTPS2*) (Table 1) and a third  
587 was identified as a canonical (*E,E*)-FPP synthase (*HhFPPS*).<sup>68</sup> *HhTPS1* was found to be a  
588 putative ortholog of *MhTPS1*, which shares more than 80% amino acid sequence with the *M.*  
589 *histrionica* enzyme and converts (*E,E*)-FPP to the same (1*S*,6*S*,7*R*)-sesquipiperitol intermediate  
590 (**80**) as in the *M. histrionica* pheromone biosynthetic pathway. Similar to *M. histrionica*, *HhTPS1*  
591 is most highly expressed in mature males, in agreement with the male-specific release of the  
592 pheromone. However, its tissue-specific expression is highest in the fat body, suggesting a  
593 different localization of the pheromone-specific enzymes in *H. halys* and *M. histrionica*. *HhTPS2*,  
594 which encodes a multi-sesquiterpene synthase, shows comparatively low expression in males  
595 and females, and the function of this gene remains unclear.<sup>68</sup> The downstream enzymatic steps  
596 from sesquipiperitol to 10,11-epoxy-1-bisabolen-3-ol are not yet identified but presumably include  
597 a conversion to a zingiberenol isomer (**81**) and an epoxidation at C10/C11 by a cytochrome P450  
598 monooxygenase enzyme (Fig. 12A). Zingiberenol and sesquipiperitol have also been identified  
599 as pheromone constituents of other stink bugs such as the rice stink bugs *Oebalus poecilus* and  
600 *Mormidea v-luteum*, and the rice stalk stink bug *Tibraca limbativentris* (tribe Carpocorini), all of  
601 which are severe pests in South America.<sup>130-132</sup> This finding and the identification of two closely  
602 related sesquipiperitol synthases in *M. histrionica* and *H. halys* suggest that the enzymatic steps  
603 in terpene pheromone formation in these species have been evolutionary conserved independent  
604 of their tribe and geographic origin.

605 [Place Fig. 12 here]

606  
607 Another pheromone biosynthetic pathway in stink bugs leads to the formation of *trans*-/*cis*-(*Z*)- $\alpha$ -  
608 bisabolene epoxide (**79**) (Fig. 12B). The epoxide isomers are released by the males of the  
609 southern green stink bug *Nezara viridula*, which is a globally invasive pest with the origin in East  
610 Africa.<sup>133,134</sup> The same isomers are emitted by the neotropical green stink bug *Chinavia*

611 *impicticornis*, a close relative in the same tribe (Nezarini) as *N. viridula*. Comparative  
612 transcriptome analyses of *N. viridula* mature males and females discovered an *N. viridula* IDS-  
613 type TPS enzyme (*NvTPS1*), which catalyzes the conversion of (*E,E*)-FPP to (+)-(*S,Z*)- $\alpha$ -  
614 bisabolene (**82**) as the likely precursor of the sesquiterpene pheromone (Fig. 12B, Table 1).<sup>135</sup>  
615 The biosynthetic pathway is presumably localized in glandular cells at the ventral abdomen of  
616 mature males, from which the pheromone is emitted.<sup>136</sup> Unexpectedly, a functionally active  
617 sesquiperitol synthase gene (*NvTPS2*) was identified in *N. viridula* genome and transcriptome  
618 resources, which encodes a protein with approximately 80% amino acid sequence identity to the  
619 corresponding enzymes of *M. histrionica* and *H. halys*.<sup>68</sup> This finding is surprising since *N. viridula*  
620 neither releases sesquiperitol or any other compound of the murgantiol pheromone complex nor  
621 stores non-volatile derivatives of sesquiperitol. The conserved status of the sesquiperitol  
622 synthase independent of its role in pheromone biosynthesis suggests that the enzyme had  
623 additional or other functions in the common progenitor of these pentatomid species.

624

### 625 3.3.1.3 Genomic organization and evolution of stink bug and hemipteran IDS-like genes

626 The availability of quality genomes of *H. halys* and *N. viridula* (<sup>128</sup>,  
627 <https://www.ncbi.nlm.nih.gov/bioproject/PRJEB47893/>) has allowed a more detailed investigation  
628 of the architecture and genomic position of the IDS-type TPS genes in these species. The six  
629 IDS-like genes of *H. halys* were found to be organized in two separate clusters, each of which  
630 most likely emerged by gene duplication (Fig. 13A, B).<sup>68</sup> The canonical *HhFPPS* gene shares only  
631 low sequence identity with the IDS-like genes and is positioned independently of the IDS-like  
632 clusters, which suggests that this gene is derived from a more ancient duplication event. Genes  
633 of the *N. viridula* IDS-like family are organized in a similar fashion, although in the form of a single  
634 three-gene cluster (*NvTPS1* and two uncharacterized IDS genes), with *NvTPS2* and *NvFPPS*  
635 being positioned separately on two other chromosomes (Fig. 13A).<sup>68</sup>

636 [place Fig. 13 here]

637 A closer analysis of the architecture of the *H. halys* canonical FPPS and IDS-like genes revealed  
638 an identical composition of seven introns and eight exons and identical positions of nearly all  
639 intron phases. The intron phases are conserved in FPPS genes of other hemipteran and  
640 blattodean insects, which provides evidence for a shared ancestral exon-intron structure of IDS  
641 genes in these lineages.<sup>68</sup> Interestingly, the coleopteran FPPS genes and the IDS-type TPS  
642 genes of *P. striolata* have a reduced number of three introns.<sup>120</sup> This difference in gene  
643 architecture between representatives of the Hemiptera and Coleoptera supports the hypothesis  
644 of an independent emergence of TPS genes from FPPS progenitors in these orders.

645  
646 To gain a more in-depth understanding of the evolution of IDS-like genes in Pentatomids and  
647 Hemiptera that are known to release volatile terpenes as pheromones or defense secretions,  
648 Rebholz et al. mined hemipteran IDS-like genes from NCBI nucleotide and transcriptome  
649 assembly databases using *H. halys* FPPS and IDS-like protein sequences as search queries.<sup>68</sup>  
650 The search resulted in the identification of nearly 300 unique sequences, with 80% classified as  
651 canonical type FPPSs and the remaining 20% classified as IDS (FPPS)-like proteins with the  
652 potential to function as TPSs (Fig. 14). Phylogenetic analysis of the IDS-like proteins indicated a  
653 paralogous division of the pentatomid sequences in two clades, named TPS-a and TPS-b clades,  
654 in agreement with the position of the corresponding genes in two different clusters (Fig. 13, 14).  
655 The clades most likely evolved from a common ancestor of the Pentatomidae or possibly the  
656 Pentatomoidea superfamily approximately more than 100 million years ago. In agreement with  
657 the neofunctionalization of IDS-like genes and in contrast to the conserved clade of canonical  
658 FPPS genes, the pentatomid TPS clades evolved under positive selective pressure.<sup>68</sup> However,  
659 genes within both TPS clades have undergone limited inter- and intraspecific diversification  
660 following clade-specific divergence, which is evidenced by the conservation of sesquipiperitol  
661 synthases in different species (Fig. 12). Thus, Pentatomids seemingly have maintained small-size  
662 gene families that generate a limited number of terpene intermediates. Limited steps of  
663 derivatizations and combinations with other metabolites such as fatty acid derivatives are  
664 sufficient to generate species-specific pheromone blends.<sup>41</sup> A cross-kingdom comparison shows  
665 that pentatomid TPS gene families are notably smaller than those of flowering plants<sup>15,137-139</sup>,  
666 despite similar evolutionary time spans of more than 100 million years (Fig. 13B). The  
667 diversification of plant TPSs into several subfamilies is associated with the synthesis of complex  
668 terpene mixtures, which are believed to have multiple functions in attraction and defense and  
669 presumably target a larger number of organisms than the smaller compound mixtures released  
670 by insects.<sup>8</sup> Therefore, it can be assumed that the diversification of TPS genes in Pentatomids  
671 and other insects is directed by more specific chemical interactions and perhaps other constraints.  
672 [place Fig. 14 here]

673  
674 Several currently uncharacterized IDS-like genes, which may be associated with the formation of  
675 volatile terpenes, have been identified in other species of Pentatomids and the two main  
676 suborders of the Hemiptera, the Heteroptera and the Sternorrhyncha. For instance, IDS-like  
677 genes with close similarity to sesquipiperitol synthases were found to be expressed in the  
678 predatory spined soldier bug, *Podisus maculiventris*, which releases monoterpene alcohol

679 pheromones including *trans*-piperitol (**83**), the C10 analog of sesquiperitol (Fig. 14).<sup>140</sup> Several  
680 other families in the heteropteran infraorders Pentatomomorpha and Cimicomorpha are known to  
681 secrete monoterpenes for defense or attraction: Acanthosomatidae (shield bugs), Cimicidae (bed  
682 bugs), Cydnidae (burrowing bugs), Miridae (plant bugs), Lygaeidae (seed bugs), Pyrrhocoridae  
683 (red bugs), Tingidae (lace bugs) and others.<sup>68</sup> In conjunction with these findings, IDS-like gene  
684 transcripts have been identified in the burrower bug *Sehirus cinctus* and the boxelder bug *Boisea*  
685 *trivittata*, which are known to release monoterpenes such as  $\alpha$ -pinene (**29**) and 3-carene (**84**),  
686 respectively (Fig. 14).<sup>141-143</sup> In the hemipteran suborder Sternorrhyncha, IDS-like transcripts are  
687 present in scale insects (infraorder Coccoomorpha) including the lac insect *Kerria lacca*, which  
688 excretes cyclic sesquiterpene acids as lac components.<sup>144</sup> Another group of insects in which IDS-  
689 like transcripts have been identified are mealybugs (infraorder Coccoomorpha) such as the cotton  
690 mealybug, *Phenacoccus solenopsis*, which releases a methylbutenoate ester of the cyclobutane  
691 monoterpene *R*-maconelliol as a sex pheromone (**85**) (Fig. 14).<sup>65</sup> Related irregular monoterpenes  
692 (e.g. lavandulol, **33**) are also made by plant TPSs that catalyze an irregular coupling between  
693 isoprenoid units.<sup>145</sup> On the other hand, IDS-like genes are absent in aphids despite the presence  
694 of  $\beta$ -farnesene as a well-known alarm pheromone in this hemipteran group.<sup>42</sup> Moreover, no IDS-  
695 like genes were identified in the suborders Auchenorrhyncha (including cicadas and plant  
696 hoppers) and Coleorrhyncha (including moss bugs) in agreement with the lack of terpenes in  
697 these orders.<sup>68</sup> Overall, the large-scale phylogeny of IDS-like sequences in the Hemiptera  
698 supports an ancient emergence of these genes from canonical FPP synthases, possibly in a  
699 shared progenitor more than 350 million years ago (Fig. 14), and suggests that volatile terpenes  
700 are synthesized *de novo* in several hemipteran lineages. Further functional characterization of  
701 IDS-like genes will provide more insight into the extent of terpene biosynthetic evolution in  
702 hemipteran insects.

703

### 704 3.3.2 GGPP synthase type terpene synthases

705 IDS-like TPSs in insects are not only derived from FPPS enzymes but also evolved from  
706 geranylgeranyl diphosphate synthases (GGPPSs). Evidence for GGPPS-derived TPSs was  
707 provided by a recent finding of two GGPPS-like proteins with monoterpene synthase activity in  
708 the butterfly *Heliconius melpomene*.<sup>146</sup> Males of *H. melpomene* transfer the monoterpene  $\beta$ -  
709 ocimene (**37**) as an anti-aphrodisiac pheromone to females during mating to prevent subsequent  
710 mating attempts by other males.<sup>147</sup> However, the formation of the pheromone varies between the  
711 *Heliconius* species.<sup>148</sup> In order to determine the genetic origin of  $\beta$ -ocimene synthesis, Darragh et  
712 al. generated genetic mapping families between the  $\beta$ -ocimene producing *H. melpomene* and the

713 non-producing closely related species *H. cydno*.<sup>146</sup> The authors identified a QTL region with eight  
714 GGPPS-like genes derived from repeated gene duplications. One of these genes was found to  
715 encode a functional  $\beta$ -ocimene synthase (Fig. 15, Table 1), while the enzyme encoded by a  
716 second gene (HMEL037108g1) converted GPP and FPP to the monoterpene and sesquiterpene  
717 alcohols (*S*)- and (*R*)-linalool (**63**) and nerolidol (**31**, stereoisomer not determined), respectively.  
718 The in vivo function of the latter enzyme is unknown. Both proteins exhibit residual IDS activity,  
719 which may indicate their evolution through subfunctionalization from a bifunctional IDS/TPS  
720 progenitor. In agreement with the absence of  $\beta$ -ocimene in *H. cydno*, the  $\beta$ -ocimene synthase  
721 ortholog in this species was found to be functionally inactive, probably due to several non-  
722 synonymous mutations in the coding sequence. Similar to FPPS-like TPSs and in contrast to the  
723 conserved canonical GGPPSs, the evolution of the two monoterpene synthases occurred under  
724 relaxed selection constraints. Several pseudogenes were identified in the GGPPS-like family,  
725 indicating loss-of-function events.

726 [place Fig. 15 here]

727 It is possible that other insects which use  $\beta$ -ocimene as a pheromone, such as bumble bees and  
728 honey bees, have GGPPS-like or perhaps FPPS-like proteins that make  $\beta$ -ocimene. It is curious  
729 to note that the *H. melpomene*  $\beta$ -ocimene synthase is unable to convert GGPP to a diterpene  
730 product, which suggests possible constraints in accommodating GGPP as a substrate. This  
731 scenario might be different in other insect lineages. For instance, a family of GGPPS-like genes  
732 was found to be expressed in soldiers of the nasute termite *Nasutitermes takasagoensis*.<sup>149</sup> The  
733 defensive secretions of these termites contain a mixture of diterpenes and monoterpenes, which  
734 may be produced by the GGPPS-like enzymes. Based on these findings, it can be assumed that  
735 GGPPS-type genes with TPS function have emerged independently multiple times throughout  
736 insect evolution. This is supported by the identification of a GGPPS-like TPS in the green tea  
737 leafhopper *Empoasca onukii*, which converts GPP into geraniol (**39**).<sup>150</sup> The authors suggest that  
738 geraniol synthase activity is also present in other lepidopteran and coleopteran species.

739

#### 740 **4 Structural and mechanistic evolution of insect IDS-like TPSs**

741 Phylogenetic evidence for the emergence of IDS-like TPS enzymes from IDS progenitors in  
742 insects raises the question of which mutations and structural modifications facilitated this  
743 evolutionary transition. While extensive experimental proof is still missing, O'Maille and co-  
744 workers have developed a structural and mechanistic model for the change in catalytic function  
745 from IDS to TPS proteins.<sup>151</sup> IDS and TPS enzymes generally share a common alpha-helical  
746 protein domain ( $\alpha$ -domain) fold<sup>152,153</sup>, suggesting an ancient common origin. Whereas TPS protein

747 sequences in microbes and plants no longer have a close evolutionary relationship with IDS  
748 proteins of these organisms, the recruitment of IDS to TPS proteins seems to have occurred more  
749 recently in insects.

750  
751 IDS proteins carry two conserved aspartate rich motifs (DDxxD) called FARM and SARM, which  
752 are positioned on the opposite sides of the active site. These motifs facilitate the cleavage of the  
753 diphosphate moiety of the allylic substrate via coordination of Mg<sup>2+</sup> ions. Insect IDS-like TPS  
754 proteins possess the same motifs but show more frequent substitutions of the first and third  
755 aspartate of the SARM (Table 2, Fig. 16). In addition, aromatic amino acid residues in positions  
756 4 and 5 upstream of the FARM of bonafide FPPS proteins are substituted by non-aromatic  
757 residues in IDS-like sesqui-TPSs.<sup>120,123,135</sup> Molecular docking of (*E,E*)-FPP in the active-site cavity  
758 of a *M. histrionica* TPS homology model showed that these residue changes appear to be critical  
759 for the positioning of the FPP prenyl side chain into the cavity to facilitate subsequent  
760 cyclization.<sup>123</sup> Substitutions of the non-aromatic residues in *Mh*TPS with aromatic amino acids led  
761 to the loss of TPS activity, confirming this assumption.<sup>123</sup> The reciprocal substitutions in the  
762 *Mh*FPPS protein did not abolish IDS activity but caused the formation of GGPP instead of FPP  
763 due to the ability of the enzyme to accommodate an extended prenyl chain. The loss of aromatic  
764 residues upstream of the FARM is typical for long-chain *trans*-IDS enzymes ( $\geq$ C20), including  
765 insect GGPPSs.<sup>146,154</sup> By contrast, the bifunctional GPPS/TPS enzyme from *I. pini* maintains  
766 aromatic amino acids in this position because a presumably smaller-size cavity of this protein is  
767 sufficient to accommodate the short chain GPP product/substrate.<sup>118,119</sup>

768 [place Table 2 here]

769  
770 While residues upstream of the FARM seem to be critical for a proper positioning of the substrate  
771 in insect TPSs, Rebholz et al. hypothesize that the transition from IDS to TPS catalytic function  
772 largely depends on a change in the binding or position of the IPP substrate relative to DMAPP.<sup>151</sup>  
773 To test this hypothesis, a set of 20 IPP-binding residues positioned  $\leq$  5Å away from IPP were  
774 identified in the crystal structure of a *Homo sapiens* FPPS in complex with IPP by using the RING  
775 web server in combination with residue network interaction analysis in Cytoscape and structural  
776 analysis in Chimera.<sup>155-157</sup> The identified amino acids comprise basic residues that bind to the  
777 diphosphate moiety of IPP, a ring of residues encircling the isoprenyl tail, and residues that  
778 interact with both of these moieties. The residues are organized into six IPP binding motifs (IBMs)  
779 and were found to be conserved across IDS sequences from diverse organisms including  
780 animals, plants and fungi (Table 2, Fig. 16).

781 [place Fig. 16 here]

782 The residues that bind IPP orient the substrate and its prenyl tail in a way that allows condensation  
783 with the nascent carbocation formed from DMAPP. Modifications of these critical residues in IDS-  
784 type TPS proteins lead to alterations of the electrostatic nature of the IPP binding pocket (Fig.  
785 17). These changes may misalign IPP and DMAPP and disrupt their condensation, which likely  
786 allows competing TPS reactions of allylic substrates to occur. In agreement with this assumption,  
787 90% of the IBM residues were found to be modified in characterized insect TPS enzymes. For  
788 example, in the first IBM of hemipteran TPSs, the basic diphosphate binding residues are  
789 substituted with large aromatic residues (Table 2, Fig. 16), one of which is also conserved in its  
790 equivalent position in plant TPSs. Furthermore, several substitutions of residues interacting with  
791 the isoprenyl tail of IPP occur in the fourth IBM motif (Table 2, Fig. 16). Multiple residue  
792 substitutions are also present in the TPSs of Coleoptera and Lepidoptera; however, the  
793 substitution patterns are unique among the different taxonomic lineages, supporting independent  
794 events of TPS evolution.<sup>151</sup>

795 [place Fig. 17 here]

796

797 Lancaster et al. tested whether a re-introduction of the motif KKxR in the IBM1 of *Mh*TPS (Table  
798 2) through replacement of the corresponding SDAW sequence could convert the TPS enzyme  
799 into an IDS.<sup>123</sup> While no IDS activity was gained, the protein lost TPS activity, indicating an  
800 essential role of the SDAW residues in TPS function. A reciprocal substitution in the *Mh*FPPS  
801 also caused a loss of IDS activity, which further supports the critical role of the KKxR residues in  
802 IPP binding. To fully identify the residues that control the transition between IDS and TPS,  
803 O'Maille and co-workers currently perform combinatorial mutations paired with the identification  
804 of epistatic residue networks. A similar strategy was applied by Salmon et al. to determine residue  
805 substitutions in the transition from linear to cyclic TPSs in plants.<sup>158</sup> To probe evolutionary  
806 pathways of terpene cyclization, the authors of this study focused on the amino acid substitutions  
807 between a  $\beta$ -farnesene (**86**) synthase (BFS) from *Artemisia annua* and an amorphadiene  
808 synthase (ADS), which produces the bicyclic sesquiterpene amorpho-4,11-diene (**87**), a precursor  
809 of artemisinin (**88**) (Fig. 18). Structure-based combinatorial protein engineering (SCOPE)<sup>159</sup> was  
810 employed to construct two libraries of soluble and biochemically active mutant enzymes with ADS  
811 substitutions within 6 Angstroms of the BFS active site. Biochemical characterization resulted in  
812 the identification of multiple enzyme variants with the ability to generate cyclic terpene products.  
813 Chief among these products was the cyclic sesquiterpene alcohol, alpha-bisabolol (**89**).  
814 Quantitative determination of product-specific kinetic rates ( $k_{cat,i}$ ) for over 100 unique enzymes

815 allowed to train quantitative models for Michaelis-Menten enzymatic free energies. These models  
816 were used to construct a family of biophysical fitness landscapes describing enzyme evolution.<sup>160</sup>  
817 It was found that most mutations leading to the formation of alpha-bisabolol tended to have  
818 adverse effects on the overall magnitudes of product-specific reaction rates except for a  
819 previously identified critical gateway mutation (Y402L) that also unlocks the cyclization reaction.<sup>158</sup>  
820 Overall, this microevolutionary exploration of sequence space allowed the identification of key  
821 residues in terpene cyclization.

822 [place Fig. 18 here]

823  
824 To gain a broader understanding of the emergence of TPS proteins among all insects, Rebholz  
825 et al. predicted the presence of putative IDS-like TPSs in several insect lineages beyond the  
826 pentatomids.<sup>151</sup> To this end, UniProt sequences for polyprenyl synthetases (PFAM id PF00348)  
827 in the taxonomic class Insecta were screened for IDS and IDS-like TPS sequences based on  
828 distinct residue substitutions in the IBMs. The search identified more than 300 canonical IDS  
829 sequences and more than 125 putative TPSs, of which approximately 65% were found to be  
830 derived from FPPSs, nearly 23% from GGPPSs, with the rest derived from decaprenyl  
831 diphosphate (**6**) synthase (DPPS) like sequences (Fig. 19). Predicted TPS sequences were  
832 present in six insect orders: Blattodea, Hemiptera, Hymenoptera, Coleoptera, Lepidoptera, and  
833 Diptera.<sup>151</sup>

834 [place Fig. 19 here]

835  
836 The analysis revealed a family of FPPS-like TPS proteins specific to the lepidopteran genus  
837 *Papilio* (swallowtail butterflies), which may be responsible for the formation of linear mono- and  
838 sesquiterpenes that are released as defense compounds by *Papilio* larvae. Within the  
839 Lepidoptera, additional GGPPS-like sequences were found in the genome of the monarch  
840 butterfly *Danaus plexippus plexippus*, which emits terpene derivatives from male hairpencils.<sup>161</sup>  
841 These sequences are monophyletic to the characterized mono-TPSs in *H. melpomene*. Moreover,  
842 GGPPS-like TPSs were predicted within blattodean termites such as *Nasutitermes*  
843 *takasagoensis*. It is likely that genes of these expanded GGPPS-like families are involved in the  
844 production of the described termite terpene metabolites (see 2.1.3). Overall, the findings by  
845 Rebholz et al.<sup>151</sup> support the notion that insect TPSs originated from IDS genes and that this  
846 transition likely occurred via gene duplication and divergence through mutational drift and/or  
847 selection. Surprisingly, the study further indicates an independent emergence of TPS function not  
848 only within insect orders but also among the IDS subfamilies (FPPS, GGPPS, DPPS) of single

849 species within the same order. This suggests that parallel functionalization of IDS enzymes for  
850 volatile and specialized terpene biosynthesis is widespread in insects.

851

## 852 **5 Terpene synthases in corals**

853 Soft corals (octocorals) are known for their accumulation of diterpenes; however, the biosynthetic  
854 origin remained unknown until recently, when a new family of terpene cyclases was discovered  
855 in these organisms. Burkhardt et al.<sup>162</sup> and Scesa et al.<sup>163</sup> identified more than 15 TPSs from  
856 genomes and transcriptomes of several octocoral genera. Most of these enzymes were  
857 characterized as diterpene cyclases and a few others were found to function as sesquiterpene  
858 cyclases, which make the sesquiterpene hydrocarbons nepthene (**90**),  $\alpha$ -muuroladiene (**91**), and  
859 capnellene (**92**) (Fig. 20, Table 1). The enzymatic diterpene products such as cembrene A and C  
860 (**25**, **93**), elisabethatriene (**94**), klysimplexin R (**95**), and xeniaphyllene (**96**) (Fig. 20, Table 1) carry  
861 the scaffolds for large groups of coral-specific diterpenes and thus represent important semi-  
862 volatile precursors in generating the tremendous chemical diversity of diterpene-mediated  
863 defenses in corals.

864 [place Fig. 20 here]

865 Crystal structure analysis of a cembrene A synthase from the coral *Eleutherobia rubra* revealed  
866 that the protein carries the  $\alpha$ -helical fold typical of class I TPSs but has three additional helices  
867 (Fig. 21).<sup>162</sup> The analysis further showed the presence of conserved substrate-binding motifs and  
868 residues including the conserved aspartate rich motif, the NSE/DTE motif, and the previously  
869 identified arginine “pyrophosphate sensor” involved in carbocation formation (Fig. 21).<sup>164</sup> In  
870 contrast to plant TPS, the coral protein carries an RY motif that is conserved in microbial TPSs  
871 (e.g. Li et al.<sup>165</sup>). Its overall closer structural resemblance with microbial TPSs led the authors to  
872 speculate that an ancestral gene of the monophyletic coral TPS family might have been acquired  
873 by a common progenitor from microbial sources via horizontal gene transfer.

874 [place Fig. 21 here]

875 The evolution of these proteins predates the emergence of land plants. Interestingly, the coral  
876 TPS genes were found to cluster with P450, acyltransferase, and dehydrogenase genes among  
877 others, which presumably encode enzymes involved in secondary reactions of the TPS  
878 products.<sup>163</sup> These are the first biosynthetic gene clusters found in animals raising questions about  
879 their evolution and the potential presence of such clusters in other animals.

880

## 881 **6 Conclusions and Outlook**

882 In the past five to six years, substantial progress has been made in the identification of TPS genes  
883 in animals and in our understanding of how volatile terpenes and their derivatives are synthesized  
884 de novo in these organisms. The findings of IDS-type TPSs in insect genomes and the recent  
885 discovery of TPSs in soft corals may indicate that endogenous terpene biosynthetic pathways  
886 could be more common in animals than previously thought. The ways in which TPS gene functions  
887 have been recruited, whether through evolutionary transition from IDS precursors or by ancestral  
888 horizontal gene transfer from microbes (Fig. 21), appear to be as versatile as the diverse nature  
889 of terpene metabolites and their functions in chemical communication and defense. The finding  
890 that the formation of terpenes in corals is associated with assemblies of biosynthetic gene clusters  
891 similar to those found in microbes and plants may lead to the discovery of clusters in other animal  
892 genomes, especially when longer pathways have been established. To what extent these  
893 clustered genes are co-regulated in tissues or single cells in response to social or environmental  
894 signals will be of interest for future investigations. Another key question is whether terpene  
895 pheromones released from vertebrates such as amphibians and reptiles are the products of  
896 endogenous TPS enzymes or perhaps derived from microbial symbionts. Mining of high-quality  
897 genomes and gland-specific transcriptomes of these organisms should facilitate the elucidation  
898 of the biosynthetic origin of volatile terpene compounds. Finally, the discovery of terpene  
899 biosynthetic pathways in animals opens new possibilities for the biotechnological production of  
900 volatile terpenes such as species-specific pheromones in the development of alternative pest  
901 management strategies. For example, recent advances have been made in the metabolic  
902 engineering of fatty acid-derived moth sex pheromones in oilseed crops.<sup>166</sup> Similarly, the  
903 discovery of terpene biosynthetic gene clusters found in corals will undoubtedly accelerate efforts  
904 in the production of bioactive and pharmaceutically valuable compounds.

905

## 906 **7 Acknowledgements**

907 This work was supported by the National Science Foundation (MCB1920914, MCB1920922,  
908 MCB1920925) to AVM, PEO, and DT, respectively, and the US Department of Agriculture  
909 National Institute of Food and Agriculture (2016-67013-24759) to DT.

910 Credit to authors and licensing of the images used in modified form in Figures 6, 8, 9, 12-14 are  
911 listed in Supplementary Table 1.

912

## 913 **8 Author Contributions**

914 ZR participated in the conceptualization of the article, designed figures, wrote parts of the  
915 manuscript, and edited the manuscript. PEO designed figures and edited the manuscript. AVM

916 wrote parts of the manuscript and edited the manuscript. DT conceptualized the article, designed  
917 figures, and wrote the manuscript. All authors read and approved the final manuscript.

918

## 919 **9 Conflicts of interest**

920 There are no conflicts to declare.

921

## 922 **10 References**

- 923 1. R. R. Junker, J. Kuppler, L. Amo, J. D. Blande, R. M. Borges, N. M. van Dam, M. Dicke,  
924 S. Dötterl, B. K. Ehlers, F. Etl, J. Gershenzon, R. Glinwood, R. Gols, A. T. Groot, M. Heil,  
925 M. Hoffmeister, J. K. Holopainen, S. Jarau, L. John, A. Kessler, J. T. Knudsen, C. Kost,  
926 A. A. C. Larue-Kontic, S. D. Leonhardt, D. Lucas-Barbosa, C. J. Majetic, F. Menzel, A. L.  
927 Parachnowitsch, R. S. Pasquet, E. H. Poelman, R. A. Raguso, J. Ruther, F. P. Schiestl,  
928 T. Schmitt, D. Tholl, S. B. Unsicker, N. Verhulst, M. E. Visser, B. T. Weldegergis and T.  
929 G. Köllner, *New Phytol.*, 2018, **220**, 739-749.
- 930 2. D. W. Christianson, *Chemical Reviews*, 2017, **117**, 11570-11648.
- 931 3. M. C. Lemfack, B. O. Gohlke, S. M. T. Toguem, S. Preissner, B. Piechulla and R.  
932 Preissner, *Nucleic Acids Res.*, 2018, **46**, D1261-D1265.
- 933 4. J. T. Knudsen and J. Gershenzon, in *Biology of plant volatiles*, eds. E. Pichersky and N.  
934 Dudareva, CRC Press, Boca Raton, 2nd edn., 2020, ch. 4, 57-78.
- 935 5. E. Pichersky and N. Dudareva, *Biology of plant volatiles*, CRC Press, Boca Raton, 2nd  
936 edn., 2020.
- 937 6. M. Morita and E. W. Schmidt, *Nat. Prod. Rep.*, 2018, **35**, 357-378.
- 938 7. T. Reichstein, J. von Euw, J. A. Parsons and M. Rothschild, *Science*, 1968, **161**, 861-  
939 866.
- 940 8. D. Tholl, in *Biotechnology of isoprenoids*, eds. J. Schrader and J. Bohlmann, Springer  
941 International Publishing, 2015, vol. 148, pp. 63-106.
- 942 9. H. K. Lichtenthaler, in *Chloroplast: Basics and Applications*, eds. C. A. Rebeiz, C.  
943 Benning, H. J. Bohnert, H. Daniell, J. K. Hooper, H. K. Lichtenthaler, A. R. Portis and B.  
944 C. Tripathy, 2010, vol. 31, pp. 95-118.
- 945 10. M. Rohmer, *Nat. Prod. Rep.*, 1999, **16**, 565-574.
- 946 11. H. M. Miziorko, *Arch. Biochem. Biophys.*, 2011, **505**, 131-143.
- 947 12. H. Tao, L. Lauterbach, G. Bian, R. Chen, A. Hou, T. Mori, S. Cheng, B. Hu, L. Lu, X. Mu,  
948 M. Li, N. Adachi, M. Kawasaki, T. Moriya, T. Senda, X. Wang, Z. Deng, I. Abe, J. S.  
949 Dickschat and T. Liu, *Nature*, 2022, **606**, 414-419.
- 950 13. R. Saiki, A. Nagata, T. Kainou, H. Matsuda and M. Kawamukai, *The FEBS Journal*,  
951 2005, **272**, 5606-5622.
- 952 14. J. D. Rudolf and C. Y. Chang, *Nat. Prod. Rep.*, 2020, **37**, 425-463.
- 953 15. F. Chen, D. Tholl, J. Bohlmann and E. Pichersky, *Plant J.*, 2011, **66**, 212-229.
- 954 16. M. B. Quin, C. M. Flynn and C. Schmidt-Dannert, *Nat. Prod. Rep.*, 2014, **31**, 1449-1473.
- 955 17. J. S. Dickschat, *Nat. Prod. Rep.*, 2016, **33**, 87-110.
- 956 18. Q. Jia, G. Li, T. G. Köllner, J. Fu, X. Chen, W. Xiong, B. J. Crandall-Stotler, J. L.  
957 Bowman, D. J. Weston, Y. Zhang, L. Chen, Y. Xie, F.-W. Li, C. J. Rothfels, A. Larsson,  
958 S. W. Graham, D. W. Stevenson, G. K.-S. Wong, J. Gershenzon and F. Chen, *Proc.*  
959 *Natl. Acad. Sci. USA*, 2016, **113**, 12328-12333.
- 960 19. Z. H. Chen, Y. W. Guo and X. W. Li, *Nat. Prod. Rep.*, 2022, DOI: 10.1039/d2np00021k.
- 961 20. R. B. Pereira, P. B. Andrade and P. Valentão, *Mar. Drugs*, 2016, **14**.

- 962 21. C. Lyu, T. Chen, B. Qiang, N. Liu, H. Wang, L. Zhang and Z. Liu, *Nucleic Acids Res.*,  
963 2021, **49**, D509-d515.
- 964 22. W. Fenical, R. K. Okuda, M. M. Bandurraga, P. Culver and R. S. Jacobs, *Science*, 1981,  
965 **212**, 1512-1514.
- 966 23. H. Ben, Y. Sawall, A. Al-Sofyani and M. Wahl, *Aquat. Biol.*, 2015, **23**, 129–137.
- 967 24. G. Giordano, M. Carbone, M. L. Ciavatta, E. Silvano, M. Gavagnin, M. J. Garson, K. L.  
968 Cheney, I. W. Mudianta, G. F. Russo, G. Villani, L. Magliozzi, G. Polese, C. Zidorn, A.  
969 Cutignano, A. Fontana, M. T. Ghiselin and E. Mollo, *Proc. Natl. Acad. Sci. USA*, 2017,  
970 **114**, 3451-3456.
- 971 25. S.-M. Shen, Z.-Y. Zhang, L.-G. Yao, J.-R. Wang, Y.-W. Guo and X.-W. Li, *Chin. J.*  
972 *Chem.*, 2022, **40**, 235-246.
- 973 26. E. P. McCauley, I. C. Piña, A. D. Thompson, K. Bashir, M. Weinberg, S. L. Kurz and P.  
974 Crews, *J. Antibiot.*, 2020, **73**, 504-525.
- 975 27. L. D. Mydlarz, R. S. Jacobs, J. Boehnlein and R. G. Kerr, *Chem. Biol.*, 2003, **10**, 1051-  
976 1056.
- 977 28. C. Avila, *Mar. Drugs*, 2020, **18**, 162.
- 978 29. W. A. Shear, *Biochem. Syst. Ecol.*, 2015, **61**, 78-117.
- 979 30. K. Mori and Y. Takagi, *Tetrahedron Lett.*, 2000, **41**, 6623-6625.
- 980 31. J. Smolanoff, A. F. Kluge, J. Meinwald, A. McPhail, R. W. Miller, K. Hicks and T. Eisner,  
981 *Science*, 1975, **188**, 734-736.
- 982 32. M. F. Hassler, D. P. Harrison, T. H. Jones, C. H. Richart and R. A. Saporito, *J. Nat.*  
983 *Prod.*, 2020, **83**, 2764-2768.
- 984 33. Y. Kuwahara, Y. Ichiki, M. Morita, T. Tanabe and Y. Asano, *J. Chem. Ecol.*, 2015, **41**,  
985 15-21.
- 986 34. F. Jüttner and S. B. Watson, *Appl. Environ. Microbiol.*, 2007, **73**, 4395-4406.
- 987 35. O. Ekpa, J. W. Wheeler, J. C. Cokendolpher and R. M. Duffield, *Tetrahedron Lett.*, 1984,  
988 **25**, 1315-1318.
- 989 36. G. Raspotnig, V. Leutgeb, G. Krisper and H. J. Leis, *Exp. Appl. Acarol.*, 2011, **54**, 211-  
990 224.
- 991 37. A. C. Skelton, M. M. Cameron, J. A. Pickett and M. A. Birkett, *J. Med. Entomol.*, 2010,  
992 **47**, 798-804.
- 993 38. N. Shimizu, N. Mori and Y. Kuwahara, *Biosci. Biotechnol. Biochem.*, 2003, **67**, 1732-  
994 1736.
- 995 39. X. Dong, K. Chaisiri, D. Xia, S. D. Armstrong, Y. Fang, M. J. Donnelly, T. Kadowaki, J.  
996 W. McGarry, A. C. Darby and B. L. Makepeace, *Gigascience*, 2018, **7**.
- 997 40. M. Müller and G. Buchbauer, *Flavour Fragr. J.*, 2011, **26**, 357-377.
- 998 41. D. C. Weber, A. Khimian, M. C. Blassioli-Moraes and J. G. Millar, in *Invasive stink bugs*  
999 *and related species (Pentatomoidea): Biology, higher systematics, semiochemistry, and*  
1000 *management*, ed. J. E. McPherson, CRC Press, Boca Raton, Florida, 2018, pp. 677-  
1001 725.
- 1002 42. J. A. Pickett, R. K. Allemann and M. A. Birkett, *Nat. Prod. Rep.*, 2013, **30**, 1277-1283.
- 1003 43. C. I. Keeling, C. Tittiger, M. MacLean and G. J. Blomquist, in *Insect pheromone*  
1004 *biochemistry and molecular biology*, eds. G. J. Blomquist and R. G. Vogt, Academic  
1005 Press, San Diego, 2021, pp. 123-154.
- 1006 44. J. Havlíčková, K. Dolejšová, M. Tichý, V. Vrkoslav, B. Kalinová, P. Kyjaková and R.  
1007 Hanus, *Z. Naturforsch. C*, 2019, **74**, 251-264.
- 1008 45. T. A. K. Prescott, J. Bramham, O. Zompro and S. K. Maciver, *Biochem. Syst. Ecol.*,  
1009 2009, **37**, 759-760.
- 1010 46. R. M. Smith, J. J. Brophy, G. W. K. Cavill and N. W. Davies, *J. Chem. Ecol.*, 1979, **5**,  
1011 727-735.

- 1012 47. D. B. Weibel, N. J. Oldham, B. Feld, G. Glombitza, K. Dettner and W. Boland, *Insect*  
1013 *Biochem. Mol. Biol.*, 2001, **31**, 583-591.
- 1014 48. C. J. Persoons, P. E. J. Verwiël, F. J. Ritter, E. Talman, P. J. F. Nooijen and W. J.  
1015 Nooijen, *Tetrahedron Lett.*, 1976, **24**, 2055-2058.
- 1016 49. E. Talman, P. E. J. Verwiël, F. J. Ritter and C. J. Persoons, *Tetrahedron Lett.*, 1978, **17**,  
1017 227-235.
- 1018 50. H. Takegawa and S. Takahashi, *Appl. Entomol. Zoolog.*, 1989, **24**, 435-440.
- 1019 51. S. Takahashi, K. Watanabe, S. Saito and Y. Nomura, *Appl. Entomol. Zoolog.*, 1995, **30**,  
1020 357-360.
- 1021 52. T. Hirukawa, T. Shudo and T. Kato, *J. Chem. Soc.-Perkin Trans. 1*, 1993, **2**, 217-225.
- 1022 53. J. C. Braekman, D. Dalozé, A. Dupont, J. M. Pasteels and R. Ottinger, *Bulletin des*  
1023 *Sociétés Chimiques Belges*, 1984, **93**, 291-298.
- 1024 54. G. D. Prestwich, B. A. Solheim, J. Clardy, F. G. Pilkiewicz, I. Miura, S. P. Tanis and K.  
1025 Nakanishi, *J. Am. Chem. Soc.*, 1977, **99**, 8082-8083.
- 1026 55. J. Sobotnik, A. Jirosova and R. Hanus, *J. Insect Physiol.*, 2010, **56**, 1012-1021.
- 1027 56. G. D. Prestwich, *Annu. Rev. Entomol.*, 1984, **29**, 201-232.
- 1028 57. C. Everaerts, O. Bonnard, J. M. Pasteels, Y. Roisin and W. A. König, *Experientia*, 1990,  
1029 **46**, 227-230.
- 1030 58. M. R. Tarver, E. A. Schmelz, J. R. Rocca and M. E. Scharf, *J. Chem. Ecol.*, 2009, **35**,  
1031 256-264.
- 1032 59. D. Sillam-Dusses, B. Kalinova, P. Jiros, A. Brezinova, J. Cvacka, R. Hanus, J. Sobotnik,  
1033 C. Bordereau and I. Valterova, *J. Insect Physiol.*, 2009, **55**, 751-757.
- 1034 60. K. Haga, T. Suzuki, S. Kodama and Y. Kuwahara, *Appl. Entomol. Zoolog.*, 1990, **25**,  
1035 138-139.
- 1036 61. T. Suzuki, K. Haga, W. S. Leal, S. Kodama and Y. Kuwahara, *Appl. Entomol. Zool.*,  
1037 1989, **24**, 222-228.
- 1038 62. J. G. C. Hamilton, D. R. Hall and W. D. J. Kirk, *J. Chem. Ecol.*, 2005, **31**, 1369-1379.
- 1039 63. S. Niassy, A. Tamiru, J. G. C. Hamilton, W. D. J. Kirk, R. Mumm, C. Sims, W. J. de  
1040 Kogel, S. Ekesi, N. K. Maniania, K. Bandi, F. Mitchell and S. Subramanian, *J. Chem.*  
1041 *Ecol.*, 2019, **45**, 348-355.
- 1042 64. Z. A. Demissie, L. A. E. Erland, M. R. Rheault and S. S. Mahmoud, *J. Biol. Chem.*, 2013,  
1043 **288**, 6333-6341.
- 1044 65. J. Tabata and R. T. Ichiki, *J. Chem. Ecol.*, 2016, **42**, 1193-1200.
- 1045 66. J. Einhorn, A. Guerrero, P. H. Ducrot, F. D. Boyer, M. Gieselmann and W. Roelofs, *Proc.*  
1046 *Natl. Acad. Sci. U S A*, 1998, **95**, 9867-9872.
- 1047 67. A. M. El-Sayed, C. R. Unelius, A. Twidle, V. Mitchell, L.-A. Manning, L. Cole, D. M.  
1048 Suckling, M. F. Flores, T. Zaviezo and J. Bergmann, *Tetrahedron Lett.*, 2010, **51**, 1075-  
1049 1078.
- 1050 68. Z. Rebholz, J. Lancaster, H. Larose, A. Khimian, K. Luck, M. E. Sparks, K. L. Gendreau,  
1051 L. Shewade, T. G. Köllner, D. C. Weber, D. E. Gundersen-Rindal, P. O'Maille, A. V.  
1052 Morozov and D. Tholl, *Insect Biochem. Mol. Biol.*, 2023, **152**, 103879.
- 1053 69. M. Inwood and P. Morgan, *Myrmecological News*, 2008, **11**, 79-90.
- 1054 70. R. K. Vander Meer, F. Alvarez and C. S. Lofgren, *J. Chem. Ecol.*, 1988, **14**, 825-838.
- 1055 71. S. J. Keegans, J. Billen, E. D. Morgan and O. A. Gokcen, *J. Chem. Ecol.*, 1993, **19**,  
1056 2705-2719.
- 1057 72. J. A. Pickett, I. H. Williams, M. C. Smith and A. P. Martin, *J. Chem. Ecol.*, 1981, **7**, 543-  
1058 554.
- 1059 73. R. Kubo and M. Ono, *Entomol. Sci.*, 2010, **13**, 167-173.
- 1060 74. M. Kaib and H. Dittebrand, *Chemoecology*, 1990, **1**, 3-11.
- 1061 75. B. Fassotte, C. Fischer, D. Durieux, G. Lognay, E. Haubruge, F. Francis and F. J.  
1062 Verheggen, *Plos One*, 2014, **9**, e115011.

- 1063 76. R. J. Bartelt, A. A. Cossé, B. W. Zilkowski, D. Weisleder and F. A. Momany, *J. Chem. Ecol.*, 2001, **27**, 2397-2423.  
1064
- 1065 77. N. Mitlin and P. A. Hedin, *J. Insect Physiol.*, 1974, **20**, 1825-1831.  
1066 78. J. E. Carrel, M. H. McCairel, A. J. Slagle, J. P. Doom, J. Brill and J. P. McCormick, *Experientia*, 1993, **49**, 171-174.  
1067
- 1068 79. K. Honda, *J. Chem. Ecol.*, 1981, **7**, 1089-1113.  
1069 80. H. Omura, K. Honda and P. Feeny, *J. Chem. Ecol.*, 2006, **32**, 1999-2012.  
1070 81. S. Schulz, C. Estrada, S. Yildizhan, M. Boppré and L. E. Gilbert, *J. Chem. Ecol.*, 2008, **34**, 82-93.  
1071
- 1072 82. R. Baker, R. H. Herbert and G. G. Grant, *J. Chem. Soc.-Chem. Commun.*, 1985, **12**, 824-825.  
1073
- 1074 83. P. E. A. Teal, J. A. Meredith and Y. Gomez-Simuta, *Arch. Insect Biochem. Physiol.*, 1999, **42**, 225-232.  
1075
- 1076 84. C. N. Spiegel, P. Jeanbourquin, P. M. Guerin, A. M. Hooper, S. Claude, R. Tabacchi, S. Sano and K. Mori, *J. Insect Physiol.*, 2005, **51**, 1366-1375.  
1077
- 1078 85. M. Menke, K. Melnik, P. S. Peram, I. Starnberger, W. Hödl, M. Vences and S. Schulz, *Eur. J. Org. Chem.*, 2018, **2018**, 2651-2656.  
1079
- 1080 86. M. Menke, P. S. Peram, I. Starnberger, W. Hödl, G. F. Jongsma, D. C. Blackburn, M. O. Rödel, M. Vences and S. Schulz, *Beilstein J. Org. Chem.*, 2016, **12**, 2731-2738.  
1081
- 1082 87. I. Starnberger, D. Poth, P. S. Peram, S. Schulz, M. Vences, J. Knudsen, M. F. Barej, M. O. Rodel, M. Walzl and W. Hodl, *Biol. J. Linn. Soc.*, 2013, **110**, 828-838.  
1083
- 1084 88. C. Nowack, P. S. Peram, S. Wenzel, A. Rakotoarison, F. Glaw, D. Poth, S. Schulz and M. Vences, *J. Zool.*, 2017, **303**, 72-81.  
1085
- 1086 89. S. Yildizhan, J. van Loon, A. Sramkova, M. Ayasse, C. Arsene, C. ten Broeke and S. Schulz, *ChemBioChem*, 2009, **10**, 1666-1677.  
1087
- 1088 90. T. Chuman, J. Sivinski, R. R. Heath, C. O. Calkins, J. H. Tumlinson, M. A. Battiste, R. L. Wydra, L. Strekowski and J. L. Nation, *Tetrahedron Lett.*, 1988, **29**, 6561-6563.  
1089
- 1090 91. D. Vanderwel, B. Johnston and A. C. Oehlschlager, *Insect Biochem. Mol. Biol.*, 1992, **22**, 875-883.  
1091
- 1092 92. B. P. C. Smith, Y. Hayasaka, M. J. Tyler and B. D. Williams, *Aust. J. Zool.*, 2004, **52**, 521-530.  
1093
- 1094 93. D. L. Mattern, W. D. Scott, C. A. McDaniel, P. J. Weldon and D. E. Graves, *J. Nat. Prod.*, 1997, **60**, 828-831.  
1095
- 1096 94. J. W. Avery, A. Shafagati, A. Turner, J. W. Wheeler and P. J. Weldon, *Biochem. Syst. Ecol.*, 1993, **21**, 533-534.  
1097
- 1098 95. S. Schulz, K. Krückert and P. J. Weldon, *J. Nat. Prod.*, 2003, **66**, 34-38.  
1099 96. S. García-Rubio, A. B. Attygalle, P. J. Weldon and J. Meinwald, *J. Chem. Ecol.*, 2002, **28**, 769-781.  
1100
- 1101 97. B. V. Burger, M. Leroux, H. S. C. Spies, V. Truter and R. C. Bigalke, *Z. Naturforsch. C Bio. Sci.*, 1981, **36**, 340-343.  
1102
- 1103 98. J. W. Wheeler, L. E. Rasmussen, F. Ayorinde, I. O. Buss and G. L. Smuts, *J. Chem. Ecol.*, 1982, **8**, 821-835.  
1104
- 1105 99. T. E. Goodwin, F. D. Brown, R. W. Counts, N. C. Dowdy, P. L. Fraley, R. A. Hughes, D. Z. Liu, C. D. Mashburn, J. D. Rankin, R. S. Roberson, K. D. Wooley, E. L. Rasmussen, S. W. Riddle, H. S. Riddle and S. Schulz, *J. Nat. Prod.*, 2002, **65**, 1319-1322.  
1106  
1107
- 1108 100. C. G. Faulkes, J. S. Elmore, D. A. Baines, B. Fenton, N. B. Simmons and E. L. Clare, *PeerJ*, 2019, **7**, e7734.  
1109
- 1110 101. E. J. Elwell, D. Walker and S. Vaglio, *Animals*, 2021, **11**, 2091.  
1111 102. M. Salamon and N. W. Davies, *J. Chem. Ecol.*, 1998, **24**, 1659-1676.  
1112 103. D. J. King, R. M. Gleadow and I. E. Woodrow, *New Phytol.*, 2006, **172**, 440-451.  
1113 104. C. S. Mathela and N. Joshi, *Nat. Prod. Commun.*, 2008, **3**, 945-950.

- 1114 105. M. A. Birkett, A. Hassanali, S. Hoglund, J. Pettersson and J. A. Pickett, *Phytochemistry*,  
1115 2011, **72**, 109-114.
- 1116 106. B. R. Lichman, G. T. Godden, J. P. Hamilton, L. Palmer, M. O. Kamileen, D. Zhao, B.  
1117 Vaillancourt, J. C. Wood, M. Sun, T. J. Kinser, L. K. Henry, C. Rodriguez-Lopez, N.  
1118 Dudareva, D. E. Soltis, P. S. Soltis, C. R. Buell and S. E. O'Connor, *Sci. Adv.*, 2020, **6**,  
1119 eaba0721.
- 1120 107. K. Miettinen, L. M. Dong, N. Navrot, T. Schneider, V. Burlat, J. Pollier, L. Woittiez, S. van  
1121 der Krol, R. Lugan, T. Ilc, R. Verpoorte, K. M. Oksman-Caldentey, E. Martinoia, H.  
1122 Bouwmeester, A. Goossens, J. Memelink and D. Werck-Reichhart, *Nat. Commun.*,  
1123 2014, **5**, 4175.
- 1124 108. F. Geu-Flores, N. H. Sherden, V. Courdavault, V. Burlat, W. S. Glenn, C. Wu, E. Nims,  
1125 Y. Cui and S. E. O'Connor, *Nature*, 2012, **492**, 138-142.
- 1126 109. R. R. Bodemann, P. Rahfeld, M. Stock, M. Kunert, N. Wielsch, M. Groth, S. Frick, W.  
1127 Boland and A. Burse, *Proc. R. Soc. B-Biol. Sci.*, 2012, **279**, 4126-4134.
- 1128 110. S. Frick, R. Nagel, A. Schmidt, R. R. Bodemann, P. Rahfeld, G. Pauls, W. Brandt, J.  
1129 Gershenzon, W. Boland and A. Burse, *Proc. Natl. Acad. Sci. USA*, 2013, **110**, 56-61.
- 1130 111. P. Rahfeld, W. Haeger, R. Kirsch, G. Pauls, T. Becker, E. Schulze, N. Wielsch, D. Wang,  
1131 M. Groth, W. Brandt, W. Boland and A. Burse, *Insect Biochem. Mol. Biol.*, 2015, **58**, 28-  
1132 38.
- 1133 112. P. Rahfeld, R. Kirsch, S. Kugel, N. Wielsch, M. Stock, M. Groth, W. Boland and A.  
1134 Burse, *Proc. R. Soc. B-Biol. Sci.*, 2014, **281**, 20140842.
- 1135 113. N. Fu, Z. L. Yang, Y. Pauchet, C. Paetz, W. Brandt, W. Boland and A. Burse, *Insect*  
1136 *Biochem. Mol. Biol.*, 2019, **113**, 103212.
- 1137 114. T. G. Köllner, A. David, K. Luck, F. Beran, G. Kunert, J. J. Zhou, L. Caputi and S. E.  
1138 O'Connor, *Proc. Natl. Acad. Sci. U S A*, 2022, **119**, e2211254119.
- 1139 115. T. Laursen, J. Borch, C. Knudsen, K. Bavishi, F. Torta, H. J. Martens, D. Silvestro, N. S.  
1140 Hatzakis, M. R. Wenk, T. R. Dafforn, C. E. Olsen, M. S. Motawia, B. Hamberger, B. L.  
1141 Moller and J. E. Bassard, *Science*, 2016, **354**, 890-893.
- 1142 116. M. Sugumaran, K. Nellaiappan, C. Amaratunga, S. Cardinale and T. Scott, *Arch.*  
1143 *Biochem. Biophys.*, 2000, **378**, 393-403.
- 1144 117. S. J. Seybold, D. R. Quilici, J. A. Tillman, D. Vanderwel, D. L. Wood and G. J. Blomquist,  
1145 *Proc. Natl. Acad. Sci. U S A*, 1995, **92**, 8393-8397.
- 1146 118. A. B. Gilg, J. C. Bearfield, C. Tittiger, W. H. Welch and G. J. Blomquist, *Proc. Natl. Acad.*  
1147 *Sci. USA*, 2005, **102**, 9760-9765.
- 1148 119. A. B. Gilg, C. Tittiger and G. J. Blomquist, *Naturwissenschaften*, 2009, **96**, 731-735.
- 1149 120. F. Beran, P. Rahfeld, K. Luck, R. Nagel, H. Vogel, N. Wielsch, S. Irmisch, S. Ramasamy,  
1150 J. Gershenzon, D. G. Heckel and T. G. Köllner, *Proc. Natl. Acad. Sci. USA*, 2016, **113**,  
1151 2922-2927.
- 1152 121. A. Čokl, A. M. Dias, M. C. B. Moraes, M. Borges and R. A. Laumann, *J. Insect Behav.*,  
1153 2017, **30**, 741-758.
- 1154 122. D. Tholl, in *Insect pheromone biochemistry and molecular biology*, eds. G. J. Blomquist  
1155 and R. G. Vogt, Academic Press, San Diego, 2nd edn., 2021, pp. 269-281.
- 1156 123. J. Lancaster, A. Khimian, S. Young, B. Lehner, K. Luck, A. Wallingford, S. K. B. Ghosh,  
1157 P. Zerbe, A. Muchlinski, P. E. Marek, M. E. Sparks, J. G. Tokuhsa, C. Tittiger, T. G.  
1158 Köllner, D. C. Weber, D. E. Gundersen-Rindal, T. P. Kuhar and D. Tholl, *Proc. Natl.*  
1159 *Acad. Sci. USA*, 2018, **115**, E8634-E8641.
- 1160 124. A. Khimian, S. Shirali, K. E. Vermillion, M. A. Siegler, F. Guzman, K. Chauhan, J. R.  
1161 Aldrich and D. C. Weber, *J. Chem. Ecol.*, 2014, **40**, 1260-1268.
- 1162 125. M. E. Sparks, J. H. Rhoades, D. R. Nelson, D. Kuhar, J. Lancaster, B. Lehner, D. Tholl,  
1163 D. C. Weber and D. E. Gundersen-Rindal, *Insects*, 2017, **8**, 55.
- 1164 126. L. K. Sy and G. D. Brown, *Phytochemistry*, 1997, **45**, 537-544.

- 1165 127. A. Khimian, A. J. Zhang, D. C. Weber, H. Y. Ho, J. R. Aldrich, K. E. Vermillion, M. A.  
1166 Sieglar, S. Shirali, F. Guzman and T. C. Leskey, *J. Nat. Prod.*, 2014, **77**, 1708-1717.
- 1167 128. M. E. Sparks, R. Bansal, J. B. Benoit, M. B. Blackburn, H. Chao, M. Chen, S. Cheng, C.  
1168 Childers, H. Dinh, H. V. Doddapaneni, S. Dugan, E. N. Elpidina, D. W. Farrow, M.  
1169 Friedrich, R. A. Gibbs, B. Hall, Y. Han, R. W. Hardy, C. J. Holmes, D. S. T. Hughes, P.  
1170 Ioannidis, A. M. Cheatle Jarvela, J. S. Johnston, J. W. Jones, B. A. Kronmiller, F. Kung,  
1171 S. L. Lee, A. G. Martynov, P. Masterson, F. Maumus, M. Munoz-Torres, S. C. Murali, T.  
1172 D. Murphy, D. M. Muzny, D. R. Nelson, B. Oppert, K. A. Panfilio, D. P. Paula, L. Pick, M.  
1173 F. Poelchau, J. Qu, K. Reding, J. H. Rhoades, A. Rhodes, S. Richards, R. Richter, H. M.  
1174 Robertson, A. J. Rosendale, Z. J. Tu, A. S. Velamuri, R. M. Waterhouse, M. T.  
1175 Weirauch, J. T. Wells, J. H. Werren, K. C. Worley, E. M. Zdobnov and D. E. Gundersen-  
1176 Rindal, *BMC Genomics*, 2020, **21**.
- 1177 129. M. E. Sparks, K. S. Shelby, D. Kuhar and D. E. Gundersen-Rindal, *Plos One*, 2014, **9**.
- 1178 130. M. Borges, M. Birkett, J. R. Aldrich, J. E. Oliver, M. Chiba, Y. Murata, R. A. Laumann, J.  
1179 A. Barrigossi, J. A. Pickett and M. C. B. Moraes, *J. Chem. Ecol.*, 2006, **32**, 2749-2761.
- 1180 131. M. W. M. de Oliveira, M. Borges, C. K. Z. Andrade, R. A. Laumann, J. A. F. Barrigossi  
1181 and M. C. Blassioli-Moraes, *J. Agric. Food Chem.*, 2013, **61**, 7777-7785.
- 1182 132. A. A. C. Moliterno, D. J. De Melo and P. H. G. Zarbin, *J. Chem. Ecol.*, 2021, **47**, 1-9.
- 1183 133. J. R. Aldrich, J. E. Oliver, W. R. Lusby, J. P. Kochansky and J. A. Lockwood, *J. Exp.*  
1184 *Zool.*, 1987, **244**, 171-175.
- 1185 134. V. E. Harris and J. W. Todd, *Entomol. Exp. Appl.*, 1980, **27**, 117-126.
- 1186 135. J. Lancaster, B. Lehner, A. Khimian, A. Muchlinski, K. Luck, T. G. Köllner, D. C. Weber,  
1187 D. E. Gundersen-Rindal and D. Tholl, *J. Chem. Ecol.*, 2019, **45**, 187-197.
- 1188 136. B. W. Cribb, K. N. Siriwardana and G. H. Walter, *J. Morphol.*, 2006, **267**, 831-840.
- 1189 137. S. Kumar, C. Kempinski, X. Zhuang, A. Norris, S. Mafu, J. C. Zi, S. A. Bell, S. E. Nybo,  
1190 S. E. Kinison, Z. D. Jiang, S. Goklany, K. B. Linscott, X. L. Chen, Q. D. Jia, S. D. Brown,  
1191 J. L. Bowman, P. C. Babbitt, R. J. Peters, F. Chen and J. Chappell, *Plant Cell*, 2016, **28**,  
1192 2632-2650.
- 1193 138. G. L. Li, T. G. Köllner, Y. B. Yin, Y. F. Jiang, H. Chen, Y. Xu, J. Gershenzon, E.  
1194 Pichersky and F. Chen, *Proc. Natl. Acad. Sci. USA*, 2012, **109**, 14711-14715.
- 1195 139. A. Muchlinski, X. Chen, J. T. Lovell, T. G. Köllner, K. A. Pelot, P. Zerbe, M. Ruggiero, L.  
1196 Callaway, S. Laliberte, F. Chen and D. Tholl, *Front. Plant Sci.*, 2019, **10**, 1144.
- 1197 140. J. R. Aldrich, W. R. Lusby and J. P. Kochansky, *Experientia*, 1986, **42**, 583-585.
- 1198 141. J. P. Farine, O. Bonnard, R. Brossut and J. L. Le Quere, *J. Chem. Ecol.*, 1992, **18**, 1673-  
1199 1682.
- 1200 142. B. S. Krall, B. W. Zilkowski, S. L. Kight, R. J. Bartelt and D. W. Whitman, *J. Chem. Ecol.*,  
1201 1997, **23**, 1951-1962.
- 1202 143. J. R. Aldrich, S. P. Carroll, J. E. Oliver, W. R. Lusby, A. A. Rudmann and R. M. Waters,  
1203 *Biochem. Syst. Ecol.*, 1990, **18**, 369-376.
- 1204 144. E. D. Morgan, *Biosynthesis in insects*, RSC Publishing, Cambridge, UK, Advanced  
1205 Edition edn., 2010.
- 1206 145. S. B. Rivera, B. D. Swedlund, G. J. King, R. N. Bell, C. E. Hussey, D. M. Shattuck-  
1207 Eidens, W. M. Wrobel, G. D. Peiser and C. D. Poulter, *Proc. Natl. Acad. Sci. USA*, 2001,  
1208 **98**, 4373-4378.
- 1209 146. K. Darragh, A. Orteu, D. Black, K. J. R. P. Byers, D. Szczerbowski, I. A. Warren, P.  
1210 Rastas, A. Pinharanda, J. W. Davey, S. Fernanda Garza, D. Abondano Almeida, R. M.  
1211 Merrill, W. O. McMillan, S. Schulz and C. D. Jiggins, *PLoS Biol.*, 2021, **19**, e3001022.
- 1212 147. S. Schulz, C. Estrada, S. Yildizhan, M. Boppre and L. E. Gilbert, *J. Chem. Ecol.*, 2008,  
1213 **34**, 82-93.

- 1214 148. K. Darragh, G. Montejó-Kovacevich, K. M. Kozak, C. R. Morrison, C. M. E. Figueiredo, J.  
1215 S. Ready, C. Salazar, M. Linares, K. Byers, R. M. Merrill, W. O. McMillan, S. Schulz and  
1216 C. D. Jiggins, *Ecol. Evol.*, 2020, **10**, 3895-3918.
- 1217 149. M. Hojo, S. Shigenobu, K. Maekawa, T. Miura and G. Tokuda, *Insect Biochem. Mol.*  
1218 *Biol.*, 2019, **111**.
- 1219 150. Y. Zhou, X. Liu and Z. Yang, *Biomolecules*, 2019, **9**, 808.
- 1220 151. Z. Rebholz, L. Shewade, K. Kaler, H. Larose, F. Schubot, D. Tholl, A. V. Morozov and P.  
1221 E. O'Maille, **in revision**.
- 1222 152. C. M. Starks, K. W. Back, J. Chappell and J. P. Noel, *Science*, 1997, **277**, 1815-1820.
- 1223 153. C. A. Lesburg, G. Zhai, D. E. Cane and D. W. Christianson, *Science*, 1997, **277**, 1820-  
1224 1824.
- 1225 154. F. H. Wallrapp, J.-J. Pan, G. Ramamoorthy, D. E. Almonacid, B. S. Hillerich, R. Seidel,  
1226 Y. Patskovsky, P. C. Babbitt, S. C. Almo, M. P. Jacobson and C. D. Poulter, *Proc. Natl.*  
1227 *Acad. Sci. USA*, 2013, **110**, E1196-E1202.
- 1228 155. E. F. Pettersen, T. D. Goddard, C. C. Huang, G. S. Couch, D. M. Greenblatt, E. C. Meng  
1229 and T. E. Ferrin, *J. Comput. Chem.*, 2004, **25**, 1605-1612.
- 1230 156. D. Piovesan, G. Minervini and S. C. Tosatto, *Nucleic Acids Res.*, 2016, **44**, W367-374.
- 1231 157. P. Shannon, A. Markiel, O. Ozier, N. S. Baliga, J. T. Wang, D. Ramage, N. Amin, B.  
1232 Schwikowski and T. Ideker, *Genome Res.*, 2003, **13**, 2498-2504.
- 1233 158. M. Salmon, C. Laurendon, M. Vardakou, J. Cheema, M. Defernez, S. Green, J. A.  
1234 Faraldos and P. E. O'Maille, *Nat. Commun.*, 2015, **6**, 6143.
- 1235 159. M. Dokarry, C. Laurendon and P. E. O'Maille, in *Natural product biosynthesis by*  
1236 *microorganisms and plants, part A*, ed. D. A. Hopwood, 2012, vol. 515, pp. 21-42.
- 1237 160. A. Ballal, C. Laurendon, M. Salmon, M. Vardakou, J. Cheema, M. Defernez, P. E.  
1238 O'Maille and A. V. Morozov, *Mol. Biol. Evol.*, 2020, **37**, 1907-1924.
- 1239 161. J. Meinwald, A. M. Chalmers, T. E. Pliske and T. Eisner, *Tetrahedron Lett.*, 1968, **9**,  
1240 4893-4896.
- 1241 162. I. Burkhardt, T. De Rond, P. Y.-T. Chen and B. S. Moore, *Nat. Chem. Biol.*, 2022, **18**,  
1242 664-669.
- 1243 163. P. D. Scesa, Z. Lin and E. W. Schmidt, *Nat. Chem. Biol.*, 2022, **18**, 659-663.
- 1244 164. P. Baer, P. Rabe, K. Fischer, C. A. Citron, T. A. Klapschinski, M. Groll and J. S.  
1245 Dickschat, *Angew. Chem.-Int. Edit.*, 2014, **53**, 7652-7656.
- 1246 165. R. Li, W. K. Chou, J. A. Himmelberger, K. M. Litwin, G. G. Harris, D. E. Cane and D. W.  
1247 Christianson, *Biochemistry*, 2014, **53**, 1155-1168.
- 1248 166. Y. H. Xia, H. L. Wang, B. J. Ding, G. P. Svensson, C. Jarl-Sunesson, E. B. Cahoon, P.  
1249 Hofvander and C. Löfstedt, *J. Chem. Ecol.*, 2021, **47**, 950-967.
- 1250 167. B. Misof, S. Liu, K. Meusemann, R. S. Peters, A. Donath, C. Mayer, P. B. Frandsen, J.  
1251 Ware, T. Flouri, R. G. Beutel, O. Niehuis, M. Petersen, F. Izquierdo-Carrasco, T.  
1252 Wappler, J. Rust, A. J. Aberer, U. Aspöck, H. Aspöck, D. Bartel, A. Blanke, S. Berger, A.  
1253 Böhm, T. R. Buckley, B. Calcott, J. Chen, F. Friedrich, M. Fukui, M. Fujita, C. Greve, P.  
1254 Grobe, S. Gu, Y. Huang, L. S. Jermin, A. Y. Kawahara, L. Krogmann, M. Kubiak, R.  
1255 Lanfear, H. Letsch, Y. Li, Z. Li, J. Li, H. Lu, R. Machida, Y. Mashimo, P. Kapli, D. D.  
1256 McKenna, G. Meng, Y. Nakagaki, J. L. Navarrete-Heredia, M. Ott, Y. Ou, G. Pass, L.  
1257 Podsiadlowski, H. Pohl, B. M. von Reumont, K. Schütte, K. Sekiya, S. Shimizu, A.  
1258 Slipinski, A. Stamatakis, W. Song, X. Su, N. U. Szucsich, M. Tan, X. Tan, M. Tang, J.  
1259 Tang, G. Timelthaler, S. Tomizuka, M. Trautwein, X. Tong, T. Uchifune, M. G. Walz, B.  
1260 M. Wiegmann, J. Wilbrandt, B. Wipfler, T. K. Wong, Q. Wu, G. Wu, Y. Xie, S. Yang, Q.  
1261 Yang, D. K. Yeates, K. Yoshizawa, Q. Zhang, R. Zhang, W. Zhang, Y. Zhang, J. Zhao,  
1262 C. Zhou, L. Zhou, T. Ziesmann, S. Zou, Y. Li, X. Xu, Y. Zhang, H. Yang, J. Wang, J.  
1263 Wang, K. M. Kjer and X. Zhou, *Science*, 2014, **346**, 763-767.
- 1264 168. G. Giribet, *Org. Divers. Evol.*, 2016, **16**, 419-426.

1265

1266 **Tables**

1267 **Table 1** Functionally characterized terpene biosynthetic genes/proteins and their products in  
1268 insects and soft corals. Insect species are grouped by orders. Sesquitrps, sesquiterpenes; \*  
1269 indicates that no accession numbers were listed based on cDNA amplification and gene  
1270 sequences were synthesized. Nucleotide sequences are listed in the respective publications.

1271

1272

1273 **Table 2** Structure-based sequence alignment of IDS IBM motifs and substitutions in IDS-type  
1274 TPS proteins in hemipteran species.

1275 Extract of IBM motifs from a structure-based sequence alignment of characterized hemipteran  
1276 IDS and TPS proteins. Residue positions that are  $\leq 5 \text{ \AA}$  away from IPP and highly conserved are  
1277 colored according to their interaction with the diphosphate moiety (dark blue) or the isoprenyl tail  
1278 (light blue), respectively. TPS residue substitutions of the diphosphate binding residues and  
1279 prenyl tail binding residues are marked in purple and green, respectively. Substitutions that  
1280 deviate from the IBM regular expressions in other insects and animals are shaded in grey.  
1281 Unshaded residues (white) correspond to variable residues and positions outside of the  
1282 consensus IPP binding residues.

1283

1284

1285 **Figures**

1286 **Figure 1** Enzymatic steps in the biosynthesis of volatile or semi-volatile monoterpenes,  
1287 sesquiterpenes, and diterpenes. IPP, isopentenyl diphosphate; DMAPP, dimethylallyl  
1288 diphosphate; GPP, geranyl diphosphate; FPP, farnesyl diphosphate; GGPP, geranylgeranyl  
1289 diphosphate; GPPS, geranyl diphosphate synthase; FPPS, farnesyl diphosphate synthase;  
1290 GGPPS, geranylgeranyl diphosphate synthase; IDS, isoprenyl diphosphate synthase; TPS,  
1291 terpene synthase; MEP, methylerythritol phosphate; MVA, mevalonic acid.

1292

1293 **Figure 2** Examples of sesquiterpenes and diterpenes in marine gastropods and octocorals.

1294

1295 **Figure 3** Examples of volatile terpene semiochemicals and derivatives in millipedes and  
1296 arachnids. No stereochemical configuration is shown in case the stereochemistry was not  
1297 determined.

1298

1299 **Figure 4** Examples of volatile terpene semiochemicals in stick insects (Phasmatodea), roaches  
1300 and termites (Blattodea), thrips (Thysanoptera), and hemipteran (true bug) insects.

1301  
1302 **Figure 5** Examples of volatile terpene semiochemicals in the orders Hymenoptera (including ants  
1303 and bees), Coleoptera (beetles), Lepidoptera (including butterflies), and Diptera (flies). No  
1304 stereochemical configuration is shown in case the stereochemistry was not determined.

1305  
1306 **Figure 6** Phylogeny of insect orders and occurrence of volatile terpenes in select insect species  
1307 representing these orders. Phylogeny of insects adapted from Misof et al.<sup>167</sup> with modifications.

1308  
1309 **Figure 7** Examples of volatile terpene semiochemicals in amphibians, reptiles, and mammals. No  
1310 stereochemical configuration is shown in case the stereochemistry was not determined.

1311  
1312 **Figure 8** Animal phylogeny and occurrence of volatile terpenes (or volatile precursors in the case  
1313 of corals) in select species representing different phyla or clades. Species shown from top to  
1314 bottom are: octocoral species; mantellid frog species, *Mantella aurantiaca*; alligatorid species;  
1315 springbok, *Antidorcas marsupialis*; sea hare, *Aplysia brasiliana*; dust mite species; polyzoniidan  
1316 millipede species; pentatomid species, *Murgantia histrionica* (harlequin bug). Phylogeny adapted  
1317 from Giribet<sup>168</sup> with modifications.

1318  
1319 **Figure 9** Biosynthetic pathway of the monoterpene iridoid sex pheromone components  
1320 nepetalactol and nepetalactone, in females of the pea aphid, *Acyrtosiphon pisum* (*Ap*). *Ap*GES,  
1321 geraniol synthase; *Ap*G8H, geraniol-8-hydroxylase; *Ap*HGO, hydroxygeraniol oxidase; *Ap*ISY,  
1322 iridoid synthase; *Ap*NEPO, nepetalactol oxidase.

1323 **Figure 10** IDS-type TPSs in monoterpene and sesquiterpene pheromone biosynthesis in beetles.  
1324 (A) A bifunctional IDS/TPS enzyme catalyzing the formation of  $\beta$ -myrcene (**32**), the precursor of  
1325 the male aggregation pheromone ipsdienol (**45**); (B) terpene cyclase converting (*Z,E*)-FPP (**73**),  
1326 made by an unusual IDS enzyme (*Ps*IDS3), to (*6R,7S*)-himachala-9,11-diene (**74**), a major  
1327 constituent of the male aggregation pheromone of the striped flea beetle, *Phyllotreta striolata* (*Ps*),  
1328 and other sesquiterpene products.

1329  
1330 **Figure 11** Bisabolane-type sesquiterpene pheromones in stink bug (pentatomid) species.

1331

1332 **Figure 12** IDS-type sesqui-TPSs involved in de novo biosynthesis of male aggregation  
1333 pheromones in stink bugs (Pentatomids). (A) proposed reaction mechanism of TPSs converting  
1334 (*E,E*)-FPP (**4**) to the pheromone precursor (1*S*,6*S*,7*R*)-sesquipiperitol (**80**) in three different  
1335 pentatomid species; (B) proposed reaction mechanism of a TPS catalyzing the formation of the  
1336 pheromone precursor (*S,Z*)- $\alpha$ -bisabolene (**82**) in *Nezara viridula*. Unidentified steps beyond the  
1337 TPS-catalyzed reaction are indicated with question marks. *Hh*, *H. halys*; *Mh*, *M. histrionica*; *Nv*,  
1338 *N. viridula*.

1339  
1340 **Figure 13** Pentatomid IDS-type gene family organization and size. (A) Genomic organization of  
1341 IDS-type gene families in *Halyomorpha halys* (*Hh*) and *Nezara viridula* (*Nv*). Two different clusters  
1342 represent IDS-type genes of the two pentatomid TPS-a and TPS-b clades (modified from Rebholz  
1343 et al.<sup>68</sup>); (B) schematic comparison of TPS gene family sizes in Pentatomids and plants  
1344 (*Arabidopsis thaliana*) in conjunction with volatile terpene product diversity and function.

1345  
1346 **Figure 14** Evolution of putative IDS-like TPSs in hemipteran insects. The phylogram shows  
1347 canonical FPPS and IDS (FPPS)-like proteins mined from hemipteran genomes and  
1348 transcriptomes. Branch lengths represent the number of amino acid substitutions per site. A  
1349 monophyletic clade (purple) of IDS-like sequences clustering with characterized IDS-type TPSs  
1350 was found to be distributed across species within the terpene-emitting hemipteran suborders  
1351 Heteroptera and Sternorrhyncha. Select terpene-emitting species are displayed alongside  
1352 representative terpene compounds. Figure modified from Rebholz et al.<sup>68</sup>.

1353  
1354 **Figure 15** Biosynthesis of the male anti-aphrodisiac (*E*)- $\beta$ -ocimene (**37**) and terpene alcohols by  
1355 GGPPS-type TPSs in the butterfly *Heliconius melpomene* (*Hmel*). *Os*, ocimene synthase.

1356  
1357 **Figure 16** Structural analysis of IPP binding residues in *H. halys* FPPS and a homology model of  
1358 the IDS-type *H. halys* TPS1 enzyme. (A) Structural model of *Hh*FPPS with the IPP binding pocket  
1359 rendered as a colored surface; (B) *Hh*FPPS residues binding the diphosphate moiety of IPP are  
1360 marked in dark blue and labeled; (C) *Hh*FPPS residues binding the prenyl tail of IPP are marked  
1361 in light blue and labeled; (D) structural model of *Hh*TPS1 with a modified IPP binding pocket  
1362 rendered as a colored surface; (E) *Hh*TPS1 residue substitutions of the diphosphate binding  
1363 residues in (B) are marked in purple and labeled. Aromatic substitutions in this region favor  
1364 interactions with the isoprenyl tail of the docked FPP substrate; (F) *Hh*TPS1 residue substitutions  
1365 of the prenyl tail binding residues in (C) are marked in green and labeled. Substitutions alter the

1366 substrate binding region to accommodate a larger isoprenyl diphosphate substrate. Figure  
1367 adapted from Rebholz et al.<sup>151</sup>.

1368  
1369 **Figure 17** Substitutions in the pyrophosphate-binding region is a common feature of insect TPS  
1370 enzymes. (A) The structure of a canonical FPPS from the eastern spruce budworm *Choristoneura*  
1371 *fumiferana* (*CfFPPS*) (PDBid 6b04; light purple ribbons) is shown with IBM-1, 2, 3, and 6 depicted  
1372 as an atomic surface. The electrostatic character of *CfFPPS* IBM motifs was mapped onto the  
1373 surface using UCSF Chimera (blue = basic; red = acidic; grey = neutral/aliphatic); (B) IBM-1, 2,  
1374 3, and 6 of *CfFPPS* and selected insect TPS models was depicted as atomic surfaces for  
1375 comparison. Figure adapted from Rebholz et al.<sup>151</sup>.

1376  
1377 **Figure 18** Acyclic and cyclic sesquiterpenes (**86-88**) in *Artemisia annua* and example of a cyclic  
1378 sesquiterpene product (**89**) generated by an *Artemisia*  $\beta$ -farnesene synthase mutant variant.

1379  
1380 **Figure 19** Phylogram of insect IDS and characterized and predicted IDS-like TPS proteins  
1381 adapted from Rebholz et al.<sup>151</sup>. Branch lengths are proportional to amino acid substitutions per  
1382 site. Previously characterized and predicted IDS-like TPS sequences are labeled with dark blue  
1383 and white squares at branch tips, respectively. Insect orders, from which protein sequences  
1384 originated, are indicated by colors of the circular perimeter and branches. Results suggest a  
1385 recurring parallel emergence of TPS activity in IDS-like enzymes within and between insect  
1386 lineages and IDS enzyme subfamilies.

1387  
1388 **Figure 20** Sesquiterpene and diterpene products of TPS enzymes identified in octocorals.

1389  
1390 **Figure 21** Structures and motif comparison of an insect IDS (*C. fumiferana*) and IDS-like TPS (*H.*  
1391 *halys*) with TPSs from a mite (*L. delicense*) and an octocoral (*E. rubra*). (A) The *C. fumiferana*  
1392 FPPS (*CfFPPS*) structure (PDBid 6b04; light grey ribbons) is shown with FARM and SARM motifs  
1393 colored orange and green, respectively; (B) Active site zoom up of *CfFPPS* showing catalytic  
1394 residues of the FARM and SARM motifs; (C) Homology models of *H. halys* TPS1 and *L. delicense*  
1395 TPS and protein structure of *E. rubra* TPS (PDBid; 7S5L) depicted as ribbons with motifs colored  
1396 according to the scheme in panels A and B. NSE/DTE and DDXXD represent the metal binding  
1397 motifs. *Er*TPS and *Ld*TPS contain two additional motifs: the phosphate sensor R colored in  
1398 magenta, and the RY motif (substituted by RF in *Ld*TPS) unique to bacterial TPS sequences.

1399

Table 1

Species	Gene/ protein	Accession number	Product	Reference
<i>Acyrtosiphon pisum</i>	GES	*	<b>39</b>	114
	G8H	ON862918	<b>70</b>	
	HGO	*	<b>71</b>	
	ISY	ON862920	<b>72, 69</b>	
	NEPO	*	<b>23</b>	
<i>Ips pini</i>	GPP Myrcene s.	AY953508	<b>3, 32</b>	118, 119
<i>Phyllotreta striolata</i>	TPS1	KT959248	<b>74, 75, 76</b>	120
	TPS2	KT959251	Unidentified sesquitrps	
	TPS3	KT959254	<b>31</b> (main product)	
	TPS4	KT959257	<b>31</b> (main product)	
<i>Halyomorpha halys</i>	TPS1	MG917093	<b>80</b>	68
	TPS2	MG870388	Sesquitrp.	
<i>Murgantia histrionica</i>	TPS1	MG662378	<b>80</b> Sesquitrps	123
<i>Nezara viridula</i>	TPS1	MG748543	<b>82</b>	68, 135
	TPS2	ON934605	<b>80</b>	
<i>Heliconius melpomene</i>	Ocimene s.	*	<b>37</b>	146
	HMEL037108g1	*	<b>31, 63</b>	
<i>Empoasca onukii</i>	TPS	MH383159	<b>39</b>	150
<i>Briareum asbestinum</i>	TC-1	*	<b>25</b>	162
	TC-2	*	<b>95</b>	
<i>Capnella imbricata</i>	TC-1	*	<b>92</b>	162
<i>Dendronephthya gigantea</i>	TC-1	*	<b>90</b>	162
	TC-2	*	<b>25</b>	
<i>Eleutherobia rubra</i>	TC-1	*	Unidentified sesquitrps	162
	TC-2	*	<b>25</b>	
<i>Erythropodium caribaeorum</i>	TPS1	OK081311	<b>95</b>	163
	TPS6	OK081312	<b>93</b>	
<i>Heliopora coerulea</i>	TC-1	*	Unidentified sesquitrps	162
	TC-2	*	<b>90</b>	
	TC-3	*	<b>94</b>	
<i>Paramuricea biscaya</i>	TC-1	*	<b>94</b>	162
<i>Renilla muelleri</i>	TC-1	*	<b>93</b>	162
<i>Tubipora musica</i>	TC-1	*	<b>25</b>	162
	TC-2	*	<b>91</b>	
<i>Xenia</i> sp.	TC-1	*	<b>96</b>	162

Table 2

Name	Type	IBM-1			IBM-2			IBM-3			IBM-4			SARM			IBM-5			IBM-6																																	
		G109	K110	R113	E146	O149	L153	R165	R166	T251	Y254	S255	F289	Q290	D293				K307		R401	K403																															
Aphis gossypii_DPPS	DPPS	G	K	A	L	R	F	E	M	I	H	S	A	S	L	S	D	F	R	R	G	K	T	A	S	L	I	A	F	Q	L	V	D	D	L	L	D	M	G	K	P	T	A	A	D	I	V	N	R	M	K		
Empoasca onukii_FPPS2	FPPS	G	K	K	N	R	G	E	M	L	Q	A	F	F	L	S	E	T	R	R	R	G	K	T	S	Y	Y	S	F	Y	Q	A	Q	S	D	D	F	F	N	T	M	K	-	P	G	H	D	I	Y	K	R	E	S
Empoasca onukii_FPPS1		S	S	N	L	R	G	E	L	L	H	T	S	L	V	A	D	E	R	R	R	G	R	S	A	Y	H	T	F	F	Q	V	Q	D	D	Y	M	D	N	G	K	-	V	G	T	D	M	T	G	R	-	L	
Halyomorpha halys_FPPS		G	K	K	V	R	G	E	M	L	Q	G	F	F	V	S	V	T	R	R	R	G	K	T	A	Y	Y	T	F	F	Q	V	Q	D	D	Y	L	D	S	G	K	-	K	G	T	D	M	Y	G	R	K	Q	
Murgantia histrionica_FPPS		G	K	K	V	R	G	E	M	L	Q	G	F	F	V	S	V	T	R	R	R	G	K	T	A	Y	Y	T	Y	F	Q	V	Q	D	D	Y	L	D	I	G	K	-	K	G	T	D	L	Y	G	R	K	Q	
Myzus persicae_FPPS2		G	K	K	N	R	G	E	I	L	Q	A	Y	Q	L	A	I	T	R	R	R	G	K	T	A	Y	Y	S	F	F	Q	I	Q	D	D	Y	L	D	T	G	K	-	I	G	T	D	I	Y	K	R	T	L	
Nezara viridula_FPPS		G	K	K	V	R	G	E	M	L	Q	G	F	F	L	S	I	T	R	R	R	G	K	T	S	Y	Y	T	F	F	Q	V	Q	D	D	Y	L	D	I	G	K	-	I	G	T	D	T	Y	G	R	K	Q	
Rhopalosiphum padi_FPPS		G	K	K	N	R	G	E	I	L	Q	A	Y	Q	L	A	I	T	R	R	R	G	K	T	A	Y	Y	S	F	F	Q	V	Q	D	D	F	L	D	M	G	K	-	I	G	T	D	I	Y	K	R	T	L	
Rhopalosiphum padi_FPPS		G	K	K	N	R	G	E	I	L	Q	A	Y	Q	L	A	I	T	R	R	R	G	K	T	A	Y	Y	S	F	F	Q	V	Q	D	D	F	L	D	T	G	K	-	I	G	T	D	I	Y	K	R	T	L	
Acyrtosiphon pisum_GGPPS		GGPPS	G	K	Q	I	R	A	Q	M	L	H	N	S	S	L	S	V	L	R	R	R	G	K	T	G	-	G	L	F	F	Q	I	R	D	D	Y	C	N	E	N	K	S	Y	C	E	D	L	R	T	W	S	Y
Acyrtosiphon pisum_FPPS/GPPS		GPPS/FPPS	G	K	K	N	R	G	E	I	L	Q	A	Y	Q	L	A	I	T	R	R	R	G	K	T	A	Y	Y	S	F	F	Q	V	Q	D	D	Y	L	D	T	G	K	-	I	G	T	D	I	Y	K	R	T	L
Acyrtosiphon pisum_FPPS/GPPS1	G		K	K	N	R	G	E	I	L	Q	A	Y	Q	L	A	I	T	R	R	R	G	K	T	A	Y	Y	S	F	F	Q	V	Q	D	D	F	L	D	M	G	K	-	I	G	T	D	I	Y	K	R	T	L	
Acyrtosiphon pisum_FPPS/GPPS2	G		K	K	N	R	G	E	I	L	Q	A	Y	Q	L	A	I	T	R	R	R	G	K	T	A	Y	Y	S	F	F	Q	V	Q	D	D	F	L	D	M	G	K	-	I	G	T	D	I	Y	K	R	T	L	
Aphis gossypii_FPPS/GPPS1	G		K	K	N	R	G	E	I	L	Q	A	Y	Q	L	A	I	T	R	R	R	G	K	T	A	Y	Y	S	F	F	Q	V	Q	D	D	F	L	D	M	G	K	-	I	G	T	D	I	Y	K	R	T	L	
Aphis gossypii_FPPS/GPPS2	G		K	K	N	R	G	E	I	L	Q	A	Y	Q	L	A	I	T	R	R	R	G	K	T	A	Y	Y	S	F	F	Q	V	Q	D	D	F	L	D	M	G	K	-	I	G	T	D	I	Y	K	R	T	L	
Myzus persicae_FPPS/GPPS	G		K	K	N	R	G	E	I	L	Q	A	Y	Q	L	A	I	T	R	R	R	G	K	T	A	Y	Y	S	F	F	Q	V	Q	D	D	F	L	D	M	G	K	-	I	G	T	D	I	Y	K	R	T	L	
Empoasca onukii_TPS	TPS		G	K	Q	I	R	A	Q	M	L	H	N	S	S	L	S	I	L	R	R	R	G	K	T	G	-	G	L	F	F	Q	I	R	D	D	Y	C	N	E	N	K	S	F	C	E	D	L	L	N	W	D	R
Halyomorpha halys_TPS1			F	S	D	A	W	N	D	L	L	F	T	M	S	A	S	D	S	R	R	R	G	K	A	G	Q	F	V	A	I	Q	T	W	D	D	F	N	D	N	G	K	-	P	S	C	D	L	V	I	R	E	B
Halyomorpha halys_TPS2			C	Y	E	G	W	N	D	M	S	H	S	M	H	F	A	E	F	R	Q	G	K	S	R	N	T	M	C	F	Q	V	W	N	D	F	M	D	S	G	K	-	G	N	Y	D	L	H	G	N	G	H	
Murgantia histrionica_TPS			F	S	D	A	W	N	D	L	L	F	T	M	S	A	S	E	F	R	K	G	K	A	G	Q	F	V	A	I	Q	T	W	D	D	F	N	D	N	G	K	-	L	S	C	D	L	V	V	R	E	F	
Nezara viridula_TPS		Y	F	E	G	W	A	D	M	S	Y	A	M	A	G	G	E	F	R	R	G	K	A	A	N	T	V	F	F	Q	V	W	D	D	F	M	E	S	G	K	-	G	A	P	D	L	L	V	B	P	P		

Fig. 1




Fig. 2




**Marine gastropods**brasudol **8****Octocorals**lophotoxin **9**isofuranodiene **10**

Fig. 3




















**Millipedes****(S)-(+)-polyzonimine 11****(4S,5R,6S)-(+)-nitropolyzonamine 12****gosodesmine 13****limonene 14****(-)-2-methylisoborneol 15****(-)-geosmin 16****Arachnids****bornyl acetate 17****(3S,8S)-chrysolidial 18****β-springene 19****neryl formate 20****β-(E)-acaridial 21**

Fig. 4



## Phasmatodea

actinidine **22***cis,trans*-nepetalactone **23**

## Blattodea

periplanone B **24**kempene-type diterpene **28**neocembrene  
(cembrene A) **25***(+)*- $\alpha$ -pinene **29**secotrinervitane-  
type diterpene **26***(E,E)*- $\alpha$ -farnesene **30**trinervitane-  
type diterpene **27***(3R,6E)*-nerolidol **31**

## Thysanoptera

 $\beta$ -myrcene **32***(R)*-lavandulol **33**

## Hemiptera


















oleanderlure **34**murgantiol **35**

Fig. 5




## Hymenoptera

(Z,E)- $\alpha$ -farnesene **36**(E)- $\beta$ -ocimene **37**(Z)-citral **38**geraniol **39**(E,E)-farnesol **40**(S)-citronellol **41** $\alpha$ -phellandrene **42**

## Coleoptera

(+)-*trans*-verbenol **43**verbenone **44**ipsdienol **45**sulcatone **46**frontalin **47**(E)- $\beta$ -caryophyllene **48**(+)-ar-himachalene **49**(+)- $\gamma$ -cadinene **50**

## Lepidoptera

grandisol **51**cantharidin **52**germacrene A **53**

## Diptera


(1S,3S,7R)-3-methyl-himachalene **54**

Fig. 6









Fig. 7

## Amphibians

frogolide **55**cucujolide I **56**suspensolide **57**

## Reptiles

11,12-dihydrocembren-10-one **58**citronellyl acetate **59**

## Mammals










 $\alpha$ -springene **60**farnesol hydrate **61**drima-8 $\alpha$ ,11-diol **62**linalool **63**1,8-cineole **64**piperitone **65** $\delta$ -cadinene **66**aromadendrene **67**viridiflorol **68**

Fig. 8




Fig. 9




Fig. 10




Fig. 11




Fig. 12




Fig. 13




Fig. 14




Fig. 15




Fig. 16




Fig. 17







Fig. 18




86



87



88



89

Fig. 19

Tree scale: 1

- Insect Order**
- Blattodea
  - Phthiraptera
  - Hemiptera
  - Hymenoptera
  - Coleoptera
  - Lepidoptera
  - Diptera




Fig. 20




Fig. 21

