


# ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

# Optical reading of the open and closed states of a molecular turnstile<sup>†</sup>

Cite this: DOI: 10.1039/x0xx00000x

Nicolas Zigon, Patrick Larpent, Abdelaziz Jouaiti, Nathalie Kyritsakas and Mir Wais Hosseini\*

Received 00th January 2012,  
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x


www.rsc.org/

**A molecular turnstile composed of a hydroquinone based rotor and a stator bearing a tridentate coordinating site can be reversibly switched between open and closed states. The locking and unlocking processes may be red optically.**

The control of intramolecular movements has attracted considerable efforts over the last two decades and still is of current interest.<sup>1-3</sup> Among many dynamic systems such as motors, rotors and machines, molecular turnstiles form an interesting ensemble.<sup>4</sup> Over the last years, we have investigated a series of porphyrins<sup>5</sup> or Pt(II) organometallic<sup>6</sup> turnstiles.

Herein, we report on a new system for which the open and closed states display different luminescent properties.

The turnstile **1** (Scheme 1) is composed of a rotor based on a hydroquinone moiety equipped with two divergently oriented benzonitrile units as monodentate coordinating sites and a stator bearing a tridentate chelating unit. The terphenyl group was chosen as the hinge for its luminescent properties. As the external effector leading to the closed state of the turnstile **1**-Pd, Pd(II) was used.




**Scheme 1** Structures of compounds **1-3** (L = DMAP or CN).

The rationale behind the choice of the latter was its square planar coordination geometry allowing thus its simultaneous binding by the

rotor and stator and its propensity to behave as a heavy atom and thus to quench the emission of the terphenyl moiety.

For the open state of the turnstile **1** (Fig. 1 O1), the rotor should freely rotate around the stator, whereas in the presence of Pd(II), the turnstile would be locked in its closed state (Fig. 1 C).



**Fig. 1** Schematic representations of the turnstile in its open (O1 and O2) and closed (C) states.

The synthesis of **1** (see ESI) was achieved in 8 steps. The precursor macrocyclic compound **3** was obtained in 90 % yield upon condensation of the handle **2**, prepared in 6 steps,<sup>59</sup> with 2,5-dibromohydroquinone in DMF in the presence of Cs<sub>2</sub>CO<sub>3</sub>. The turnstile **1** was obtained in 68 % yield by coupling 4-benzonitrile boronic acid with the dibromo compound **3** in the presence of Pd(PPh<sub>3</sub>)<sub>4</sub>. The closed state of the turnstile (**1**-Pd) was generated quantitatively upon metallation of the compound **1** by Pd(OAc)<sub>2</sub>.

In addition to solution characterizations, the solid-state structure of **1**-Pd was studied by X-Ray diffraction (Fig. 2, see ESI). Single crystals of **1**-Pd were obtained at 25 °C upon slow diffusion of Et<sub>2</sub>O into a CHCl<sub>3</sub> solution of **1**-Pd. The latter crystallizes (monoclinic P2<sub>1</sub>/n) without any solvent. The Pd(II) cation, adopting a strongly distorted square planar geometry (NPdN<sub>cis</sub> angle in the 80.7 - 99.8° range and NPdN<sub>trans</sub> angles of 161.6 and 175.9°), is coordinated to the tridentate chelating moiety of the stator (Pd-N distance in the 1.92-2.02 Å range) and to the N atom of the benzonitrile unit of the rotor (d<sub>PdN</sub> = 2.01 Å).



with a hyperchromic effect. Interestingly, whereas for the turnstile **1** in its open state, excitation at 350 nm, assigned to  $\pi$ - $\pi^*$  transitions, leads to strong emission in the blue domain of the visible spectrum ( $\lambda_{em} = 434$  nm,  $\Phi_F = 0.96$ ), the closed state **1**-Pd ( $\lambda_{em} = 436$  nm,  $\Phi_F = 0.03$ ) is far less emissive (Fig. 7). For the determination of the quantum yields, quinine sulphate was used as reference. Lifetimes of the excited state of **3.2** and **3.1** ns for **1** and **1**-Pd respectively have been determined. A plausible hypothesis for the observed decrease of luminescence resulting by the binding of Pd(II) could be that, upon excitation of the  $\pi$ - $\pi^*$  transition of the rotor, a  $S_1$  excited state with a lifetime of ca. 3 ns is generated. For the closed state of the turnstile, the heavy atom effect of Pd(II) promotes the transition to a  $T_1$  excited state which is efficiently quenched by solvent or  $O_2$  under aerated condition.




Fig. 6 UV-visible spectra ( $CH_2Cl_2$ , 298 K) of **1** (black) and **1**-Pd (red).

Using *iso*-absorbing solutions of **1** and **1**-Pd and under the same conditions and identical recording parameters, the excitation at 350 nm leads to a strong signal for **1**, whereas for the closed state of the turnstile **1**-Pd, the emission is below the detection limits of the spectrometer used (Fig. 7).




Fig. 7 Picture of **1** (left) and **1**-Pd (right) in  $CH_2Cl_2$  (298 K, aerated,  $\lambda_{exc} = 365$  nm) and emission spectra for *iso*-absorbing solutions of the open **1** (blue) and closed **1**-Pd (red) states of the turnstiles ( $CH_2Cl_2$ , 298 K, aerated,  $\lambda_{exc} = 350$  nm).

In conclusion, the purely organic turnstile **1**, composed of a rotor based on a hydroquinone derivative bearing two peripheral coordinating sites of the benzonitrile type connected to a stator equipped with a chelating moiety undergoes a reversible locking/unlocking process between an open and a closed state. The closed state is generated upon binding of Pd(II) as an effector. The switch between the locked and unlocked states is achieved either by a competitive external ligand such as  $CN^-$  or DMAP or in the latter case by an acid ( $MsOH$ ) and a base ( $Et_3N$ ). Interestingly, owing to the emissive nature of the turnstile, whereas the open state  $O_1$  is strongly luminescent, for the closed state, the emission is quenched by the presence of Pd(II) leading thus to an optical reading of the two states.

Financial supports by the University of Strasbourg, the International Centre for Frontier Research in Chemistry (PhD fellowship to P.L.), Strasbourg, the Institut Universitaire de France, the CNRS and the Ministry of Education and Research (PhD fellowship to N. Z.) are acknowledged.

## Notes and references

Molecular Tectonic Laboratory, UMR UDS-CNRS 7140, icFRC, University of Strasbourg, Institut Le Bel, 4, rue Blaise Pascal, F-67000 Strasbourg, France. hosseini@unistra.fr.

† This contribution is dedicated to J.-P. Sauvage on the occasion of his 70<sup>th</sup> birthday.

Electronic Supplementary Information (ESI) available: [experimental details, characterisation of new compounds, additional 2D-NMR and UV/luminescence spectra]. Experimental crystallographic data: CCDC 985992 See DOI: 10.1039/c000000x/

- (a) J.-P. Sauvage, *Science*, 2001, **291**, 2105; (b) J.-P. Sauvage, *Acc. Chem. Res.*, 1998, **31**, 611; (c) J.-P. Sauvage, in *Molecular machines and motors, Structure and bonding*, Vol. 99, (Ed.: J.-P. Sauvage), Springer, Berlin, Heidelberg, 2001, pp. 1-282; (d) V. Balzani, M. Venturi and A. Credi, in *Molecular devices and machines: a journey into the nanoworld*, Wiley-VCH, Weinheim, 2003, pp. 1-457; (e) T. R. Kelly, in *Molecular machines, Topics in Current Chemistry*, Vol. 262, (Ed.: T. R. Kelly), Springer, Berlin, Heidelberg, 2005, pp. 1-227; (f) Y. Shirai, J.-F. Morin, T. Sasaki, J. M. Guerrero and J. M. Tour., *Chem. Soc. Rev.*, 2006, **35**, 1043.
- (a) N. Koumura, R. W. J. Zijlstra, R. A. van Delden, N. Harada and B. L. Feringa, *Nature*, 1999, **401**, 152; (b) V. Balzani, A. Credi, F. M. Raymo and J. F. Stoddart, *Angew. Chem. Int. Ed.*, 2000, **39**, 3348; (c) J. Michl and E. C. H. Sykes, *ACS Nano*, 2009, **3**, 1042; (d) S. Sengupta, M. E. Ibele and A. Sen, *Angew. Chem. Int. Ed.*, 2012, **51**, 8434.
- E. R. Kay, D. A. Leigh and F. Zerbetto, *Angew. Chem. Int. Ed.*, 2007, **46**, 72.
- (a) T. C. Bedard and J. S. Moore, *J. Am. Chem. Soc.*, 1995, **117**, 10662; (b) A. Carella, J. Jaud, G. Rapenne and J.-P. Launay, *Chem. Commun.*, 2003, 2434; (c) N. Weibel, A. Mishchenko, T. Wandlowski, M. Neuburger, Y. Leroux and M. Mayor, *Eur. J. Org. Chem.*, 2009, **35**, 6140; (d) K. Skopek, M. C. Hershberger and J. A. Gladysz, *Coord. Chem. Rev.*, 2007, **251**, 1723.
- (a) A. Guenet, E. Graf, N. Kyritsakas, L. Allouche and M. W. Hosseini, *Chem. Commun.*, 2007, 2935; (b) A. Guenet, E. Graf, N. Kyritsakas and M. W. Hosseini, *Inorg. Chem.*, 2010, **49**, 1872; (c) T. Lang, A. Guenet, E. Graf, N. Kyritsakas and M. W. Hosseini, *Chem. Commun.*, 2010, **46**, 3508; (d) A. Guenet, E. Graf, N. Kyritsakas and M. W. Hosseini, *Chem. Eur. J.*, 2011, **17**, 6443; (e) T. Lang, E. Graf, N. Kyritsakas and M. W. Hosseini, *Dalton Trans.*, 2011, **40**, 3517; (f) T. Lang, E. Graf, N. Kyritsakas and M. W. Hosseini, *Dalton Trans.*, 2011, **40**, 5244; (g) T. Lang, E. Graf, N. Kyritsakas and M. W. Hosseini, *Chem. Eur. J.*, 2012, **18**, 10419; (h) T. Lang, E. Graf, N. Kyritsakas and M. W. Hosseini, *New J. Chem.*, 2013, **37**, 112.
- (a) N. Zigon, A. Guenet, E. Graf and M. W. Hosseini, *Chem. Commun.*, 2013, **49**, 3637; (b) N. Zigon, A. Guenet, E. Graf, N. Kyritsakas and M. W. Hosseini, *Dalton Trans.*, 2013, **42**, 9740; (c) N. Zigon, N. Kyritsakas and M. W. Hosseini, *Dalton Trans.*, 2013, **43**, 152.

Graphical Abstract:

The reversible switching between the open and closed states of a molecular turnstile can be addressed by emission spectroscopy.

