

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/chemcomm

ARTICLE TYPE

Gold-catalyzed intermolecular oxidation of *o*-alkynylbiaryls: an easy and practical access to functionalized fluorenes

Fei Pan, Shuang Liu, Chao Shu, Rong-Kun Lin, Yong-Fei Yu, Jin-Mei Zhou and Long-Wu Ye*

Received (in XXX, XXX) Xth XXXXXXXXXX 200X, Accepted Xth XXXXXXXXXX 200X

DOI: 10.1039/b000000x


A novel gold-catalyzed intermolecular oxidation of *o*-alkynylbiaryls has been developed. A variety of functionalized fluorenes are readily accessed by utilizing this non-diazo approach, thus providing a viable alternative to synthetically useful fluorenes.

Among the polycyclic aromatic hydrocarbons (PAH), fluorenes are core structures employed in a variety of practical applications in materials chemistry as dyes and optical brightening agents.¹ In addition, fluorenyl functions have also played important roles in the field of organic synthesis, mainly as unique protecting groups in peptide synthesis.² Somewhat surprisingly, however, only a few preparative methods have been reported.³ Therefore, the development of novel strategies, especially those based on assembling structures directly from readily available and easily diversified building blocks, is highly desirable.


In recent years, significant progress has been made in the gold-catalyzed intermolecular alkyne oxidation to make alkynes as equivalents of α -diazo ketones.⁴⁻⁶ For example, this non-diazo approach offers a viable alternative for the transition metal-catalyzed intramolecular aromatic carbenoid insertion of diazo compounds.⁷⁻⁸ In this regard, L. Zhang and co-workers reported an elegant protocol for the synthesis of variously substituted chroman-3-ones involving a gold-catalyzed intermolecular oxidation of propargyl aryl ethers.^{7a} Later, J. Zhang and co-workers realized a gold-catalyzed oxidation/C–H functionalization reaction of electron-deficient *N*-arylpropionamides for the construction of 3-acyloxindoles.^{7b} The group of Gagosz disclosed a gold-catalyzed oxidative cyclization of propynyl arenes, allowing rapid and practical access to a variety of functionalized indan-2-ones.^{7c} Recently, Li and co-workers reported that the preparation of oxindoles could be achieved via a gold-catalyzed intermolecular oxidation of *N*-arylnamides.^{7d} Despite these significant achievements, these oxidative cyclizations, presumably involving the intramolecular trapping of the α -oxo gold carbenoids by the aryl ring, are limited to terminal alkynes or electron-deficient internal alkynes, as electron-rich internal alkynes often suffers the competing overoxidation of the carbene center by the very oxidant.^{9,7d} Recently, Chang and co-workers described a general route for the synthesis of fluorenes that involves a rhodium- or copper-catalyzed intramolecular aromatic carbenoid insertion of biaryldiazoacetates (Scheme 1a).¹⁰ Inspired by these works and our recent findings on the gold-catalyzed synthesis of polycyclic

aromatic hydrocarbons,¹¹ we envisioned that these synthetically useful fluorene skeletons might be accessed directly from *o*-alkynylbiaryls through a gold-catalyzed intermolecular alkyne oxidation (Scheme 1b). In this communication, we describe herein the realization of such a gold-catalyzed oxidative protocol, providing a flexible and alternative way for the preparation of versatile fluorene derivatives.

(a) Rh- or Cu-catalyzed aromatic carbenoid insertion of biaryldiazoacetates (Chang)



(b) Au-catalyzed intermolecular oxidation of *o*-alkynylbiaryls (this work)




Scheme 1 Formation of functionalized fluorenes through gold-catalyzed intermolecular alkyne oxidation.

Ynamide **1a** was chosen as a model substrate for our initial study and some of the results are outlined in Table 1.¹² The screening of different gold catalysts was first performed (Table 1, entries 1-8). In most cases, diketone compound **3a** was detected as a major product (Table 1, entries 1-6). To our delight, the use of the bulky gold catalysts could dramatically suppress the undesired overoxidation (Table 1, entries 7-8) and the desired fluorene **2a** was obtained in 78% yield when using BrettPhosAuNTf₂ as the gold catalyst (Table 1, entry 7). Here, the bulky ligand likely prevents the overoxidation of the carbene center by the very oxidant via intermolecular reactions as the carbene center is sterically shielded.^{7a} Attempts to improve the yield of this reaction by the screening of other quinoline *N*-oxides and pyridine *N*-oxides was unsuccessful (Table 1, entries 9-13). As water could assist some gold-catalyzed oxidative reactions,^{6a} water was used as the solvent or mixture solvent to further improve this reaction but resulted in a decrease of the yield (Table 1, entries 14-15). Of note, without a gold catalyst, the reaction failed to give even a trace of fluorene **2a**, and PtCl₂ and AgNTf₂ could not catalyze this reaction. Finally, it should be specially mentioned that the corresponding phenanthrene **3a'** was


obtained in 85% yield without the addition of *N*-oxide (Table 1, entry 16), indicating that the gold-catalyzed 6-*endo-dig* cycloisomerization can be suppressed by the oxidant.¹³

Table 1 Optimization of reaction conditions^a



Entry	L	Oxidant (R)	Yield ^b (%)	
			2a	3a
1	Et ₃ P	4a (Et)	<5	>95
2	IPr	4a (Et)	21	65
3	PPh ₃	4a (Et)	<5	>95
4	P(4-CF ₃ C ₆ H ₄) ₃	4a (Et)	<5	>95
5	Cy-JohnPhos	4a (Et)	<5	>95
6	XPhos	4a (Et)	18	75
7	BrettPhos	4a (Et)	78	<2
8	Me ₄ ^t BuXPhos	4a (Et)	62	<2
9	BrettPhos	4b (Me)	50	25
10	BrettPhos	4c (ⁱ Pr)	60	<2
11	BrettPhos	5a (R' = 2-Br)	<5 ^c	12
12	BrettPhos	5b (R' = 2,6-Br ₂)	<5 ^d	10
13	BrettPhos	5c (R' = 3,5-Cl ₂)	30	14
14 ^e	BrettPhos	4a (Et)	47	<2
15 ^f	BrettPhos	4a (Et)	62	<2
16 ^g	BrettPhos	/	0	0

^a Reaction conditions: [1a] = 0.05 M, oxidant (2.0 equiv.); DCE: 1, 2-dichloroethane. ^b Estimated by ¹H NMR using diethyl phthalate as internal reference. ^c 70% of 1a remained unreacted. ^d 75% of 1a remained unreacted. ^e Reaction was run in H₂O. ^f Reaction was run in H₂O/DCE = 1:1. ^g The corresponding phenanthrene 3a' was obtained in 85% yield (see SI for details).



With the optimized reaction conditions in hand (Table 1, entry 15 7), we subsequently investigated the substrate scope of this oxidative cyclization. Different functional groups such as fluoro, chloro and methyl on the 4'-methyl-1,1'-biaryl ring were tolerated in this mild transformation, providing the corresponding fluorene derivatives 2b-2f in moderate to good yields (Table 2, entries 2-20 6). Notably, the reaction of methoxy-substituted ynamide 1g only afforded spiro compound 2g¹⁴ through a gold-catalyzed 5-*endo-dig* cyclization and no fluorene formation could be observed (Table 2, entry 7). When the substrate bearing methyl substituent at the meta-position (1h) was employed, a 3/1 regioselectivity

Table 2 Reaction scope study^a

Entry	Substrate	1	Product	2	Yield (%)
1		1a		2a	77%
2		1b		2b	75%
3		1c		2c	56%
4		1d		2d	67%
5		1e		2e	58%
6		1f		2f	64%
7		1g		2g'	86%
8 ^b		1h		2h	72%
9		1i		2i	88%
10		1j		2j	54%

Table 2 (continued)

11		1k		2k	<10% ^c
12		1l		2l	<10% ^c
13		1m		2m	71%

^a Reactions run in toluene; [1] = 0.05 M; isolated yields are reported. ^b 2h/2h' = 3/1, determined by ¹H NMR integration of crude mixture. ^c Estimated by ¹H NMR using diethyl phthalate as internal reference.

and 72% combined yield of **2h** and **2h'** could be achieved (Table 2, entry 8). In case of more electron-rich 3',5'-dimethyl-1,1'-biphenyl **1i**, the reaction could give the desired product **2i** in 88% yield (Table 2, entry 9). In addition, ynamide **1j** was also a suitable substrate for this oxidative cyclization to furnish the corresponding **2j** in a serviceable yield (Table 2, entry 10) while no reactivity was observed with the electron-deficient ynamide **1k** (Table 2, entry 11). In spite of this limitation, the electron-deficient fluorene can be obtained from another aryl ring of the biphenyl due to the symmetric structure of the fluorene compound (Table 2, entry 2-4). Subsequent investigation of *N*-protecting groups demonstrated that Bs (*p*-bromobenzenesulfonyl) protected substrate **1l** did not undergo this oxidative cyclization (Table 2, entry 12), but for MBS (*p*-methoxybenzenesulfonyl) protected substrate **1m**, the reaction could lead to a respectable 71% yield (Table 2, entry 13).

In summary, we have developed a gold-catalyzed intermolecular oxidation of *o*-alkynylbiaryls, allowing the convenient synthesis of functionalized fluorenes in generally moderate to good yields. In comparing our method with literature protocols, this non-diazo approach offers a convenient and viable alternative for the preparation of synthetically useful fluorenes. Other notable features of this method include widespread availability of the substrates, a simple procedure, and mild reaction conditions and, in particular, no need to exclude moisture or air ("open flask"). Further studies on the mechanism and synthetic application of the current reaction are ongoing in our group.

We are grateful for the financial support from the National Natural Science Foundation of China (No. 21102119 and 21272191), the Program for Changjiang Scholars and Innovative Research Team in University (PCSIRT) and NFFTBS (No. J1310024).

Notes and references

⁴⁰ State Key Laboratory for Physical Chemistry of Solid Surfaces, The Key Laboratory for Chemical Biology of Fujian Province and Department of

Chemistry, College of Chemistry and Chemical Engineering, Xiamen University, Xiamen, 361005, Fujian, PR China. Fax: (+86) 592-218-5833; E-mail: longwuyue@xmu.edu.cn

⁴⁵ † Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/b000000x/

1 For selected recent examples, see: (a) J. L. Zafra, J. Casado, I. I. Percepichka, M. R. Bryce, F. J. Ramirez and J. T. L. Navarrete, *J. Chem. Phys.*, 2011, **134**, 044520; (b) M. Zhu, T. Ye, C.-G. Li, X. Cao, C. Zhong, D. Ma, J. Qin and C. Yang, *J. Phys. Chem. C*, 2011, **115**, 17965; (c) K.-Y. Pu, R. Zhan and B. Liu, *Chem. Commun.*, 2010, 1470; (d) S. M. Aly, C.-L. Ho, W.-Y. Wong, D. Fortin and P. D. Harvey, *Macromolecules*, 2009, **42**, 6902; (e) H.-C. Yeh, C.-H. Chien, P.-I. Shih, M.-C. Yuan and C.-F. Shu, *Macromolecules*, 2008, **41**, 3801; (f) Y. Mo, X. Jiang, D. Cao, *Org. Lett.*, 2007, **9**, 4371.

2 For selected examples, see: (a) O. Bassas, J. Huuskonen, K. Rissanen and A. M. P. Koskinen, *Eur. J. Org. Chem.*, 2009, 1340; (b) M. R. Paleo, N. Aurrecochea, K.-Y. Jung and H. Rapoport, *J. Org. Chem.*, 2003, **68**, 130; (c) Y. Ding, J. Wang, K. A. Abboud, Y. Xu, W. R. Dolbier, Jr. and N. G. J. Richards, *J. Org. Chem.*, 2001, **66**, 6381; (d) K. D. Stigers, M. R. Koutroulis, D. M. Chung and J. S. Nowick, *J. Org. Chem.*, 2000, **65**, 3858; (e) L. A. Carpino, *J. Org. Chem.*, 1980, **45**, 4250; (f) L. A. Carpino and G. Y. Han, *J. Am. Chem. Soc.*, 1970, **92**, 5748.

3 For recent examples, see: (a) M. Itoh, K. Hirano, T. Satoh, Y. Shibata, K. Tanaka and M. Miura, *J. Org. Chem.*, 2013, **78**, 1365; (b) K. Morimoto, M. Itoh, K. Hirano, T. Satoh, Y. Shibata, K. Tanaka and M. Miura, *Angew. Chem. Int. Ed.*, 2012, **51**, 5359; (c) X.-C. Wang, R.-L. Yan, M.-J. Zhong and Y.-M. Liang, *J. Org. Chem.*, 2012, **77**, 2064; (d) T.-P. Liu, C.-H. Xing and Q.-S. Hu, *Angew. Chem. Int. Ed.*, 2010, **49**, 2909; (e) F.-L. Sun, M. Zeng, Q. Gu and S.-L. You, *Chem. Eur. J.*, 2009, **15**, 8709; (f) D. J. Gorin, I. D. G. Watson and F. D. Toste, *J. Am. Chem. Soc.*, 2008, **130**, 3736; (g) K. Fuchibe and T. Akiyama, *J. Am. Chem. Soc.*, 2006, **128**, 1434.

4 For reviews, see: (a) H.-S. Yeom and S. Shin, *Acc. Chem. Res.*, 2014, **47**, 966; (b) L. Zhang, *Acc. Chem. Res.*, 2014, **47**, 877; (c) J. Xiao and X. Li, *Angew. Chem. Int. Ed.*, 2011, **50**, 7226.

5 For recent examples, see: (a) S. N. Karad and R.-S. Liu, *Angew. Chem. Int. Ed.*, 2014, **53**, 5444; (b) T. Wang, S. Shi, M. M. Hansmann, E. Rettenmeier, M. Rudolph and A. S. K. Hashmi, *Angew. Chem. Int. Ed.*, 2014, **53**, 3715; (c) M. D. Santos and P. W. Davies, *Chem. Commun.*, 2014, 6001; (d) J. Li, K. Ji, R. Zheng, J. Nelson and L. Zhang, *Chem. Commun.*, 2014, 4130; (e) G. Wu, R. Zheng, J. Nelson and L. Zhang, *Adv. Synth. Catal.*, 2014, **356**, 1229; (f) K. Ji, L. Zhang, *Org. Chem. Front.*, 2014, **1**, 34; (g) P. Nösel, L. N. dos Santos Comprido, T. Lauterbach, M. Rudolph, F. Rominger and A. S. K. Hashmi, *J. Am. Chem. Soc.*, 2013, **135**, 15662; (h) L. Wang, X. Xie and Y. Liu, *Angew. Chem. Int. Ed.*, 2013, **52**, 13302; (i) S. K. Pawar, C.-D. Wang, S. Bhunia, A. M. Jadhav and R.-S. Liu, *Angew. Chem. Int. Ed.*, 2013, **52**, 7559; (j) K. Ji, Y. Zhao and L. Zhang, *Angew. Chem. Int. Ed.*, 2013, **52**, 6508; (k) S. Ghorpade, M.-D. Su and R.-S. Liu, *Angew. Chem. Int. Ed.*, 2013, **52**, 4229; (l) J. Fu, H. Shang, Z. Wang, L. Chang, W. Shao, Z. Yang and Y. Tang, *Angew. Chem. Int. Ed.*, 2013, **52**, 4198; (m) D. B. Hupple, S. Ghorpade and R.-S. Liu, *Chem. Eur. J.*, 2013, **19**, 12965; (n) S. Shi, T. Wang, W. Yang, M. Rudolph and A. S. K. Hashmi, *Chem. Eur. J.*, 2013, **19**, 6576; (o) R. K. Kawade and R.-S. Liu, *Org. Lett.*, 2013, **15**, 4094; (p) H.-H. Hung, Y.-C. Liao and R.-S. Liu, *J. Org. Chem.*, 2013, **78**, 7970.

6 (a) L. Li, C. Shu, B. Zhou, Y.-F. Yu, X.-Y. Xiao and L.-W. Ye, *Chem. Sci.*, 2014, **5**, DOI: 10.1039/C4SC00983E; (b) C. Shu, L. Li, X.-Y. Xiao, Y.-F. Yu, Y.-F. Ping, J.-M. Zhou and L.-W. Ye, *Chem. Commun.*, 2014, 8689; (c) C. Shu, L. Li, Y.-F. Yu, S. Jiang and L.-W. Ye, *Chem. Commun.*, 2014, 2522.

7 (a) Y. Wang, K. Ji, S. Lan and L. Zhang, *Angew. Chem. Int. Ed.*, 2012, **51**, 1915; (b) D. Qian and J. Zhang, *Chem. Commun.*, 2012, 7082; (c) G. Henrion, T. E. J. Chava, X. Le Goff and F. Gagosz, *Angew. Chem. Int. Ed.*, 2013, **52**, 6277; (d) L.-Q. Yang, K.-B. Wang and C.-Y. Li, *Eur. J. Org. Chem.*, 2013, 2775.

- 8 For a metal-free oxidative cyclization of alkynyl aryl ethers to benzofuranones, see: K. Graf, C. L. Rühl, M. Rudolph, F. Rominger and A. S. K. Hashmi, *Angew. Chem. Int. Ed.*, 2013, **52**, 12727.
- 9 (a) K.-B. Wang, R.-Q. Ran, S.-D. Xiu and C.-Y. Li, *Org. Lett.*, 2013, **15**, 2374; (b) R. B. Dateer, K. Pati and R.-S. Liu, *Chem. Commun.*, 2012, 7200; (c) A. Mukherjee, R. B. Dateer, R. Chaudhuri, S. Bhunia, S. N. Karad and R.-S. Liu, *J. Am. Chem. Soc.*, 2011, **133**, 15372; (d) D. Vasu, H.-H. Hung, S. hunia, S. A. Gawade, A. Das and R.-S. Liu, *Angew. Chem. Int. Ed.*, 2011, **50**, 6911; (e) C.-F. Xu, M. Xu, Y.-X. Jia and C.-Y. Li, *Org. Lett.*, 2011, **13**, 1556. (f) C.-W. Li, K. Pati, G.-Y. Lin, S. M. A. Sohel, H.-H. Hung and R.-S. Liu, *Angew. Chem. Int. Ed.*, 2010, **49**, 9891
- 10 J. Kim, Y. Ohk, S. H. Park, Y. Jung and S. Chang, *Chem. Asian J.*, 2011, **6**, 2040.
- 15 11 (a) C. Shu, L. Li, C.-B. Chen, H.-C. Shen and L.-W. Ye, *Chem. Asian J.*, 2014, **9**, 1525; (b) C. Shu, C.-B. Chen, W.-X. Chen and L.-W. Ye, *Org. Lett.*, 2013, **15**, 5542.
- 12 For recent reviews on ynamide reactivity, see: (a) X.-N. Wang, H.-S. Yeom, L.-C. Fang, S. He, Z.-X. Ma, B. L. Kedrowski and R. P. Hsung, *Acc. Chem. Res.*, 2014, **47**, 560; (b) K. A. DeKorver, H. Li, A. G. Lohse, R. Hayashi, Z. Lu, Y. Zhang and R. P. Hsung, *Chem. Rev.*, 2010, **110**, 5064; (c) G. Evano, A. Coste and K. Jouvin, *Angew. Chem. Int. Ed.*, 2010, **49**, 2840.
- 20 13 For the synthesis of phenanthrenes via transition metal-catalyzed 6-endo cyclization of *o*-alkynylbiaryls, see: (a) K. Komeyama, R. Igawa and K. Takaki, *Chem. Commun.*, 2010, 1748; (b) A. Fürstner and J. W. J. Kennedy, *Chem. Eur. J.*, 2006, **12**, 7398; (c) V. Mamane, P. Hannen and A. Fürstner, *Chem. Eur. J.*, 2004, **10**, 4556; (d) A. Fürstner and V. Mamane, *J. Org. Chem.*, 2002, **67**, 6264.
- 30 14 CCDC-1015243 (**2g'**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.